O. Broche E. Jespers C. Polcino Milies M. Ruiz

Abstract

Let R be a commutative ring, G a group and RG its group ring. Let φ : RG \rightarrow RG denote the R-linear extension of an involution φ defined on G. An element x in RG is said to be φ antisymmetric if $\varphi(x) = -x$. A characterization is given of when the φ -antisymmetric elements of RG commute. This is a completion of earlier work.

keywords: Involution; group ring; antisymmetric elements. keywords: 2000 Mathematics Subject Classification: 16S34, 16W10, 20C07.

1 Introduction.

Throughout this paper R is a commutative ring with identity, G is a group and φ is an involution on G. Clearly φ can be extended linearly to an involution $\varphi : RG \to RG$ of the group ring RG. Set $R_2 = \{r \in \mathbb{R} \mid 2r = 0\}$. We denote by $(RG)_{\varphi}^-$ the Lie algebra consisting of the φ -antisymmetric elements of RG, that is

$$
(RG)_{\varphi}^- = {\alpha \in RG | \varphi(\alpha) = -\alpha}.
$$

For general algebras A with an involution φ , we recall some important results that show that crucial information of the algebraic structure of A can be determined by that of $(A)_{\varphi}^-$ and the latter has information that is determined by the φ -unitary unit group $U_{\varphi}(A) = \{u \in A \mid u\varphi(u) = \varphi(u)u =$ 1}. By $U(A)$ we denote the unit group of A. Amitsur in [\[1\]](#page-9-0) proves that if A_{φ}^- satisfies a polynomial identity (in particular when A_{φ}^- is commutative) then A satisfies a polynomial identity. Gupta and Levin in [\[10\]](#page-10-0) proved that for all $n \geq 1$ $\gamma_n(\mathcal{U}(A)) \leq 1 + L_n(A)$. Here $\gamma_n(G)$ denotes the *n*th term in the lower central series of the group G and $L_n(A)$ denotes the two sided ideal of A generated by all Lie elements of the form $[a_1, a_2, ..., a_n]$ with $a_i \in A$ and $[a_1] = a_1$, $[a_1, a_2] = a_1 a_2 - a_2 a_1$ and inductively $[a_1, a_2, \ldots, a_n] = [[a_1, a_2, \ldots, a_{n-1}], a_n]$. Smirnov and Zalesskii in [\[14\]](#page-10-1), proved that, for example, if the Lie ring generated by the elements of the form $g + g^{-1}$ with $g \in \mathcal{U}(A)$ is Lie nilpotent then A is Lie nilpotent. In [\[5\]](#page-9-1) Giambruno and Polcino Milies show that if A is a finite dimensional semisimple algebra over an algebraically closed field F with $char(F) \neq 2$ then $\mathcal{U}_{\varphi}(A)$ satisfies a group identity if and only if $(A)^{-}_{\varphi}$ is commutative. Furthermore, if F is a nonabsolute field then $\mathcal{U}_{\varphi}(A)$ does not contain a free group of rank 2 if and only if $(A)^{-}_{\varphi}$ is commutative. Giambruno and Sehgal, in [\[6\]](#page-9-2), showed that if B is a semiprime ring with involution φ , $B = 2B$ and $(B)_{\varphi}^-$ is Lie nilpotent then $(B)_{\varphi}^-$ is commutative and B satisfies a polynomial identity of degree 4.

Special attention has been given to the classical involution $*$ on RG, that is, the R-linear map defined by mapping $g \in G$ onto g^{-1} . In case R is a field of characteristic 0 and G is a periodic group, Giambruno and Polcino Milies in [\[5\]](#page-9-1) described when $\mathcal{U}_*(RG)$ satisfies a group identity. Gonçalves and Passman in [\[8\]](#page-9-3) characterized when $\mathcal{U}_*(RG)$ does not contain non abelian free groups when G is a finite group and R is a nonabsolute field. Giambruno and Sehgal, in [\[7\]](#page-9-4), characterized when $(RG)^{-}_{*}$ is Lie nilpotent provided R is a field of characteristic $p \geq 0$, with $p \neq 2$.

Motivated by all these connections, in this paper we deal with the question of when $(RG)_{\varphi}^-$ is commutative for an arbitrary involution φ on G. Let $G_{\varphi} = \{g \in G \mid \varphi(g) = g\}$ be the subset of φ -symmetric elements of G, i.e. the set of elements of G fixed by φ . The following complete answer is obtained.

Theorem 1.1 Let R be a commutative ring. Suppose G is a non-abelian group and φ is an invo*lution on G. Then,* $(RG)_{\varphi}^-$ *is commutative if and only if one of the following conditions holds:*

- *1.* $K = \langle g \in G | g \notin G_{\varphi} \rangle$ *is abelian (and thus* $G = K \cup Kx$ *, where* $x \in G_{\varphi}$ *, and* $\varphi(k) = xkx^{-1}$ *for all* $k \in K$ *)* and $R_2^2 = \{0\}$.
- 2. $R_2 = \{0\}$ and G contains an abelian subgroup of index 2 that is contained in G_φ .
- *3.* char(R) = 4, $|G'| = 2$, $G/G' = (G/G')_{\varphi}$, $g^2 \in G_{\varphi}$ for all $g \in G$, and G_{φ} is commutative in $case \ R_2^2 \neq \{0\}.$
- 4. char(R) = 3, $|G'| = 3$, $G/G' = (G/G')_{\varphi}$ and $g^3 \in G_{\varphi}$ for all $g \in G$.

Clearly, as an R-module, $(RG)_{\varphi}^-$ is generated by the set

$$
S = \{ g - \varphi(g) \mid g \in G \setminus G_{\varphi} \} \cup \{ rg \mid g \in G_{\varphi}, \ r \in R_2 \}
$$

Therefore $(RG)_{\varphi}^-$ is commutative if and only if the elements in S commute.

This work is a continuation of the work started in [\[4\]](#page-9-5) (for the classical involution), [\[11\]](#page-10-2) and [\[3\]](#page-9-6). In the latter one considers the involutions η on RG introduced by Novikov in [\[13\]](#page-10-3): $\eta(\sum)$ g∈G $\alpha_g g) =$ \sum g∈G $\alpha_g \sigma(g) g^{-1}$, where $\sigma : G \to {\pm 1}$ is a group homomorphism. Unfortunately, in [\[4,](#page-9-5) [11\]](#page-10-2) the set $S_1 = \{rg \mid r \in R_2, g \in G_{\varphi}\}\$ was not included in the set S. Therefore, the results given in [\[4,](#page-9-5) [11\]](#page-10-2) only deal with commuting of elements in the set $S \setminus S_1$. Hence, provided $R_2 = \{0\}$, there is a complete characterization of when $(RG)_{\varphi}^-$ is commutative in [\[11\]](#page-10-2) when $char(R) \neq 2, 3$ and in [\[4\]](#page-9-5) when φ is the classical involution and char(R) \neq 2. The case char(R) = 3 was left as an open problem in [\[11\]](#page-10-2), and the case char(R) = 2 has been dealt with in [\[2,](#page-9-7) [12\]](#page-10-4) because then $(RG)_{\varphi}$ coincides with the set of φ -symmetric elements of RG.

So, throughout the paper we assume $char(R) \neq 2$. The center of G is denoted by $Z(G)$, the additive commutator $\alpha\beta - \beta\alpha$ of $\alpha, \beta \in RG$ is denoted $[\alpha, \beta]$, and the multiplicative commutator $ghg^{-1}h^{-1}$ of $g, h \in G$ is denoted by (g, h) .

As mentioned above, the Theorem has been proved in [\[11\]](#page-10-2) provided $R_2 = \{0\}$ and $char(R) \neq 3$. Theorem [2.5](#page-2-0) shows the result holds in case $R_2 \neq 0$ and Theorem [3.8](#page-7-0) shows that it also holds if $char(R) = 3.$

2 Rings with elements of additive order 2

We begin with recalling some technical results from [\[11\]](#page-10-2). The first lemma shows that the group generated by the non-fixed elements has index at most 2.

Lemma 2.1 [\[11,](#page-10-2) Lemma 2.3] *If* φ *is non-trivial then the subgroup* $K = \langle g \in G \mid g \notin G_{\varphi} \rangle$ *has index at most* 2 *in* G*.*

Lemma 2.2 [\[11,](#page-10-2) Lemma 1.1 and Lemma 1.2] Let R be a commutative ring with char(R) $\neq 2, 3$. Let $g, h \in G \setminus G_\varphi$ be two non-commuting elements. If $(RG)_{\varphi}^-$ is commutative then one of the following *conditions holds*

1. $gh \in G_{\varphi}, hg \in G_{\varphi}, h\varphi(g) = g\varphi(h)$ and $\varphi(h)g = \varphi(g)h$. 2. $gh = h\varphi(g) = \varphi(h)g = \varphi(g)\varphi(h)$ *and* $char(R) = 4$ *.*

Note that, if non-commuting elements $g, h \in G \setminus G_{\varphi}$ satisfy condition (2) in the lemma then $h^{-1}gh = \varphi(g)$. Non-commutative groups G with an involution φ such that $h^{-1}gh \in \{g, \varphi(g)\}\$ for all $g, h \in G$ have been described in [\[9,](#page-10-5) Theorem III.3.3]. These are precisely the groups G with a unique non-trivial commutator and that satisfy the *lack of commutativity property* ("LC" for short). The latter means that for any pair of elements $g, h \in G$ it is the case that $gh = hg$ if and only if either $g \in Z(G)$ or $h \in Z(G)$ or $gh \in Z(G)$. It turns out [\[9,](#page-10-5) Proposition III.3.6] that such groups are precisely those non-commutative groups G with $G/Z(G) \cong C_2 \times C_2$, where C_2 denotes the cyclic group of order 2.

In the next lemma we give the structure of the group generated by two elements $g, h \in G \setminus G_\varphi$ satisfying (2) of Lemma [2.2.](#page-1-0)

Lemma 2.3 [\[11,](#page-10-2) Lemma 3.1] *Let* R *be a commutative ring with* char(R) = 4*. Suppose* $g, h \in G \backslash G_\varphi$ *are non-commuting elements that satisfy* (2) *of Lemma [2.2.](#page-1-0) If* $(RG)_{\varphi}^-$ *is commutative then the group* $H = \langle g, h, \varphi(g), \varphi(h) \rangle = \langle g, h \rangle$ *satisfies the LC-property and has a unique non-trivial commutator* s *and the involution restricted to* H *is given by* $\varphi(h) = sh$ *if* $h \in H \setminus Z(H)$ *and* $\varphi(h) = h$ *if* $h \in Z(H)$ *.*

From the next lemma it follows that if $R_2 \neq \{0\}$ and char(R) $\neq 4$ then any two elements of $G \setminus G_{\varphi}$ that satisfy condition (1) of Lemma [2.2](#page-1-0) must commute.

Lemma 2.4 *Assume* $R_2 \neq \{0\}$ *and* $(RG)_{\varphi}^-$ *is commutative. Let* $g, h \in G$ *and suppose* $(g, h) \neq 1$ *.*

- *1. If* $g \in G_{\varphi}$ *and* $h \notin G_{\varphi}$ *then* $gh = \varphi(h)g$ *and* $hg = g\varphi(h)$ *.*
- 2. If $g, h \in G \setminus G_\varphi$ then $gh \notin G_\varphi$ (in particular, g and h do not satisfy condition (1) of *Lemma [2.2\)](#page-1-0).*
- *3. If* $g, h \in G \setminus G_\varphi$ *then* $char(R) = 4$, $\langle g, h \rangle$ *is LC with a unique non-trivial commutator and* $\varphi(g) = (g, h)g$ and $\varphi(h) = (g, h)h$. In particular, if $char(R) \neq 4$ then $K = \langle g \in G \mid g \notin G_{\varphi} \rangle$ *is abelian.*

Proof. Let $0 \neq r \in R_2$.

(1) Since $(RG)_{\varphi}^-$ is commutative, we have that $0 = [rg, h - \varphi(h)] = r(gh + g\varphi(h) + hg + \varphi(h)g)$. As $(g, h) \neq 1$ and $h \notin G_{\varphi}$, it follows that $gh = \varphi(h)g$ and $hg = g\varphi(h)$.

(2) Suppose $g, h \in G \setminus G_{\varphi}$. Assume $gh \in G_{\varphi}$. Then, by (1), $\varphi(h)\varphi(g)h = ghh = \varphi(h)gh$ and thus $g \in G_{\varphi}$, a contradiction.

(3) This follows at once from Lemma [2.2,](#page-1-0) (2) and Lemma [2.3](#page-2-1). \blacksquare

We now give a complete characterization of when $(RG)_{\varphi}^-$ is commutative provided $R_2 \neq \{0\}$ (and thus $char(R) \neq 3$).

Theorem 2.5 *Let* R *be a commutative ring with elements of additive order* 2*. Assume* G *is a non-abelian group and* φ *is an involution on* G. Then, $(RG)_{\varphi}^-$ *is commutative if and only if one of the following conditions holds:*

(a) $K = \langle g \in G | g \notin G_{\varphi} \rangle$ *is abelian (and thus* $G = K \cup Kx$ *, where* $x \in G_{\varphi}$ *, and* $\varphi(k) = xkx^{-1}$ *for all* $k \in K$ *)*, and $R_2^2 = \{0\}$.

(b) char(R) = 4, $|G'| = 2$, $G/G' = (G/G')_{\varphi}$, $g^2 \in G_{\varphi}$ for all $g \in G$, and G_{φ} is commutative in $case \ R_2^2 \neq \{0\}.$

Proof. Let G be a non-abelian group and φ an involution on G. Assume $(RG)_{\varphi}^-$ is commutative. Notice that Lemma [2.1](#page-1-1) implies that if $K = \langle g \in G | g \notin G_{\varphi} \rangle$ is abelian (and thus $K \neq G$) then $G = K \cup Kx$ for some $x \in G_{\varphi}$. Furthermore, one gets that $\varphi(k) = xkx^{-1}$ for all $k \in K$. Indeed, since $x \notin K$ it follows that $x \notin K$ and hence $xk = \varphi(xk) = \varphi(k)x$ and therefore $\varphi(k) = xkx^{-1}$. Also, since x is not central, we get that $x_k \neq kx$ for some $k \in K$. Now, for any $r_1, r_2 \in R_2$ we have that $x, kx \in G_\varphi$ and thus, by assumption, $r_1r_2(kx - kx^2) = [r_1x, r_2kx] = 0$. Since $xkx \neq kx^2$, it follows that $r_1r_2 = 0$. Consequently, $R_2^2 = \{0\}$. So, condition (a) follows.

If char(R) \neq 4 then it follows from Lemma [2.4.](#page-2-2)(3) that K is abelian. Hence, by the above, condition (a) follows.

So, to prove the necessity of the mentioned conditions, we are left to deal with the case that $char(R) = 4$ and K is not abelian. We need to prove that condition (b) holds. Because of Lemma [2.4](#page-2-2) (3), we also know that $H = \langle x, y \rangle$ is LC with a unique non-trivial commutator and $\varphi(h) = (x, y)h$ if $h \in H \setminus Z(H)$ and $\varphi(h) = h$ if $h \in Z(H)$.

Now we claim that for all $g \in G \setminus G_\varphi$ we have that $g^2 \in G_\varphi$ and $g^{-1}\varphi(g) = (x, y)$ (in particular, $G/G' = (G/G')_{\varphi}$). Indeed, let $g \in G \setminus G_{\varphi}$. If $(g, x) \neq 1$ then by Lemma [2.4](#page-2-2) (3) $g^2 \in G_{\varphi}$ and $g^{-1}\varphi(g) = x^{-1}\varphi(x) = (x, y)$. Similarly, if $(g, y) \neq 1$ then $g^{-1}\varphi(g) = y^{-1}\varphi(y) = (x, y)$. Assume now that $(g, x) = (g, y) = 1$. If $gx \in G_\varphi$ then $g^2x^2 = (gx)^2 = \varphi((gx)^2) = \varphi(g^2x^2) = \varphi(g^2)x^2$ and hence $g^2 \in G_{\varphi}$. Moreover, in this case, $gx = \varphi(g)\varphi(x)$ and hence $g^{-1}\varphi(g) = x\varphi(x^{-1}) =$ $x^{-1}\varphi(x) = (x, y)$ as desired. If $gx \notin G_{\varphi}$ then, by Lemma [2.4](#page-2-2) (3) and since $(gx, y) = (x, y) \neq 1$, we get that $(gx)^{-1}\varphi(gx) = (gx, y) = (x, y) = x^{-1}\varphi(x)$. Hence, $x^{-1}g^{-1}\varphi(x)\varphi(g) = x^{-1}\varphi(x)$ and thus $g^{-1}\varphi(g)\varphi(x) = g^{-1}\varphi(x)\varphi(g) = \varphi(x)$. So, $g = \varphi(g)$, a contradiction. This finishes the proof of the claim.

Next we show that $G' = \langle (x, y) \rangle = \{1, (x, y)\}\$ (and thus $G' \subseteq G_{\varphi}$). Indeed, let $g, h \in G$ such that $(g, h) \neq 1$. If $g, h \notin G_{\varphi}$ then by the previous claim and Lemma [2.4](#page-2-2) (3) it follows that (g, h) = $g^{-1}\varphi(g) = x^{-1}\varphi(x) = (x, y)$, as desired. If $g \in G_{\varphi}$ and $h \notin G_{\varphi}$ then, by Lemma [2.4](#page-2-2) (1), $gh = \varphi(h)g$ and hence by the previous claim we get that $(x, y) = \varphi(x)x^{-1} = \varphi(h)h^{-1} = ghg^{-1}h^{-1} = (g, h)$. Finally if $g, h \in G_\varphi$ then $hg \notin G_\varphi$ (because otherwise $(g, h) = 1$), and hence by the previous claim $(x, y) = \varphi(x)x^{-1} = \varphi((hg))(hg)^{-1} = ghg^{-1}h^{-1} = (g, h)$, as desired.

To finish the prove of the necessity, we remark that if $R_2^2 \neq \{0\}$ then G_φ is commutative. Indeed, let $r_1, r_2 \in R_2$ be so that $r_1r_2 \neq 0$ and let $g_1, g_2 \in G_{\varphi}$. Since $(RG)_{\varphi}^-$ is commutative, we have that $r_1r_2(g_1g_2 - g_2g_1) = [r_1g_1, r_2g_2] = 0$. Hence $(g_1, g_2) = 1$.

In order to prove the sufficiency we need to show that the elements in

$$
\mathcal{S} = \{ g - \varphi(g) \mid g \in G, g \notin G_{\varphi} \} \cup \{ rg \mid g \in G_{\varphi}, r \in R_2 \}
$$

commute.

First assume G satisfies condition (a). So $G = K \cup Kx$ with $x \in G_{\varphi}$ and K abelian. We need to show that $[g - \varphi(g), r_1h_1] = 0$ and $[r_1h_1, r_2h_2] = 0$ for $g \in G \setminus G_\varphi$, $h_1, h_2 \in G_\varphi$ and $r_1, r_2 \in R_2$ with $(g, h_1) \neq 1$ and $(h_1, h_2) \neq 1$. The later equality is obviously satisfied because of the assumptions. To prove the former equality, we note that, by Lemma [2.4](#page-2-2) (1), $h_1g = \varphi(g)h_1$ and $gh_1 = h_1\varphi(g)$. Hence,

$$
[g - \varphi(g), r_1 h_1] = r_1 gh_1 - r_1 \varphi(g) h_1 - r_1 h_1 g + r_1 h_1 \varphi(g)
$$

= $r_1 gh_1 + r_1 h_1 g + r_1 h_1 g + r_1 gh_1$
= $2r_1 gh_1 + 2r_1 h_1 g$
= 0,

as desired.

Second, assume G satisfies (b) and that char(R) = 4. Notice that in this case if $g \notin G_{\varphi}$ then $g^{-1}\varphi(g) = g\varphi(g^{-1})$ is central and equal to the unique commutator of G. Let $g, h \in G$ with $(g, h) \neq 1$ and let $r_1, r_2 \in R_2$. If $g, h \in G_\varphi$, then the assumptions imply that $R_2^2 = \{0\}$ and thus $[r_1g, r_2h] = 0$, as desired. If $g \notin G_{\varphi}$ and $h \in G_{\varphi}$ then

$$
[g - \varphi(g), r_1 h] = r_1 gh - r_1 \varphi(g) h - r_1 hg + r_1 h \varphi(g)
$$

= $r_1 gh + r_1 h g \varphi(g^{-1}) \varphi(g) + r_1 hg - r_1 \varphi(g) \varphi(g^{-1}) gh$
= $r_1 gh + r_1 hg + r_1 hg + r_1 gh$
= $2r_1 hg + 2r_1 gh = 0$.

Finally, if $g, h \in G \setminus G_{\varphi}$ then

$$
[g - \varphi(g), h - \varphi(h)] = gh - g\varphi(h) - \varphi(g)h + \varphi(g)\varphi(h) - hg + h\varphi(g) + \varphi(h)g - \varphi(h)\varphi(g)
$$

= gh - hg - hg + gh - hg + gh - hg = 4gh - 4hg = 0

which finishes the proof of the theorem.

For the classical involution $*$ on G we get the following consequence.

Corollary 2.6 *Let* R *be a commutative ring with elements of additive order* 2*. Let* G *be a nonabelian group. Denote by* $*$ *the classical involution. Then* $(RG)_{*}^{-}$ *is commutative if and only if one of the following conditions holds:*

- *1.* $G = K \rtimes \langle x \rangle$ where $K = \langle g | g^2 \neq 1 \rangle$, K is abelian, $x^2 = 1$, $x k x = k^{-1}$ for all $k \in K$ and $R_2^2 = \{0\}.$
- 2. char $(R) = 4$, G has exponent 4, G' is a cyclic group of order 2, G/G' is an elementary abelian 2-subgroup and elements of order 2 commute if $R_2^2 \neq \{0\}$.

3 Rings of characteristic three

In this section we determine when $(RG)_{\varphi}^-$ is commutative if $char(R) = 3$ (and thus $R_2 = \{0\}$). Again we begin by recalling two technical lemmas from [\[11\]](#page-10-2).

Lemma 3.1 [\[11,](#page-10-2) Lemma 1.3] *Let* R *be a commutative ring with char(R)* \neq *2 <i>and let* $g \in G \setminus G_{\varphi}$ *. If* $(RG)_{\varphi}^-$ is commutative then one of the following conditions holds:

- *1.* $g\varphi(g) = \varphi(g)g$.
- 2. $g^2 \in G_{\varphi}$.

Lemma 3.2 [\[11,](#page-10-2) Lemma 1.1] *Let* R *be a commutative ring with* char(R) = 3*. Let* $g, h \in G \backslash G_\varphi$ *be two non-commuting elements.* If $(RG)_{\varphi}^-$ *is commutative then one of the following conditions holds*

1.
$$
gh \in G_{\varphi}
$$
, $hg \in G_{\varphi}$, $h\varphi(g) = g\varphi(h)$ and $\varphi(h)g = \varphi(g)h$.

2. $gh \in G_{\varphi}, hg \in G_{\varphi}, h\varphi(g) = \varphi(g)h$.

3.
$$
gh \in G_{\varphi}
$$
, $hg = \varphi(g)h = g\varphi(h)$.

4. $gh = h\varphi(g) = \varphi(g)\varphi(h), \varphi(h)g = \varphi(g)h.$

- 5. $gh = \varphi(h)g = \varphi(g)\varphi(h)$, $h\varphi(g) = g\varphi(h)$.
- 6. $hg \in G_{\varphi}, gh = \varphi(h)g = h\varphi(g)$.

The following lemma was proved in [\[11\]](#page-10-2) in the case when $char(R)$ is distinct from both 2 and 3.

Lemma 3.3 Let R be a commutative ring. Let $g, h \in G \setminus G_\varphi$ be non-commuting elements that *satisfy* (2) *of Lemma [3.2.](#page-4-0) If* $char(R) = 3$ *then* g, h *also satisfy* (1) *of Lemma 3.2.*

Proof. Consider the element $\varphi(g)hg \in G$. Since $h \notin G_{\varphi}$ we have that $\varphi(g)hg \notin G_{\varphi}$. Also $\varphi(g)hg$ and h do not commute because, by assumption, $h\varphi(g) = \varphi(g)h$ and $gh \neq hg$. Assume that g and h satisfy (2) of Lemma [3.2.](#page-4-0) We claim that then $\varphi(h)g = \varphi(g)h$.

We deal with two mutually exclusive cases. First, assume that $\varphi(g)hgh \in G_{\varphi}$, i.e., $\varphi(g)hgh =$ $\varphi(\varphi(g)hgh) = \varphi(h)\varphi(g)\varphi(h)g = \varphi(h)hg^2$ since $hg \in G_{\varphi}$. If $(g^2, h) = 1$ we obtain that $\varphi(g)h =$ $\varphi(h)g$, as desired. So, to deal with this case, we may assume that $(g^2, h) \neq 1$. If $g^2 \in G_{\varphi}$ then, using (2), we observe that $\varphi(h)hg^2 = \varphi(g)hgh = \varphi(g^2)\varphi(h)h = g^2\varphi(h)h = \varphi(h)g^2h$. Hence, we get that $(g^2, h) = 1$, a contradiction. In the rest of the proof we will several times use (without referring to this) that gh , $hg \in G_{\varphi}$ and $(g, \varphi(h)) = 1 = (h, \varphi(g))$. Moreover, since $g^2 \notin G_{\varphi}$, we also have that $g\varphi(g) = \varphi(g)g$, by Lemma [3.1.](#page-4-1)

So, if $\varphi(g)hgh \in G_{\varphi}$ then we may assume that $g^2 \notin G_{\varphi}$ and $(g^2, h) \neq 1$. Hence, g^2 and h satisfy one of the six conditions of Lemma [3.2.](#page-4-0) We now show that this situation can not occur. Assume first that $g^2h \in G_{\varphi}$. Then, $g^2h = \varphi(g^2h) = \varphi(h)\varphi(g^2) = gh\varphi(g) = g\varphi(g)h$ and thus $g \in G_{\varphi}$, a contradiction. Therefore g^2 and h do not satisfy conditions (1) – (3) of Lemma [3.2.](#page-4-0) If g^2 and h satisfy (4) of Lemma [3.2](#page-4-0) then $g^2h = h\varphi(g^2) = \varphi(g^2)h$ and hence $g^2 \in G_{\varphi}$, a contradiction. Finally, if g^2 and h satisfy either (5) or (6) of Lemma [3.2](#page-4-0) then $g^2h = \varphi(h)g^2 = g^2\varphi(h)$ and thus $h \in G_{\varphi}$, a contradiction. This finishes the proof of the first case.

Second, assume that $\varphi(g)hgh \notin G_{\varphi}$. Then $\varphi(g)hg$ and h satisfy one of the conditions $(4)-(6)$ of Lemma [3.2.](#page-4-0) We show that all these lead to a contradiction and hence that this case also can not occur. If $\varphi(q)hg$ and h satisfy (4) of Lemma [3.2,](#page-4-0) then $\varphi(q)hqh = h\varphi(\varphi(q)hq) = h\varphi(q)\varphi(h)q =$ $h\varphi(hg)g = h^2g^2$ and thus $\varphi(g)gh = hg^2 = \varphi(g)\varphi(h)g$; so $gh = \varphi(h)g$ and hence $g \in G_{\varphi}$, a contradiction.

Suppose that $\varphi(g)hg$ and h satisfy (5) or (6) of Lemma [3.2.](#page-4-0) Then

$$
\varphi(g)hgh = \varphi(h)\varphi(g)hg \tag{1}
$$

First assume that $g^2 \in G_\varphi$ then we have that $\varphi(h)hg^2 = g^2\varphi(h)h = \varphi(g^2)\varphi(h)h = \varphi(g)hgh$. On the other hand $\varphi(h)\varphi(g)hg = \varphi(h)h\varphi(g)g$. Thus, by [\(1\)](#page-5-0) we get that $g \in G_{\varphi}$, a contradiction. Therefore $g^2 \notin G_{\varphi}$ and hence, by Lema [3.1](#page-4-1) we get that $(g, \varphi(g)) = 1$. If also $(h, \varphi(h)) = 1$ then $\varphi(g)hgh = hg\varphi(g)h$ and on the other hand, $\varphi(h)\varphi(g)hg = hg\varphi(h)\varphi(g)$. Then, by [\(1\)](#page-5-0), we get that $\varphi(g)h = \varphi(h)\varphi(g) = gh$ and thus $g \in G_{\varphi}$, a contradiction. So, again by Lemma [3.1](#page-4-1) we have that $h^2 \in G_{\varphi}$. Therefore $\varphi(h)\varphi(g)hg = gh^2g = h^2g^2$ and on the other hand $\varphi(g)hgh = h\varphi(g)gh$. Thus, by [\(1\)](#page-5-0), we have that $\varphi(g)gh = hg^2 = \varphi(g)\varphi(h)g$. Therefore $\varphi(h)\varphi(g) = gh = \varphi(h)g$ and hence $g \in G_{\varphi}$, again a contradiction. So $\varphi(g)hg$ and h do not satisfy neither (5) nor (6) of Lemma [3.2.](#page-4-0)

So, we have proved that if (2) of Lemma [3.2](#page-4-0) holds for non-commuting elements $g, h \in G \backslash G_\varphi$ then $\varphi(h)g = \varphi(g)h$. Since $(h, \varphi(g)) = 1 = (g, \varphi(h))$ it also follows that $h\varphi(g) = g\varphi(h)$. Consequently, we have shown that (1) of Lemma [3.2](#page-4-0) holds for g and h. \blacksquare

Lemma 3.4 *Let* R *be a commutative ring. Let* $g, h \in G \setminus G_\varphi$ *be non-commuting elements.*

- *1. If* g and h satisfy (1) *(or* (2)*)* of Lemma [3.2](#page-4-0) then $g^3, h^3 \notin G_\varphi$
- 2. If g and h satisfy one of the conditions $(3) (6)$ of Lemma [3.2](#page-4-0) then g^3 , $h^3 \in G_\varphi$ and g^3 , $h^3 \in G_\varphi$ $Z(\langle q, h, \varphi(q), \varphi(h)\rangle).$

Proof. 1. Let $g, h \in G \setminus G_\varphi$ be non-commuting elements. Assume that g and h satisfy (1) of Lemma [3.2.](#page-4-0) We prove by contradiction that $g^3 \notin G_\varphi$. So, suppose that $g^3 \in G_\varphi$. Since $g \notin G_\varphi$, it follows that $g^2 \notin G_{\varphi}$. Also by (1) of Lemma [3.2](#page-4-0) we have that

$$
g^{3}h = g^{2}\varphi(h)\varphi(g) = gh\varphi(g^{2}) = \varphi(h)\varphi(g^{3}) = \varphi(h)g^{3}
$$
\n(2)

Notice that by [\(2\)](#page-6-0) it follows that $(g^2, h) \neq 1$, because otherwise $gh = \varphi(h)g$, a contradiction. Therefore g^2 and h satisfy one of the conditions $(1) - (6)$ of Lemma [3.2.](#page-4-0) Assume first that $g^2 h \in G_\varphi$. Then by [\(2\)](#page-6-0) we have that $g\varphi(h)\varphi(g^2) = gg^2h = \varphi(h)\varphi(g^3)$. Hence $g\varphi(h) = \varphi(h)\varphi(g) = gh$ and thus $h \in G_{\varphi}$, a contradiction. Therefore g^2 and h do not satisfy conditions $(1) - (3)$ of Lemma [3.2.](#page-4-0) Second, assume that $g^2h = \varphi(h)g^2$. Then, by [\(2\)](#page-6-0), it follows that $\varphi(h)g^3 = gg^2h = g\varphi(h)g^2$ and thus $(g, \varphi(h)) = 1$. Therefore, again by [\(2\)](#page-6-0), we get that $h \in G_{\varphi}$, a contradiction. So, g^2 and h do not satisfy conditions $(5) - (6)$. Hence, g^2 and h satisfy (4). Then, since $(g, \varphi(g)) = 1$ by Lemma [3.1,](#page-4-1) we have that $\varphi(g^3)h = gg^2h = gh\varphi(g^2) = g\varphi(g^2)\varphi(h) = \varphi(g^2)g\varphi(h)$. Hence $\varphi(g)h = g\varphi(h) = h\varphi(g)$. Consequently, by [\(2\)](#page-6-0), we get that $h \in G_{\varphi}$, a contradiction. This finishes the proof of the fact that $g^3 \notin G_\varphi$. Because of the symmetry in g and h in condition (1) of Lemma [3.2,](#page-4-0) we thus also obtain that $h^{3'} \notin G_{\varphi}$.

2. Notice that if in (3) of Lemma [3.2](#page-4-0) we interchange the roles of q and h then we obtain (6), if we change h by $\varphi(h)$ we have (5) and finally if we change g by $\varphi(g)$ we have (4). Therefore it is enough to show the result for (3).

So, assume that $g, h \in G \backslash G_\varphi$ are non-commuting elements that satisfy (3) of Lemma [3.2.](#page-4-0) Then $g^3h = g^2\varphi(h)\varphi(g) = g\varphi(g)h\varphi(g) = g\varphi(g^2)\varphi(h)$ and therefore, since $h \notin G_{\varphi}$, it follows that $g^2 \notin G_{\varphi}$. Thus, by Lemma [3.1,](#page-4-1) we get that $(g, \varphi(g)) = 1$. Consequently, $g^3 h = \varphi(g^2) g \varphi(h) = \varphi(g^3) h$ and therefore $g^3 \in G_\varphi$. Analogously we obtain that $h^3 \in G_\varphi$. Moreover, $g^3 h = g^2 \varphi(h)\varphi(g) = ghg\varphi(g)$ $\varphi(h)\varphi(g)g\varphi(g) = \varphi(h)g\varphi(g)\varphi(g) = h\varphi(g^3) = hg^3$. So, $g^3h = hg^3$ and thus also $\varphi(h)g^3 = g^3\varphi(h)$, as desired. Similarly we get that $h^3 \in Z(\langle g, h, \varphi(g), \varphi(h) \rangle)$.

Remark 3.5 *Notice that Lemma [3.4](#page-5-1) implies that if* $g, h, x, y \in G \setminus G_\varphi$ *are such that* g *and* h *are non-commuting elements satisfying condition* (1) *of Lemma [3.2](#page-4-0) and* x *and* y *are non-commuting elements satisfying one of the conditions* (3) − (6) *of Lemma [3.2](#page-4-0) then* x *and* y *commute with both* g *and* h*.*

Lemma 3.6 Let R be a commutative ring with $char(R) = 3$ and assume RG_{φ}^- is commutative. *If there exist non-commuting* $g, h \in G \setminus G_\varphi$ *so that* g and h *satisfy* (1) of Lemma [3.2](#page-4-0) then all $x, y \in G \setminus G_{\varphi}$ *satisfy* (1) *of Lemma [3.2.](#page-4-0)*

Proof. Let $g, h \in G \backslash G_\varphi$ be non-commuting elements so that g and h satisfy (1) of Lemma [3.2,](#page-4-0) that is, gh, hg, $g\varphi(h)$ and $\varphi(g)h$ are elements of G_{φ} . Also, by Lemma [3.4,](#page-5-1) we have that g^3 , $h^3 \notin G_{\varphi}$.

Let $x \in G \setminus G_\varphi$. Then, by Lemma [3.2](#page-4-0) and Lemma [3.4,](#page-5-1) it follows that $(g, x) = 1$ or $gx \in G_\varphi$. We claim that $gx \in G_{\varphi}$. In order to prove this claim suppose that $gx \notin G_{\varphi}$ and thus $(g, x) = 1$. Again, by Lemma [3.2](#page-4-0) and Lemma [3.4,](#page-5-1) it follows that $(h, x) = 1$ or $xh \in G_{\varphi}$; and $(gx, h) = 1$ or $gxh \in G_{\varphi}$. Assume first that $(gx, h) = 1$, that is, $gxh = hgx$. Since $gh \neq gh$ we get that $hx \neq xh$ and thus $xh \in G_{\varphi}$. Therefore, $hgx = gxh = g\varphi(h)\varphi(x) = h\varphi(g)\varphi(x) = h\varphi(gx)$ and thus $gx \in G_{\varphi}$, a contradiction. Second assume that $gxh \in G_{\varphi}$. Then $gxh = \varphi(h)\varphi(x)\varphi(g) = \varphi(h)\varphi(g)\varphi(x) =$ $gh\varphi(x)$ and hence $xh = h\varphi(x)$. Therefore, and since $x \notin G_{\varphi}$, we get that $xh \neq hx$ and thus $xh \in G_{\varphi}$. Then $\varphi(h)\varphi(x) = xh = h\varphi(x)$ and hence $h \in G_{\varphi}$, again a contradiction. This finishes the proof of the claim.

Now, let $x, y \in G \setminus G_{\varphi}$. We need to prove that x and y satisfy (1) of Lemma [3.2.](#page-4-0) First we deal with the case that $(x, y) \neq 1$. Because of Lemma [3.3,](#page-5-2) we only have to show that it is impossible that x and y satisfy one of the conditions $(3) - (6)$ of Lemma [3.2.](#page-4-0) So suppose the contrary. Then, by Remark [3.5,](#page-6-1) $(g, x) = 1 = (g, y)$. Also, by Lemma [3.4,](#page-5-1) $x^3, y^3 \in G_\varphi$. By the previous claim we have that $gx \in G_{\varphi}$. Consequently, $g^3x^3 = (gx)^3 = \varphi(gx)^3 = \varphi(g^3)\varphi(x^3) = \varphi(g^3)x^3$, and thus $g^3 \in G_\varphi$, a contradiction. So, if $xy \neq yx$ then x and y satisfy (1) of Lemma [3.2.](#page-4-0)

Finally, assume $x, y \in G \backslash G_\varphi$ and $(x, y) = 1$. Then $xy \in G_\varphi$. Indeed, suppose the contrary, that is assume $xy \notin G_{\varphi}$. Hence, by the above claim, $gxy \in G_{\varphi}$. Thus $gyx = gxy = \varphi(y)\varphi(x)\varphi(y)$ $\varphi(y)gx$, because $gx \in G_{\varphi}$. Therefore $gy = \varphi(y)g$. Since $gy \in G_{\varphi}$, it follows that $\varphi(y)g = gy =$ $\varphi(y)\varphi(y)$ and thus $g \in G_{\varphi}$, a contradiction. Hence, indeed $yx = xy \in G_{\varphi}$. Replacing y by $\varphi(y)$ we thus also get that $x\varphi(y) \in G_{\varphi}$ if $(x, \varphi(y)) = 1$. If, on the other hand, $(x, \varphi(y)) \neq 1$ then the previous implies that again $x\varphi(y) \in G_{\varphi}$. Similarly, $\varphi(x)y \in G_{\varphi}$. Consequently, we have shown that x and y satisfy (1) of Lemma [3.2.](#page-4-0) \blacksquare

Lemma 3.7 Let R be a commutative ring with $char(R) = 3$. Let $g, h \in G \setminus G_{\varphi}$ be non-commuting *elements satisfying any of the conditions* (3) – (6) *of Lemma [3.2.](#page-4-0) Then* $\langle g^{-1}\varphi(g) \rangle = \langle h^{-1}\varphi(h) \rangle =$ $\langle (g, h) \rangle$ and $(g, h)^3 = 1$.

Proof. Let $g, h \in G \backslash G_\varphi$ be as in the statement of the Lemma. Because of Lemma [3.4,](#page-5-1) $g^3, h^3 \in G_\varphi$. Therefore $g^2, h^2 \notin G_\varphi$, because $g, h \notin G_\varphi$. Hence, by Lemma [3.1,](#page-4-1) it follows that $(g, \varphi(g)) = 1$ $(h, \varphi(h)).$

First, assume that g and h satisfy (3) of Lemma [3.2.](#page-4-0) Hence, $\varphi(g) = hgh^{-1}$ and $\varphi(h) = g^{-1}hg$. Therefore $g^{-1}\varphi(g) = g^{-1}hgh^{-1} = \varphi(h)h^{-1} = h^{-1}\varphi(h)$. Thus, $h^{-1}\varphi(h) = g^{-1}\varphi(g) = \varphi(g)g^{-1} =$ $hgh^{-1}g^{-1} = (g,h)^{-1}$, as desired.

Second, assume that g and h satisfy (4) of Lemma [3.2.](#page-4-0) Then $\varphi(g) = h^{-1}gh$ and $\varphi(h) =$ $\varphi(g^{-1})gh = h^{-1}g^{-1}hgh$. Therefore $(g, \varphi(g)) = (g, h^{-1}gh) = 1 = (h, \varphi(h)) = (h, g^{-1}hg)$. Thus, $g^{-1}\varphi(g) = g^{-1}h^{-1}gh = h^{-1}ghg^{-1} = ghg^{-1}h^{-1} = (g,h)$ and $h^{-1}\varphi(h) = h^{-1}g^{-1}hg = (g^{-1}\varphi(g))^{-1}$, again as desired.

Third, assume that g and h satisfy (5) of Lemma [3.2.](#page-4-0) Then $\varphi(h) = ghg^{-1}$ and $\varphi(g) =$ $gh\varphi(h)^{-1} = ghgh^{-1}g^{-1}$. Therefore $(h, \varphi(h)) = (h, ghg^{-1}) = 1 = (g, \varphi(g)) = (g, hgh^{-1})$. Thus, $g^{-1}\varphi(g) = hgh^{-1}g^{-1} = (g,h)^{-1}$ and $h^{-1}\varphi(h) = h^{-1}ghg^{-1} = ghg^{-1}h^{-1} = (g^{-1}\varphi(g))^{-1}$, again as desired.

Fourth, assume that g and h satisfy (6) of Lemma [3.2.](#page-4-0) Then $\varphi(h) = ghg^{-1}$ and $\varphi(g) = h^{-1}gh$. Therefore $(h, \varphi(h)) = (h, ghg^{-1}) = 1 = (g, \varphi(g)) = (g, h^{-1}gh)$. Thus, $g^{-1}\varphi(g) = g^{-1}h^{-1}gh =$ $h^{-1}ghg^{-1} = ghg^{-1}h^{-1} = (g,h)$ and $h^{-1}\varphi(h) = h^{-1}ghg^{-1} = ghg^{-1}h^{-1} = g^{-1}\varphi(g)$, as desired.

To finish the proof of the lemma notice that since $g^3 \in G_\varphi$ and $(g, \varphi(g)) = 1$ it follows that $(g^{-1}\varphi(g))^3 = 1$ and therefore $(g,h)^3 = 1$.

Theorem 3.8 Let R be a commutative ring with $char(R) = 3$. Suppose G is a non-abelian group and φ *is an involution on* G. Then $(RG)_{\varphi}^-$ *is commutative if and only if one of the following conditions holds:*

(a)
$$
K = \langle g \in G | g \notin G_{\varphi} \rangle
$$
 is abelian, $G = K \cup Kx$ where $x \in G_{\varphi}$ and $\varphi(k) = xkx^{-1}$ for all $k \in K$.

(b) G contains an abelian subgroup of index 2 that is contained in G_{φ} .

(c)
$$
|G'| = 3
$$
, $(G/G') = (G/G')_{\varphi}$ and $g^3 \in G_{\varphi}$ for all $g \in G$.

Proof. Assume that there exist non-commuting elements $g, h \in G \setminus G_\varphi$ so that g and h satisfy (1) of Lemma [3.2.](#page-4-0) Then, by Lemma [3.6,](#page-6-2) all $x, y \in G \setminus G_\varphi$ satisfy (1) of Lemma 3.2. Because of all the stated Lemmas, one now obtains, exactly as in the proof of Theorem 2.1 in [\[11\]](#page-10-2) that condition (a) or (b) holds.

So now suppose that there do not exist non-commuting elements $g, h \in G \setminus G_{\varphi}$ satisfying condition (1) (and thus not (2) , by Lemma [3.3](#page-5-2)) of Lemma [3.2.](#page-4-0) Then, all non-commuting elements $x, y \in G\setminus G_{\varphi}$ satisfying one of the conditions $(3) - (6)$ of Lemma [3.2.](#page-4-0) In particular, by Lemma [3.4,](#page-5-1) $x^3, y^3 \in G_\varphi$. Since $x \notin G_\varphi$, we thus have that $x^2 \notin G_\varphi$ and thus, by Lemma [3.1,](#page-4-1) $(\varphi(x), x) = 1$.

We claim that $g^3 \in G_\varphi$ for all $g \in G$. So, let $g \in G$. In case $(g, x) \neq 1$ then it follows at once from Lemma [3.4](#page-5-1) that $g^3 \in G_\varphi$. If, on the other hand, $(g, x) = 1$, then we consider two mutually exclusive cases. First, assume $gx \notin G_{\varphi}$. Then, again by Lemma [3.4,](#page-5-1) $g^3x^3 = (gx)^3 \in G_{\varphi}$ and thus $g^3x^3 = \varphi(g^3)\varphi(x^3) = \varphi(g^3)x^3$ and thus $g^3 \in G_{\varphi}$. Second, assume $gx \in G_{\varphi}$. Thus $xg = gx = \varphi(x)\varphi(g) = \varphi(g)\varphi(x)$. Hence, by Lemma [3.7,](#page-7-1) $g^{-1}\varphi(g) = x\varphi(x)^{-1} = \varphi(x)^{-1}x = \varphi(g)g^{-1}$ is an element of order 3. Thus $1 = (g^{-1}\varphi(g))^3 = g^{-3}\varphi(g^3)$. So $g^3 \in G_\varphi$, as claimed.

Next, we claim that if $g \notin G_{\varphi}$ then $g^{-1}\varphi(g) \in \langle (x, y) \rangle$ (in particular $G/G' = (G/G')_{\varphi}$). Indeed, if $(g, x) \neq 1$, $(g, \varphi(x)) \neq 1$, $(g, \varphi(y)) \neq 1$ or $(g, y) \neq 1$ the result follows from Lemma [3.7.](#page-7-1) So assume that $(g, x) = (g, \varphi(x)) = (g, y) = (g, \varphi(y)) = 1$. If $gx \in G_{\varphi}$ then $gx = \varphi(gx) = \varphi(x)\varphi(g) =$ $\varphi(g)\varphi(x)$ and hence $g^{-1}\varphi(g) = x\varphi(x)^{-1} \in \langle (x,y) \rangle$ by Lemma [3.7](#page-7-1) as desired. Finally if $gx \notin G_{\varphi}$ then $(gx, y) = (x, y) \neq 1$. Then, by Lemma [3.7,](#page-7-1) we have that $(gx)^{-1}\varphi(gx) = g^{-1}\varphi(g)x^{-1}\varphi(x) \in \langle (x, y) \rangle$ and therefore, since $x^{-1}\varphi(x) \in \langle (x, y) \rangle$ we get that $g^{-1}\varphi(g) \in \langle (x, y) \rangle$, which finishes the proof of the claim.

To finish the proof of the necessity, we have to prove that $G' = \langle (x, y) \rangle$ and thus, by Lemma [3.7,](#page-7-1) $|G'| = 3$. In order to prove this, let $g, h \in G$ with $(g, h) \neq 1$. If $g, h \notin G_{\varphi}$ then by Lemma [3.7](#page-7-1) and the previous claim we have that $(g, h) \in \langle g^{-1}\varphi(g) \rangle = \langle (x, y) \rangle$. Next assume that $g \in G_{\varphi}$ and $h \notin G_{\varphi}$. If $gh \notin G_{\varphi}$ then, by Lemma [3.7](#page-7-1) and the previous claim, we have that $(g, h) = (gh, h) \in$ $\langle h^{-1}\varphi(h)\rangle = \langle (x, y)\rangle$, as desired. If $gh \in G_{\varphi}$ we have that $gh = \varphi(h)g$ and thus, by the previous claim, $ghg^{-1}h^{-1} = \varphi(h)h^{-1} \in \langle (x, y) \rangle$, as desired. Finally, assume that $g, h \in G_{\varphi}$. Then, since $(g, h) \neq 1$, it follows that $h^{-1}g^{-1} \notin G_{\varphi}$. Therefore, by the previous claim, $(g, h) = ghg^{-1}h^{-1} =$ $(h^{-1}g^{-1})^{-1}\varphi(h^{-1}g^{-1}) \in \langle (x,y) \rangle$ and the proof of the necessity concludes.

In order to prove the sufficiency, we need to show that the elements in $S = \{g - \varphi(g) \mid g \in G, g \notin G\}$ G_{φ} commute. If G satisfies conditions (a) or (b), then the proof is the same as the sufficiency proof of Theorem 2.1 in [\[11\]](#page-10-2). So, assume that G satisfies condition (c). Then, since $g^3 \in G_\varphi$, we have that $g^2 \notin G_{\varphi}$ for all $g \in G \setminus G_{\varphi}$ and hence, by Lemma [3.1,](#page-4-1) $(g, \varphi(g)) = 1$. Moreover, it follows from (c) that $\varphi(g) = t^i g$, where $G' = \langle t \rangle$, $i \in {\pm 1}$ for $g \notin G_{\varphi}$. Then, clearly, $(g, t) = 1$.

Let now $g, h \in G \setminus G_{\varphi}$. Then $\varphi(g) = t^i g$ and $\varphi(h) = t^j h$ with $i, j \in \{\pm 1\}$ and

$$
[g - \varphi(g), h - \varphi(h)] = gh - g\varphi(h) - \varphi(g)h + \varphi(g)\varphi(h) - hg + h\varphi(g) + \varphi(h)g - \varphi(h)\varphi(g)
$$

= gh - t^jgh - tⁱgh + tⁱt^jgh - hg + tⁱhg + t^jhg - tⁱt^jhg.

If $(g, h) = 1$ then clearly $[g - \varphi(g), h - \varphi(h)] = 0$. So, assume that $1 \neq (g, h) = t$. If $i = j$ then, since $char(R) = 3$,

$$
[g - \varphi(g), h - \varphi(h)] = gh - t^{i}gh - t^{i}gh + t^{-i}gh - hg + t^{i}hg + t^{i}hg - t^{-i}hg
$$

= $(1 - 2t^{i} + t^{-i})gh - (1 - 2t^{i} + t^{-i})hg$
= $(1 + t^{i} + t^{-i})(t - 1)hg$
= $(1 + t + t^{2})(t - 1)gh$
= 0

On the other hand if $i \neq j$ then, again since char(R) = 3,

$$
[g - \varphi(g), h - \varphi(h)] = gh - t^{j}gh - t^{i}gh + t^{i}t^{j}gh - hg + t^{i}hg + t^{j}hg - t^{i}t^{j}hg
$$

\n
$$
= 2gh - t^{j}gh - t^{i}gh - 2hg + t^{i}hg + t^{j}hg
$$

\n
$$
= (2 - t^{j} - t^{-i})gh - (2 - t^{i} - t^{j})hg
$$

\n
$$
= (2 - t^{j} - t^{i})(t - 1)hg
$$

\n
$$
= 2(1 + t + t^{2})(t - 1)hg
$$

\n
$$
= 0
$$

Similarly, if $(g, h) = t^{-1}$ one gets that $[g - \varphi(g), h - \varphi(h)] = 0$, which finishes the proof of the theorem. \blacksquare

Acknowledgements

Research supported by Onderzoeksraad of Vrije Universiteit Brussel, Fonds voor Wetenschappelijk Onderzoek (Vlaanderen), Flemish-Polish bilateral agreement BIL 01/31, FAPEMIG and CNPq. Proc. 300243/79-0(RN) of Brazil, D.G.I. of Spain and Fundación Séneca of Región de Murcia.

References

- [1] S. A. Amitsur, Identities in rings with involutions, *Israel J. Math.* 7 (1969) 63–68.
- [2] O. Broche Cristo, Commutativity of symmetric elements in group rings, *J. Group Theory*, 9, 5 (2006), 673–683
- [3] O. Broche, E. Jespers, M. Ruiz, Antisymmetric elements in group rings with an orientation morphism, *Forum Mathematicum*, to appear.
- [4] O. Broche Cristo and C. Polcino Milies, Commutativity of skew symmetric elements in group rings, *Proc. Edinburgh Math. Soc.* (2) 50 (2007), no. 1, 37-47.
- [5] A. Giambruno, C.Polcino Milies, Unitary units and skew elements in group algebras. *Manuscripta Math.* 111 (2003), no. 2, 195–209.
- [6] A. Giambruno, S.K. Sehgal, Lie nilpotence of group rings, *Comm. Algebra* 21 (1993), 4253– 4261.
- [7] A. Giambruno and S.K. Sehgal, Groups algebras whose Lie algebra of skew-symmetric elements is nilpotent, Groups, rings and algebras, 113–120, *Contemp. Math.*, 420, Amer. Math. Soc., Providence, RI, 2006.
- [8] J.Z. Gonçalves and D.S. Passman, Unitary units in group algebras, *Israel J. Math.* **125** (2001), 131–155.
- [9] E.G. Goodaire, E. Jespers, C. Polcino Milies, *Alternative Loop Rings*, North Holland Math. Studies 184, Elsevier, Amsterdam, 1996.
- [10] N. D. Gupta and F. Levin, On the Lie ideals of a ring, *Journal of Algebra*, 81 (1), (1993), 225–231.
- [11] E. Jespers and M. Ruiz, Antisymmetric elements in group rings, *J. Algebra and its Appl.*, 4, 4 (2005), 341–353.
- [12] E. Jespers, M. Ruiz, On symmetric elements and symmetric units in group rings, *Comm. Algebra*, 34, (2) (2006), 727-736.
- [13] S. P. Novikov, Algebraic construction and properties of hermitian analogues of K-theory over rings with involution from the viewpoint of Hamiltonian formalism, Applications to differential topology and the theory of characteristitic classes II, Izv. Akad. Nauk SSSR Ser. Mat. 34 (1970), 475-500; English transl. in Math. USSR Izv., 4 (1970).
- [14] Zalesskiı̆, A. E.; Smirnov, M. B. Lie algebra associated with linear group. *Comm. Algebra* 9 (1981), 20, 2075–2100.

C. Polcino Milies Manuel Ruiz Instituto de Matemática y Estatística Dep. Métodos Cuantitativos e Informáticos Universidaded de São Paulo Universidad Politécnica de Cartagena Caixa Postal 66281 Paseo Alfonso XIII, 50 05311-970 São Paulo, Brazil 30.203 Cartagena, Spain polcino@ime.usp.br Manuel.Ruiz@upct.es