SPECTRAL GAPS FOR PERIODIC SCHRÖDINGER OPERATORS WITH HYPERSURFACE MAGNETIC WELLS

B. HELFFER AND Y. A. KORDYUKOV

ABSTRACT. We consider a periodic magnetic Schrödinger operator on a noncompact Riemannian manifold M such that $H^1(M, \mathbb{R}) = 0$ endowed with a properly discontinuous cocompact isometric action of a discrete group. We assume that there is no electric field and that the magnetic field has a periodic set of compact magnetic wells. We review a general scheme of a proof of existence of an arbitrary large number of gaps in the spectrum of such an operator in the semiclassical limit, which was suggested in our previous paper, and some applications of this scheme. Then we apply these methods to establish similar results in the case when the wells have regular hypersurface pieces.

INTRODUCTION

Let M be a noncompact oriented manifold of dimension $n \geq 2$ equipped with a properly discontinuous action of a finitely generated, discrete group Γ such that M/Γ is compact. Suppose that $H^1(M, \mathbb{R}) = 0$, i.e. any closed 1-form on M is exact. Let g be a Γ -invariant Riemannian metric and \mathbf{B} a real-valued Γ -invariant closed 2-form on M. Assume that \mathbf{B} is exact and choose a real-valued 1-form \mathbf{A} on M such that $d\mathbf{A} = \mathbf{B}$.

Thus, one has a natural mapping

$$u \mapsto ih \, du + \mathbf{A}u$$

from $C_c^{\infty}(M)$ to the space $\Omega_c^1(M)$ of smooth, compactly supported oneforms on M. The Riemannian metric allows to define scalar products in these spaces and consider the adjoint operator

$$(ih d + \mathbf{A})^* : \Omega^1_c(M) \to C^\infty_c(M).$$

A Schrödinger operator with magnetic potential **A** is defined by the formula

$$H^h = (ih \, d + \mathbf{A})^* (ih \, d + \mathbf{A}).$$

Here h > 0 is a semiclassical parameter, which is assumed to be small.

Key words and phrases. magnetic Schrödinger operator; magnetic well; spectral gaps; Riemannian manifolds; semiclassical limit; quasimodes.

B.H. was partially supported by the ESF programme SPECT. Y.K. was partially supported by the Russian Foundation of Basic Research (grant 06-01-00208).

Choose local coordinates $X = (X_1, \ldots, X_n)$ on M. Write the 1-form **A** in the local coordinates as

$$\mathbf{A} = \sum_{j=1}^{n} A_j(X) \, dX_j,$$

the matrix of the Riemannian metric g as

$$g(X) = (g_{j\ell}(X))_{1 \le j,\ell \le n}$$

and its inverse as

$$g(X)^{-1} = (g^{j\ell}(X))_{1 \le j, \ell \le n}.$$

Denote $|g(X)| = \det(g(X))$. Then the magnetic field **B** is given by the following formula

$$\mathbf{B} = \sum_{j < k} B_{jk} \, dX_j \wedge dX_k, \quad B_{jk} = \frac{\partial A_k}{\partial X_j} - \frac{\partial A_j}{\partial X_k}.$$

Moreover, the operator H^h has the form

$$\begin{split} H^{h} &= \frac{1}{\sqrt{|g(X)|}} \sum_{1 \leq j, \ell \leq n} \left(ih \frac{\partial}{\partial X_{j}} + A_{j}(X) \right) \\ & \times \left[\sqrt{|g(X)|} g^{j\ell}(X) \left(ih \frac{\partial}{\partial X_{\ell}} + A_{\ell}(X) \right) \right]. \end{split}$$

For any $x \in M$, denote by B(x) the anti-symmetric linear operator on the tangent space $T_x M$ associated with the 2-form **B**:

$$g_x(B(x)u, v) = \mathbf{B}_x(u, v), \quad u, v \in T_x M.$$

Recall that the intensity of the magnetic field is defined as

$$\operatorname{Tr}^{+}(B(x)) = \sum_{\substack{\lambda_{j}(x) > 0\\ i\lambda_{j}(x) \in \sigma(B(x))}} \lambda_{j}(x) = \frac{1}{2} \operatorname{Tr}([B^{*}(x) \cdot B(x)]^{1/2}).$$

It turns out that in many problems the function $x \mapsto h \cdot \operatorname{Tr}^+(B(x))$ can be considered as a magnetic potential, that is, as a magnetic analogue of the electric potential V in a Schrödinger operator $-h^2\Delta + V$.

We will also use the trace norm of B(x):

$$|B(x)| = [\operatorname{Tr}(B^*(x) \cdot B(x))]^{1/2}.$$

It coincides with the norm of B(x) with respect to the Riemannian metric on the space of linear operators on $T_x M$ induced by the Riemannian metric g on M.

In this paper we will always assume that the magnetic field has a periodic set of compact potential wells. More precisely, put

$$b_0 = \min\{\operatorname{Tr}^+(B(x)) : x \in M\}$$

and assume that there exist a (connected) fundamental domain \mathcal{F} and a constant $\epsilon_0 > 0$ such that

(1)
$$\operatorname{Tr}^+(B(x)) \ge b_0 + \epsilon_0, \quad x \in \partial \mathcal{F}$$

For any $\epsilon_1 \leq \epsilon_0$, put

$$U_{\epsilon_1} = \{ x \in \mathcal{F} : \operatorname{Tr}^+(B(x)) < b_0 + \epsilon_1 \}.$$

Thus U_{ϵ_1} is an open subset of \mathcal{F} such that $U_{\epsilon_1} \cap \partial \mathcal{F} = \emptyset$ and, for $\epsilon_1 < \epsilon_0$, $\overline{U_{\epsilon_1}}$ is compact and included in the interior of \mathcal{F} . Any connected component of U_{ϵ_1} with $\epsilon_1 < \epsilon_0$ and also any its translation under the action of an element of Γ can be understood as a magnetic well. These magnetic wells are separated by potential barriers, which are getting higher and higher when $h \to 0$ (in the semiclassical limit).

For any linear operator T in a Hilbert space, we will denote by $\sigma(T)$ its spectrum. By a gap in the spectrum of a self-adjoint operator T we will mean any connected component of the complement of $\sigma(T)$ in \mathbb{R} , that is, any maximal interval (a, b) such that

$$(a,b) \cap \sigma(T) = \emptyset$$
.

The problem of existence of gaps in the spectra of second order periodic differential operators has been extensively studied recently. Some related results on spectral gaps for periodic magnetic Schrödinger operators can be found for example in [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13] (see also the references therein).

In this paper, we consider the magnetic Schrödinger operator H^h as an unbounded self-adjoint operator in the Hilbert space $L^2(M)$ and will study gaps in the spectrum of this operator, which are located below the top of potential barriers, that is, on the interval $[0, h(b_0 + \epsilon_0)]$. In this case, the important role is played by the tunneling effect, that is, by the possibility for the quantum particle described by the Hamiltonian H^h with such an energy to pass through a potential barrier. Using the semiclassical analysis of the tunneling effect, we showed in [2] that the spectrum of the magnetic Schrödinger operator H^h on the interval is localized in an exponentially small neighborhood of the spectrum of its Dirichlet realization inside the wells. This result reduces the investigation of gaps in the spectrum of the operator H^h to the study of the eigenvalue distribution for a "one-well" operator and leads us to suggest a general scheme of a proof of existence of spectral gaps in [3]. We review this scheme and some of its applications in Section 1. Then, in Section 2, we will apply these methods to prove the existence of an arbitrary large number of gaps in the spectrum of the operator H^h , as $h \to 0$, under the assumption that $b_0 = 0$ and the zero set of **B** has regular codimension one pieces.

1. Quasimodes and spectral gaps

In this section, we review a general scheme of a proof of existence of gaps in the spectrum of the magnetic Schrödinger operator H^h on the interval $[0, h(b_0 + \epsilon_0)]$ and some of its applications obtained in [3].

1.1. A general scheme. For any domain W in M, denote by H_W^h the unbounded self-adjoint operator in the Hilbert space $L^2(W)$ defined by the operator H^h in \overline{W} with the Dirichlet boundary conditions. The operator H_W^h is generated by the quadratic form

$$u \mapsto q_W^h[u] := \int_W |(ih\,d + \mathbf{A})u|^2\,dx$$

with the domain

$$Dom(q_W^h) = \{ u \in L^2(W) : (ih \, d + \mathbf{A}) u \in L^2\Omega^1(W), u |_{\partial W} = 0 \},\$$

where $L^2\Omega^1(W)$ denotes the Hilbert space of L^2 differential 1-forms on W, dx is the Riemannian volume form on M.

Assume now that the operator H^h satisfies the condition (1). Fix $\epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $\epsilon_1 < \epsilon_2 < \epsilon_0$, and consider the operator H^h_D associated with the domain $D = \overline{U_{\epsilon_2}}$. The operator H^h_D has discrete spectrum.

The following result is a slight generalization of Theorem 2.1 in [3], which is concerned with the case when N_h is independent of h. It permits to get a more precise information on the number of gaps as $h \to 0$.

Theorem 1.1. Suppose that there exist $h_0 > 0$, c > 0, $M \ge 1$ and that, for $h \in (0, h_0]$, there exists N_h and a subset $\mu_0^h < \mu_1^h < \ldots < \mu_{N_h}^h$ of an interval $I(h) \subset [0, h(b_0 + \epsilon_1))$ such that

$$\mu_j^h - \mu_{j-1}^h > ch^M, \quad j = 1, \dots, N_h,$$

$$\operatorname{dist}(\mu_0^h, \partial I(h)) > ch^M, \quad \operatorname{dist}(\mu_{N_h}^h, \partial I(h)) > ch^M,$$

and, for each $j = 0, 1, ..., N_h$, there exists some non trivial $v_j^h \in C_c^{\infty}(D)$ such that

$$||H_D^h v_j^h - \mu_j^h v_j^h|| \le \frac{c}{3} h^M ||v_j^h||.$$

Then there exists $h_1 \in (0, h_0]$ such that the spectrum of H^h on the interval I(h) has at least N_h gaps for $h \in (0, h_1)$.

1.2. A generic situation. As a first application of Theorem 1.1, we show in [3] that the spectrum of the Schrödinger operator H^h , satisfying the assumption (1), always has gaps (moreover, an arbitrarily large number of gaps) on the interval $[0, h(b_0 + \epsilon_0)]$ in the semiclassical limit $h \to 0$. Under some additional generic assumption, this result was obtained in [2]. Indeed, slightly modifying the arguments of [3], one can show the following theorem.

Theorem 1.2. Under the assumption (1), for any interval $[\alpha, \beta] \subset [b_0, b_0 + \epsilon_0]$ and for any natural N, there exists $h_0 > 0$ such that, for any $h \in (0, h_0]$, the spectrum of H^h in the interval $[h\alpha, h\beta]$ has at least N gaps.

The proof of this theorem can be given by a straightforward repetition of the proof of Theorem 3.1 in [3] with the only difference — one should choose $\mu_0 < \mu_1 < \ldots < \mu_N$ in the interval (α, β) instead of $(b_0, b_0 + \epsilon_0)$.

Indeed, using Theorem 1.1 with N_h dependent on h and a continuous family of quasimodes constructed in the proof of Proposition 2.3 in [2], we can get an estimate for the number of gaps in the constant rank case. Denote by [a] the integer part of a (the largest integer n satisfying $n \leq a$).

Theorem 1.3. Under the assumption (1), suppose that the rank of **B** is constant in an open set $U \subset M$. Then, for any interval $[\alpha, \beta] \subset \text{Tr}^+B(U)$, there exists $h_0 > 0$ and C > 0 such that, for any $h \in (0, h_0]$, the spectrum of H^h in the interval $[h\alpha, h\beta]$ has at least $[Ch^{-1/3}]$ gaps.

1.3. The case of discrete wells. A more precise information on location and asymptotic behavior of gaps in the spectrum of the magnetic Schrödinger operator H^h , satisfying the assumption (1), can be obtained, if we impose additional hypotheses on the bottoms of magnetic wells. In this section, we consider a case when the bottom of the magnetic well contains zero-dimensional components, that is, isolated points, and, moreover, the magnetic field behaves regularly near these points. More precisely, we will assume that $b_0 = 0$ and that there is at least one zero x_0 of B such that, for some integer k > 0, there exists a positive constant C such that for all x in some neighborhood of x_0 the estimate holds:

(2)
$$C^{-1}d(x,x_0)^k \le |B(x)| \le Cd(x,x_0)^k$$

(here d(x, y) denotes the geodesic distance between x and y). In this case, the important role is played by a differential operator $K_{\bar{x}_0}^h$ in \mathbb{R}^n , which is in some sense an approximation to the operator H^h near x_0 . Recall its definition (see [14]).

Let \bar{x}_0 be a zero of B. Choose local coordinates $f : U(\bar{x}_0) \to \mathbb{R}^n$ on M, defined in a sufficiently small neighborhood $U(\bar{x}_0)$ of \bar{x}_0 . Suppose that $f(\bar{x}_0) = 0$, and the image $f(U(\bar{x}_0))$ is a ball B(0,r) in \mathbb{R}^n centered at the origin.

Write the 2-form \mathbf{B} in the local coordinates as

$$\mathbf{B}(X) = \sum_{1 \le \ell < m \le n} b_{\ell m}(X) \, dX_{\ell} \wedge dX_m, \quad X = (X_1, \dots, X_n) \in B(0, r).$$

Let \mathbf{B}^0 be the closed 2-form in \mathbb{R}^n with polynomial components defined by the formula

$$\mathbf{B}^{0}(X) = \sum_{1 \le \ell < m \le n} \sum_{|\alpha| = k} \frac{X^{\alpha}}{\alpha!} \frac{\partial^{\alpha} b_{\ell m}}{\partial X^{\alpha}}(0) \, dX_{\ell} \wedge dX_{m}, \quad X \in \mathbb{R}^{n}.$$

One can find a 1-form \mathbf{A}^0 on \mathbb{R}^n with polynomial components such that

$$d\mathbf{A}^0(X) = \mathbf{B}^0(X), \quad X \in \mathbb{R}^n.$$

Let $K^h_{\bar{x}_0}$ be a self-adjoint differential operator in $L^2(\mathbb{R}^n)$ with polynomial coefficients given by the formula

$$K^{h}_{\bar{x}_{0}} = (ih \, d + \mathbf{A}^{0})^{*} (ih \, d + \mathbf{A}^{0}),$$

where the adjoints are taken with respect to the Hilbert structure in $L^2(\mathbb{R}^n)$ given by the flat Riemannian metric $(g_{\ell m}(0))$ in \mathbb{R}^n . If \mathbf{A}^0 is written as

$$\mathbf{A}^0 = A_1^0 \, dX_1 + \ldots + A_n^0 \, dX_n,$$

then $K^h_{\bar{x}_0}$ is given by the formula

$$K_{\bar{x}_0}^h = \sum_{1 \le \ell, m \le n} g^{\ell m}(0) \left(ih \frac{\partial}{\partial X_\ell} + A_\ell^0(X) \right) \left(ih \frac{\partial}{\partial X_m} + A_m^0(X) \right).$$

The operators $K_{\bar{x}_0}^h$ have discrete spectrum (cf, for instance, [15, 16]). Using the simple dilation $X \mapsto h^{\frac{1}{k+2}}X$, one can show that the operator $K_{\bar{x}_0}^h$ is unitarily equivalent to $h^{\frac{2k+2}{k+2}}K_{\bar{x}_0}^1$. Thus, $h^{-\frac{2k+2}{k+2}}K_{\bar{x}_0}^h$ has discrete spectrum, independent of h.

Theorem 1.4 ([3]). Suppose that the operator H^h satisfies the condition (1) with some $\epsilon_0 > 0$ and that there exists a zero \bar{x}_0 of B, satisfying the assumption (2) for some integer k > 0. Denote by $\lambda_1 < \lambda_2 < \lambda_3 < \ldots$ the eigenvalues of the operator $K^1_{\bar{x}_0}$ (not taking into account multiplicities). Then, for any natural N and any $C > \lambda_{N+1}$, there exists $h_0 > 0$ such that the spectrum of H^h in the interval $[0, Ch^{\frac{2k+2}{k+2}}]$ has at least N gaps for any $h \in (0, h_0)$.

2. Hypersurface wells

In this section, we consider the case when $b_0 = 0$ and the zero set of the magnetic field has regular codimension one pieces. More precisely, suppose that there exists $x_0 \in M$ such that $\mathbf{B}(x_0) = 0$ and in a neighborhood U of x_0 the zero set of \mathbf{B} is a smooth oriented hypersurface S, and, moreover, there are constants $k \in \mathbb{N}$ and C > 0 such that for all $x \in U$ we have:

(3)
$$C^{-1}d(x,S)^k \le |B(x)| \le Cd(x,S)^k$$

On compact manifolds, this model was introduced for the first time by Montgomery [17] and was further studied in [14, 18, 19].

Let

$$\omega_{0.0} = i_S^* \mathbf{A}$$

be the closed one form on S induced by **A**, where i_S is the embedding of S to M.

Denote by N the external unit normal vector to S and by \tilde{N} an arbitrary extension of N to a smooth vector field on U.

Let $\omega_{0,1}$ be the smooth one form on S defined, for any vector field V on S, by the formula

$$\langle V, \omega_{0,1} \rangle(y) = \frac{1}{k!} \tilde{N}^k(\mathbf{B}(\tilde{N}, \tilde{V}))(y), \quad y \in S,$$

where \tilde{V} is a C^{∞} extension of V to U. By (3), it is easy to see that $\omega_{0,1}(x) \neq 0$ for any $x \in S$. Denote

$$\omega_{\min}(B) = \inf_{x \in S} |\omega_{0,1}(x)| > 0.$$

For any $\alpha \in \mathbb{R}$ and $\beta \in \mathbb{R}, \beta \neq 0$, consider the self-adjoint second order differential operator in $L^2(\mathbb{R})$ given by

$$Q(\alpha,\beta) = -\frac{d^2}{dt^2} + \left(\frac{1}{k+1}\beta t^{k+1} - \alpha\right)^2.$$

In the context of magnetic bottles, this family of operators (for k = 1) first appears in [17] (see also [14]). Denote by $\lambda_0(\alpha, \beta)$ the bottom of the spectrum of the operator $Q(\alpha, \beta)$.

Let us recall some properties of $\lambda_0(\alpha, \beta)$, which were established in [17, 14, 18]. First of all, remark that $\lambda_0(\alpha, \beta)$ is a continuous function of $\alpha \in \mathbb{R}$ and $\beta \in \mathbb{R} \setminus \{0\}$. One can see by scaling that, for $\beta > 0$,

(4)
$$\lambda_0(\alpha,\beta) = \beta^{\frac{2}{k+2}} \lambda_0(\beta^{-\frac{1}{k+2}}\alpha,1) .$$

A further discussion depends on k odd or k even.

When k is odd, $\lambda_0(\alpha, 1)$ tends to $+\infty$ as $\alpha \to -\infty$ by monotonicity. For analyzing its behavior as $\alpha \to +\infty$, it is suitable to do a dilation $t = \alpha^{\frac{1}{k+1}}s$, which leads to the analysis of

$$\alpha^2 \left(-h^2 \frac{d^2}{ds^2} + \left(\frac{s^{k+1}}{k+1} - 1 \right)^2 \right)$$

with $h = \alpha^{-(k+2)/(k+1)}$ small. Semi-classical analysis is relevant, and it is easy to show, using harmonic approximation, that

$$\lambda_0(\alpha, 1) \sim (k+1)^{\frac{2k}{k+1}} \alpha^{\frac{k}{k+1}}$$
, as $\alpha \to +\infty$.

In particular, we see that $\lambda_0(\alpha, 1)$ tends to $+\infty$.

When k is even, we have $\lambda_0(\alpha, 1) = \lambda_0(-\alpha, 1)$, and, therefore, it is sufficient to consider the case $\alpha \ge 0$. As $\alpha \to +\infty$, semi-classical analysis again shows that $\lambda_0(\alpha, 1)$ tends to $+\infty$.

So in both cases, it is clear that the continuous function $\lambda_0(\alpha, 1)$ is lower semi-bounded:

$$\hat{\nu} := \inf_{\alpha \in \mathbb{R}} \lambda_0(\alpha, 1) > -\infty,$$

and there exists (at least one) $\alpha_{\min} \in \mathbb{R}$ such that $\lambda_0(\alpha, 1)$ is minimal:

$$\lambda_0(\alpha_{\min}, 1) = \hat{\nu}.$$

For k odd, one can show that the minimum α_{\min} is strictly positive. One can indeed compute the derivative of $\lambda_0(\alpha, 1)$ at $\alpha = 0$ and find that

$$\frac{\partial \lambda_0}{\partial \alpha}(0,1) < 0$$

In the case k = 1, it has been shown that this minimum is unique (see [18]). Numerical computations show (see [17, 14]) that, in this case, $\hat{\nu} \approx 0.5698$.

Theorem 2.1. For any *a* and *b* such that

$$\hat{\nu}\,\omega_{\min}(B)^{\frac{2}{k+2}} < a < b$$

and for any natural N, there exists $h_0 > 0$ such that, for any $h \in (0, h_0]$, the spectrum of H^h in the interval

$$[h^{\frac{2k+2}{k+2}}a,h^{\frac{2k+2}{k+2}}b]$$

has at least N gaps.

Proof. Let g_0 be the Riemannian metric on S induced by g. Without loss of generality, we can assume that U coincides with an open tubular neighborhood of S and choose a diffeomorphism

$$\Theta: I \times S \to U,$$

where I is an open interval $(-\varepsilon_0, \varepsilon_0)$ with $\varepsilon_0 > 0$ small enough, such that $\Theta|_{\{0\}\times S} = \text{id}$ and

$$\left(\Theta^*g - \tilde{g}_0\right)\Big|_{\{0\} \times S} = 0,$$

where \tilde{g}_0 is a Riemannian metric on $I \times S$ given by

$$\tilde{g}_0 = dt^2 + g_0.$$

By adding to **A** the exact one form $d\phi$, where ϕ is the function satisfying

$$N(x)\phi(x) = -\langle N, \mathbf{A} \rangle(x), \quad x \in U,$$

$$\phi(x) = 0, \quad x \in S,$$

we may assume that

$$\langle N, \mathbf{A} \rangle(x) = 0, \quad x \in U.$$

As above, denote by H_D^h the unbounded self-adjoint operator in $L^2(D)$ given by the operator H^h in the domain $D = \overline{U}$ with the Dirichlet boundary conditions.

For any $t \in \mathbb{R}$, let $P_S^h\left(\omega_{0,0} + \frac{1}{k+1}t^{k+1}\omega_{0,1}\right)$ be a formally self-adjoint operator in $L^2(S, dx_{q_0})$ defined by

$$P_{S}^{h}\left(\omega_{0,0} + \frac{1}{k+1}t^{k+1}\omega_{0,1}\right) = \left(ihd + \omega_{0,0} + \frac{1}{k+1}t^{k+1}\omega_{0,1}\right)^{*} \times \left(ihd + \omega_{0,0} + \frac{1}{k+1}t^{k+1}\omega_{0,1}\right).$$

Consider the self-adjoint operator $H^{h,0}$ in $L^2(\mathbb{R} \times S, dt \, dx_{g_0})$ defined by the formula

$$H^{h,0} = -h^2 \frac{\partial^2}{\partial t^2} + P_S^h \left(\omega_{0,0} + \frac{1}{k+1} t^{k+1} \omega_{0,1} \right)$$

with the Dirichlet boundary conditions. By Theorem 2.7 of [14], the operator $H^{h,0}$ has discrete spectrum. Moreover, it can be seen from the proof of this theorem that if $\lambda^0(h)$ is an approximate eigenvalue of $H^{h,0}$ with the corresponding approximate eigenfunction $w^h \in C_c^{\infty}(\mathbb{R} \times S)$ such that

$$\lambda^0(h) < Dh^{(2k+2)/(k+2)}$$

and

$$\|(H^{h,0} - \lambda^0(h))w^h\| \le Ch^{(2k+3)/(k+2)} \|w^h\|,$$

then $\lambda^0(h)$ is an approximate eigenvalue of H_D^h with the corresponding approximate eigenfunction $v^h = (\Theta^{-1})^* w^h \in C_c^\infty(U)$:

$$||(H_D^h - \lambda^0(h))v^h|| \le Ch^{(2k+3)/(k+2)}||v^h||.$$

So it remains to construct approximate eigenvalues of $H^{h,0}$.

Lemma 2.1. For any $\lambda \geq \hat{\nu} \omega_{\min}(B)^{2/(k+2)}$, there exists $\Phi \in C_c^{\infty}(\mathbb{R} \times S)$ such that

$$\|(H^{h,0} - \lambda h^{\frac{2k+2}{k+2}})\Phi\| \le Ch^{\frac{6k+8}{3(k+2)}} \|\Phi\|.$$

Proof. Take $x_1 \in S$ such that $|\omega_{0,1}(x_1)| = \omega_{\min}(B)$. Consider $\alpha_1 \in \mathbb{R}$ such that $\lambda_0(\alpha_1, 1) = \lambda \omega_{\min}(B)^{-2/(k+2)} \geq \hat{\nu}$. Let $\psi \in L^2(\mathbb{R})$ be a normalized eigenfunction of $Q(\alpha_1, 1)$, corresponding to $\lambda_0(\alpha_1, 1)$:

$$\left[-\frac{d^2}{dt^2} + \left(\frac{1}{k+1}t^{k+1} - \alpha_1\right)^2\right]\psi(t) = \lambda\omega_{\min}(B)^{-\frac{2}{k+2}}\psi(t), \quad \|\psi\|_{L^2(\mathbb{R})} = 1.$$

Then the function

$$\Psi(t) = \omega_{\min}(B)^{\frac{1}{2(k+2)}} h^{-\frac{1}{2(k+2)}} \psi(\omega_{\min}(B)^{\frac{1}{k+2}} h^{-\frac{1}{k+2}} t)$$

satisfies

$$\left(-h^2 \frac{d^2}{dt^2} + \left(\frac{1}{k+1}\omega_{\min}(B)t^{k+1} - \alpha_1\omega_{\min}(B)^{\frac{1}{k+2}}h^{\frac{k+1}{k+2}}\right)^2\right)\Psi(t)$$
$$= \lambda h^{\frac{2k+2}{k+2}}\Psi(t), \quad \|\Psi\|_{L^2(\mathbb{R})} = 1.$$

Take normal coordinates $f: U(x_1) \subset S \to \mathbb{R}^{n-1}$ on S defined in a neighborhood $U(x_1)$ of x_1 , where $f(U(x_1)) = B(0,r)$ is a ball in \mathbb{R}^{n-1} centered at the origin and $f(x_1) = 0$. Choose a function $\phi \in C^{\infty}(B(0,r))$ such that $d\phi = \omega_{0,0}$. Write $\omega_{0,1} = \sum_{j=1}^{n-1} \omega_j(s) \, ds_j$. Note that

$$\omega_{\min}(B) = \left(\sum_{j=1}^{n-1} |\omega_j(0)|^2\right)^{1/2}.$$

Consider the function $\Phi \in C^{\infty}(B(0,r) \times \mathbb{R})$ given by

$$(5) \quad \Phi(s,t) = ch^{-\beta/2(n-1)}\chi(s)\exp\left(-i\frac{\phi(s)}{h}\right)\exp\left(\frac{\alpha_1\sum\limits_{j=1}^{n-1}\omega_j(0)s_j}{\omega_{\min}(B)^{\frac{k+1}{k+2}}h^{\frac{1}{k+2}}}\right) \\ \times \exp\left(-\frac{|s|^2}{2h^{2\beta}}\right)\Psi(t), \quad s \in B(0,r), \quad t \in \mathbb{R},$$

with some β , where $\chi \in C_c^{\infty}(B(0,r))$ is a cut-off function, c is chosen in such a way that $\|\Phi\|_{L^2(S \times \mathbb{R})} = 1$.

Put

$$\begin{aligned} H^{h,1} &= -h^2 \frac{\partial^2}{\partial t^2} + P_S^h \left(\frac{1}{k+1} t^{k+1} \omega_{0,1} - \alpha_1 \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_{0,1}(0) \right), \\ E(s) &= c h^{-\beta/2(n-1)} \chi(s) \exp\left(-\frac{|s|^2}{2h^{2\beta}}\right), \\ \Phi_1(s,t) &= E(s) \Psi(t). \end{aligned}$$

Then we have

$$H^{h,0}\Phi(s,t) = \exp\left(-i\frac{\phi(s)}{h}\right) \exp\left(i\frac{\alpha_1\sum_{j=1}^{n-1}\omega_j(0)s_j}{\omega_{\min}(B)^{\frac{k+1}{k+2}}h^{\frac{1}{k+2}}}\right) H^{h,1}\Phi_1(s,t).$$

Next, we have

$$\begin{split} P_{S}^{h} \left(\frac{1}{k+1} t^{k+1} \omega_{0,1} - \alpha_{1} \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_{0,1}(0) \right) \\ &= \sum_{j,\ell} \frac{1}{\sqrt{g_{0}}} \left(ih \frac{\partial}{\partial s_{j}} + \frac{1}{k+1} t^{k+1} \omega_{j}(s) - \alpha_{1} \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_{j}(0) \right) \\ &\times \left(g_{0}^{j\ell} \sqrt{g_{0}} \left(ih \frac{\partial}{\partial s_{\ell}} + \frac{1}{k+1} t^{k+1} \omega_{\ell}(s) - \alpha_{1} \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_{\ell}(0) \right) \right) \\ &= \sum_{j,\ell} g_{0}^{j\ell} \left(ih \frac{\partial}{\partial s_{j}} + \frac{1}{k+1} t^{k+1} \omega_{j}(s) - \alpha_{1} \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_{j}(0) \right) \\ &\times \left(ih \frac{\partial}{\partial s_{\ell}} + \frac{1}{k+1} t^{k+1} \omega_{\ell}(s) - \alpha_{1} \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_{\ell}(0) \right) \\ &+ \sum_{\ell} ih \Gamma^{\ell}(s) \left(ih \frac{\partial}{\partial s_{\ell}} + \frac{1}{k+1} t^{k+1} \omega_{\ell}(s) - \alpha_{1} \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_{\ell}(0) \right) \\ &= -h^{2} \sum_{j,\ell} g_{0}^{j\ell} \frac{\partial^{2}}{\partial s_{j} \partial s_{\ell}} + 2ih \sum_{j,\ell} g_{0}^{j\ell} \frac{1}{k+1} t^{k+1} \frac{\partial \omega_{\ell}}{\partial s_{j}}(s) \end{split}$$

$$+ 2ih \sum_{j,\ell} g_0^{j\ell} \left(\frac{1}{k+1} t^{k+1} \omega_\ell(s) - \alpha_1 \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_\ell(0) \right) \frac{\partial}{\partial s_j} \\ + \sum_{j,\ell} g_0^{j\ell} \left(\frac{1}{k+1} t^{k+1} \omega_j(s) - \alpha_1 \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_j(0) \right) \\ \times \left(\frac{1}{k+1} t^{k+1} \omega_\ell(s) - \alpha_1 \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_\ell(0) \right) - h^2 \sum_{\ell} \Gamma^\ell(s) \frac{\partial}{\partial s_\ell} \\ + ih \sum_{\ell} \Gamma^\ell(s) \left(\frac{1}{k+1} t^{k+1} \omega_\ell(s) - \alpha_1 \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_\ell(0) \right),$$

where

$$\Gamma^{\ell} = \sum_{j} \frac{1}{\sqrt{g_0}} \frac{\partial}{\partial s_j} \left(g_0^{j\ell} \sqrt{g_0} \right).$$

By a well-known property of normal coordinates, we have $\partial_j g_0^{\ell m}(0) = 0$. So we get $\Gamma^{\ell}(0) = 0$, and

(6)
$$g_0^{\ell m}(s) = \delta^{\ell m} + O(|s|^2), \quad \Gamma^{\ell}(s) = O(|s|), \quad s \to 0.$$

We get

$$\begin{split} H^{h,1}\Phi_{1}(s,t) &= \lambda h^{\frac{2k+2}{k+2}}\Phi_{1}(s,t) - h^{2}\sum_{j,\ell}g_{0}^{j\ell}\frac{\partial^{2}E}{\partial s_{j}\partial s_{\ell}}(s)\Psi(t) \\ &+ 2ih\sum_{j,\ell}g_{0}^{j\ell}\frac{1}{k+1}\frac{\partial\omega_{\ell}}{\partial s_{j}}E(s)t^{k+1}\Psi(t) \\ &+ 2ih\sum_{j,\ell}g_{0}^{j\ell}\frac{\partial E}{\partial s_{j}}(s)\left(\frac{1}{k+1}t^{k+1}\omega_{\ell}(s) - \alpha_{1}\omega_{\min}(B)^{-\frac{k+1}{k+2}}h^{\frac{k+1}{k+2}}\omega_{\ell}(0)\right)\Psi(t) \\ &+ R(s,t)E(s)\Psi(t) - h^{2}\sum_{\ell}\Gamma^{\ell}(s)\frac{\partial}{\partial s_{\ell}}E(s)\Psi(t) \\ &+ ih\sum_{\ell}\Gamma^{\ell}(s)\left(\frac{1}{k+1}t^{k+1}\omega_{\ell}(s) - \alpha_{1}\omega_{\min}(B)^{-\frac{k+1}{k+2}}h^{\frac{k+1}{k+2}}\omega_{\ell}(0)\right)E(s)\Psi(t), \end{split}$$

where

$$R(s,t) = \sum_{j,\ell} g_0^{j\ell} \left(\frac{1}{k+1} t^{k+1} \omega_j(s) - \alpha_1 \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_j(0) \right)$$
$$\times \left(\frac{1}{k+1} t^{k+1} \omega_\ell(s) - \alpha_1 \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_\ell(0) \right)$$
$$- \sum_j \left(\frac{1}{k+1} t^{k+1} \omega_{\min}(B)^2 - \alpha_1 \omega_{\min}(B)^{\frac{1}{k+2}} h^{\frac{k+1}{k+2}} \right)^2$$

$$= \frac{1}{(k+1)^2} \left(\sum_{j,\ell} g_0^{\ell}(s)\omega_j(s)\omega_\ell(s) - \sum_j (\omega_j(0))^2 \right) t^{2(k+1)} - \frac{2}{k+1} t^{k+1} \sum_j (\omega_j(s) - \omega_j(0))\alpha_1 \omega_{\min}(B)^{-\frac{k+1}{k+2}} h^{\frac{k+1}{k+2}} \omega_j(0) + O(|s|^2) \sum_j \left(\frac{1}{k+1} t^{k+1} \omega_{\min}(B) - \alpha_1 \omega_{\min}(B)^{\frac{1}{k+2}} h^{\frac{k+1}{k+2}} \right)^2.$$

We have

(7)
$$|||s|^m E(s)|| = \left(h^{-\beta(n-1)} \int_{\mathbb{R}^{n-1}} |s|^{2m} \exp\left(-\frac{|s|^2}{h^{2\beta}}\right) ds\right)^{1/2} = C_1 h^{\beta m},$$

and, furthermore,

(8)
$$||s|^m \frac{\partial E}{\partial s_j}(s)|| = C_2 h^{\beta(m-1)}, \quad ||s|^m \frac{\partial^2 E}{\partial s_j \partial s_\ell}(s)|| = C_3 h^{\beta(m-2)}.$$

We also have

(9)
$$||t^{k+1}\Psi(t)|| \le C_4 h^{\frac{k+1}{k+2}}, \quad ||t^{2(k+1)}\Psi(t)|| \le C_5 h^{\frac{2k+2}{k+2}}.$$

Since s = 0 is a minimum of $|\omega_{0,1}(s)|^2$, we have

(10)
$$|\omega_{0,1}(s)|^2 - \omega_{\min}(B)^2 = \sum_{j,\ell} g_0^{j\ell}(s)\omega_j(s)\omega_\ell(s) - (\omega_j(0))^2 \le C_6|s|^2$$

and

$$\left(\frac{\partial}{\partial s_r}|\omega_{0,1}|^2\right)(0) = 2\sum_j \frac{\partial \omega_j}{\partial s_r}(0)\omega_j(0) = 0,$$

that implies

(11)
$$\left|\sum_{j} (\omega_j(s) - \omega_j(0))\omega_j(0)\right| \le C_7 |s|^2.$$

Using (6), (7), (8), (9), (10) and (11) and putting $\beta = \frac{1}{3(k+2)}$, one can easily get that

$$\|H^{h,0}\Phi - \lambda h^{\frac{2k+2}{k+2}}\Phi\| \le Ch^{\frac{6k+8}{3(k+2)}}.$$

Given a and b such that $\hat{\nu} \,\omega_{\min}(B)^{2/(k+2)} < a < b$ and some natural N, choose some finite sequence $\{\nu_j\}_{j=0,\dots,N}$ such that

$$a < \nu_0 < \nu_1 < \ldots < \nu_N < b.$$

By Lemma 2.1, for any $m = 0, 1, \ldots, N$,

$$\mu_m^h = \nu_m h^{\frac{2k+2}{k+2}} \in [h^{(2k+2)/(k+2)}a, h^{(2k+2)/(k+2)}b]$$

is an approximate eigenvalue of the operator H_D^h : for some $\Phi_m^h \in C_c^{\infty}(D)$

$$\|(H^{h,0} - \mu_m^h)\Phi_m^h\| \le Ch^{\frac{6k+8}{3(k+2)}} \|\Phi_m^h\|.$$

Using Theorem 1.1 with $N_h = N$ independent of h, we complete the proof.

Remark 2.1. Using the methods of the proof of Theorem 2.1, one can construct much more approximate eigenvalues of the operator H^h on the interval $[h^{(2k+2)/(k+2)}a, h^{(2k+2)/(k+2)}b]$ with some a and b such that $\hat{\nu} \omega_{\min}(B)^{2/(k+2)} < a < b$. Applying then Theorem 1.1 with N_h dependent on h, one can get the following theorem.

Theorem 2.2. Under the assumptions of Theorem 2.1, for any a and b such that

$$\hat{\nu}\,\omega_{\min}(B)^{\frac{2}{k+2}} < a < b,$$

there exist $h_0 > 0$ and C > 0 such that, for any $h \in (0, h_0]$, the spectrum of H^h in the interval

$$[h^{\frac{2k+2}{k+2}}a, h^{\frac{2k+2}{k+2}}b]$$

has at least $[Ch^{-\frac{2}{3(k+2)}}]$ gaps.

3. Concluding remarks

1. Suppose that the operator H^h satisfies the condition (1) with some $\epsilon_0 > 0$, and the zero set of the magnetic field **B** is a smooth oriented hypersurface S. Moreover, assume that there are constants $k \in \mathbb{N}$ and C > 0 such that for all x in a neighborhood of S we have:

$$C^{-1}d(x,S)^k \le |B(x)| \le Cd(x,S)^k.$$

It is interesting to determine the bottom $\lambda_0(H^h)$ of the spectrum of the operator H^h in $L^2(M)$. By Theorem 2.1 in [2] and Theorem 2.7 in [14], $\lambda_0(H^h)$ is asymptotically equal to the bottom $\lambda_0(H^{h,0})$ of the spectrum of the operator $H^{h,0}$. From the construction of approximate eigenvalues of $H^{h,0}$ given in Lemma 2.1, one can see that, in order to find $\lambda_0(H^{h,0})$, it is natural to consider a self-adjoint second order differential operator $P(\mathbf{v}, \mathbf{w})$, $\mathbf{v}, \mathbf{w} \in \mathbb{R}^{n-1}$, in $L^2(\mathbb{R})$ given by

$$P(\mathbf{v}, \mathbf{w}) = -\frac{d^2}{dt^2} + \left|\frac{1}{k+1}\mathbf{w}t^{k+1} - \mathbf{v}\right|^2$$

and minimize the bottoms $\lambda_0(\mathbf{v}, \mathbf{w})$ of the spectrum of the operator $P(\mathbf{v}, \mathbf{w})$ over $\mathbf{v} \in \mathbb{R}^{n-1}$ and $\mathbf{w} \in K$, where $K = \{\omega_{0,1}(s) : s \in \overline{S}\}$ is a compact subset of $\mathbb{R}^{n-1} \setminus \{0\}$.

The identity

$$P(\mathbf{v}, \mathbf{w}) = \left(-\frac{d^2}{dt^2} + \left(\frac{1}{k+1}|\mathbf{w}|t^{k+1} - \frac{\mathbf{v} \cdot \mathbf{w}}{|\mathbf{w}|}\right)^2\right) + \left|\mathbf{v} - \frac{\mathbf{v} \cdot \mathbf{w}}{|\mathbf{w}|^2}\mathbf{w}\right|^2$$

shows that, for determining the minimum of $\lambda_0(\mathbf{v}, \mathbf{w})$ over $\mathbf{v} \in \mathbb{R}^{n-1}$ and $\mathbf{w} \in K$, it is sufficient to assume that \mathbf{v} is parallel to \mathbf{w} . For such \mathbf{v} and \mathbf{w} , we obtain $P(\mathbf{v}, \mathbf{w}) = Q(\alpha, \beta)$ with $\alpha = \pm |\mathbf{v}|, \beta = |\mathbf{w}|$. By (4), it follows that, for determining the minimum of $\lambda_0(\alpha, \beta)$ over $\alpha \in \mathbb{R}$ and $\beta \in \{|\omega_{0,1}(s)| : s \in \overline{S}\}$, we should first minimize over β , that is, take $s_1 \in S$ such that

$$|\omega_{0,1}(s_1)| = \min\{|\omega_{0,1}(s)| : s \in \bar{S}\},\$$

and then, for the minimal β , minimize over α .

This observation provides some explanations of our construction of approximate eigenvalues of the operator $H^{h,0}$ in Lemma 2.1, in particular, of our choice for the exponent in (5). It also motivates us to formulate the following conjecture:

Conjecture 3.1. Under current assumptions, for the bottom $\lambda_0(H^h)$ of the spectrum of the operator H^h in $L^2(M)$, we have

$$\lim_{h \to 0} h^{-\frac{2k+2}{k+2}} \lambda_0(H^h) = \hat{\nu} \,\omega_{\min}(B)^{\frac{2}{k+2}}.$$

Observe that a similar result was obtained by Pan and Kwek [18] for the bottom of the spectrum of the Neumann realization of the operator H^h in a bounded domain in the case k = 1.

2. In the setting of Section 2, one can assume that the function $|\omega_{0,1}(x)|$ has a non-degenerate minimum at some $x_1 \in S$. In some sense this is the "miniwells case" analyzed in [20] in comparison with the "uniform case" analyzed in [21], which in this setting was studied in [3]. Then we can obtain a more precise information about gaps located near the bottom of the spectrum of H^h (see some relevant calculations in [19]). This will be discussed elsewhere.

3. The similar results both like in Section 2 and like in the "miniwells case" mentioned in the previous remark can be obtained when $b_0 \neq 0$ (for the "miniwells case", see some relevant results in [22]). We will consider these problems in a future publication (see [23]).

References

- J. Brüning, S. Yu. Dobrokhotov and K. V. Pankrashkin. The spectral asymptotics of the two-dimensional Schrödinger operator with a strong magnetic field. I. Russ. J. Math. Phys. 9 (2002), no. 1, 14–49; II. Russ. J. Math. Phys. 9 (2002), no. 4, 400–416 (see also e-print version math-ph/0411012).
- [2] B. Helffer and Yu. A. Kordyukov, Semiclassical asymptotics and gaps in the spectra of periodic Schrödinger operators with magnetic wells, *Trans. Amer. Math. Soc.* 360 (2008), 1681-1694.
- [3] B. Helffer and Yu. A. Kordyukov, The periodic magnetic Schrödinger operators: spectral gaps and tunneling effect, Preprint math.SP/0702776; to appear in *Proc. Steklov Inst. Math.*
- [4] B. Helffer and J. Sjöstrand, Analyse semi-classique pour l'équation de Harper (avec application à l'équation de Schrödinger avec champ magnétique). Mém. Soc. Math. France (N.S.) 34 (1988)

15

- [5] B. Helffer and J. Sjöstrand, Équation de Schrödinger avec champ magnétique et équation de Harper, In: *Schrödinger operators (Sønderborg, 1988)*, Lecture Notes in Phys., 345 (Springer, Berlin, 1989), pp. 118–197.
- [6] R. Hempel and I. Herbst, Strong magnetic fields, Dirichlet boundaries, and spectral gaps. Commun. Math. Phys. 169 (1995), 237–259.
- [7] R. Hempel and O. Post, Spectral gaps for periodic elliptic operators with high contrast: an overview, In: *Progress in analysis, Vol. I, II (Berlin, 2001).* (World Sci. Publishing, River Edge, NJ, 2003), pp. 577–587.
- [8] I. Herbst and S. Nakamura. Schrodinger operators with strong magnetic fields: quasiperiodicity of spectral orbits and topology. In: *Differential operators and spectral theory*, Amer. Math. Soc. Transl. Ser. 2, 189 (Amer. Math. Soc., Providence, RI, 1999), pp. 105–123.
- [9] Yu. A. Kordyukov, Spectral gaps for periodic Schrödinger operators with strong magnetic fields. Commun. Math. Phys. 253 (2005), 371–384.
- [10] Yu. A. Kordyukov, Semiclassical asymptotics and spectral gaps for periodic magnetic Schrödinger operators on covering manifolds, In: "C^{*}-algebras and elliptic theory", Trends in Mathematics (Birkhäuser, Basel, 2006), pp. 129 – 150.
- [11] Yu. A. Kordyukov, V. Mathai and M. Shubin, Equivalence of projections in semiclassical limit and a vanishing theorem for higher traces in K-theory. J. Reine Angew. Math. 581 (2005), 193–236.
- [12] V. Mathai and M. Shubin, Semiclassical asymptotics and gaps in the spectra of magnetic Schrödinger operators. *Geometriae Dedicata* 91 (2002), 155–173.
- [13] S. Nakamura, Band spectrum for Schrödinger operators with strong periodic magnetic fields. In: *Partial differential operators and mathematical physics (Holzhau, 1994)*, Operator Theory: Advances and Applications. vol. 78 (Birkhäuser, Basel, 1995), pp. 261–270.
- [14] B. Helffer and A. Mohamed, Semiclassical analysis for the ground state energy of a Schrödinger operator with magnetic wells. J. Funct. Anal. 138 (1996), 40–81.
- [15] B. Helffer and J. Nourrigat, Hypoellipticité maximale pour des opérateurs polynômes de champs de vecteurs, (Birkhäuser, Boston, 1985)
- [16] B. Helffer and A. Mohamed, Caractérisation du spectre essentiel de l'opérateur de Schrödinger avec un champ magnétique. Ann. Inst. Fourier 38 (1988), 95–112.
- [17] R. Montgomery, Hearing the zero locus of a magnetic field. Comm. Math. Phys. 168 (1995), 651–675.
- [18] X-B. Pan and K-H. Kwek, Schrödinger operators with non-degenerately vanishing magnetic fields in bounded domains. *Trans. Amer. Math. Soc.* **354** (2002), 4201– 4227.
- [19] B. Helffer, Introduction to semi-classical methods for the Schrödinger operator with magnetic fields. To appear in: Séminaires et Congrès, Cours du CIMPA, SMF.
- [20] B. Helffer and J. Sjöstrand, Puits multiples en mécanique semi-classique. V. Étude des minipuits. *Current topics in partial differential equations* (Kinokuniya, Tokyo, 1986), pp. 133–186.
- [21] B. Helffer and J. Sjöstrand, Puits multiples en mécanique semi-classique. VI. Cas des puits sous-variétés. Ann. Inst. H. Poincaré Phys. Théor. 46 (1987), 353–372.
- [22] B. Helffer and A. Morame, Magnetic bottles in connection with superconductivity. J. Funct. Anal. 185 (2001), 604–680.
- [23] B. Helffer and Yu. A. Kordyukov, Spectral gaps for periodic Schrödinger operators with hypersurface magnetic wells : Analysis near the bottom. In preparation.

Département de Mathématiques, Bâtiment 425, Univ Paris-Sud et CNRS,, F-91405 Orsay Cédex, France

 $E\text{-}mail\ address:\ \texttt{Bernard}.\texttt{Helffer}\texttt{Qmath}.\texttt{u-psud}.\texttt{fr}$

Institute of Mathematics, Russian Academy of Sciences, 112 Chernyshevsky str., 450077 Ufa, Russia

E-mail address: yurikor@matem.anrb.ru