

The strict and relaxed stochastic maximum principle for optimal control problem of backward systems

Seid Bahlali*

Abstract

We consider a stochastic control problem where the set of controls is not necessarily convex and the system is governed by a nonlinear backward stochastic differential equation. We establish necessary as well as sufficient conditions of optimality for two models. The first concerns the strict (classical) controls. The second is an extension of the first to relaxed controls, who are a measure valued processes.

Keywords. Backward stochastic differential equation, strict control, relaxed control, maximum principle, adjoint equation, variational inequality, variational principle.

AMS Subject Classification. 93 Exx

1 Introduction

In this paper we study a stochastic control problem where the system is governed by a nonlinear backward stochastic differential equation (BSDE for short) of the type

$$\begin{cases} dy_t^v = b(t, y_t^v, z_t^v, v_t) dt + z_t^v dW_t, \\ y_T^v = \xi, \end{cases}$$

*Laboratory of applied mathematics, University Med Khider, P.O. Box 145, Biskra 07000, Algéria.

where b is given function, ξ is the terminal data and $W = (W_t)_{t \geq 0}$ is a standard d -dimensional Brownian motion, defined on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathcal{P})$ satisfying the usual conditions. The control variable $v = (v_t)$, called strict (classical) control, is an \mathcal{F}_t -adapted process with values in some set U of \mathbb{R}^k . We denote by \mathcal{U} the class of all strict controls.

The criteria to be minimized, over the set \mathcal{U} , has the form

$$J(v) = \mathbb{E} \left[g(y_0^v) + \int_0^T h(t, y_t^v, z_t^v, v_t) dt \right],$$

where g and h are given maps, and (y_t^v, z_t^v) is the trajectory of the system controlled by v .

A control $u \in \mathcal{U}$ is called optimal if it satisfies

$$J(u) = \inf_{v \in \mathcal{U}} J(v).$$

Stochastic control problems for the backward and forward-backward systems have been studied by many authors. The first contribution of control problems of forward-backward systems is made by Peng [30], he obtained the maximum principle with the control domain being convex. Xu [34] established the maximum principle for this kind of problem in the case where the control domain is not necessary convex, with uncontrolled diffusion coefficient and a restricted functional cost. The work of Peng [30] (convex control domain) is generalized by Wu [33], where the system is governed by a fully coupled forward-backward stochastic differential equation. Shi and Wu [32] extend the result of Xu [34] to the fully coupled forward-backward systems, with convex control domain and uncontrolled diffusion coefficient. Ji and Zhou [22] use the Ekeland variational principle and establish a maximum principle of controlled forward-backward systems, while the forward state is constrained in a convex set at the terminal time, and apply the result to state constrained stochastic linear-quadratic control models and a recursive utility optimization problem are investigated. All the cited previous works on stochastic control of forward-backward systems are obtained by introducing two adjoint equations. In the recent works on the subject, Bahlali and Labeled [3] and Bahlali [6] introduce three adjoint equations to establish necessary as well as sufficient optimality conditions. In [3] the authors establish the results in the case where the control domain being nonconvex and uncontrolled diffusion coefficient. The results of [6], are obtained while the control domain

is convex and with controlled diffusion coefficient, moreover the author apply his theory to solve the financial model of cash flow valuation.

On the other hand, stochastic maximum principle of backward systems was studied by El-Karoui et al [14], where the linear case is solved and some applications in finance are treated. Dokuchaev and Zhou [9] established necessary as well as sufficient optimality conditions, where the control domain is not convex.

Our objective in this paper is to establish necessary as well as sufficient optimality conditions, of the Pontryagin maximum principle type, for two models.

Firstly, we derive necessary as well as sufficient optimality conditions for strict controls. Since the set of strict controls is nonconvex, the classical way to use, is the spike variation method. More precisely, if u is an optimal strict control and v is arbitrary, then with a sufficiently small $\theta > 0$, we define a perturbed control as follows

$$u_t^\theta = \begin{cases} v & \text{if } t \in [\tau, \tau + \theta], \\ u_t & \text{otherwise.} \end{cases}$$

We then derive the variational equation from the state equation, and the variational inequality from the fact that

$$0 \leq J(u^\theta) - J(u).$$

The major difficulty in doing this is that the state of a backward system and the functional cost depends on two variables y_t and z_t . Then, we can't derive directly the variational inequality, because z_t is hard to handle, there is no convenient pointwise (in t) estimation for it, as opposed to the first variable y_t . To overcome this difficulty, we introduce a new method which consist to transform the initial control problem to a restricted problem without integral cost, by adding an unidimensional BSDE. We establish then necessary optimality conditions for the restricted control problem and by an adequate transformation on the adjoint process and the adjoint equation associated with the restricted problem, we reformulate necessary optimality conditions for the initial control problem.

To achieve this part of the paper, we study when these necessary optimality conditions becomes sufficient.

The second main result in this paper concerns necessary as well as sufficient optimality conditions for relaxed controls. In the relaxed model, the

controller chooses at time t a probability measure $q_t(da)$ on the control set U , rather than an element v_t of U . The system is then governed by the BSDE

$$\begin{cases} dy_t^q = \int_U b(t, y_t^q, z_t^q, a) q_t(da) dt + z_t^q dW_t, \\ y_T^q = \xi. \end{cases}$$

The criteria to be minimized, over the set \mathcal{R} of relaxed controls, has the form

$$\mathcal{J}(q) = \mathbb{E} \left[g(y_0^q) + \int_0^T \int_U h(t, y_t^q, z_t^q, a) q_t(da) dt \right].$$

A control $\mu \in \mathcal{R}$ is called optimal if it satisfies

$$\mathcal{J}(\mu) = \inf_{q \in \mathcal{R}} \mathcal{J}(q).$$

The relaxed control problem is an extension of the previous model of strict controls. Indeed, if $q_t(da) = \delta_{v_t}(da)$ is a Dirac measure concentrated at a single point v_t , then we get a strict control problem as a particular case of the relaxed one.

By using the Ekeland's variational principle, we are able to establish necessary optimality conditions for near optimal strict controls converging in some sense to the relaxed optimal control, by the so called chattering lemma. The relaxed necessary optimality conditions are then derived by using some stability properties of the trajectories and the adjoint process with respect to the control variable.

We note that necessary optimality conditions for relaxed controls, where the systems are governed by a stochastic differential equation, were studied by Mezerdi and Bahlali [27], Bahlali, Djehiche and Mezerdi [4].

The paper is organized as follows. In Section 2, we formulate the problem and give the various assumptions used throughout the paper. Section 3 is devoted to restrict the initial control problem to a problem without integral cost and we derive a restricted necessary optimality conditions. In Section 4, we give our first main result, the necessary optimality conditions for the initial control problem and under additional hypothesis, we prove that these conditions becomes sufficient. Finally, in the last Section, we give necessary optimality conditions for near optimal controls and from this we derive our second main result in this paper, necessary as well as sufficient optimality conditions for relaxed controls.

Along this paper, we denote by C some positive constant, $\mathcal{M}_{n \times d}(\mathbb{R})$ the space of $n \times d$ real matrix and $\mathcal{M}_{n \times n}^d(\mathbb{R})$ the linear space of vectors $M = (M_1, \dots, M_d)$ where $M_i \in \mathcal{M}_{n \times n}(\mathbb{R})$. We use the standard calculus of inner and matrix product.

2 Formulation of the problem

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathcal{P})$ be a probability space equipped with a filtration satisfying the usual conditions, on which a d -dimensional Brownian motion $W = (W_t)_{t \geq 0}$ is defined. We assume that (\mathcal{F}_t) is the \mathcal{P} - augmentation of the natural filtration of $(W_t)_{t \geq 0}$.

Let T be a strictly positive real number and U a non empty subset of \mathbb{R}^k .

Definition 1 *An admissible control is an \mathcal{F}_t - adapted process with values in U such that*

$$\mathbb{E} \left[\sup_{t \in [0, T]} |v_t|^2 \right] < \infty.$$

We denote by \mathcal{U} the set of all admissible controls.

For any $v \in \mathcal{U}$, we consider the following BSDE

$$\begin{cases} dy_t^v = b(t, y_t^v, z_t^v, v_t) dt + z_t^v dW_t, \\ y_T^v = \xi, \end{cases} \quad (1)$$

where

$$b : [0, T] \times \mathbb{R}^n \times \mathcal{M}_{n \times d}(\mathbb{R}) \times U \longrightarrow \mathbb{R}^n,$$

and ξ is an n -dimensional \mathcal{F}_T -measurable random variable such that

$$\mathbb{E} |\xi|^2 < \infty.$$

The expected cost is defined from \mathcal{U} into \mathbb{R} by

$$J(v) = \mathbb{E} \left[g(y_0^v) + \int_0^T h(t, y_t^v, z_t^v, v_t) dt \right], \quad (2)$$

where

$$\begin{aligned} g &: \mathbb{R}^n \longrightarrow \mathbb{R}, \\ h &: [0, T] \times \mathbb{R}^n \times \mathcal{M}_{n \times d}(\mathbb{R}) \times U \longrightarrow \mathbb{R}. \end{aligned}$$

A control $u \in \mathcal{U}$ is called optimal, if that solves

$$J(u) = \inf_{v \in \mathcal{U}} J(v). \quad (3)$$

Our goal is to establish necessary as well as sufficient optimality conditions for controls in the form of stochastic maximum principle.

The following assumptions will be in force throughout this paper

The functions b, g and h are continuous in (y, z, v) , they are differentiable with respect to (y, z) , and their derivatives b_y, b_z, g_y, h_y and h_z are continuous in (y, z, v) and uniformly bounded. b and h are bounded by $C(1 + |y| + |v|)$ and bounded in z . (4)

Under the above hypothesis, for every $v \in \mathcal{U}$, equation (1) has a unique strong $(\mathcal{F}_t)_t$ -adapted solution and the functional cost J is well defined from \mathcal{U} into \mathbb{R} .

3 Problem with restricted cost

Since the function h of the cost depends explicitly on z_t , we can't treat our problem directly. Thus, let us in this section restrict the initial control problem $\{(1), (2), (3)\}$ to a problem without integral cost. For this end, consider the following unidimensional BSDE

$$\begin{cases} dx_t^v = h(t, y_t^v, z_t^v, v_t) dt + k_t^v dW_t, \\ x_T^v = \eta, \end{cases}$$

where k^v is a $(1 \times d)$ matrix, (y_t^v, z_t^v) is the solution of equation (1) and η is a one-dimensional \mathcal{F}_T -measurable random variable such that

$$\mathbb{E} |\eta|^2 < \infty.$$

The above equation admits a unique strong $(\mathcal{F}_t)_t$ -adapted solution.

We put

$$\tilde{y}_t = \begin{pmatrix} y_t^v \\ x_t^v \end{pmatrix},$$

and consider now the following $(n + 1)$ -dimensional BSDE

$$\begin{cases} d\tilde{y}_t = \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, v_t) dt + \tilde{z}_t dW_t, \\ \tilde{y}_T = \begin{pmatrix} \xi \\ \eta \end{pmatrix}, \end{cases} \quad (5)$$

where the functions \tilde{b} is defined from $[0, T] \times \mathbb{R}^{n+1} \times \mathcal{M}_{(n+1) \times d}(\mathbb{R}) \times U$ into \mathbb{R}^{n+1} by

$$\tilde{b}(t, \tilde{y}_t, \tilde{z}_t, v_t) = \begin{pmatrix} b(t, y_t^v, z_t^v, v_t) \\ h(t, y_t^v, z_t^v, v_t) \end{pmatrix},$$

and \tilde{z}_t is a $(n + 1) \times d$ real matrix given by

$$\tilde{z}_t = \begin{pmatrix} z_t^v \\ k_t^v \end{pmatrix} = \begin{pmatrix} z_{11}^v & z_{12}^v & \dots & z_{1d}^v \\ z_{21}^v & z_{22}^v & \dots & z_{2d}^v \\ \vdots & & & \vdots \\ z_{n1}^v & z_{n2}^v & \dots & z_{nd}^v \\ k_1^v & k_2^v & \dots & k_d^v \end{pmatrix}.$$

From (4), \tilde{b} is uniformly Lipschitz in $(\tilde{y}_t, \tilde{z}_t)$, then equation (1) admits a unique strong solution $(\tilde{y}_t, \tilde{z}_t)$ adapted to the filtration $(\mathcal{F}_t)_t$.

Define now the function \tilde{g} from \mathbb{R}^{n+1} into \mathbb{R} by

$$\tilde{g}(\tilde{y}_t) = g(y_t^v) - x_t^v,$$

and the new functional cost from \mathcal{U} into \mathbb{R} by

$$\tilde{J}(v) = \mathbb{E}[\tilde{g}(\tilde{y}_0)] + \mathbb{E}[\eta]. \quad (6)$$

It's easy to see that

$$\tilde{J}(v) = J(v).$$

Consequently, it's sufficient to minimize the restricted cost \tilde{J} over \mathcal{U} . If $u \in \mathcal{U}$ is an optimal solution, that is

$$\tilde{J}(u) = \inf_{v \in \mathcal{U}} \tilde{J}(v). \quad (7)$$

From this transformation, we have reduce our initial problem $\{(1), (2), (3)\}$ to a new problem without integral cost. We can now study the restricted

problem $\{(5), (6), (7)\}$ by using a classical way of spike variation method. We establish necessary optimality conditions for a restricted problem and by an adequate transformation on the adjoint process and the adjoint equation associated with the restricted problem, we reformulate necessary optimality conditions for the initial control problem $\{(1), (2), (3)\}$.

3.1 Preliminary results

Suppose that $u \in \mathcal{U}$ is an optimal control and denote by $(\tilde{y}_t, \tilde{z}_t)$ the solution of (5) corresponding to u . Introduce the following perturbation (spike variation) of the optimal control u

$$u_t^\theta = \begin{cases} v & \text{if } t \in [\tau, \tau + \theta], \\ u_t & \text{otherwise,} \end{cases} \quad (8)$$

where $0 \leq \tau \leq T$ is fixed, $\theta > 0$ is sufficiently small and v is an arbitrary \mathcal{F}_τ -measurable random variable with values in U such that $\mathbb{E}[|v|^2] < \infty$.

The control u^θ is admissible and let $(\tilde{y}_t^\theta, \tilde{z}_t^\theta)$ be the solution of (5) associated with u_t^θ .

Since u is optimal, the variational inequality will be derived from the fact that

$$0 \leq \tilde{J}(u^\theta) - \tilde{J}(u). \quad (9)$$

For this end, we need the following lemmas.

Lemma 2 *Under assumptions (4), we have*

$$\mathbb{E} \left[\sup_{t \in [0, T]} |\tilde{y}_t^\theta - \tilde{y}_t|^2 \right] \leq C\theta^2, \quad (10)$$

$$\mathbb{E} \int_0^T |\tilde{z}_t^\theta - \tilde{z}_t|^2 dt \leq C\theta^2. \quad (11)$$

Proof. We have

$$\begin{cases} d(\tilde{y}_t^\theta - \tilde{y}_t) = \begin{bmatrix} \tilde{b}(t, \tilde{y}_t^\theta, \tilde{z}_t^\theta, u_t^\theta) - \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t) \\ \tilde{b}(t, \tilde{y}_t, \tilde{z}_t^\theta, u_t^\theta) - \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t) \\ \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t^\theta) - \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t) \end{bmatrix} dt \\ \quad + (\tilde{z}_t^\theta - \tilde{z}_t) dW_t, \\ (\tilde{y}_T^\theta - \tilde{y}_T) = 0. \end{cases}$$

Put

$$\begin{aligned} Y_t^\theta &= \tilde{y}_t^\theta - \tilde{y}_t, \\ Z_t^\theta &= \tilde{z}_t^\theta - \tilde{z}_t, \end{aligned}$$

and

$$\begin{aligned} \varphi^\theta(t, Y_t^\theta, Z_t^\theta) &= \int_0^1 \tilde{b}_y(t, \tilde{y}_t + \lambda(\tilde{y}_t^\theta - \tilde{y}_t), \tilde{z}_t + \lambda(\tilde{z}_t^\theta - \tilde{z}_t), u_t^\theta) Y_t^\theta d\lambda \\ &+ \int_0^1 \tilde{b}_z(t, \tilde{y}_t + \lambda(\tilde{y}_t^\theta - \tilde{y}_t), \tilde{z}_t + \lambda(\tilde{z}_t^\theta - \tilde{z}_t), u_t^\theta) Z_t^\theta d\lambda \\ &+ \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t^\theta) - \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t). \end{aligned} \quad (12)$$

Then

$$\begin{cases} dY_t^\theta = \varphi^\theta(t, Y_t^\theta, Z_t^\theta) dt + Z_t^\theta dW_t, \\ Y_T^\theta = 0. \end{cases} \quad (13)$$

The above equation is a linear BSDE with bounded coefficients and with terminal condition $Y_T^\theta = 0$. Then by applying a priori estimates (see Briand et al [8, Proposition 3.2, Page 7]), we get

$$\mathbb{E} \left[\sup_{t \in [0, T]} |Y_t^\theta|^2 + \int_0^T |Z_t^\theta|^2 dt \right] \leq C \mathbb{E} \left| \int_0^T |\varphi^\theta(t, 0, 0)| dt \right|^2.$$

From (12), we get

$$\mathbb{E} \left[\sup_{t \in [0, T]} |Y_t^\theta|^2 + \int_0^T |Z_t^\theta|^2 dt \right] \leq C \mathbb{E} \left| \int_0^T \left| \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t^\theta) - \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t) \right| dt \right|^2.$$

By the definition of u^θ , we have

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} |Y_t^\theta|^2 + \int_0^T |Z_t^\theta|^2 dt \right] &\leq C \mathbb{E} \left| \int_\tau^{\tau+\theta} \left| \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, v) - \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t) \right| dt \right|^2 \\ &\leq C \mathbb{E} \left| \sup_{t \in [0, T]} \left| \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, v) - \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t) \right| \int_\tau^{\tau+\theta} dt \right|^2. \end{aligned}$$

By (4), b is with linear growth with respect to (y, v) and bounded in z , then \tilde{b} satisfy the same properties, and we get

$$\mathbb{E} \left[\sup_{t \in [0, T]} |Y_t^\theta|^2 + \int_0^T |Z_t^\theta|^2 dt \right] \leq C\theta^2.$$

The lemma is proved. ■

3.2 Necessary optimality conditions for restricted problem

We can now state necessary optimality conditions for a restricted control problem $\{(5), (6), (7)\}$.

Theorem 3 (*necessary optimality conditions for restricted problem*) *Let $(u, \tilde{y}, \tilde{z})$ be an optimal solution of the restricted control problem $\{(5), (6), (7)\}$. Then there exists a unique adapted process*

$$\tilde{p} \in \mathcal{L}^2([0, T]; \mathbb{R}^{n+1}),$$

which is solution of the following forward stochastic differential equation

$$\begin{cases} -d\tilde{p}_t = \tilde{H}_y(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) dt + \tilde{H}_z(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) dW_t, \\ \tilde{p}_0 = \tilde{g}_y(\tilde{y}_0), \end{cases} \quad (14)$$

such that

$$\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) = \max_{v \in U} \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, v); \text{ a.e., a.s.} \quad (15)$$

where the Hamiltonian \tilde{H} is defined from $[0, T] \times \mathbb{R}^{n+1} \times \mathcal{M}_{(n+1) \times d}(\mathbb{R}) \times \mathbb{R}^{n+1} \times U$ into \mathbb{R} by

$$\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) = \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t) \tilde{p}_t.$$

Proof. For simplicite, we put

$$\Lambda_t^\theta = (t, \tilde{y}_t + \lambda(\tilde{y}_t^\theta - \tilde{y}_t), \tilde{z}_t + \lambda(\tilde{z}_t^\theta - \tilde{z}_t), u_t^\theta).$$

Since u minimizes the cost \tilde{J} over \mathcal{U} , then ■

$$\begin{aligned} 0 &\leq \tilde{J}(u^\theta) - \tilde{J}(u) \\ &\leq \mathbb{E} [\tilde{g}(\tilde{y}_0^\theta) - \tilde{g}(\tilde{y}_0)] \\ &\leq \mathbb{E} \int_0^1 \tilde{g}_y[\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)] (\tilde{y}_0^\theta - \tilde{y}_0) d\lambda \\ &\leq \mathbb{E} [\tilde{g}_y(\tilde{y}_0) (\tilde{y}_0^\theta - \tilde{y}_0)] + \mathbb{E} \int_0^1 [\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)] (\tilde{y}_0^\theta - \tilde{y}_0) d\lambda. \end{aligned}$$

We remark from (14) that

$$\tilde{p}_0 = \tilde{g}_y(\tilde{y}_0).$$

Then

$$0 \leq \mathbb{E} [\tilde{p}_0 (\tilde{y}_0^\theta - \tilde{y}_0)] + \mathbb{E} \int_0^1 [\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)] (\tilde{y}_0^\theta - \tilde{y}_0) d\lambda. \quad (16)$$

By applying Itô's formula to $\tilde{p}_t(\tilde{y}_t^\theta - \tilde{y}_t)$, we get

$$\begin{aligned} \mathbb{E} [\tilde{p}_0 (\tilde{y}_0^\theta - \tilde{y}_0)] &= \mathbb{E} \int_0^T \int_0^1 [\tilde{b}_y(\Lambda_t^\theta) - \tilde{b}_y(t, \tilde{y}_t, \tilde{z}_t, u_t)] (\tilde{y}_t - \tilde{y}_t^\theta) \tilde{p}_t d\lambda dt \\ &\quad + \mathbb{E} \int_0^T \int_0^1 [\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t)] (\tilde{z}_t - \tilde{z}_t^\theta) \tilde{p}_t d\lambda dt \\ &\quad + \mathbb{E} \int_0^T [\tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t) - \tilde{b}(t, \tilde{y}_t, \tilde{z}_t, u_t^\theta)] \tilde{p}_t dt. \end{aligned}$$

Then (16) becomes

$$\begin{aligned} 0 &\leq \mathbb{E} \int_0^T [\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t^\theta)] dt \quad (17) \\ &\quad - \mathbb{E} \int_0^1 [\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)] (\tilde{y}_0 - \tilde{y}_0^\theta) d\lambda \\ &\quad + \mathbb{E} \int_0^T \int_0^1 [\tilde{b}_y(\Lambda_t^\theta) - \tilde{b}_y(t, \tilde{y}_t, \tilde{z}_t, u_t)] (\tilde{y}_t - \tilde{y}_t^\theta) \tilde{p}_t d\lambda dt \\ &\quad + \mathbb{E} \int_0^T \int_0^1 [\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t)] (\tilde{z}_t - \tilde{z}_t^\theta) \tilde{p}_t d\lambda dt \end{aligned}$$

Let us show that

$$\mathbb{E} \int_0^T \int_0^1 [\tilde{b}_y(\Lambda_t^\theta) - \tilde{b}_y(t, \tilde{y}_t, \tilde{z}_t, u_t)] (\tilde{y}_t - \tilde{y}_t^\theta) \tilde{p}_t d\lambda dt \leq C\theta^{3/2}, \quad (18)$$

and

$$\mathbb{E} \int_0^T \int_0^1 [\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t)] (\tilde{z}_t - \tilde{z}_t^\theta) \tilde{p}_t d\lambda dt \leq C\theta^{3/2}. \quad (19)$$

Indeed, by using the Cauchy-Schwartz inequality to term in the left hand side of (19), we get

$$\begin{aligned} & \mathbb{E} \int_0^T \int_0^1 \left[\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t) \right] (\tilde{z}_t - \tilde{z}_t^\theta) \tilde{p}_t d\lambda dt \\ & \leq \left(\mathbb{E} \int_0^T \int_0^1 \left| \left[\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t) \right] \tilde{p}_t \right|^2 d\lambda dt \right)^{1/2} \left(\mathbb{E} \int_0^T |\tilde{z}_t - \tilde{z}_t^\theta|^2 dt \right)^{1/2}. \end{aligned}$$

By (11), we obtain

$$\begin{aligned} & \mathbb{E} \int_0^T \int_0^1 \left[\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t) \right] (\tilde{z}_t - \tilde{z}_t^\theta) \tilde{p}_t d\lambda dt \\ & \leq C\theta \left(\mathbb{E} \int_0^T \int_0^1 \left| \left[\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t) \right] \tilde{p}_t \right|^2 d\lambda dt \right)^{1/2}. \end{aligned}$$

By the definition of u^θ , we have

$$\begin{aligned} & \mathbb{E} \int_0^T \int_0^1 \left[\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t) \right] (\tilde{z}_t - \tilde{z}_t^\theta) \tilde{p}_t d\lambda dt \\ & \leq C\theta \left(\mathbb{E} \int_\tau^{\tau+\theta} \int_0^1 \left| \left[\tilde{b}_z(\Lambda_t^v) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t) \right] \tilde{p}_t \right|^2 d\lambda dt \right)^{1/2}. \end{aligned}$$

Since \tilde{b}_y is bounded, we get

$$\begin{aligned} & \mathbb{E} \int_0^T \int_0^1 \left[\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t) \right] (\tilde{z}_t^\theta - \tilde{z}_t) \tilde{p}_t d\lambda dt \\ & \leq C\theta \left(\int_\tau^{\tau+\theta} \mathbb{E} |\tilde{p}|^2 dt \right)^{1/2}. \end{aligned}$$

Since $\tilde{p} \in \mathcal{L}^2([0, T]; \mathbb{R}^{n+1})$, we obtain

$$\begin{aligned} & \mathbb{E} \int_0^T \int_0^1 \left[\tilde{b}_z(\Lambda_t^\theta) - \tilde{b}_z(t, \tilde{y}_t, \tilde{z}_t, u_t) \right] (\tilde{z}_t^\theta - \tilde{z}_t) \tilde{p}_t d\lambda dt \\ & \leq \left(C \int_\tau^{\tau+\theta} dt \right)^{1/2} C\theta = C\theta^{3/2}. \end{aligned}$$

Relation (19) is proved.

(18) is proved by the same method and by using (10) and the fact that b_y is bounded.

Now, by (17), (18) and (19) we get

$$\begin{aligned} 0 &\leq \mathbb{E} \int_0^T \left[\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t^\theta) \right] dt \\ &\quad + \mathbb{E} \int_0^1 [\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)] (\tilde{y}_0^\theta - \tilde{y}_0) d\lambda \\ &\quad + C\theta^{3/2}. \end{aligned}$$

By applying the Cauchy-Schwartz inequality to the second term in the right hand side of the above inequality, we get

$$\begin{aligned} 0 &\leq \mathbb{E} \int_0^T \left[\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t^\theta) \right] dt \\ &\quad + \left(\int_0^1 \mathbb{E} |\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)|^2 d\lambda \right)^{1/2} \left(\mathbb{E} |\tilde{y}_0^\theta - \tilde{y}_0|^2 \right)^{1/2} \\ &\quad + C\theta^{3/2}. \end{aligned}$$

By (10), we deduce

$$\begin{aligned} 0 &\leq \mathbb{E} \int_0^T \left[\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t^\theta) \right] dt \\ &\quad + C\theta \left(\int_0^1 \mathbb{E} |\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)|^2 d\lambda \right)^{1/2} \\ &\quad + C\theta^{3/2}. \end{aligned}$$

From the definition of u_t^θ , we have

$$\begin{aligned} 0 &\leq \mathbb{E} \int_\tau^{\tau+\theta} \left[\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, v) \right] dt \\ &\quad + C\theta \left(\int_0^1 \mathbb{E} |\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)|^2 d\lambda \right)^{1/2} \\ &\quad + C\theta^{3/2}. \end{aligned}$$

Dividing by θ , we get

$$\begin{aligned}
0 &\leq \frac{1}{\theta} \mathbb{E} \int_{\tau}^{\tau+\theta} \left[\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, v) \right] dt \quad (20) \\
&+ C \left(\int_0^1 \mathbb{E} |\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)|^2 d\lambda \right)^{1/2} \\
&+ C\theta^{1/2}.
\end{aligned}$$

Since \tilde{g}_y is continuous and bounded, then by (10) and the dominated convergence theorem, we have

$$\lim_{\theta \rightarrow 0} C \int_0^1 \left(\mathbb{E} |\tilde{g}_y(\tilde{y}_0 + \lambda(\tilde{y}_0^\theta - \tilde{y}_0)) - \tilde{g}_y(\tilde{y}_0)|^2 \right)^{1/2} d\lambda = 0.$$

Then, by taking the limit as $\theta \rightarrow 0$ in (20), we obtain

$$0 \leq \lim_{\theta \rightarrow 0} \frac{1}{\theta} \mathbb{E} \int_{\tau}^{\tau+\theta} \left[\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, v) \right] dt.$$

This implies that

$$0 \leq \mathbb{E} \left[\tilde{H}(\tau, \tilde{y}_\tau, \tilde{z}_\tau, \tilde{p}_\tau, u_\tau) - \tilde{H}(\tau, \tilde{y}_\tau, \tilde{z}_\tau, \tilde{p}_\tau, v) \right], \quad d\tau - a.e.$$

Now, let $a \in U$ be a deterministic element and F be an arbitrary element of the σ -algebra \mathcal{F}_t , and set

$$w_t = a \mathbf{1}_F + u_t \mathbf{1}_{\Omega - F}.$$

It is obvious that w is an admissible control.

Since $0 \leq \tau \leq T$, then for every bounded U -valued, \mathcal{F}_t -measurable random variable v such that $\mathbb{E}|v|^2 < +\infty$, we get

$$0 \leq \mathbb{E} \left[\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, v) \right], \quad dt - a.e.,$$

Applying the above inequality with w , we get

$$0 \leq \mathbb{E}[\mathbf{1}_F(\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, a))], \quad \forall F \in \mathcal{F}_t,$$

which implies that

$$0 \leq \mathbb{E}[\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) - \tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, a) \mid \mathcal{F}_t].$$

The quantity inside the conditional expectation is \mathcal{F}_t -measurable, and thus the result follows immediately. This prove theorem 3.

4 Necessary and sufficient optimality conditions for strict controls

Starting from the results of the last section, we can now reformulate the restricted necessary optimality conditions given by theorem 3, and state necessary as well as sufficient optimality conditions for the initial control problem $\{(1), (2), (3)\}$.

4.1 Necessary optimality conditions

Theorem 4 (*necessary optimality conditions for strict controls*) *Let (u, y^u, z^u) be an optimal solution of the initial control problem $\{(1), (2), (3)\}$. Then there exists a unique adapted processes*

$$p^u \in \mathcal{L}^2([0, T]; \mathbb{R}^n),$$

which are solution of the following forward stochastic differential equation

$$\begin{cases} -dp_t^u = H_y(t, y_t^u, z_t^u, p_t^u, u_t) dt + H_z(t, y_t^u, z_t^u, p_t^u, u_t) dW_t, \\ p_0^u = g_y(y_0^u), \end{cases} \quad (21)$$

such that

$$H(t, y_t^u, z_t^u, p_t^u, u_t) = \max_{v \in U} H(t, y_t^u, z_t^u, p_t^u, v) ; \text{ a.e. , a.s.} \quad (22)$$

where the Hamiltonian H is defined from $[0, T] \times \mathbb{R}^n \times \mathcal{M}_{n \times d}(\mathbb{R}) \times \mathbb{R}^n \times U$ into \mathbb{R} by

$$H(t, y, z, p, v) = pb(t, y, z, v) - h(t, y, z, v).$$

Proof. We put

$$\tilde{p}_t = \begin{pmatrix} p_t^u \\ -1 \end{pmatrix}.$$

From the definition of \tilde{H} , \tilde{p} , \tilde{b} and \tilde{z} , we have

$$\tilde{H}(t, \tilde{y}_t, \tilde{z}_t, \tilde{p}_t, u_t) = H(t, y_t^u, z_t^u, p_t^u, u_t), \quad (23)$$

and from the adjoint equation (14), we can easily deduce (21). Finally (22) is derived immediately from (23) and (15). ■

4.2 Sufficient optimality conditions

Theorem 5 (*Sufficient optimality conditions for strict controls*). *If we assume that, U is convex and for every $v \in \mathcal{U}$ and for all $t \in [0, T]$, the function g is convex and $(y_t, z_t, v_t) \rightarrow H(t, y_t, z_t, p_t, v_t)$ is concave. Then u is an optimal control of the problem $\{(1), (2), (3)\}$ if it satisfies (22).*

Proof. Let u be an arbitrary admissible control (candidate to be optimal) and (y_t^u, z_t^u) the solution of (1) associated with u . For any admissible control v , with associated trajectory (y_t^v, z_t^v) , we have

$$\begin{aligned} J(v) - J(u) &= \mathbb{E}[g(y_0^v) - g(y_0^u)] \\ &\quad + \mathbb{E} \int_0^T [h(t, y_t^v, z_t^v, v_t) - h(t, y_t^u, z_t^u, u_t)] dt. \end{aligned}$$

Since g is convex, then

$$g(y_0^v) - g(y_0^u) \geq g_y(y_0^u)(y_0^v - y_0^u).$$

Then

$$\begin{aligned} J(v) - J(u) &\geq \mathbb{E}[g_y(y_0^u)(y_0^v - y_0^u)] \\ &\quad + \mathbb{E} \int_0^T [h(t, y_t^v, z_t^v, v_t) - h(t, y_t^u, z_t^u, u_t)] dt. \end{aligned}$$

We remark from (21) that

$$p_0^u = g_y(y_0^u).$$

Then, we have

$$\begin{aligned} J(v) - J(u) &\geq \mathbb{E}[p_0^u(y_0^v - y_0^u)] \\ &\quad + \mathbb{E} \int_0^T [h(t, y_t^v, z_t^v, v_t) - h(t, y_t^u, z_t^u, u_t)] dt. \end{aligned}$$

By applying Itô's formula to $p_t^u(y_t^v - y_t^u)$, we obtain

$$\begin{aligned} &J(v) - J(u) \\ &\geq \mathbb{E} \int_0^T [H_y(t, y_t^u, z_t^u, p_t^u, u_t)(y_t^v - y_t^u) + H_z(t, y_t^u, z_t^u, p_t^u, u_t)(z_t^v - z_t^u)] dt \\ &\quad + \mathbb{E} \int_0^T [H(t, y_t^u, z_t^u, p_t^u, u_t) - H(t, y_t^v, z_t^v, p_t^u, v_t)] dt. \end{aligned}$$

Since H is concave in (y, z, u) , then

$$\begin{aligned} & H(t, y_t^v, z_t^v, p_t^u, v_t) - H(t, y_t^u, z_t^u, p_t^u, u_t) \\ & \leq H_y(t, y_t^u, z_t^u, p_t^u, u_t)(y_t^v - y_t^u) \\ & \quad + H_z(t, y_t^u, z_t^u, p_t^u, u_t)(z_t^v - z_t^u) + H_v(t, y_t^u, z_t^u, p_t^u, u_t)(v_t - u_t). \end{aligned}$$

Or equivalently

$$\begin{aligned} & H_v(t, y_t^u, z_t^u, p_t^u, u_t)(u_t - v_t) \\ & \leq H(t, y_t^u, z_t^u, p_t^u, u_t) - H(t, y_t^v, z_t^v, p_t^u, v_t) \\ & \quad + H_y(t, y_t^u, z_t^u, p_t^u, u_t)(y_t^v - y_t^u) + H_z(t, y_t^u, z_t^u, p_t^u, u_t)(z_t^v - z_t^u). \end{aligned}$$

Then, we get

$$J(v) - J(u) \geq \mathbb{E} \int_0^T H_v(t, y_t^u, z_t^u, p_t^u, u_t)(u_t - v_t) dt. \quad (24)$$

We know that $H(t, y_t^u, z_t^u, p_t^u, \cdot)$ is concave, then $-H(t, y_t^u, z_t^u, p_t^u, \cdot)$ is convex from U into \mathbb{R} . Furthermore U is convex and $-H(t, y_t^u, z_t^u, p_t^u, \cdot)$ is continuous, Gâteaux-differentiable, with differential continuous, then from the convex optimization principle (see Ekeland-Temam [11, prop 2.1, page 35]), we have

$$-H(t, y_t^u, z_t^u, p_t^u, u_t) = \inf_{v_t \in U} -H(t, y_t^u, z_t^u, p_t^u, v_t) \iff -H_v(t, y_t^u, z_t^u, p_t^u, u_t)(v_t - u_t) \geq 0.$$

Or equivalently

$$H(t, y_t^u, z_t^u, p_t^u, u_t) = \max_{v_t \in U} H(t, y_t^u, z_t^u, p_t^u, v_t) \iff H_v(t, y_t^u, z_t^u, p_t^u, u_t)(u_t - v_t) \geq 0.$$

Then from the necessary condition of optimality (22), we deduce that

$$H_v(t, y_t^u, z_t^u, p_t^u, u_t)(u_t - v_t) \geq 0.$$

And from (24), we have

$$J(v) - J(u) \geq 0.$$

The theorem is proved. ■

5 The relaxed model

In this section, we generalize the results of the above section to a relaxed control problem. The idea for relaxed the strict control problem defined above is to embed the set U of strict controls into a wider class which gives a more suitable topological structure. In the relaxed model, the U -valued process v is replaced by a $\mathbb{P}(U)$ -valued process q , where $\mathbb{P}(U)$ denotes the space of probability measure on U equipped with the topology of stable convergence.

Let V the set of positive random measures on $[0, T] \times U$ whose projection on $[0, T]$ coincide with the Lebesgue measure dt . Equipped with the topology of stable convergence of measures, V is a compact metrizable space. The stable convergence is required for bounded measurable functions $f(t, a)$ such that for each fixed $t \in [0, T]$, $h(t, \cdot)$ is continuous. The space V is equipped with its Borel σ -field, which is the smallest σ -field such that the mapping $q \mapsto \int f(s, a) q(ds, da)$ are measurable for any bounded measurable function f , continuous with respect to a (Instead of functions bounded and continuous with respect to the pair (t, a) for the weak topology).

For more details, see Jacod-Memin [18, page 629-630] and El Karoui et al [9, Page 4-5].

Definition 6 *A relaxed control $(q_t)_t$ is a $\mathbb{P}(U)$ -valued process, progressively measurable with respect to $(\mathcal{F}_t)_t$ and such that for each t , $1_{]0,t[} \cdot q$ is \mathcal{F}_t -measurable.*

We denote by \mathcal{R} the set of all relaxed controls.

Every relaxed control q may be desintegrated as $q(dt, da) = q(t, da) dt = q_t(da) dt$, where $q_t(da)$ is a progressively measurable process with value in the set of probability measures $\mathbb{P}(U)$.

The set U is embedded into the set \mathcal{R} of relaxed process by the mapping

$$f : v \in U \mapsto f_v(dt, da) = \delta_{v_t}(da) dt \in \mathcal{R}$$

where δ_v is the atomic measure concentrated at a single point v .

For more details on relaxed controls, see [2], [4], [5], [12], [16], [26], [27].

For any $q \in \mathcal{R}$, we consider the following relaxed BSDE

$$\begin{cases} dy_t^q = \int_U b(t, y_t^q, z_t^q, a) q_t(da) dt + z_t^q dW_t, \\ y_T^q = \xi. \end{cases} \quad (25)$$

The expected cost associated to a relaxed control q is defined as follows

$$\mathcal{J}(q) = \mathbb{E} \left[g(y_0^q) + \int_0^T \int_U h(t, y_t^q, z_t^q, a) q_t(da) dt \right]. \quad (26)$$

Our objective is to minimize the functional \mathcal{J} over \mathcal{R} . If $\mu \in \mathcal{R}$ is an optimal relaxed control, that is

$$\mathcal{J}(\mu) = \inf_{q \in \mathcal{R}} \mathcal{J}(q). \quad (27)$$

Throughout this section we suppose moreover that

$$\begin{aligned} &U \text{ is compact,} \\ &b \text{ and } h \text{ are bounded,} \\ &b_y, h_y, b_z \text{ and } h_z \text{ are Lipschitz continuous in } z. \end{aligned} \quad (28)$$

Remark 7 *If we put*

$$\begin{aligned} \bar{b}(t, y_t^q, z_t^q, q_t) &= \int_U b(t, y_t^q, z_t^q, a) q_t(da), \\ \bar{h}(t, y_t^q, z_t^q, q_t) &= \int_U h(t, y_t^q, z_t^q, a) q_t(da), \end{aligned}$$

then equation (25) becomes

$$\begin{cases} dy_t^q = \bar{b}(t, y_t^q, z_t^q, q_t) dt + z_t^q dW_t, \\ y^q(T) = \xi, \end{cases}$$

with a functional cost given by

$$\mathcal{J}(q) = \mathbb{E} \left[g(y_0^q) + \int_0^T \bar{h}(t, y_t^q, z_t^q, q_t) dt \right].$$

Hence by introducing relaxed controls, we have replaced U by a larger space $\mathbb{P}(U)$. We have gained the advantage that $\mathbb{P}(U)$ is both compact and convex, the new drift and the integral coefficient of \mathcal{J} are linear in q .

On the other hand, the coefficients \bar{b} (defined above) check the same assumptions as b . Then, under assumptions (4), \bar{b} is uniformly Lipschitz and with linear growth. Then, by classical results on BSDEs (The Pardoux-Peng theorem, see : Pardoux-Peng [28]), for every $q \in \mathcal{R}$, equation (25) has a unique solution.

Moreover, It is easy to see that \bar{h} checks the same assumptions as h . Then, the functional cost \mathcal{J} is well defined from \mathcal{R} into \mathbb{R} .

Remark 8 If $q_t = \delta_{v_t}$ is an atomic measure concentrated at a single point v_t , then for each $t \in [0, T]$ we have

$$\begin{aligned} \int_U b(t, y_t^q, z_t^q, a) q_t(da) &= \int_U b(t, y_t^q, z_t^q, a) \delta_{v_t}(da) = b(t, y_t^q, z_t^q, v_t), \\ \int_U h(t, y_t^q, z_t^q, a) q_t(da) &= \int_U h(t, y_t^q, z_t^q, a) \delta_{v_t}(da) = h(t, y_t^q, z_t^q, v_t). \end{aligned}$$

In this case $(y^q, z^q) = (y^v, z^v)$, $J(v) = \mathcal{J}(q)$ and we get an ordinary admissible control problem. So the problem of strict controls defined in the section 2 is a particular case of the problem of relaxed one.

5.1 Approximation of trajectories

The next lemma, known as the Chattering Lemma, tells us that any relaxed control is a stable limit of a sequence of strict controls. This lemma was first proved for deterministic measures and then extended to random measures in [12] and [16].

Lemma 9 (*Chattering Lemma*). Let q_t be a predictable process with values in the space of probability measures on U . Then there exists a sequence of predictable processes $(u^n)_n$ with values in U such that

$$dtq_t^n(da) = dt\delta_{u_t^n}(da) \xrightarrow[n \rightarrow \infty]{} dtq_t(da) \text{ stably, } \mathcal{P} - a.s. \quad (29)$$

Proof. See El Karoui et al [12]. ■

Lemma 10 Let q be a relaxed control and $(u^n)_n$ be a sequence of strict controls such that (29) holds. Then for any bounded function $f : [0, T] \times U \rightarrow \mathbb{R}$, measurable in t and continuous in a , we have

$$\int_U f(t, a) \delta_{u_t^n}(da) \xrightarrow[n \rightarrow \infty]{} \int_U f(t, a) q_t(da). \quad (30)$$

Proof. By the Chattering lemma and the definition of the stable convergence (see Jacod-Memin [21, definition 1.1, page 529], we have

$$\int_0^T \int_U f(t, a) \delta_{u_t^n}(da) dt \xrightarrow{n \rightarrow \infty} \int_0^T \int_U f(t, a) q_t(da) dt.$$

Put

$$g(s, a) = 1_{[0, t]}(s) f(s, a).$$

It's clear that

$$\int_0^T \int_U g(s, a) \delta_{u_s^n}(da) ds \xrightarrow{n \rightarrow \infty} \int_0^T \int_U g(s, a) q_s(da) ds.$$

Then

$$\int_0^t \int_U f(s, a) \delta_{u_s^n}(da) ds \xrightarrow{n \rightarrow \infty} \int_0^t \int_U f(s, a) q_s(da) ds.$$

The set $\{(s, t) ; 0 \leq s \leq t \leq T\}$ generate $\mathcal{B}_{[0, T]}$. Then $\forall B \in \mathcal{B}_{[0, T]}$, we have

$$\int_B \int_U f(s, a) \delta_{u_s^n}(da) ds \xrightarrow{n \rightarrow \infty} \int_B \int_U f(s, a) q_s(da) ds.$$

This implies that

$$\int_U f(s, a) \delta_{u_s^n}(da) \xrightarrow{n \rightarrow \infty} \int_U f(s, a) q_s(da) , \quad dt - a.e.$$

The lemma is proved. ■

The next lemma gives the stability of the controlled stochastic differential equation with respect to the control variable.

Lemma 11 *Let $q_t \in \mathcal{R}$ be a relaxed control and (y^q, z^q) the corresponding trajectory. Then there exists a sequence $(u^n)_n \subset \mathcal{U}$ such that*

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{t \in [0, T]} |y_t^n - y_t^q|^2 \right] = 0, \quad (31)$$

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^T |z_t^n - z_t^q|^2 dt = 0, \quad (32)$$

$$\lim_{n \rightarrow \infty} J(u^n) = \mathcal{J}(q), \quad (33)$$

where (y^n, z^n) denotes the solution of equation (1) associated with u^n .

Proof. We have

$$\begin{aligned} d(y_t^n - y_t^q) &= [b(t, y_t^n, z_t^n, u_t^n) - b(t, y_t^q, z_t^q, u_t^n)] dt \\ &\quad + \left[b(t, y_t^q, z_t^q, u_t^n) - \int_U b(t, y_t^q, z_t^q, a) q_t(da) \right] dt \\ &\quad + (z_t^n - z_t^q) dW_t \end{aligned}$$

Put

$$\begin{aligned} Y_t^n &= y_t^n - y_t^q, \\ Z_t^n &= z_t^n - z_t^q, \end{aligned}$$

and

$$\begin{aligned} \varphi^n(t, Y_t^n, Z_t^n) &= b(t, y_t^q, z_t^q, u_t^n) - \int_U b(t, y_t^q, z_t^q, a) q_t(da) \\ &\quad + \int_0^1 b_y(t, y_t^q + \lambda(y_t^n - y_t^q), z_t^q + \lambda(z_t^n - z_t^q), u_t^n) Y_t^n d\lambda \\ &\quad + \int_0^1 b_z(t, y_t^q + \lambda(y_t^n - y_t^q), z_t^q + \lambda(z_t^n - z_t^q), u_t^n) Z_t^n d\lambda. \end{aligned} \quad (34)$$

Then

$$\begin{cases} dY_t^n = \varphi^n(t, Y_t^n, Z_t^n) dt + Z_t^n dW_t, \\ Y_T^n = 0. \end{cases} \quad (35)$$

The above equation is a linear BSDE with bounded coefficients and with terminal condition $Y_T^n = 0$, then by applying a priori estimates (see Briand et al [8]), we get

$$\mathbb{E} \left[\sup_{t \in [0, T]} |Y_t^n|^2 + \int_0^T |Z_t^n|^2 dt \right] \leq C \mathbb{E} \left| \int_0^T |\varphi^n(t, 0, 0)| dt \right|^2.$$

From (34), we get

$$\begin{aligned} &\mathbb{E} \left[\sup_{t \in [0, T]} |Y_t^n|^2 + \int_0^T |Z_t^n|^2 dt \right] \\ &\leq C \mathbb{E} \int_0^T \left| \int_U b(t, y_t^q, z_t^q, a) \delta_{u_t^n}(da) - \int_U b(t, y_t^q, z_t^q, a) q_t(da) \right|^2 dt. \end{aligned} \quad (36)$$

By (30) and the dominated convergence theorem, the term in the right hand side of the above inequality tends to zero as n tends to infinity. This prove (31) and (32).

Let us prove (33)

Since g and h are Lipschitz continuous in (y, z) , then by using the Cauchy-Schwartz inequality, we have

$$\begin{aligned} & |J(u^n) - \mathcal{J}(q)| \\ & \leq C \left(\mathbb{E} |y_0^n - y_0^q|^2 \right)^{1/2} + C \left(\int_0^T \mathbb{E} |y_t^n - y_t^q|^2 ds \right)^{1/2} + C \left(\mathbb{E} \int_0^T |z_t^n - z_t^q|^2 dt \right)^{1/2} \\ & + C \left(\mathbb{E} \int_0^T \left| \int_U b(t, y_t^q, z_t^q, a) \delta_{u_t^n}(da) - \int_U h(t, y_t^q, z_t^q, a) q_t(da) \right|^2 dt \right)^{1/2}. \end{aligned}$$

From (31) and (32) the first, the second and the third terms in the right hand side converge to zero, and by (30) and the dominated convergence theorem, the fourth term in the right hand side tends to zero. ■

Remark 12 *As a consequence, it is easy to see that the strict and relaxed optimal control problems have the same value function.*

5.2 necessary optimality conditions for near controls

In this section we derive necessary optimality conditions for near optimal controls. This result is based on Ekeland's variational principle which is given by the following.

Lemma 13 (*Ekeland's variational principle*). *Let (E, d) be a complete metric space and $f : E \rightarrow \overline{\mathbb{R}}$ be lower-semicontinuous and bounded from below. Given $\varepsilon > 0$, suppose $u^\varepsilon \in E$ satisfies $f(u^\varepsilon) \leq \inf(f) + \varepsilon$. Then for any $\lambda > 0$, there exists $v \in E$ such that*

1. $f(v) \leq f(u^\varepsilon)$.
2. $d(u^\varepsilon, v) \leq \lambda$.
3. $f(v) < f(w) + \frac{\varepsilon}{\lambda} d(v, w)$, $\forall w \neq v$.

Proof. See Ekeland [10]. ■

To apply Ekeland's variational principle, we have to endow the set \mathcal{U} of strict controls with an appropriate metric. For any $u, v \in \mathcal{U}$, we set

$$d(u, v) = \mathcal{P} \otimes dt \{(\omega, t) \in \Omega \times [0, T], u(t, \omega) \neq v(t, \omega)\},$$

where $\mathcal{P} \otimes dt$ is the product measure of \mathcal{P} with the Lebesgue measure dt .

Let us summarize some of the properties satisfied by d .

Lemma 14 1. (\mathcal{U}, d) is a complete metric space.

2. The cost functional J is continuous from \mathcal{U} into \mathbb{R} .

Proof. See Mezerdi [25]. ■

Now let $\mu \in \mathcal{R}$ be an optimal relaxed control and denote by (y^μ, z^μ) the trajectory of the system controlled by μ . From lemmas 9, 10 and 11, there exists a sequence $(u^n)_n$ of strict controls such that

$$\begin{aligned} dt\mu_t^n(da) &= dt\delta_{u_t^n}(da) \xrightarrow[n \rightarrow \infty]{} dt\mu_t(da) \text{ Stably, } \mathcal{P}\text{-a.s.}, \\ \mathbb{E} \left[\sup_{t \in [0, T]} |y_t^n - y_t^\mu|^2 \right] &\xrightarrow[n \rightarrow \infty]{} 0, \\ \mathbb{E} \int_0^T |z_t^n - z_t^\mu|^2 dt &\xrightarrow[n \rightarrow \infty]{} 0. \end{aligned}$$

where (y_t^n, z_t^n) is the solution of equation (25) controlled by μ^n .

According to the optimality of μ and (29), there exists a sequence $(\varepsilon_n)_n$ of positive real numbers with $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ such that

$$J(u^n) = \mathcal{J}(\mu^n) \leq \mathcal{J}(\mu) + \varepsilon_n.$$

A suitable version of lemma 13 implies that, given any $\varepsilon_n > 0$, there exists $(u^n)_n \in \mathcal{U}$ such that

$$\begin{aligned} J(u^n) &\leq \inf_{u \in \mathcal{U}} J(u) + \varepsilon_n, \\ J(u^n) &\leq J(u) + \varepsilon_n d(u^n, u); \quad \forall u \in \mathcal{U}. \end{aligned} \tag{37}$$

Let us define the perturbation

$$u_t^{n,\theta} = \begin{cases} v & \text{if } t \in [\tau, \tau + \theta], \\ u_t^n & \text{Otherwise.} \end{cases} \quad (38)$$

From (37) we have

$$0 \leq J(u_t^{n,\theta}) - J(u^n) + \varepsilon_n d(u^{n,\theta}, u_t^n).$$

From the definition of the metric d , we obtain

$$0 \leq J(u_t^{n,\theta}) - J(u^n) + \varepsilon_n C\theta. \quad (39)$$

From these above inequalities, we shall establish necessary optimality conditions for near optimal controls.

Theorem 15 (*Necessary optimality conditions for near controls*). *For each $\varepsilon_n > 0$, there exists $(u^n)_n \in \mathcal{U}$ such that there exists a unique adapted processes*

$$p^n \in \mathcal{L}^2([0, T]; \mathbb{R}^n),$$

solution of the following forward stochastic differential equation

$$\begin{cases} -dp_t^n = H_y(t, y_t^n, z_t^n, p_t^n, u_t^n) dt + H_z(t, y_t^n, z_t^n, p_t^n, u_t^n) dW_t, \\ p_0^n = g_y(y_0^n), \end{cases} \quad (40)$$

such that for all $v \in \mathcal{U}$,

$$0 \leq [H(t, y_t^n, z_t^n, p_t^n, u_t^n) - H(t, y_t^n, z_t^n, p_t^n, v)] + C\varepsilon_n. \quad (41)$$

Proof. From inequality (39), we use the same method as in the last sections with index n . ■

5.3 Necessary and sufficient optimality conditions for relaxed controls

In this subsection, we will state and prove necessary as well as sufficient optimality conditions for relaxed controls. For this end, let us summarize and prove some of lemmas that we will use thereafter.

Introduce the following adjoint equation in the relaxed form

$$\begin{cases} -dp_t^\mu = H_y^\mu(t, y_t^\mu, z_t^\mu, p_t^\mu, \mu_t) dt + H_z^\mu(t, y_t^\mu, z_t^\mu, p_t^\mu, \mu_t) dW_t, \\ p_0^\mu = g_y(y_0^\mu), \end{cases} \quad (42)$$

$$p^\mu \in \mathcal{L}^2([0, T]; \mathbb{R}^n),$$

where the Hamiltonian H^μ in the relaxed form is defined from $[0, T] \times \mathbb{R}^n \times \mathcal{M}_{n \times d}(\mathbb{R}) \times \mathbb{R}^n \times \mathbb{P}(U)$ into \mathbb{R} by

$$H^\mu(t, y_t^\mu, z_t^\mu, p_t^\mu, \mu_t) = p_t^\mu \int_U b(t, y_t^\mu, z_t^\mu, a) \mu_t(a) - \int_U h(t, y_t^\mu, z_t^\mu, a) \mu_t(a).$$

For simplicity of notation, we denote

$$\begin{aligned} f^n(t) &= f(t, y_t^n, z_t^n, u_t^n), \\ f^\mu(t) &= \int_U f(t, y_t^\mu, z_t^\mu, a) \mu_t(a), \end{aligned}$$

where f stands for one of the functions b_y, b_z, h_y, h_z .

Lemma 16 *The following estimations hold*

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |b_y^n(s) - b_y^\mu(s)|^2 ds = 0, \quad (43)$$

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |b_z^n(s) - b_z^\mu(s)|^2 ds = 0, \quad (44)$$

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |h_y^n(s) - h_y^\mu(s)|^2 ds = 0, \quad (45)$$

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |h_z^n(s) - h_z^\mu(s)|^2 ds = 0. \quad (46)$$

Proof. We have

$$\begin{aligned} \mathbb{E} \int_0^t |b_y^n(s) - b_y^\mu(s)|^2 ds &= \mathbb{E} \int_0^t \left| b_y(s, y_s^n, z_s^n, u_s^n) - \int_U b_y(s, y_s^\mu, z_s^\mu, a) \mu_s(a) \right|^2 ds \\ &\leq \mathbb{E} \int_0^t |b_y(s, y_s^n, z_s^n, u_s^n) - b_y(s, y_s^\mu, z_s^\mu, u_s^n)|^2 ds \\ &\quad + \mathbb{E} \int_0^t |b_y(s, y_s^\mu, z_s^\mu, u_s^n) - b_y(s, y_s^\mu, z_s^\mu, u_s^\mu)|^2 ds \\ &\quad + \mathbb{E} \int_0^t \left| b_y(s, y_s^\mu, z_s^\mu, u_s^\mu) - \int_U b_y(s, y_s^\mu, z_s^\mu, a) \mu_s(a) \right|^2 ds. \end{aligned}$$

Since b_y is Lipschitz continuous in z , then

$$\begin{aligned} \mathbb{E} \int_0^t |b_y^n(s) - b_y^\mu(s)|^2 ds &\leq \mathbb{E} \int_0^t |b_y(s, y_s^n, z_s^n, u_s^n) - b_y(s, y_s^\mu, z_s^\mu, u_s^\mu)|^2 ds \\ &\quad + C \mathbb{E} \int_0^t |z_s^n - z_s^\mu|^2 ds \\ &\quad + \mathbb{E} \int_0^t \left| b_y(s, y_s^\mu, z_s^\mu, u_s^n) - \int_U b_y(s, y_s^\mu, z_s^\mu, a) \mu_s(a) \right|^2 ds. \end{aligned} \quad (47)$$

From (32), we have

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |z_s^n - z_s^\mu|^2 ds = 0.$$

Since b_y is bounded and continuous, then by (31) and the dominate convergence theorem, we have

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |b_y(s, y_s^n, z_s^n, u_s^n) - b_y(s, y_s^\mu, z_s^\mu, u_s^n)|^2 ds = 0.$$

On the other hand, by the chattering lemma and the dominate convergence theorem, we have

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t \left| \int_U b_y(s, y_s^\mu, z_s^\mu, a) \delta_{u_s^n}(da) - \int_U b_y(s, y_s^\mu, z_s^\mu, a) \mu_s(a) \right|^2 ds = 0.$$

By (47) and these above three limits, we deduce (43). Using the same method and arguments, we prove (44), (45) and (46). ■

Lemma 17 *Let p^n and p^μ respectively the solutions of (40) and (42), then we have*

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{t \in [0, T]} |p_t^n - p_t^\mu|^2 \right] = 0. \quad (48)$$

Proof. From (40) and (42), we have

$$\begin{aligned} p_t^n &= g_y(y_0^n) - \int_0^t H_y^n(s) ds - \int_0^t H_z^n(s) dW_s, \\ p_t^\mu &= g_y(y_0^\mu) - \int_0^t H_y^\mu(s) ds - \int_0^t H_z^\mu(s) dW_s, \end{aligned}$$

where

$$\begin{aligned} H_y^n(t) &= H_y(t, y_t^n, z_t^n, p_t^n, u_t^n) \quad ; \quad H_y^\mu(t) = \int_U H_y(t, y_t^\mu, z_t^\mu, p_t^\mu, a) \mu_t(a), \\ H_z^n(t) &= H_z(t, y_t^n, z_t^n, p_t^n, u_t^n) \quad ; \quad H_z^\mu(t) = \int_U H_z(t, y_t^\mu, z_t^\mu, p_t^\mu, a) \mu_t(a). \end{aligned}$$

Then

$$\begin{aligned} \mathbb{E} |p_t^n - p_t^\mu|^2 &\leq C \mathbb{E} |g_y(y_0^n) - g_y(y_0^\mu)|^2 + C \mathbb{E} \int_0^t |H_y^n(s) - H_y^\mu(s)|^2 ds \\ &\quad + C \mathbb{E} \int_0^t |H_z^n(s) - H_z^\mu(s)|^2 ds \\ &\leq C \mathbb{E} \int_0^t |b_y^n(s) (p_s^n - p_s^\mu)|^2 ds + C \mathbb{E} \int_0^t |b_z^n(s) (p_s^n - p_s^\mu)|^2 ds + C \alpha_t^n, \end{aligned}$$

where

$$\begin{aligned} \alpha_t^n &= \mathbb{E} |g_y(y_0^n) - g_y(y_0^\mu)|^2 + \mathbb{E} \int_0^t |h_y^n(s) - h_y^\mu(s)|^2 ds \quad (49) \\ &\quad + \mathbb{E} \int_0^t |(b_y^n(s) - b_y^\mu(s)) p_s^\mu|^2 ds + \mathbb{E} \int_0^t |h_z^n(s) - h_z^\mu(s)|^2 ds \\ &\quad + \mathbb{E} \int_0^t |(b_z^n(s) - b_z^\mu(s)) p_s^\mu|^2 ds. \end{aligned}$$

Since b_y and b_z are bounded then

$$\mathbb{E} |p_t^n - p_t^\mu|^2 \leq 2C \mathbb{E} \int_0^t |p_s^n - p_s^\mu|^2 ds + C \alpha_t^n. \quad (50)$$

Let us prove that $\lim_{n \rightarrow \infty} \alpha_t^n = 0$

Since g_y is bounded and continuous, then by (31) and the dominated convergence theorem, we have

$$\lim_{n \rightarrow \infty} \mathbb{E} |g_y(y_0^n) - g_y(y_0^\mu)|^2 = 0. \quad (51)$$

On the other hand, since b_y is bounded, then

$$|[b_y^n(s) - b_y^\mu(s)] p_s^n| \leq 2C |p_s^n|. \quad (52)$$

Hence by the Cauchy-Schwartz inequality we get,

$$\mathbb{E} \int_0^t |[b_y^n(s) - b_y^\mu(s)] p_s^\mu| ds \leq \left(\mathbb{E} \int_0^t |b_y^n(s) - b_y^\mu(s)|^2 ds \right)^{1/2} \left(\mathbb{E} \int_0^t |p_s^\mu|^2 ds \right)^{1/2}.$$

Since $p^\mu \in \mathcal{L}^2([0, T]; \mathbb{R}^n)$, then

$$\mathbb{E} \int_0^t |[b_y^n(s) - b_y^\mu(s)] p_s^\mu| ds \leq C \left(\mathbb{E} \int_0^t |b_y^n(s) - b_y^\mu(s)|^2 ds \right)^{1/2}.$$

By (43), we have

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |b_y^n(s) - b_y^\mu(s)|^2 ds = 0.$$

Then, we deduce that

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |[b_y^n(s) - b_y^\mu(s)] p_s^\mu| ds = 0. \quad (53)$$

By using the dominated convergence theorem we obtain

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |[b_y^n(s) - b_y^\mu(s)] p_s^\mu|^2 ds = 0. \quad (54)$$

Similarly, using (44), the boundness of b_z and the dominated convergence theorem, it follows that

$$\lim_{n \rightarrow \infty} \mathbb{E} \int_0^t |[b_z^n(s) - b_z^\mu(s)] p_s^\mu|^2 ds = 0. \quad (55)$$

From (45), (46), (51), (54) and (55), it is easy to see that

$$\lim_{n \rightarrow \infty} \alpha_t^n = 0. \quad (56)$$

Finally from (50), (56), Gronwall's lemma and Burkholder-Davis-Gundy inequality, we have the desired result. ■

Theorem 18 (*Necessary optimality conditions for relaxed controls*). Let μ be an optimal relaxed control minimizing the cost \mathcal{J} over \mathcal{R} and (y_t^μ, z_t^μ) the corresponding optimal trajectory. Then there exists a unique adapted processes

$$p^\mu \in \mathcal{L}^2([0, T]; \mathbb{R}^n),$$

solution of the stochastic forward differential equation (42), such that for all $q \in \mathcal{R}$, we have

$$H^\mu(t, y_t^\mu, z_t^\mu, p_t^\mu, \mu_t) = \max_{q \in \mathbb{P}(U)} H^\mu(t, y_t^\mu, z_t^\mu, p_t^\mu, q). \quad (57)$$

Proof. Let μ be an optimal relaxed control. By the necessary condition for near controls (Theorem 15), there exists a sequence $(u^n)_n \subset \mathcal{U}$ such that for all $v \in \mathcal{U}$

$$0 \leq [H(t, y_t^n, z_t^n, p_t^n, u_t^n) - H(t, y_t^n, z_t^n, p_t^n, v)] + C\varepsilon_n,$$

where $\lim_{n \rightarrow \infty} \varepsilon_n = 0$.

According to (29), (31), (31) and (48), the result follows immediately by letting n going to infinity in the last inequality. ■

Remark 19 If $\mu_t(da) = \delta_{u(t)}(da)$, we recover the strict necessary optimality conditions (Theorem 4).

Theorem 20 (*Sufficient optimality conditions for relaxed controls*). We know that the set \mathcal{R} of relaxed controls is convex and the function $H^q(t, y_t^q, z_t^q, p_t^q, q_t)$ is linear in q_t . If we assume that for every $q \in \mathcal{R}$ and for all $t \in [0, T]$, the functions g is convex and $(y_t^q, z_t^q) \rightarrow H^q(t, y_t^q, z_t^q, p_t^q, q_t)$ is concave, then μ is an optimal relaxed control if it satisfies (57).

Proof. The proof is the same that in theorem 5. ■

References

- [1] F. Antonelli, *Backward-forward stochastic differential equations*, Annals of Applied Probability, 1993, 3, pp. 777–793.
- [2] S. Bahlali, B. Mezerdi and B. Djehiche, *Approximation and optimality necessary conditions in relaxed stochastic control problems*, Journal of Applied Mathematics and Stochastic Analysis, Volume 2006, pp 1-23.

- [3] S. Bahlali and B. Labeled, *Necessary and sufficient conditions of optimality for optimal control problem with initial and terminal costs*, Random Operators and Stochastic Equations, 2006, Vol 14, No3, pp 291-301.
- [4] S. Bahlali, B. Djehiche and B. Mezerdi, *The relaxed maximum principle in singular control of diffusions*, SIAM J. Control and Optim, 2007, Vol 46, Issue 2, pp 427-444.
- [5] S. Bahlali, *Necessary and sufficient conditions of optimality for relaxed and strict control problems*, SIAM J. Control and Optim, 2008, Vol. 47, No. 4, pp. 2078–2095.
- [6] S. Bahlali, *Necessary and sufficient condition of optimality for optimal control problem of forward and backward systems*, Probability Theory and It's Application. In revision.
- [7] A. Bensoussan, *Lecture on stochastic control. in non linear filtering and stochastic control*, Lecture notes in mathematics, 972. Proc. Cortona, Springer Verlag, 1981.
- [8] Ph. Briand, B. Delyon, Y. Hu, E. Pardoux and L. Stoica, *L^p Solutions of backward stochastic differential equations*, Sochastic Process and their Applications, No 108 (2003), pp 109-129.
- [9] N. Dokuchaev and X.Y. Zhou, *Stochastic controls with terminal contingent conditions*, Journal Of Mathematical Analysis And Applications, 1999, 238, pp 143-165.
- [10] I. Ekeland, *On the variational principle*, J. Math. Anal. Appl, 1974, Vol. 47, pp 324-353.
- [11] I. Ekeland and R. Temam, *Analyse convexe et problème variationnel*, Dunod. 1974.
- [12] N. El Karoui, N. Huu Nguyen and N. Jeanblanc Piqué, *Compactification methods in the control of degenerate diffusions*, Stochastics, 1987, Vol. 20, pp 169-219.
- [13] N. El Karoui and M. Mazliak, *Backward stochastic differential equations*, Addison Wesley, Longman, 1997.

- [14] N. El-Karoui, S. Peng and M.C. Quenez, *Backward stochastic differential equations in finance*, Math. finance 7, 1997.
- [15] N. El-Karoui, S. Peng and M.C. Quenez, *A dynamic maximum principle for the optimization of recursive utilities under constraints*, Annals of Applied Probability, 11(2001), pp 664-693.
- [16] W.H. Fleming, *Generalized solutions in optimal stochastic control*, Differential games and control theory 2, (Kingston conference 1976), Lect. Notes in Pure and Appl. Math.30, 1978.
- [17] N.F. Framstad, B. Oksendal and A. Sulem, *A sufficient stochastic maximum principle for optimal control of jump diffusions and applications to finance*, J. Optim. Theory and applications, 2004, 121, pp 77-98.
- [18] M. Fuhrman and G. Tessitore, *Existence of optimal stochastic controls and global solutions of forward-backward stochastic differential equations*, SIAM Jour.Cont. Optim, 2004, Vol 43, N° 3, pp 813-830.
- [19] U.G. Haussmann, *General necessary conditions for optimal control of stochastic systems*, Math. Programming Studies 6, 1976, pp 30-48.
- [20] U.G. Haussmann, *A Stochastic maximum principle for optimal control of diffusions*, Pitman Research Notes in Math, 1986, Series 151.
- [21] J. Jacod and J. Mémin, *Sur un type de convergence intermédiaire entre la convergence en loi et la convergence en probabilité*, Sem. Proba.XV, Lect. Notes in Math, 851, 1980, Springer Verlag.
- [22] S.Ji and X. Y. Zhou, *A maximum principle for stochastic optimal control with terminal state constraints, and its applications*. Commun. Inf. Syst, 2006, 6(4), pp 321-338.
- [23] H.J. Kushner, *Necessary conditions for continuous parameter stochastic optimization problems*, SIAM J. Control Optim, Vol. 10, 1973, pp 550-565.
- [24] J. Ma and J. Zhang, *Representation theorems for backward stochastic differential equations*, Ann. Appl. Probab, 2002, 12(4) :1390-1418.

- [25] B. Mezerdi, *Necessary conditions for optimality for a diffusion with a non smooth drift*, Stochastics And Stoch. Reports, 1988, Vol. 24, pp 305-326.
- [26] B. Mezerdi and S. Bahlali, *Approximation in optimal control of diffusion processes*, Rand. Operat. and Stoch. Equ, 2000, Vol.8, No 4, pp 365-372.
- [27] B. Mezerdi and S. Bahlali, *Necessary conditions for optimality in relaxed stochastic control problems*, Stochastics And Stoch. Reports, 2002, Vol 73 (3-4), pp 201-218.
- [28] E. Pardoux and S. Peng, *Adapted solutions of backward stochastic differential equations*, Sys. Control Letters, 1990, Vol. 14, pp 55-61.
- [29] S. Peng, *A general stochastic maximum principle for optimal control problems*, SIAM Jour.Cont. Optim, 1990, 28, N° 4, pp 966-979.
- [30] S. Peng, *Backward stochastic differential equations and application to optimal control*, Appl. Math. Optim, 1993, 27, pp 125-144.
- [31] S. Peng and Z.Wu, *Fully coupled forward-backward stochastic differential equations and applications to optimal control*, SIAM J. Control Optim, 1999, 37, no. 3, pp. 825–843.
- [32] J. T. Shi and Z. Wu, *The maximum principle for fully coupled forward-backward stochastic control system*, Acta Automatica Sinica, Vol 32, No 2, 2006, pp 161-169.
- [33] Z. Wu, *Maximum Principle for Optimal Control Problem of Fully Coupled Forward-Backward Stochastic Systems*, Systems Sci. Math. Sci, A998, 11, No.3, pp 249-259.
- [34] W. Xu, *Stochastic maximum principle for optimal control problem of forward and backward system*, J. Austral. Math. Soc. Ser. B 37, 1995, pp 172-185.
- [35] J. Yong and X.Y. Zhou, *Stochastic controls : Hamilton systems and HJB equations*, vol 43, Springer, New York, 1999.
- [36] X.Y. Zhou, *Sufficient conditions of optimality for stochastic systems with controllable diffusions*. IEEE Trans. on Automatic Control, 1996, 41, pp 1176-1179.