

The Relativistic Stern-Gerlach Force

C. Tschalaer

1. Introduction

For over a decade, speculation on the Stern Gerlach (SG) force acting on a particle with spin moving at a relativistic velocity in an electromagnetic field have been put forward and experiments proposed. Using the Hamiltonian-Lagrangian framework in close analogy to the well-known derivation of the Lorentz Force [1], the SG interaction including the effect of the Thomas Precession is derived from the well-established non-relativistic SG force.

In the true rest frame of a particle of charge e , mass M , spin \vec{s} , and magnetic moment

$$\vec{\mu} = (e/M)(g/2)\vec{s} \quad (1)$$

where g is the gyromagnetic factor, the Hamiltonian H_{rest}^* of the particle is the sum of its kinetic and potential energy, i.e. the total energy

$$H_{rest}^* = Mc^2 + S^* + e\phi^*.$$

ϕ^* is the electromagnetic field potential, and S^* the SG interaction potential in a magnetic field \vec{B}^* :

$$S^* = -\vec{\mu} \cdot \vec{B}^*. \quad (2)$$

2. Thomas precession

If the particle is accelerated by an external force perpendicular to its velocity $\vec{\beta}c$, the true rest frame of the particle rotates with respect to the laboratory frame by the well-established Thomas precession frequency $\vec{\omega}_T^*$ (see ref. 1)

$$\vec{\omega}_T^* = (e/M)(\gamma/(\gamma+1))(\vec{\beta} \times \vec{E}^*/c) \quad (3)$$

with

$$1/\gamma \equiv \sqrt{1 - \vec{\beta}^2}$$

if the external force is the Lorentz force ($=e\vec{E}^*$ in the rest frame). In preparation for a Lorentz transformation to the laboratory frame, we first transform from the true rest frame of the particle to a rest frame rotating with $-\vec{\omega}_r^*$. While the particle's mass and the electromagnetic potential do not change, the rotating spin creates an additional potential

$$T^* = -\vec{s} \cdot \vec{\omega}_r^* \quad (4)$$

which is added to H_{rest}^* to form the rotating rest frame Hamiltonian

$$H_{rot}^* = Mc^2 + e\phi^* + G^* \quad (5)$$

where G^* is the sum of S^* and T^* .

3. Canonical Momentum

Since the rotating rest frame RF is an accelerated frame, we define an instantaneous rest frame IRF which is moving with respect to the laboratory frame LF with a constant velocity $c\vec{\beta}_0$ which for simplicity we assume to be in the z -direction. The particle is moving with velocity $c\vec{\beta}^*$ in the IRF where $\vec{\beta}^*$ is zero at time t_0 for which we want to determine the forces on the particle. At times near t_0 , the velocity $c\vec{\beta}^*$ is small and we can obtain the Lagrangian L_0^* from the classical definition that L_0^* is $2K^* - E^*$ where E^* is the total energy in the RF i.e. H_{rot}^* , and K^* is the kinetic energy i.e. $v^2/2$ times the total mass including the contributions of the electromagnetic and Stern-Gerlach-Thomas potentials ϕ^* and G^* . Thus

$$\begin{aligned} L_0^* &= 2K^* - E^* = \vec{\beta}^{*2} H_{rot}^* - \gamma^* H_{rot}^* \\ &= -(1/\gamma^*) (Mc^2 + e\phi^* + G^*) \\ &= -(Mc^2 + G^*)/\gamma^* - e(\gamma^* \phi_0^* - \gamma^* \vec{\beta}^* \cdot \vec{A}_0^*)/\gamma^* \end{aligned}$$

or

$$L_0^* = -(Mc^2 + G^*)/\gamma^* - e\phi_0^* + e\vec{\beta}^* \cdot \vec{A}_0^* \quad (6)$$

where ϕ_0^* and \vec{A}_0^* are the electromagnetic potential and vector potential in the IRF.

The canonical momentum \vec{P}_0^* conjugate to the position coordinate \vec{x}^* is classically defined [1] as

$$c\vec{P}_0^* = \partial L_0^* / \partial \vec{\beta}^* = \gamma^* \vec{\beta}^* Mc^2 + e\vec{A}_0^* - \partial(G^*/\gamma^*) / \partial \vec{\beta}^*.$$

The last term arises from the fact that the Stern-Gerlach-Thomas potential G^* is velocity dependent.

In preparation for the transformation to the LF, we deduce the Hamiltonian H_0^* or total energy in the IRF from the relation

$$H_0^* = c\bar{P}_0^* \bar{\beta}^* - L_0^* = \gamma^* M c^2 - \bar{\beta}^* \partial(G^* / \gamma^*) / \partial \bar{\beta}^* + e\phi_0^* + G^* / \gamma^*$$

or

$$H_0^* = \gamma^* M c^2 + e\phi_0^* - \bar{\beta}^* \bar{\Gamma}^* + G^* / \gamma^*$$

and

$$c\bar{P}_0^* = \gamma^* \bar{\beta}^* M c^2 + e\bar{A}_0^* - \bar{\Gamma}^* \quad (7)$$

where $\bar{\Gamma}^*$ is defined as

$$\bar{\Gamma}^* \equiv \partial(G^* / \gamma^*) / \partial \bar{\beta}^*.$$

We can now form the energy-momentum 4-vector P_v^* in the IRF:

$$P_v^* = (\bar{P}_0^*; iH_0^* / c)$$

which may be represented as a sum of three 4-vectors representing the free-particle momentum p_v^* , the electromagnetic field A_v^* , and the Stern-Gerlach-Thomas potential G_v^* respectively, where

$$\begin{aligned} p_v^* &= (\gamma^* \bar{\beta}^* M c; i\gamma^* M c); & p_v^* p_v^* &= -M^2 c^2 \\ A_v^* &= (\bar{A}_0^*; i\phi_0^*); & A_v^* A_v^* &= \bar{A}_0^{*2} - \phi_0^{*2} \\ G_v^* &= (\bar{G}_0^*; iG_0^*); & G_v^* G_v^* &= \bar{\Gamma}^{*2} - G^{*2}(t_0) \end{aligned}$$

and

$$cP_v^* = cp_v^* + eA_v^* + G_v^*$$

with

$$\begin{aligned} \bar{G}_0^* &= -\bar{\Gamma}^* \\ G_0^* &= G^* / \gamma^* - \bar{\beta}^* \bar{\Gamma}^* = G^*(t_0) + \text{order}(\beta^{*2}). \end{aligned} \quad (8)$$

The IRF Hamiltonian can then be expressed in terms of the 4-vector components as

$$H_0^* = e\phi_0^* + G_0^* + \sqrt{M^2 c^4 + (c\bar{P}_0^* - e\bar{A}_0^* - \bar{G}_0^*)^2}.$$

The equation of motion in the IRF is then given by the Lagrange equation

$$d(\partial L_0^* / \partial \vec{\beta}^*) / dt^* = cd\vec{P}_0^* / dt^* = c\vec{V}^* L_0^*.$$

Using eqs. (6) and (7) we find

$$d(\gamma^* \vec{\beta}^* Mc) / dt^* = e\vec{E}_0^* + e(\vec{\beta}^* \times \vec{B}_0^*) - (1/\gamma^*) \vec{V}^* G^* + d\vec{\Gamma}^* / dt^* \quad (9)$$

4. Lorentz Transformation

Having established the energy-momentum 4-vector for a spinning particle in the IRF moving with velocity $c\vec{\beta}_0$ with respect to the LF, we can now perform the final transformation to the LF by a Lorentz rotation of P_ν^* to the laboratory vector P_ν using the Lorentz rotation matrix $a_{\nu\mu}$

$$P_\nu = a_{\nu\mu} P_\mu^* = (\vec{P}; iH/c)$$

to obtain the LF Hamiltonian H and momentum \vec{P} .

Since P_μ^* is a sum of three 4-vectors, we can rotate each of these component 4-vectors separately using the formulae for velocity addition in appendix A1 to obtain

$$\begin{aligned} p_\nu &= (\gamma\vec{\beta}Mc; i\gamma Mc) \\ A_\nu &= (\vec{A}; i\phi) \\ G_\nu &= (\vec{G}; iG) \end{aligned} \quad (10)$$

where

$$\begin{aligned} G &= \gamma_0 G_0^* + \gamma_0 \beta_0 \vec{G}_{0z}^* \\ \vec{G}_z &= \gamma_0 \vec{G}_{0z}^* + \gamma_0 \beta_0 G_0^*; \quad \vec{G}_y = \vec{G}_{0y}^*; \quad \vec{G}_x = \vec{G}_{0x}^*. \end{aligned}$$

and,

$$\vec{G} = (-\Gamma_x^*; -\Gamma_y^*; -\gamma_0 \Gamma_z^* - \gamma_0 \beta_0 \vec{\beta}^* \vec{\Gamma}^* + \gamma_0 \beta_0 G^* / \gamma^*) \quad (11)$$

$$G = \gamma_0 G^* / \gamma^* - \gamma_0 \vec{\beta}^* \vec{\Gamma}^* - \gamma_0 \beta_0 \Gamma_z^*.$$

Therefore, $\vec{P} = \gamma\vec{\beta}Mc + (e/c)\vec{A} + (1/c)\vec{G}$

and $H = \gamma Mc^2 + e\phi + G$

or $H = e\phi + G + \sqrt{M^2 c^4 + (c\bar{P} - e\bar{A} - \bar{G})^2}$.

5. Lagrangian

It has been shown in appendix A2 that the Lorentz transformation of the energy-momentum 4-vector conserves the canonical relationships such that \bar{P} is again the canonical conjugate momentum to \bar{x} of H . Therefore, we can derive the Lagrangian L for the spinning particle in the LF

$$L = \bar{P} \cdot \bar{\beta} c - H.$$

Using eq. (10), we find

$$\begin{aligned} L &= \gamma \beta^2 Mc^2 + e\bar{\beta}\bar{A} + \bar{\beta}\bar{G} - \gamma Mc^2 - e\phi - G \\ &= -Mc^2 / \gamma + e\bar{\beta}\bar{A} - e\phi + \bar{\beta}\bar{G} - G. \end{aligned}$$

Here, from eq. (11) and appendix A1

$$\begin{aligned} \bar{\beta}\bar{G} - G &= (\gamma_0 \beta_0 \beta_z - \gamma_0) G^* / \gamma^* + (\gamma_0 - \gamma_0 \beta_0 \beta_z) \bar{\beta}^* \bar{\Gamma}^* \\ &\quad - \beta_x \Gamma_x^* - \beta_y \Gamma_y^* - \gamma_0 \beta_z \Gamma_z^* + \gamma_0 \beta_0 \Gamma_z^* \\ &= -G^* / \gamma + (\gamma^* - 1) (\beta_x \Gamma_x^* + \beta_y \Gamma_y^*). \end{aligned}$$

The last term is of order β^{*3} and can be neglected as β^* goes to zero so that

$$L = -(Mc^2 + G^*) / \gamma + e\bar{\beta}\bar{A} - e\phi. \quad (12)$$

From eqs. (1) to (4), the Stern-Gerlach-Thomas part of the Lagrangian is

$$-(1/\gamma)G^* = -(1/\gamma)(S^* + T^*) = (e/M)\bar{s} \cdot \left[(g/2)\bar{B}^* / \gamma + (1/(\gamma+1))\bar{\beta} \times \bar{E}^* / c \right]$$

which in terms of the LF fields becomes

$$L_{SGT} = (e/M)\bar{s} \left[(g/2 - \gamma\beta^2 / (\gamma+1))\bar{B}_\perp + (g/2)\bar{B}_\parallel / \gamma - (g/2 - \gamma / (\gamma+1))\bar{\beta} \times \bar{E} / c \right]$$

where \vec{B}_\perp and \vec{B}_\parallel are the \vec{B} -field components perpendicular and parallel to \vec{B} . L_{SGT} can readily be identified with the well-known spin motion Lagrangian [2].

The partial derivative of L with respect to \vec{v} is

$$\partial L / \partial (\vec{\beta}c) = \vec{P}$$

because of the canonical relation

$$\partial H / \partial \vec{P} = \vec{\beta}c.$$

Therefore the Lagrange equation of motion in the LF is

$$d(\partial L / \partial \vec{v}) / dt = d\vec{P} / dt = \vec{\nabla}L.$$

Explicitly, using eq. (8), we obtain

$$d(\gamma\vec{\beta}Mc) / dt = e\vec{E} + e(\vec{\beta} \times \vec{B}) - (1/\gamma)\vec{\nabla}G^* - (1/c)d\vec{G} / dt. \quad (13)$$

Using relations of appendix A1, and the vector $\vec{\Gamma}$ defined as

$$\vec{\Gamma} = \partial(G^* / \gamma^*) / \partial \vec{\beta} \quad (14)$$

we find

$$\begin{aligned} \Gamma_{x,y}^* &= \Gamma_{x,y} / \gamma \\ \Gamma_z^* &= \Gamma_z / \gamma^2 \end{aligned}$$

and can rewrite \vec{G} of eq. (11) as

$$\begin{aligned} \vec{G}_{x,y} &= -\Gamma_{x,y} / \gamma \\ \vec{G}_z &= -\gamma_0 \left[\Gamma_z / \gamma^2 + \beta_0 \vec{\beta}^* \vec{\Gamma}^* - \beta_0 G^* / \gamma^* \right]. \end{aligned}$$

Since

$$\vec{\beta}^* \vec{\Gamma}^* = \beta_x \Gamma_x + \beta_y \Gamma_y + (\Gamma_z / \gamma^2)(\beta_z - \beta_0) / (1 - \beta_0 \beta_z)$$

we find

$$\begin{aligned} \beta_0 \vec{\beta}^* \vec{\Gamma}^* + \Gamma_z / \gamma^2 &= (\Gamma_z / \gamma^2)(1 - \beta_0^2) / (1 - \beta_0 \beta_z) \\ &= \Gamma_z / (\gamma \gamma_0 \gamma^*) \end{aligned}$$

and therefore

$$\bar{G}_z = -\Gamma_z / \gamma - \gamma_0 \beta_0 (\beta_x \Gamma_x + \beta_y \Gamma_y) + \gamma_0 \beta_0 G^* / \gamma^*$$

or

$$\bar{G} = -\bar{\Gamma} / \gamma + \gamma_0 \bar{\beta}_0 (G^* / \gamma^* - \beta_x \Gamma_x - \beta_y \Gamma_y). \quad (15)$$

Therefore, the equation for the LF acceleration of a spinning particle in an electromagnetic field is

$$d(\gamma \bar{\beta} M c) / dt = e \bar{E} + e(\bar{\beta} + \bar{B}) - (1/\gamma) \bar{\nabla} G^* - d(\bar{\Gamma} / \gamma) / dt + \gamma_0 \bar{\beta}_0 d(G^* / \gamma^* - \beta_x \Gamma_x - \beta_y \Gamma_y) / dt. \quad (16)$$

6. Conclusion

Starting with the total energy or Hamiltonian for a spinning charged particle in a rotating rest frame (RF) (eq. (5)) based on the classical Stern-Gerlach potential S^* (eqs. (1) and (2)) and the Thomas precession potential T^* (eqs. (3) and (4)) in this frame, we have established the classical Lagrangian L_0^* (eq. (6)) and the canonical momentum \bar{P}_0^* conjugate to the position coordinate \bar{x}_0^* (eq.(7)) in the instantaneous rest frame (IRF). This yields the correct equation of motion (eq. 9)) in the IRF, as well as the corresponding Hamiltonian H_0^* which together with \bar{P}_0^* forms the energy-momentum 4-vector P_v^* in the IRF. Transformation of P_v^* into the laboratory frame LF provides the LF 4-vector P_v from which the LF Lagrangian L (eq. 12)) is obtained which provides the LF equations of motion in eqs. (13)-(16).

From these equations, we can identify the Stern-Gerlach force \bar{F}_{SG} and the Thomas force \bar{F}_T defined as the corresponding acceleration times the rest mass:

$$\bar{F}_{SG} = -(1/\gamma) \bar{\nabla} S^* - (1/c) d(\bar{\sigma} / \gamma) / dt + (\gamma_0 \bar{\beta}_0 / c) d(S^* / \gamma^* - \beta_x \sigma_x - \beta_y \sigma_y) / dt$$

and

$$\bar{F}_T = -(1/\gamma) \bar{\nabla} T^* - (1/c) d(\bar{\tau} / \gamma) / dt + (\gamma_0 \bar{\beta}_0 / c) d(T^* / \gamma^* - \beta_x \tau_x - \beta_y \tau_y) / dt. \quad (17)$$

Here we have defined

$$\bar{\sigma} \equiv \partial(S^* / \gamma^*) / \partial \bar{\beta}; \quad \bar{\tau} \equiv \partial(T^* / \gamma^*) / \partial \bar{\beta}$$

and the S^* and T^* in terms of the LF fields are

$$S^* = -(e/M)(g/2)\bar{s}(\gamma\vec{B}_\perp + \vec{B}_\parallel - \gamma\vec{\beta} \times \vec{E}/c);$$

$$T^* = (e/M)(\gamma-1)\bar{s}(\vec{B}_\perp - (1/\beta^2)\vec{\beta} \times \vec{E}/c)$$

where again \vec{B}_\perp and \vec{B}_\parallel are the \vec{B} components perpendicular and parallel to $\vec{\beta}$ respectively.

Note that the last terms in eq. (17) do not contribute to the effective force i.e. the total momentum change of a spinning particle traversing a localized field region (zero field outside the region) since these terms are total time differentials.

A1 Relations Between Rest Frame and Laboratory Frame Velocities

Starting with the relativistic addition theorem for the IRF velocity $c\bar{\beta}_0$ (chosen in the z -direction), the velocity $c\bar{\beta}^*$ of the RF in the IRF and the velocity $c\bar{\beta}$ of the RF in the LF

$$\beta_z = (\beta_0 + \beta_z^*) / (1 + \beta_0 \beta_z^*)$$

$$\beta_x = \beta_x^* / \gamma; \beta_y = \beta_y^* / \gamma$$

we find

$$(1/\gamma^2) = 1 - \beta_x^2 - \beta_y^2 - \beta_z^2 = 1 - (\beta_x^{*2} + \beta_y^{*2}) / \gamma^2 - (\beta_0 + \beta_z^*)^2 / (1 + \beta_0 \beta_z^*)^2$$

$$= \left[\gamma^2 (1 - \beta_0^2) (1 - \beta_z^{*2}) - (\beta_x^{*2} + \beta_y^{*2}) (1 + \beta_0 \beta_z^*)^2 \right] / \left[\gamma^2 (1 + \beta_0 \beta_z^*)^2 \right]$$

or

$$\gamma = \gamma_0 (1 + \beta_0 \beta_z^*) \sqrt{1 + \beta_x^{*2} + \beta_y^{*2}} / \sqrt{1 - \beta_z^{*2}}.$$

For $\beta_x^* = \beta_y^* = 0$, this becomes

$$\gamma = \gamma^* \gamma_0 (1 + \beta_0 \beta_z^*)$$

and

$$\gamma \beta_z = \gamma^* \gamma_0 (\beta_0 + \beta_z^*).$$

The partial derivatives of γ and β with respect to $\bar{\beta}^*$ are then

$$\partial \gamma / \partial \beta_{x,y}^* = \gamma \beta_{x,y}^* / (1 + \beta_x^{*2} + \beta_y^{*2}) = \gamma \beta_{x,y}^* \text{ for } \beta_x^* = \beta_y^* = 0$$

$$\partial \gamma / \partial \beta_z^* = \gamma_0 \sqrt{1 + \beta_x^{*2} + \beta_y^{*2}} (\beta_0 - \beta_0 \beta_z^{*2} + \beta_z^* + \beta_z^{*2} \beta_0) / \sqrt{1 - \beta_z^{*2}}^3$$

$$= \gamma \beta_z^* / (1 - \beta_z^{*2}) = \gamma^{*2} \gamma \beta_z^* \text{ for } \beta_x^* = \beta_y^* = 0$$

$$\partial \beta_z / \partial \beta_z^* = \gamma_0^{-2} (1 + \beta_0 \beta_z^*)^{-2} = \gamma^{*2} / \gamma^2 \text{ for } \beta_x^* = \beta_y^* = 0.$$

The partial derivatives among γ and $\bar{\beta}$ inside a given frame are

$$\partial \gamma / \partial \bar{\beta} = \partial (1 - \beta_x^2 - \beta_y^2 - \beta_z^2)^{-1/2} / \partial \bar{\beta} = \gamma^3 \bar{\beta}$$

$$\partial (\gamma \beta_x) / \partial \beta_x = \beta_x^2 \gamma^3 + \gamma = \gamma \text{ for } \beta_x = \beta_y = 0$$

$$\partial (\gamma \beta_z) / \partial \beta_x = \beta_x \beta_z \gamma^3$$

$$\partial (\gamma \beta_z) / \partial \beta_z = \beta_z^2 \gamma^2 + \gamma = \gamma^3 \text{ for } \beta_x = \beta_y = 0.$$

A2 Lorentz Transform of Canonical Variables

Assume that in a given frame the Hamiltonian of a system depends on the position variable z^* , its canonical momentum p^* and time t^* . Then the Hamilton relations hold

$$\begin{aligned} \left(\frac{\partial H^*}{\partial z^*}\right)_{p^*, t^*} &= -dp^* / dt^* \\ \left(\frac{\partial H^*}{\partial p^*}\right)_{z^*, t^*} &= dz^* / dt^* \equiv v^*. \end{aligned}$$

We also assume the canonical momentum p^* and the total energy H^* / c to form the z - and time – component of a 4-vector whose x - and y - components are unaltered in the following Lorentz transformation in z -direction. This transformation yields the relations for the unstarred components in the new system.

$$\begin{aligned} H &= \gamma H^* + \gamma \beta c p^* ; H^* = \gamma H - \gamma \beta c p \\ p &= \gamma p^* + (\gamma \beta / c) H^* ; p^* = \gamma p - (\gamma \beta / c) H \end{aligned}$$

$$\begin{aligned} z &= \gamma z^* + \gamma \beta c t^* ; z^* = \gamma z - \gamma \beta c t \\ t &= \gamma t^* + (\gamma \beta / c) z^* ; t^* = \gamma t - (\gamma \beta / c) z. \end{aligned}$$

Therefore, we find

$$\begin{aligned} \left(\frac{\partial H}{\partial z}\right)_{p,t} &= (1/\gamma) \left(\frac{\partial H^*}{\partial z}\right)_{p,t} + \beta c \left(\frac{\partial p}{\partial z}\right)_{p,t} = (1/\gamma) \left(\frac{\partial H^*}{\partial z}\right)_{p,t} \\ &= (1/\gamma) \left[\left(\frac{\partial H^*}{\partial z^*}\right)_{p^*, t^*} \cdot \left(\frac{\partial z^*}{\partial z}\right)_{p,t} + \left(\frac{\partial H^*}{\partial p^*}\right)_{z^*, t^*} \cdot \left(\frac{\partial p^*}{\partial z}\right)_{p,t} + \left(\frac{\partial H^*}{\partial t^*}\right)_{z^*, p^*} \cdot \left(\frac{\partial t^*}{\partial z}\right)_{p,t} \right]. \end{aligned}$$

$$\text{Since } \left(\frac{\partial z^*}{\partial z}\right)_{p,t} = \gamma; \left(\frac{\partial p^*}{\partial z}\right)_{p,t} = -(\gamma \beta / c) \left(\frac{\partial H}{\partial z}\right)_{p,t}; \left(\frac{\partial t^*}{\partial z}\right)_{p,t} = -\gamma \beta / c,$$

we have

$$\begin{aligned} \left(\frac{\partial H}{\partial z}\right)_{p,t} \left[1 + (\beta / c) \left(\frac{\partial H^*}{\partial p^*}\right)_{z^*, t^*} \right] &= \left(\frac{\partial H^*}{\partial z^*}\right)_{p^*, t^*} - (\beta / c) \left(\frac{\partial H^*}{\partial t^*}\right)_{z^*, p^*} \\ &= -dp^* / dt^* - (\beta / c) dH^* / dt^* = -(1/\gamma) dp / dt^* \end{aligned}$$

$$\text{Since } dt / dt^* = \gamma (1 + \beta v^* / c)$$

$$\text{we find } \left(\frac{\partial H}{\partial z}\right)_{p,t} = -dp / dt .$$

Conversely

$$\left(\frac{\partial H}{\partial p}\right)_{z,t} = \left(\frac{\partial H}{\partial p^*}\right)_{z^*,t^*} \cdot \left(\frac{\partial p^*}{\partial p}\right)_{z,t} + \left(\frac{\partial H}{\partial z^*}\right)_{p^*,t^*} \cdot \left(\frac{\partial z^*}{\partial p}\right)_{z,t} + \left(\frac{\partial H}{\partial t^*}\right)_{p^*,z^*} \cdot \left(\frac{\partial t^*}{\partial p}\right)_{z,t}.$$

Since
$$\left(\frac{\partial p^*}{\partial p}\right)_{z,t} = \gamma \left(\frac{\partial p}{\partial p}\right)_{z,t} - (\gamma\beta/c) \left(\frac{\partial H}{\partial p}\right)_{z,t} = \gamma \left[1 - (\beta/c) \left(\frac{\partial H}{\partial p}\right)_{z,t}\right]$$

and
$$\left(\frac{\partial z^*}{\partial p}\right)_{z,t} = \left(\frac{\partial t^*}{\partial p}\right)_{z,t} = 0$$

we find

$$\begin{aligned} \left(\frac{\partial H}{\partial p}\right)_{z,t} \left[1 + (\gamma\beta/c) \left(\frac{\partial H}{\partial p^*}\right)_{z^*,t^*}\right] &= \gamma \left(\frac{\partial H}{\partial p^*}\right)_{z^*,t^*} \\ &= \gamma^2 \left(\frac{\partial H^*}{\partial p^*}\right)_{z^*,t^*} + \gamma^2 \beta/c \left(\frac{\partial p^*}{\partial p^*}\right)_{z^*,t^*} = \gamma^2 (v^* + \beta c) \end{aligned}$$

and

$$\begin{aligned} \left(\frac{\partial H}{\partial p}\right)_{z,t} &= \gamma^2 (v^* + \beta c) / \left[1 + (\gamma^2 \beta/c) (v^* + \beta c)\right] \\ &= (v^* + \beta c) / (1 + v^* \beta/c) = v = dz/dt \end{aligned}$$

according to the rule of relativistic velocity addition.

Thus, the Hamiltonian relations hold and the transformed position z and momentum p are canonical variables of the transformed Hamiltonian H .

References

[1] J.D. Jackson: “Classical Electrodynamics”, Wiley & Sons

[2] A.A. Pomeranskii and I.B. Khriplovich; Journal of Experimental and Theoretical Physics; Vol. 86 Nr 5, 1998

Acknowledgements: The author wishes to thank William Donnelly, Mario Conte, and Waldo McKay for their helpful discussions.