

# SIMPLIFIED PROOF OF THE THEOREM OF VAROPOULOS IN THE COMMUTATIVE CASE

MARCO THILL

ABSTRACT. We give continuity properties of bitraces on (possibly non-commutative) Banach  $*$ -algebras based on the Closed Graph Theorem, leading to a simplified proof of the Theorem of Varopoulos in the commutative case.

## 1. INTRODUCTION

The Theorem of Varopoulos [8] says that a positive linear functional on a Banach  $*$ -algebra with continuous involution and with a bounded one-sided approximate unit is continuous. Its proof depends on two closely related Factorisation Theorems, one by Paul J. Cohen, and the other by Nicholas Th. Varopoulos himself. We quote them here:

**Theorem 1** (Paul J. Cohen [2]). *Let  $A$  be a Banach algebra with a bounded one-sided approximate unit. Then every element of  $A$  is a product of two elements of  $A$ .*

**Theorem 2** (Nicholas Th. Varopoulos [8]). *Let  $A$  be a Banach algebra with a bounded left approximate unit. Then for a sequence  $(a_n)$  in  $A$  converging to 0, there exists a sequence  $(b_n)$  in  $A$  converging to 0 as well as  $c \in A$  with  $a_n = c b_n$  for all  $n$ .*

Proofs of the above two results can also be found in the following books: [9, § 6], [1, § 11], and [4, Section 5.2].

Satish Shirali [6] extended the Theorem of Varopoulos to the case of a Banach  $*$ -algebra with a bounded one-sided approximate unit (and with a possibly discontinuous involution). (This was non-trivial also because Varopoulos used the fact that a bounded left approximate unit in a Banach  $*$ -algebra with continuous involution gives rise to a bounded right approximate unit.) Proofs of the Theorem of Varopoulos and Shirali can also be found in the books [5, Theorem 11.3.7 pp. 1189–1190] and [7, § 27].

We shall show how to prove the Theorem of Varopoulos in the commutative case without using the more technical Factorisation Theorem of Varopoulos. This will depend on continuity properties of bitraces

---

2000 *Mathematics Subject Classification*. Primary: 46K05; Secondary: 46K10.

*Key words and phrases*. Varopoulos, commutative, positive linear functional, bitrace, continuity.

Many thanks to Torben Maack Bisgaard for his excellent comments.

on (possibly non-commutative) Banach  $*$ -algebras. These continuity properties are based in a novel way on the Closed Graph Theorem.

The Factorisation Theorem of Varopoulos heavily depends on additional bounds on the factors in Cohen's Factorisation Theorem. These additional bounds are hard to establish, and are not needed for our proof. Furthermore, a number of simpler proofs for Cohen's Factorisation Theorem without these sharp bounds have been given. See the paragraph preceding the statement of [4, Theorem 5.2.2] and the references given there.

## 2. TERMINOLOGY AND NOTATION

We use the terminology of [3, Chapter XV, Sections 6 & 7]. In the following, let  $A$  be a  $*$ -algebra, and let  $\langle \cdot, \cdot \rangle$  be a positive Hilbert form on  $A$ , that is, a positive semi-definite sesquilinear form on  $A$  such that

$$\langle ab, c \rangle = \langle b, a^*c \rangle$$

holds for all  $a, b \in A$ . For example, any positive linear functional  $\varphi$  on  $A$  induces a positive Hilbert form on  $A$  by putting

$$\langle a, b \rangle := \varphi(b^*a) \quad (a, b \in A).$$

See [3, pp. 348–349].

The isotropic subspace

$$I := \{ a \in A : \langle a, b \rangle = 0 \text{ for all } b \in A \}$$

is a left ideal. The quotient space

$$\underline{A} := A/I$$

is a pre-Hilbert space with respect to the inner product

$$\langle \underline{a}, \underline{b} \rangle := \langle a, b \rangle \quad (a, b \in A),$$

where  $\underline{a} \in \underline{A}$  shall denote the projection of  $a \in A$ . Since  $I$  is a left ideal, it follows that for  $a \in A$  an operator  $\pi(a)$  mapping  $\underline{b} \in \underline{A}$  to  $\underline{ab} \in \underline{A}$  is well defined. We denote the completion of  $\underline{A}$  by  $H$ , which is a Hilbert space.

If  $A$  is a Banach  $*$ -algebra, then the operators  $\pi(a)$  ( $a \in A$ ) are bounded, and thus have a continuation to  $H$ , which we denote by the same symbol. It follows that the GNS representation

$$\begin{aligned} \pi : A &\rightarrow B(H) \\ a &\mapsto \pi(a) \end{aligned}$$

is bounded. For proofs, see for example [7, Theorem 22.14 p. 83].

One says that  $\langle \cdot, \cdot \rangle$  is a bitrace, if furthermore

$$\langle a, b \rangle = \langle b^*, a^* \rangle$$

holds for all  $a, b \in A$ . One then also has

$$\langle ba, c \rangle = \langle b, ca^* \rangle$$

for all  $a, b, c \in A$ , and it is this property that we shall need in lemma 3. For example, if  $\langle \cdot, \cdot \rangle$  is induced by a positive linear functional  $\varphi$ , then  $\langle \cdot, \cdot \rangle$  is a bitrace if and only if

$$\varphi(ab) = \varphi(ba)$$

holds for all  $a, b \in A$ . One then says that  $\varphi$  is a trace. See [3, pp. 356–358].

As usual, let  $A^2 := \text{span} \{ ab \in A : a, b \in A \}$ .

The pair  $(A, \langle \cdot, \cdot \rangle)$  is called a Hilbert algebra, if it is a pre-Hilbert space with  $A^2$  dense, if  $\langle \cdot, \cdot \rangle$  is a bitrace, and if the operators  $\pi(a)$  ( $a \in A$ ) all are bounded, cf. [3, p. 358].

In the following, we shall let  $A$  be a Banach  $*$ -algebra, and let  $p$  denote the projection on the closed subspace  $M$  of  $H$  given by

$$M := \overline{\text{span}} \{ \pi(a)x \in H : x \in H, a \in A \} \subset H.$$

### 3. THE PROOFS

**Lemma 3.** *Let  $\langle \cdot, \cdot \rangle$  be a bitrace on a Banach  $*$ -algebra  $A$ . The mapping*

$$\begin{aligned} A &\rightarrow H \\ a &\mapsto p\underline{a} \end{aligned}$$

(with notation as above) is bounded.

*Proof.* We shall show that the mapping in question has closed graph. So let  $a_n \rightarrow 0$  in  $A$ , and  $p\underline{a_n} \rightarrow x$  in  $H$ . We have to show that  $x = 0$ . The vector  $x$  lies in  $M$ . It is thus enough to prove that  $\langle x, \pi(b)\underline{c} \rangle = 0$  for all  $b, c \in A$ . One calculates

$$\begin{aligned} \langle x, \pi(b)\underline{c} \rangle &= \lim_{n \rightarrow \infty} \langle p\underline{a_n}, \pi(b)\underline{c} \rangle = \lim_{n \rightarrow \infty} \langle \underline{a_n}, p\pi(b)\underline{c} \rangle \\ &= \lim_{n \rightarrow \infty} \langle \underline{a_n}, \pi(b)\underline{c} \rangle = \lim_{n \rightarrow \infty} \langle a_n, bc \rangle = \lim_{n \rightarrow \infty} \langle a_n c^*, b \rangle \\ &= \lim_{n \rightarrow \infty} \langle \pi(a_n)\underline{c}^*, \underline{b} \rangle = 0 \end{aligned}$$

by continuity of  $\pi$ . □

A positive Hilbert form  $\langle \cdot, \cdot \rangle$  on a normed  $*$ -algebra  $(A, |\cdot|)$  is called bounded if there exists  $\gamma \geq 0$  such that

$$|\langle a, b \rangle| \leq \gamma \cdot |a| \cdot |b|$$

holds for all  $a, b \in A$ .

**Theorem 4.** *A bitrace  $\langle \cdot, \cdot \rangle$  on a Banach  $*$ -algebra  $A$  is bounded if the corresponding GNS representation is non-degenerate. The last condition is satisfied, if  $(A, \langle \cdot, \cdot \rangle)$  is a Hilbert algebra.*

From the lemma also follows:

**Theorem 5.** *A bitrace on a Banach  $*$ -algebra  $A$  is bounded on  $A^2$ .*

**Theorem 6.** *A bitrace on a Banach  $*$ -algebra with a bounded one-sided approximate unit is bounded.*

*Proof.* This follows now from the Factorisation Theorem of Cohen (theorem 1 above).  $\square$

**Theorem 7.** *Let  $A$  be a Banach  $*$ -algebra with a bounded right approximate unit  $(e_i)_{i \in I}$ . Let  $\langle \cdot, \cdot \rangle$  be a non-zero bitrace on  $A$ . The corresponding GNS representation  $\pi$  then is cyclic. Furthermore, there exists a cyclic vector  $c$  for  $\pi$ , such that the positive linear functional  $a \mapsto \langle \pi(a)c, c \rangle$  ( $a \in A$ ) induces the bitrace  $\langle \cdot, \cdot \rangle$ . The vector  $c$  can be chosen to be any adherence point of the net  $(\underline{e}_i)_{i \in I}$  in the weak topology.*

(We give the proof here for the sake of staying self-contained. A more general result is given in [7, Theorem 27.4]. See also [7, Theorem 27.5].)

*Proof.* By boundedness of  $\langle \cdot, \cdot \rangle$  (theorem 6), the net  $(\underline{e}_i)_{i \in I}$  is bounded in  $H$ , and therefore has an adherence point in the weak topology on  $H$ . Let  $c$  be any such adherence point. By going over to a subnet, we can assume that  $(\underline{e}_i)_{i \in I}$  converges weakly to  $c$ . It shall be shown that  $\pi(a)c = \underline{a}$  for all  $a \in A$ , which implies that  $c$  is cyclic for  $\pi$ . It is enough to show that for  $b \in A$ , one has  $\langle \pi(a)c, \underline{b} \rangle = \langle \underline{a}, \underline{b} \rangle$ . So one calculates

$$\begin{aligned} \langle \pi(a)c, \underline{b} \rangle &= \langle c, \pi(a)^* \underline{b} \rangle = \lim_{i \in I} \langle \underline{e}_i, \pi(a)^* \underline{b} \rangle \\ &= \lim_{i \in I} \langle \pi(a) \underline{e}_i, \underline{b} \rangle = \lim_{i \in I} \langle a e_i, b \rangle = \langle a, b \rangle = \langle \underline{a}, \underline{b} \rangle, \end{aligned}$$

where we have used the continuity of  $\langle \cdot, \cdot \rangle$ . Consider now the positive linear functional  $\varphi$  on  $A$  defined by

$$\varphi(a) := \langle \pi(a)c, c \rangle \quad (a \in A).$$

We have

$$\varphi(a^*a) = \langle \pi(a^*a)c, c \rangle = \|\pi(a)c\|^2 = \|\underline{a}\|^2 = \langle \underline{a}, \underline{a} \rangle = \langle a, a \rangle$$

whence, after polarisation,

$$\langle a, b \rangle = \varphi(b^*a) \quad (a, b \in A),$$

that is,  $\varphi$  induces the bitrace  $\langle \cdot, \cdot \rangle$ .  $\square$

A positive linear functional  $\varphi$  on a normed  $*$ -algebra  $(A, \|\cdot\|)$  is called representable, if there exists a non-degenerate continuous representation  $\pi$  of  $(A, \|\cdot\|)$  in a Hilbert space, and a vector  $c$  such that

$$\varphi(a) = \langle \pi(a)c, c \rangle$$

holds for all  $a \in A$ . The functional  $\varphi$  then is continuous by the continuity assumption on  $\pi$ .

It is known that a positive linear functional  $\varphi$  on a Banach  $*$ -algebra  $A$  is representable, if and only if it has finite variation, in the sense that there exists  $\gamma \geq 0$  such that

$$|\varphi(a)|^2 \leq \gamma \varphi(a^*a)$$

holds for all  $a \in A$ . See for example [7, §§ 24–25].

**Theorem 8.** *A trace on a Banach  $*$ -algebra with a bounded one-sided approximate unit has finite variation. Hence it is representable, and thus also continuous.*

*Proof.* Let  $\varphi$  be a trace on a Banach  $*$ -algebra  $A$  with a bounded one-sided approximate unit. In case of a bounded right approximate unit, it follows from theorem 7 and the Factorisation Theorem of Cohen (theorem 1 above), that  $\varphi$  is representable. Then  $\varphi$  has finite variation and is Hermitian in the sense that  $\varphi(a^*) = \overline{\varphi(a)}$  for all  $a \in A$ . In case of a bounded left approximate unit, one changes the multiplication in  $A$  to  $(a, b) \mapsto ba$ , and one applies the preceding case of a bounded right approximate unit. Using the fact that  $\varphi$  turns out to be Hermitian, it follows again that  $\varphi$  has finite variation.  $\square$

**Corollary 9.** *A positive linear functional on a commutative Banach  $*$ -algebra with a bounded one-sided approximate unit has finite variation. Hence it is representable, and thus also continuous.*

#### REFERENCES

- [1] F. F. Bonsall, J. Duncan. *Complete Normed Algebras*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 80, Springer-Verlag, Berlin Heidelberg New York, 1973.
- [2] P. Cohen, Factorization in group algebras, *Duke Math. J.*, 26, pp. 199–205, 1959.
- [3] J. Dieudonné. *Treatise on Analysis*, vol. 2, enlarged and corrected printing, Academic Press, 1976.
- [4] Th. W. Palmer. *Encyclopedia of Mathematics and its Applications*, vol. 49: *Algebras and Banach Algebras*, Cambridge University Press, 1994.
- [5] Th. W. Palmer. *Encyclopedia of Mathematics and its Applications*, vol. 79: *Banach algebras and the General Theory of  $*$ -Algebras*, Cambridge University Press, 2001.
- [6] S. Shirali. Representability of positive functionals, *J. London Math. Soc.*, 2 (3), pp. 145–155, 1971.
- [7] M. Thill. *Introduction to Normed  $*$ -Algebras and their Representations*, 5th ed., arXiv:math/0701306v2 [math.RT], 2007.
- [8] N. Th. Varopoulos. Sur les formes positives d'une algèbre de Banach, *Comptes Rendus Acad. Sc. Paris*, tome 258, Groupe 1, pp. 2465–2467, 1964.
- [9] W. Żelazko. *Banach Algebras*, Polish Scientific Publishers and Elsevier Publishing Company, 1973.

BD G. - D. CHARLOTTE 53, L - 1331 LUXEMBOURG CITY, EUROPE  
E-mail address: math@pt.lu