

A SYMPLECTIC ISOTOPY OF A DEHN TWIST ON $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$

EMIKO DUPONT

ABSTRACT. The complex manifold $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ with symplectic form $\sigma_\mu = \sigma_{\mathbb{C}\mathbb{P}^n} + \mu\sigma_{\mathbb{C}\mathbb{P}^{n+1}}$, where $\sigma_{\mathbb{C}\mathbb{P}^n}$ and $\sigma_{\mathbb{C}\mathbb{P}^{n+1}}$ are normalized Fubini-Study forms, $n \in \mathbb{N}$ and $\mu > 1$ a real number, contains a natural Lagrangian sphere L^μ . We prove that the Dehn twist along L^μ is symplectically isotopic to the identity for all $\mu > 1$. This isotopy can be chosen so that it pointwise fixes a complex hypersurface in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ and lifts to the blow-up of $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ along a complex n -dimensional submanifold.

1. INTRODUCTION

Suppose L is a Lagrangian sphere in a symplectic manifold (M, ω) . The generalized Dehn twist (the Dehn twist for short) τ_L along L is a symplectomorphism on M that is compactly supported near L and restricts to L as the antipodal map. A symplectomorphism τ is said to be symplectically isotopic to the identity if there is a smooth family $(\tau_s)_{0 \leq s \leq 1}$ of symplectomorphisms such that $\tau_0 = \text{id}$ and $\tau_1 = \tau$.

In this paper we consider the (real) $4n + 2$ -dimensional manifold $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ with the product symplectic form $\sigma_\mu = \sigma_{\mathbb{C}\mathbb{P}^n} + \mu\sigma_{\mathbb{C}\mathbb{P}^{n+1}}$ for $\mu > 1$ a real number. Here $n \in \mathbb{N}$ and $\sigma_{\mathbb{C}\mathbb{P}^n}$ and $\sigma_{\mathbb{C}\mathbb{P}^{n+1}}$ are the Fubini-Study forms on $\mathbb{C}\mathbb{P}^n$ and $\mathbb{C}\mathbb{P}^{n+1}$, respectively, normalized to integrate to π on $\mathbb{C}\mathbb{P}^1$. As described in Lemma 2.5, the graph of the complex conjugate of the Hopf map $S^{2n+1} \rightarrow \mathbb{C}\mathbb{P}^n$ embeds as the Lagrangian sphere

$$(1.1) \quad L^\mu = \{([\bar{z}_0 : \cdots : \bar{z}_n], [z_0 : \cdots : z_n : \sqrt{\mu - 1}]) \mid |z|^2 = 1\}$$

in $(\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$. The main result of this paper is the following.

Theorem 1.1. *For all $\mu > 1$, the Dehn twist $\tau_{L^\mu} \in \text{Symp}(\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$ along the Lagrangian L^μ is symplectically isotopic to the identity by an isotopy whose restriction to the complex hypersurface*

$$S = \{([s_0 : \cdots : s_n], [x_0 : \cdots : x_{n+1}]) \in \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1} \mid s_0x_0 + \cdots + s_nx_n = 0\}$$

is the identity.

The isotopy of Theorem 1.1 was established in the case $n = 1$ and $\mu \gg 1$ by Corti-Smith [CS05, Section 7]. Corti-Smith construct a singular fibration with non-singular fibers isotopic to $(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^2, \sigma_\mu)$ and such that the monodromy around the only singular fiber of this fibration is symplectically isotopic to the Dehn twist τ_{L^μ} . As the monodromy is also known to be symplectically trivial, this proves the result. The proof of Theorem 1.1 uses exactly the same fibration but by examining the construction in more detail, we are able to establish an isotopy for all $\mu > 1$. Furthermore, we observe that the construction generalizes to all $n \in \mathbb{N}$ and that the complex hypersurface S is pointwise-fixed under the isotopy.

The hypersurface S is a $\mathbb{C}\mathbb{P}^n$ -bundle over $\mathbb{C}\mathbb{P}^n$ in which the base coordinates are $[s_0 : \cdots : s_n]$, and each fiber is a linearly embedded copy of $\mathbb{C}\mathbb{P}^n$ in $\mathbb{C}\mathbb{P}^{n+1}$.

Write $\mathbb{C}\mathbb{P}^{n+1}$ as $\mathbb{C}^n \sqcup D$ where \mathbb{C}^n is the coordinate chart centered at the point $p_0 = [0 : \cdots : 0 : 1]$ and $D = (x_{n+1} = 0) \cong \mathbb{C}\mathbb{P}^n$. Then S contains a section at $0 \in \mathbb{C}^n$ given by $S_0 = \mathbb{C}\mathbb{P}^n \times \{p_0\}$ and a section at infinity, namely,

$$S_\infty = \{([s_0 : \cdots : s_n], [x_0 : x_1 : 0 : \cdots : 0]) \mid s_0 x_0 + s_1 x_1 = 0\} \subset \mathbb{C}\mathbb{P}^n \times D.$$

We show that S_0 and S_∞ are not only pointwise-fixed under the isotopy but in fact we have the following result.

Corollary 1.2. *Let S_0 and S_∞ be the complex submanifolds in S , each isomorphic to $\mathbb{C}\mathbb{P}^n$, defined by*

$$\begin{aligned} S_0 &= \mathbb{C}\mathbb{P}^n \times \{p_0\}, \\ S_\infty &= \{([s_0 : \cdots : s_n], [x_0 : x_1 : 0 : \cdots : 0]) \in \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1} \mid s_0 x_0 + s_1 x_1 = 0\} \end{aligned}$$

where $p_0 = [0 : \cdots : 0 : 1]$. *The isotopy of Theorem 1.1 lifts to the blow-up of $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ along S_0 and S_∞ if the size of the blow-up is sufficiently small.*

One should note that the Dehn twist τ_{L^μ} is well-defined on the blow-up since both S_0 and S_∞ are disjoint from L^μ and the Dehn twist is supported near L^μ .

Theorem 1.1 is a consequence of Theorem 1.3. The latter constructs a Lefschetz fibration with exactly one critical point whose non-singular fibers are isotopic to $(\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$ and with vanishing cycle isotopic to the Lagrangian sphere L^μ . Thus, the monodromy around a positively-oriented loop in $\mathbb{C}\mathbb{P}^1$ that circles the critical value exactly once, is symplectically isotopic to the Dehn twist τ_{L^μ} . Since there is only one critical value, the monodromy is also symplectically isotopic to the identity.

Theorem 1.3. *Let $\mu > 1$. There is a Lefschetz fibration $(\mathcal{X}, \pi, J, \Omega^\mu)$ such that:*

- (1) *π has exactly one critical point $z_{\text{crit}} \in \mathcal{X}_0 = \pi^{-1}(0)$;*
- (2) *there is a holomorphic trivialization*

$$\Phi : \mathcal{X} \setminus \mathcal{X}_0 \rightarrow (\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}) \times \mathbb{C}$$

whose restriction to the fiber \mathcal{X}_∞ over the point at infinity in $\mathbb{C}\mathbb{P}^1$ composed with the projection to $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ is a symplectomorphism

$$\Phi_\infty : (\mathcal{X}_\infty, \Omega^\mu |_{\mathcal{X}_\infty}) \rightarrow (\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu);$$

- (3) *the map Φ_∞ takes the vanishing cycle \mathcal{V}_∞^μ in \mathcal{X}_∞ to L^μ in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$.*

For $n = 1$, Theorem 1.3 is an extended version of [CS05, Proposition 1]. The Lefschetz fibration $(\mathcal{X}, \pi, J, \Omega^\mu)$ is an extension of the fibration constructed in [CS05] from a disk $\Delta \subset \mathbb{C}$ to $\mathbb{C}\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$. Note, however, that unless $\mu \gg 1$, the full statement of Theorem 1.3 is necessary in order to identify the vanishing cycle with the Lagrangian 3-sphere L^μ in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^2$. After deforming Ω^μ to a form $(\Omega^\mu)'$ which is standard in a small neighborhood of the critical point z_{crit} , Corti-Smith compute the vanishing cycle $(\mathcal{V}_t^\mu)'$ in the fibers close to the singular fiber \mathcal{X}_0 . For each $t \in \Delta$, write $\Phi_t : \mathcal{X}_t \rightarrow \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^2$ for the restriction $\Phi |_{\mathcal{X}_t}$ of the map Φ described in Theorem 1.3 followed by projection to $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^2$. The map Φ_t is biholomorphic and $\alpha'_t = (\Phi_t^{-1})^*((\Omega^\mu)' |_{\mathcal{X}_t})$ is a symplectic form on $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^2$ in the same cohomology class as σ_μ . Hence by Moser's Theorem, α'_t is isotopic to σ_μ . If $\mu \gg 1$, the fiber \mathcal{X}_1 intersects the neighborhood in which $(\Omega^\mu)'$ is standard and Corti-Smith show that the pull-back by Φ_1 of the vanishing cycle $(\mathcal{V}_1^\mu)'$ in \mathcal{X}_1 equals the Lagrangian $L^\mu \cong S^3$. Hence the isotopy from α'_1 to σ_μ can be made to fix L^μ .

For general $\mu > 1$, however, it is not clear what happens to the vanishing cycle under the isotopy and, in particular, we cannot identify it with L^μ . On the other hand, by extending the fibration over $\mathbb{C}\mathbb{P}^1$ as in Theorem 1.3, we see that the form $\alpha_\infty = (\Phi_\infty^{-1})^*(\Omega^\mu)$ is not only isotopic to σ_μ but, in fact, $\alpha_\infty = \sigma_\mu$ and $\Phi_\infty : \mathcal{X}_\infty \rightarrow \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^2$ is a symplectomorphism. The key step in our construction is the following. By symplectically embedding a large part of the total space $(\mathcal{X}, \Omega^\mu)$ of the Lefschetz fibration into a toric manifold $(\mathbb{F}, \omega_{\mathbb{F}}^{(1, \mu)})$, we obtain a Darboux chart on $(\mathcal{X}, \Omega^\mu)$ that enables us to compute the horizontal spaces of the symplectic connection coming from Ω^μ in a large neighborhood of the critical point z_{crit} . In particular, we are able to compute the vanishing cycle \mathcal{V}_∞^μ explicitly and to see that $\Phi_\infty(\mathcal{V}_\infty^\mu)$ is the Lagrangian L^μ .

Our construction generalizes easily from the case $n = 1$ to arbitrary $n \in \mathbb{N}$. Furthermore, with $S \subset \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ as defined in Theorem 1.1, we see that the corresponding hypersurface $\Phi_t^{-1}(S)$ in the fiber \mathcal{X}_t is held fixed under symplectic parallel transport in $(\mathcal{X}, \pi, J, \Omega^\mu)$. This implies that the isotopy of Theorem 1.1 fixes the complex hypersurface S in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$.

The results in this paper are motivated by the symplectic isotopy problem which is described below. Given a symplectic manifold (M, ω) , the group of symplectomorphisms on M is an infinite-dimensional Lie subgroup of the group $\text{Diff}^+(M)$ of orientation-preserving diffeomorphisms of M . We say that a symplectomorphism τ is smoothly isotopic to the identity if there is an isotopy $(\tau_s)_{0 \leq s \leq 1}$ with $\tau_s \in \text{Diff}^+(M)$. A symplectomorphism is called essential if it is smoothly isotopic to the identity but not symplectically so.

The symplectic isotopy problem: *Given a symplectic manifold (M, ω) , does it admit essential symplectomorphisms?*¹

When the dimension of M is 2, the space of symplectic forms on M is convex and it follows from Moser's Theorem that no essential symplectomorphisms exist. In dimension 4, Gromov and Abreu-McDuff (see [MS04, p.320-321]) show that the answer to the isotopy problem is once again negative for $\mathbb{C}\mathbb{P}^2$, $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ and the one point blow-up of $\mathbb{C}\mathbb{P}^2$. In 1997, however, Seidel [Sei97] shows that under fairly weak conditions on a 4-manifold, if τ_L is the Dehn twist in a Lagrangian sphere L , then τ_L^2 is an essential symplectomorphism. Notice here that since τ_L restricts to L as the antipodal map, τ_L cannot be isotopic to the identity when the dimension of L is even. On the other hand, Seidel showed that when M is 4-dimensional, the square of the Dehn twist is always smoothly isotopic to the identity.

In dimensions 6 and above, very little is known. It is not even clear that Dehn twists (or their squares) are smoothly isotopic to the identity. Notice that in the example $(\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$ studied in this paper, the Lagrangian sphere is odd-dimensional. Thus, there is no homological obstruction for the Dehn twist to be isotopic to the identity, and indeed, Theorem 1.1 shows that it is. Although Theorem 1.1 does not provide an answer to the symplectic isotopy problem, the construction of the isotopy relies on certain symmetries specific to these examples that suggests they may be higher-dimensional analogues of the special cases $\mathbb{C}\mathbb{P}^2$ and $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ in dimension 4. (See [Dup07, Concluding remarks]).

Furthermore, by understanding how the isotopy could be destroyed, one may be able to construct examples of essential symplectomorphisms in these dimensions.

¹In this paper we consider only compact symplectic manifolds.

In contrast to Corollary 1.2, the proof of [CS05, Proposition 2] shows the following result about S_0 and a different n -dimensional submanifold S'_∞ at infinity.

Proposition 1.4. *Consider the complex submanifolds in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ given by*

$$S_0 = \mathbb{C}\mathbb{P}^n \times \{p_0\} \quad \text{and} \quad S'_\infty = \mathbb{C}\mathbb{P}^n \times \{p_\infty\},$$

where $p_0 = [0 : \cdots : 0 : 1]$ and $p_\infty = [1 : 0 : \cdots : 0]$. *There is no smooth isotopy between the Dehn twist τ_{L^μ} and the identity that simultaneously fixes S_0 and S'_∞ pointwise.*

It therefore seems likely that blowing up along the submanifolds S_0 and S'_∞ would destroy the symplectic isotopy of Theorem 1.1. This does not preclude there being no symplectic isotopies between the Dehn twist along L^μ and the identity in the blow-up. If one could nevertheless exhibit a smooth isotopy between τ_{L^μ} and the identity in the blow-up, then this example would provide a candidate for an essential symplectomorphism in dimension $4n + 2$. An alternative approach to destroying the isotopy of Theorem 1.1 would be to perform a large blow-up of the submanifolds S_0 and S_∞ of Corollary 1.2. It would be interesting to estimate the bounds of the size of the blow-ups allowed for the isotopy to persist and to see what happens when we exceed the bounds.

Acknowledgments. This paper contains the results of my Ph.D. thesis, and I thank my Ph.D. advisor, Dusa McDuff, for her help and guidance throughout my Ph.D. I am also grateful to Aleksey Zinger for his many comments and suggestions.

2. PRELIMINARIES

This section is expository and can be skipped if one is already familiar with Dehn twists and Lefschetz fibrations. The main statement is Proposition 2.7 which shows that the monodromy of a Lefschetz fibration around a loop that circles a critical value once is symplectically isotopic to the Dehn twist along the vanishing cycle corresponding to the critical value. The discussion in this section is based on [Sei97] and [Sei03, Section 1]. Note that although [Sei03] assumes exactness of Lefschetz fibrations, i.e., that each non-singular fiber is a symplectic manifold with boundary and the symplectic form on these fibers is exact, the proofs of the results that we use are easily adapted to ordinary Lefschetz fibrations. Details can be found in [Dup07].

2.1. The model Dehn twist. Consider the cotangent bundle of S^N

$$T^*S^N = \{(u, v) \in \mathbb{R}^{N+1} \times \mathbb{R}^{N+1} \mid \|u\| = 1, \langle u, v \rangle = 0\}$$

with symplectic form $\omega_{T^*S^N} = -\sum_j du_j \wedge dv_j$. The zero section L_0 is a Lagrangian submanifold of $(T^*S^N, \omega_{T^*S^N})$. The length function $h : T^*S^N \rightarrow \mathbb{R}$ given by $h(u, v) = \|v\|$ generates a Hamiltonian circle action on $T^*S^N \setminus L_0$ whose flow is given by

$$(2.1) \quad \varphi_\theta^h(u, v) = \left(\cos(\theta)v - \sin(\theta)\|v\|u, \cos(\theta)u + \sin(\theta)\frac{v}{\|v\|} \right).$$

The time- π map φ_π^h extends over the zero section by the antipodal map $A(u, 0) = (-u, 0)$. Now let $R : \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function such that $R(s) = 0$ for $s \geq s_R$ for some $s_R > 0$ and $R(-s) = R(s) - s$ for all s . Let $H = R \circ h$. The flow of H is $\varphi_\theta^H(u, v) = \varphi_{\theta R'(\|v\|)}^h(u, v)$. Since $R'(0) = \frac{1}{2}$, $\varphi_{2\pi}^H$ extends continuously to T^*S^N by the antipodal map. By [Sei03, Lemma 1.8] this extension is smooth and hence is a symplectomorphism.

Definition 2.1. Let τ be the time- 2π flow of $H = R \circ h$ on $T^*S^N \setminus L_0$ as above, extended to L_0 by the antipodal map. The symplectomorphism τ is called a *model Dehn twist*.




FIGURE 1. The model Dehn twist

Figure 1 shows the image of a fiber $F_u = \pi_{T^*S^N}^{-1}(u, 0)$ of T^*S^N under a model Dehn twist. Here $\pi_{T^*S^N} : T^*S^N \rightarrow L_0 \cong S^N$ is the natural projection. We see that $(u, v) \in T^*S^N$ with $\|v\|$ large are held fixed by τ , but as $\|v\|$ decreases, the circle action φ_θ^h is “turned on” with larger and larger $\theta \in (\pi, 0)$ until at $\|v\| = 0$, we reach the antipodal map φ_π^h . Although the definition of the model Dehn twist depends on the choice of function R , the symplectic isotopy class of τ is independent of R .

Let $\omega_{\mathbb{C}^N} = \frac{i}{2} \sum_j dz_j \wedge d\bar{z}_j$ denote the standard symplectic structure on \mathbb{C}^N . Consider the singular fibration $\pi_{\text{std}} : \mathbb{C}^N \rightarrow \mathbb{C}$ given by

$$\pi_{\text{std}}(z_1, \dots, z_N) = \sum_{j=1}^N z_j^2.$$

All fibers of π_{std} , except for $\pi_{\text{std}}^{-1}(0)$, are symplectic submanifolds of $(\mathbb{C}^N, \omega_{\mathbb{C}^N})$. For each $r > 0$,

$$\pi_{\text{std}}^{-1}(r) := \{z \in \mathbb{C}^N \mid \|\Re(z)\|^2 - \|\Im(z)\|^2 = r, \sum_j \Re(z_j)\Im(z_j) = 0\}$$

is symplectomorphic to $(T^*S^{N-1}, \omega_{T^*S^{N-1}})$ by the map

$$(2.2) \quad \Phi_r(z) = \left(\frac{\Re(z)}{\|\Re(z)\|}, -\|\Re(z)\|\Im(z) \right).$$

Away from the singular fiber, the symplectic structure $\omega_{\mathbb{C}^N}$ gives a natural connection on $\pi_{\text{std}} : \mathbb{C}^N \rightarrow \mathbb{C}$ for which parallel transport maps are symplectomorphisms. Each non-singular fiber $\pi_{\text{std}}^{-1}(t)$, $t \neq 0$, contains a Lagrangian sphere

$$(2.3) \quad \Sigma_t = \{\sqrt{r}e^{i\frac{\theta}{2}}z \in \mathbb{C}^N \mid \Im(z) = 0, \|z\|^2 = 1\}, \quad \text{where } t = re^{i\theta}.$$

These spheres are mapped to each other by parallel transport. Note that for $r > 0$, $\Phi_r(\Sigma_r)$ is the zero section L_0 . The following result is based on the proof of [Sei03, Lemma 1.10] and is proved in detail in [Dup07].

Lemma 2.2. Consider the singular symplectic fibration $\pi_{\text{std}} : \mathbb{C}^N \rightarrow \mathbb{C}$ given by $\pi_{\text{std}}(z_1, \dots, z_N) = \sum_{j=1}^N z_j^2$, where \mathbb{C}^N has the standard symplectic form $\omega_{\mathbb{C}^N}$. For $r > 0$, let $\rho_r : \pi_{\text{std}}^{-1}(r) \rightarrow \pi_{\text{std}}^{-1}(r)$ denote the monodromy around the loop $\gamma(\theta) = re^{i\theta}$, $0 \leq \theta \leq 2\pi$, and let $\Phi_r : \pi_{\text{std}}^{-1}(r) \rightarrow T^*S^{N-1}$ be defined as in (2.2). Then for all $r > 0$,

$$\tilde{\tau}_r = \Phi_r \circ \rho_r \circ (\Phi_r)^{-1} : T^*S^{N-1} \rightarrow T^*S^{N-1}$$

is symplectically isotopic to a model Dehn twist.

2.2. Dehn twists.

Definition 2.3. Let (M^{2N}, ω) be a symplectic manifold and $L \subset M$ a Lagrangian sphere with a chosen identification $\iota : S^N \rightarrow L$. By the Lagrangian Neighborhood Theorem, for some small $\lambda > 0$, we can extend ι to a symplectic embedding of the space $T_{\leq \lambda}^*S^N$ of cotangent vectors with $\|v\| \leq \lambda$ to a neighborhood $\mathcal{N}(L)$ of L . Choose a model Dehn twist τ whose support is in $T_{\leq \frac{\lambda}{2}}^*S^N$. The *generalized Dehn twist* (or simply *Dehn twist*) along L is the map

$$\tau_L(z) = \begin{cases} \iota \circ \tau \circ \iota^{-1}, & \text{if } z \in \mathcal{N}(L); \\ z & \text{otherwise.} \end{cases}$$

Remark 2.4. The Dehn twist is a symplectomorphism of M that depends on the choice of identification $\iota : S^N \rightarrow L$. We say that two identifications ι_1 and ι_2 give the same *framing* if $\iota_2^{-1} \circ \iota_1 : S^N \rightarrow S^N$ can be deformed inside the group of diffeomorphisms of S^N to an element of $O(N+1)$. Identifications that give the same framing define Dehn twists that are symplectically isotopic. However, it is unknown whether the Dehn twist can be defined as a symplectic isotopy class independent of framing. We omit the choice of framing in our notation as there is often a natural choice in the situations we describe below. \square

Now consider $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ with the product symplectic form $\sigma_\mu = \sigma_{\mathbb{C}\mathbb{P}^n} + \mu\sigma_{\mathbb{C}\mathbb{P}^{n+1}}$, where $\mu > 1$ and $\sigma_{\mathbb{C}\mathbb{P}^n}, \sigma_{\mathbb{C}\mathbb{P}^{n+1}}$ are the normalized Fubini-Study forms on $\mathbb{C}\mathbb{P}^n$ and $\mathbb{C}\mathbb{P}^{n+1}$, respectively. As shown in [CS05, Lemma 1], the graph of the complex conjugate of the Hopf map $H_{\mathbb{C}\mathbb{P}^n} : S^{2n+1} \rightarrow \mathbb{C}\mathbb{P}^n$ embeds naturally into this manifold as a Lagrangian sphere L^μ .

Lemma 2.5. If $\mu > 1$, then

$$L^\mu := \{[\bar{z}_0 : \dots : \bar{z}_n][z_0 : \dots : z_n : \sqrt{\mu-1}] \in \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1} \mid |z|^2 = 1\}$$

is a Lagrangian sphere in $(\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$.

Proof. For $\lambda > 0$, let $B(\lambda) = \{z \in \mathbb{C}^{n+1} \mid |z|^2 < \lambda^2\}$. Then the map $i : B(\sqrt{\mu}) \hookrightarrow \mathbb{C}\mathbb{P}^{n+1}$ given by

$$i : (z_0, \dots, z_n) \mapsto [z_0 : \dots : z_n : \sqrt{\mu - |z_0|^2 - \dots - |z_n|^2}]$$

is a symplectic embedding $(B(\sqrt{\mu}), \omega_{\mathbb{C}^{n+1}}) \hookrightarrow (\mathbb{C}\mathbb{P}^{n+1}, \mu\sigma_{\mathbb{C}\mathbb{P}^{n+1}})$. This follows from the fact that i factors through the Hopf map $H_{\mathbb{C}\mathbb{P}^{n+1}} : S^{2n+3} \rightarrow \mathbb{C}\mathbb{P}^{n+1}$

$$\begin{array}{ccc} & & S^{2n+3} \\ & \nearrow \tilde{i} & \downarrow H_{\mathbb{C}\mathbb{P}^{n+1}} \\ B(\sqrt{\mu}) & \xrightarrow{i} & \mathbb{C}\mathbb{P}^{n+1} \end{array}$$

where

$$\tilde{i}(z_0, \dots, z_n) = \left(\frac{z_0}{\sqrt{\mu}}, \dots, \frac{z_n}{\sqrt{\mu}}, \sqrt{1 - \left| \frac{z}{\sqrt{\mu}} \right|^2} \right).$$

Hence,

$$i^*(\mu\sigma_{\mathbb{C}\mathbb{P}^{n+1}}) = \tilde{i}^*((H_{\mathbb{C}\mathbb{P}^{n+1}})^*(\mu\sigma_{\mathbb{C}\mathbb{P}^{n+1}})) = \tilde{i}^*(\mu\omega_{\mathbb{C}^{2n+3}}|_{S^{2n+3}}) = \omega_{\mathbb{C}^{n+1}}|_{B(\sqrt{\mu})}.$$

Thus, $S^{2n+1} := \partial B(1) \subset B(\sqrt{\mu})$ embeds symplectically into $(\mathbb{C}\mathbb{P}^{n+1}, \mu\sigma_{\mathbb{C}\mathbb{P}^{n+1}})$. Therefore, the graph of the conjugate of the Hopf map $H_{\mathbb{C}\mathbb{P}^n} : S^{2n+1} \rightarrow \mathbb{C}\mathbb{P}^n$ embeds symplectically into $(\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu^n)$ as L^μ . Since $(\overline{H}_{\mathbb{C}\mathbb{P}^n})^*(\sigma_{\mathbb{C}\mathbb{P}^n}) = -\omega_{\mathbb{C}^{n+1}}|_{S^{2n+1}}$,

$$\sigma_\mu|_{L^\mu} = \omega_{\mathbb{C}^{n+1}}|_{S^{2n+1}} + (\overline{H}_{\mathbb{C}\mathbb{P}^n})^*(\sigma_{\mathbb{C}\mathbb{P}^n}) = 0,$$

i.e., L^μ is Lagrangian. \square

2.3. Lefschetz fibrations.

Definition 2.6. A *Lefschetz fibration* is a smooth fibration $\pi : E \rightarrow \mathbb{C}\mathbb{P}^1$ such that

- the set E_{crit} of critical points of π is finite and no two critical points lie in the same fiber;
- E is compact and has a closed 2-form Ω that restricts to a symplectic form on the non-singular fibers;
- there is a complex structure J defined in a neighborhood of each critical point such that Ω is J -Kähler;
- if j is the complex structure on $\mathbb{C}\mathbb{P}^1$, then π is (J, j) -holomorphic and at each critical point the Hessian of π is non-degenerate.

Note that by the Morse Lemma, the fourth condition is equivalent to the condition that for each $z_{\text{crit}} \in E_{\text{crit}}$ we can find a J -holomorphic coordinate chart Ψ in a neighborhood of z_{crit} and a chart ψ on $\mathbb{C}\mathbb{P}^1$ centered at $\pi(z_{\text{crit}})$ in $\mathbb{C}\mathbb{P}^1$ such that

$$\psi \circ \pi \circ \Psi^{-1}(z) = \sum_{i=1}^N z_i^2.$$

Such a pair of charts (Ψ, ψ) will be called a *Morse chart*. In the following we often suppress the choice of ψ in our notation.

Away from the critical points, (E, π, Ω) is a symplectic fiber bundle and hence admits a symplectic connection. Each critical point $z_{\text{crit}} \in E_{\text{crit}}$ gives rise to a Lagrangian sphere in the non-singular fibers called the vanishing cycle. Let $\gamma : [a, b] \rightarrow \mathbb{C}\mathbb{P}^1$ be an embedded path which avoids $\pi(E_{\text{crit}})$ except at the endpoint $\gamma(b) = \pi(z_{\text{crit}})$. Define

$$B_\gamma = \{z_{\text{crit}}\} \cup \bigcup_{a \leq s < b} \{z \in E_{\gamma(s)} \mid \lim_{s' \rightarrow b} \rho_{\gamma|_{[s, s']}}(z) = z_{\text{crit}}\}.$$

By [Sei03, Lemma 1.13 and 1.14], B_γ is an embedded closed N -ball in E with $\Omega|_{B_\gamma} = 0$ whose boundary

$$V_\gamma = \partial B_\gamma = B_\gamma \cap E_{\gamma(a)}$$

is a Lagrangian sphere in $(E_{\gamma(a)}, \Omega|_{E_{\gamma(a)}})$ that comes with a natural framing (see Remark 2.4). We call V_γ the *vanishing cycle* associated to γ . (See Figure 2). If γ' is path-homotopic to γ in $\mathbb{C}\mathbb{P}^1 \setminus \pi(E_{\text{crit}})$, then $V_{\gamma'}$ with its natural framing is symplectically isotopic to V_γ . Hence, for each $z_{\text{crit}} \in E_{\text{crit}}$ and $t \in \mathbb{C}\mathbb{P}^1 \setminus \pi(E_{\text{crit}})$, the symplectic isotopy class of V_γ with its natural framing depends only on the path-homotopy class.

Let (E, π, J, Ω) be a Lefschetz fibration and $\gamma : [a, b] \rightarrow \mathbb{C}\mathbb{P}^1$ an embedded path that avoids $\pi(E_{\text{crit}})$ except at the endpoint $\gamma(b) = \pi(z_{\text{crit}}) \in \pi(E_{\text{crit}})$. We say that a loop $\ell : [c, d] \rightarrow \mathbb{C}\mathbb{P}^1 \setminus \pi(E_{\text{crit}})$ *doubles* γ if $\ell(c) = \ell(d) = \gamma(a)$, ℓ is positively oriented with respect to the standard orientation of $\mathbb{C}\mathbb{P}^1$ and ℓ circles the point




FIGURE 2. The vanishing cycle

$\pi(z_{\text{crit}})$ exactly once and circles no other critical values of π . The following result is the main result of this section and is described in [Sei03, Proposition 1.15].

Proposition 2.7. *Let (E, π, J, Ω) be a Lefschetz fibration and $V_\gamma \subset E_{t_0}$ the vanishing cycle of a path $\gamma : [a, b] \rightarrow \mathbb{CP}^1$ with $\gamma(a) = t_0$ and $\gamma(b) = \pi(z_{\text{crit}}) \in E_{\text{crit}}$. Let ℓ be a loop that doubles γ . The monodromy $\rho_\ell : E_{t_0} \rightarrow E_{t_0}$ around ℓ is symplectically isotopic to the Dehn twist τ_{V_γ} along the vanishing cycle V_γ .*

Sketch proof. We deform the form Ω to a closed 2-form Ω^1 which is standard with respect to the Morse coordinates at each critical point. We can choose this deformation such that the monodromy of the resulting Lefschetz fibration agrees with that of (E, π, J, Ω) up to symplectic isotopy. In a neighborhood of each critical point, the Lefschetz fibration (E, π, J, Ω^1) agrees with a neighborhood of 0 in the standard model $\pi_{\text{std}} : \mathbb{C}^N \rightarrow \mathbb{C}$ described in Section 2.1. Here the vanishing cycles correspond to the Σ_t 's defined in (2.3). We can therefore use Lemma 2.2 to show the result. Details can be found in [Dup07, Proposition 2.2.3]. \square

3. THE LEFSCHETZ FIBRATION $\pi : \mathcal{X} \rightarrow \mathbb{CP}^1$

This section contains the essential steps in the proofs of the main results of this paper. In the first two sections we construct the Lefschetz fibration $(\mathcal{X}, \pi, J, \Omega^\mu)$ of Theorem 1.3. Its total space \mathcal{X} is a complex hypersurface in $\mathbb{F} \times \mathbb{CP}^1$, where \mathbb{F} is a \mathbb{CP}^{n+1} -bundle over \mathbb{CP}^{n+1} . The manifold \mathcal{X} fibers over \mathbb{CP}^1 as a subbundle of the bundle $\mathbb{F} \times \mathbb{CP}^1 \rightarrow \mathbb{CP}^1$. For each $\mu > 1$, the closed 2-form Ω^μ on \mathcal{X} comes from a toric symplectic structure on \mathbb{F} , and the symplectic isotopy class of the non-singular fibers of $(\mathcal{X}, \pi, J, \Omega^\mu)$ depends on the cohomology class of this toric structure. For each $\mu > 1$, it is possible to choose a toric structure $\omega_{\mathbb{F}}^{(1, \mu)}$ on \mathbb{F} such that the non-singular fibers of $(\mathcal{X}, \pi, J, \Omega^\mu)$ are isotopic to $(\mathbb{CP}^n \times \mathbb{CP}^{n+1}, \sigma_\mu)$. By symplectically embedding a large part of $(\mathcal{X}, \Omega^\mu)$ into the toric manifold $(\mathbb{F}, \omega_{\mathbb{F}}^{(1, \mu)})$, and using a Darboux chart on \mathbb{F} , we show in Section 3.3 that the vanishing cycle \mathcal{V}_∞ in the fiber at infinity is the Lagrangian L^μ defined in (1.1).

3.1. **The toric manifold** $(\mathbb{F}, \omega_{\mathbb{F}}^{(1, \mu)})$. Let \mathbb{C}^{2n+4} have coordinates $(\underline{s}, q, \underline{x}, x_{n+1}) = (s_0, \dots, s_n, q, x_0, \dots, x_{n+1})$ and consider the $(\mathbb{C}^*)^2$ action on \mathbb{C}^{2n+4} with weights

$$(3.1) \quad \begin{pmatrix} \overbrace{1 \ \cdots \ 1}^{n+1} & 1 & \overbrace{0 \ \cdots \ 0}^{n+2} \\ 0 & \cdots & 0 & 1 & 1 & \cdots & 1 \end{pmatrix}.$$

We define \mathbb{F} to be the quotient

$$\mathbb{F} = (\mathbb{C}^{n+2} \setminus \{0\} \times \mathbb{C}^{n+2} \setminus \{0\}) / (\mathbb{C}^*)^2.$$

Since the action of $(\mathbb{C}^*)^2$ on $\mathbb{C}^{n+2} \setminus \{0\} \times \mathbb{C}^{n+2} \setminus \{0\}$ is free and proper, this quotient is a smooth complex manifold.² We denote the points of \mathbb{F} by $[[\underline{s}, q, \underline{x}, x_{n+1}]]$, where $(\underline{s}, q, \underline{x}, x_{n+1}) \in \mathbb{C}^{n+2} \setminus \{0\} \times \mathbb{C}^{n+2} \setminus \{0\}$ and

$$[[\underline{s}, q, \underline{x}, x_{n+1}]] = [[\alpha \underline{s}, \alpha \beta q, \beta \underline{x}, \beta x_{n+1}]] \text{ for } (\alpha, \beta) \in (\mathbb{C}^*)^2$$

The map $\mathbb{F} \rightarrow \mathbb{C}\mathbb{P}^{n+1}$ given by $[[\underline{s}, q, \underline{x}, x_{n+1}]] \mapsto [x_0 : \dots : x_{n+1}]$ is clearly well-defined and makes \mathbb{F} a $\mathbb{C}\mathbb{P}^{n+1}$ bundle over $\mathbb{C}\mathbb{P}^{n+1}$.

We now describe a Kähler structure on \mathbb{F} by constructing it as a symplectic quotient. Let \mathbb{C}^{2n+4} have the standard symplectic form $\omega_{\mathbb{C}^{2n+4}}$ and suppose T^2 acts on \mathbb{C}^{2n+4} with weights (3.1). The moment map $\Psi : \mathbb{C}^{2n+4} \rightarrow (\mathfrak{t}^2)^* \cong \mathbb{R}^2$ of the action is given by

$$\Psi(\underline{s}, q, \underline{x}, x_{n+1}) = \left(\sum_{j=0}^n |s_j|^2 + |q|^2, |q|^2 + \sum_{j=0}^{n+1} |x_j|^2 \right).$$

Its image is the cone illustrated in Figure 3 and the regular values are the interiors of the cones C_1 and C_2 . By the standard symplectic quotient construction [MS98,




FIGURE 3. The image of Ψ

Section 5.4], for each $\kappa \in C_1$, the form $\omega_{\mathbb{C}^{2n+4}}|_{\Psi^{-1}(\kappa)}$ descends to a symplectic form on the quotient manifold $\Psi^{-1}(\kappa)/T^2$. The elements of $\Psi^{-1}(\kappa)/T^2$ are denoted $[[\underline{s}, q, \underline{x}, x_{n+1}]]$ where $(\underline{s}, q, \underline{x}, x_{n+1}) \in \Psi^{-1}(\kappa)$ and

$$[[\underline{s}, q, \underline{x}, x_{n+1}]] = [e^{i\theta_1} \underline{s}, e^{i(\theta_1 + \theta_2)} q, e^{i\theta_2} \underline{x}, e^{i\theta_2} x_{n+1}] \text{ for } (e^{i\theta_1}, e^{i\theta_2}) \in T^2.$$

² \mathbb{F} is the GIT quotient $\mathbb{F} = \mathbb{C}^{2n+4} //_{\kappa} (\mathbb{C}^*)^2$ where κ is an integral point of the chamber C_1 of the effective cone depicted in Figure 3. The effective cone is the cone in \mathbb{R}^2 generated by the columns of the matrix (3.1). It decomposes naturally into a union of 2-dimensional cones, each generated by a pair of column vectors. The interior of these cones are the chambers C_1 and C_2 . As a complex manifold, the GIT quotient is the same for different choices of κ in the same chamber [Tha94, Theorem 3.9].

By [Aud04, Theorem VII.2.1], as a complex manifold, the quotient $\Psi^{-1}(\kappa)/T^2$ equals \mathbb{F} . In fact, the map $\Psi^{-1}(\kappa)/T^2 \rightarrow \mathbb{F}$ given by

$$[\underline{s}, q, \underline{x}, x_{n+1}] \mapsto [[\underline{s}, q, \underline{x}, x_{n+1}]]$$

is biholomorphic and the quotient symplectic form is a Kähler structure $\omega_{\mathbb{F}}^{\kappa}$ on \mathbb{F} .

Remark 3.1. Points of \mathbb{F} are denoted $[\underline{s}, q, \underline{x}, x_{n+1}]$, when \mathbb{F} is thought of as a symplectic quotient $\Psi^{-1}(\kappa)/T^2$ and by $[[\underline{s}, q, \underline{x}, x_{n+1}]]$, when we are interested in the complex structure on \mathbb{F} . \square

Lemma 3.2. *The manifold \mathbb{F} is a $\mathbb{C}\mathbb{P}^{n+1}$ bundle over $\mathbb{C}\mathbb{P}^{n+1}$ with fiber coordinates $[s_0 : \cdots : s_n : q]$ and base coordinates $[x_0 : \cdots : x_{n+1}]$. For each $\kappa = (\kappa_1, \kappa_2) \in C_1$, the form $\omega_{\mathbb{F}}^{\kappa}$ integrates to $\kappa_1\pi$ on a line in the fiber and to $\kappa_2\pi$ on a line in the section $s_0 = \cdots = s_{n-1} = q = 0$. Moreover, the submanifold $\mathbb{F} \cap (q = 0)$ is isomorphic to the product $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ and the restriction $\omega_{\mathbb{F}}^{\kappa}|_{\mathbb{F} \cap (q=0)}$ is the product form $\kappa_1\sigma_{\mathbb{C}\mathbb{P}^n} + \kappa_2\sigma_{\mathbb{C}\mathbb{P}^{n+1}}$.*

Proof. We have already seen the bundle structure of \mathbb{F} .

To see the statement about $\mathbb{F} \cap (q = 0)$, note that $\mathbb{F} \cap (q = 0)$ can be thought of as $(\Psi|_{(q=0)})^{-1}(\kappa)/T^2$ which is by construction $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ with symplectic form $\kappa_1\sigma_{\mathbb{C}\mathbb{P}^n} + \kappa_2\sigma_{\mathbb{C}\mathbb{P}^{n+1}}$. More precisely, the map

$$(\mathbb{F} \cap (q = 0), \omega_{\mathbb{F}}^{\kappa}|_{(q=0)}) \rightarrow (\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \kappa_1\sigma_{\mathbb{C}\mathbb{P}^n} + \kappa_2\sigma_{\mathbb{C}\mathbb{P}^{n+1}})$$

given by

$$[\underline{s}, 0, \underline{x}, x_{n+1}] \mapsto ([s_0 : \cdots : s_n], [x_0 : \cdots : x_{n+1}])$$

is a biholomorphic symplectomorphism.

The line $(q = s_2 = \cdots = s_n = 0$ and $\underline{x} = 0)$ in the fiber $(\underline{x} = 0)$ lies in $(\mathbb{F} \cap (q = 0), \omega_{\mathbb{F}}^{\kappa}|_{(q=0)})$ and corresponds to

$$\{([s_0 : s_1 : 0 : \cdots : 0], [0 : \cdots : 0 : 1]) \in \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}\}.$$

Similarly, the line $(q = s_0 = \cdots = s_{n-1} = x_0 = \cdots = x_{n-1})$ in the section $(q = s_0 = \cdots = s_{n-1} = 0)$ corresponds to

$$\{([0 : \cdots : 0 : 1], [0 : \cdots : 0 : x_n : x_{n+1}]) \in \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}\}.$$

It follows that $\omega_{\mathbb{F}}^{\kappa}$ integrates to $\kappa_1\pi$ on $(q = s_2 = \cdots = s_n = 0$ and $\underline{x} = 0)$ and to $\kappa_2\pi$ on $(q = s_0 = \cdots = s_{n-1} = x_0 = \cdots = x_{n-1} = 0)$. \square

Remark 3.3. The manifold $(\mathbb{F}, \omega_{\mathbb{F}}^{\kappa})$ admits an action by the quotient torus $T^{2n+4}/T^2 \cong T^{2n+2}$. Hence it can be represented by a Delzant polytope Δ^{κ} of dimension $2n + 2$. If $p : \mathbb{R}^{2n+4} \rightarrow \mathbb{R}^2$ is the linear map given by the matrix (3.1), then

$$(3.2) \quad \begin{aligned} \Delta^{\kappa} &= \mathbb{R}_+^{2n+4} \cap p^{-1}(\kappa) \\ &= \left\{ (\underline{\xi}, \underline{\nu}, \underline{\eta}, \eta_{n+1}) \in \mathbb{R}_+^{2n+4} \mid \nu + \sum_{j=0}^n \xi_j = \kappa_1, \nu + \sum_{j=0}^{n+1} \eta_j = \kappa_2 \right\} \end{aligned}$$

where $\underline{\xi} = (\xi_0, \dots, \xi_n)$, $\underline{\eta} = (\eta_0, \dots, \eta_n)$ and $\kappa = (\kappa_1, \kappa_2)$. Let $A : \mathbb{C}^{2n+4} \rightarrow \mathbb{R}^{2n+4}$ be the map

$$A(\underline{s}, q, \underline{x}, x_{n+1}) = (|s_0|^2, \dots, |s_n|^2, |q|^2, |x_0|^2, \dots, |x_{n+1}|^2).$$

Then $\Psi^{-1}(\kappa) = A^{-1}(\Delta^{\kappa})$. Suppose we have a d -dimensional face of Δ^{κ} . Then it is given by the vanishing of $2n + 4 - d$ coordinates in \mathbb{R}^{2n+4} . The vanishing of the corresponding coordinates in \mathbb{C}^{2n+4} is a T^2 -invariant complex submanifold of $\Psi^{-1}(\kappa)$ and corresponds to a d -dimensional complex submanifold of $\mathbb{F} = \Psi^{-1}(\kappa)/T^2$. Two faces of Δ^{κ} intersect precisely when the corresponding submanifolds intersect in \mathbb{F} .

For $n = 1$ the polytope representing $(\mathbb{F}, \omega_{\mathbb{F}}^{\kappa})$ is illustrated in Figure 4. The polytope that represents $(\mathbb{F} \cap (x_0 = 0), \omega_{\mathbb{F}}^{\kappa}|_{x_0=0})$ can be illustrated in a 3-dimensional subspace of \mathbb{R}^6 as seen on the left hand side of Figure 4. This is a $\mathbb{C}\mathbb{P}^2$ -bundle over $\mathbb{C}\mathbb{P}^1$ with base coordinates $[x_1 : x_2]$ and fiber coordinates $[s_0 : s_1 : q]$. We get the polytope associated to $(\mathbb{F}, \omega_{\mathbb{F}}^{\kappa})$ by replacing each section $\mathbb{C}\mathbb{P}^1$ by a $\mathbb{C}\mathbb{P}^2$ as shown in the right hand side of the figure. \square




FIGURE 4. Polytope Δ^{κ} representing $(\mathbb{F}, \omega_{\mathbb{F}}^{\kappa})$ for $n = 1$

3.2. The Lefschetz fibration $\pi : \mathcal{X} \rightarrow \mathbb{C}\mathbb{P}^1$. Let

$$\mathcal{X} = \{([\underline{s}, q, \underline{x}, x_{n+1}], [t_0 : t_1]) \in \mathbb{F} \times \mathbb{C}\mathbb{P}^1 \mid t_1(s_0x_0 + \cdots + s_nx_n) = t_0q\}.$$

The holomorphic equation $t_1(s_0x_0 + \cdots + s_nx_n) = t_0q$ is invariant under the $(\mathbb{C}^*)^2$ -action on \mathbb{C}^{2n+4} and \mathcal{X} is a complex submanifold of $\mathbb{F} \times \mathbb{C}\mathbb{P}^1$. Let $t = \frac{t_0}{t_1}$ (where we write ∞ to mean the point $[1 : 0] \in \mathbb{C}\mathbb{P}^1$) and let

$$X_t = \{([\underline{s}, q, \underline{x}, x_{n+1}]) \in \mathbb{F} \mid t_1(s_0x_0 + \cdots + s_nx_n) = t_0q\}$$


and $\mathcal{X}_t = X_t \times \{t\}$. Note that for $t_1 = 0$ we get

$$X_{\infty} = \mathbb{F} \cap (q = 0).$$

Let $\pi : \mathcal{X} \rightarrow \mathbb{C}\mathbb{P}^1$ be the natural projection map. Figure 5 shows how the submanifolds X_t lie inside the polytope representing $(\mathbb{F} \cap (x_0 = 0), \omega_{\mathbb{F}}^{\kappa}|_{x_0=0})$ when $n = 1$.

Consider the submanifold

$$(3.3) \quad \mathbb{S} = \{([\underline{s}, q, \underline{x}, x_{n+1}]) \in \mathbb{F} \mid s_0x_0 + \cdots + s_nx_n = 0 \text{ and } q = 0\}$$

FIGURE 5. X_t in $\mathbb{F} \cap (x_0 = 0)$

in \mathbb{F} . We see that

$$\mathbb{S} = \bigcap_{t \in \mathbb{C}\mathbb{P}^1} X_t$$

and moreover, when \mathbb{S} is removed from the X_t 's, they are disjoint in \mathbb{F} and

$$\mathbb{F} = \mathbb{S} \sqcup \bigsqcup_{t \in \mathbb{C}} X_t \setminus \mathbb{S}.$$

More precisely, if $\pi_{\mathbb{F}} : \mathbb{F} \times \mathbb{C}\mathbb{P}^1 \rightarrow \mathbb{F}$ denotes projection to the first factor, then

$$(3.4) \quad \pi_{\mathbb{F}}|_{\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1)} : \mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1) \rightarrow \mathbb{F} \setminus \mathbb{S}$$

has inverse $(\pi_{\mathbb{F}}|_{\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1)})^{-1} : \mathbb{F} \setminus \mathbb{S} \rightarrow \mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1)$ given by

$$[[\underline{s}, q, \underline{x}, x_{n+1}]] \mapsto ([[s, q, \underline{x}, x_{n+1}], [s_0 x_0 + \cdots + s_n x_n : q]]).$$

In particular, $\pi_{\mathbb{F}}|_{\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1)}$ is biholomorphic.

Lemma 3.4. *The singular fibration $\pi : \mathcal{X} \rightarrow \mathbb{C}\mathbb{P}^1$ is holomorphic and has exactly one critical point at*

$$z_{\text{crit}} = ([[0, 1, 0, 1]], 0) \in \mathcal{X}_0.$$

In a neighborhood of z_{crit} we obtain holomorphic coordinates (z_1, \dots, z_{2n+2}) such that π is the map $(z_1, \dots, z_{2n+2}) \mapsto \sum_{j=1}^{2n+2} z_j^2$.

Moreover, when the singular fiber \mathcal{X}_0 is removed, $\pi : \mathcal{X} \rightarrow \mathbb{C}\mathbb{P}^1$ is trivialized by a biholomorphic map $\Phi : \mathcal{X} \setminus \mathcal{X}_0 \rightarrow (\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}) \times \mathbb{C}$.

Proof. Clearly, $\pi : \mathcal{X} \rightarrow \mathbb{C}$ is a holomorphic fibration with exactly one critical point z_{crit} .

For $q \neq 0$ and $x_{n+1} \neq 0$,

$$\tilde{s}_i = \frac{s_i x_{n+1}}{q}, \quad \tilde{x}_j = \frac{x_j}{x_{n+1}} \quad \text{for } 0 \leq i, j \leq n$$

are holomorphic and invariant under the $(\mathbb{C}^*)^2$ -action. The map

$$\mathbb{F} \cap (q \neq 0, x_{n+1} \neq 0) \rightarrow \mathbb{C}^{2n+2}$$

given by

$$[[\underline{s}, q, \underline{x}, x_{n+1}]] \mapsto (\tilde{s}_0, \dots, \tilde{s}_n, \tilde{x}_0, \dots, \tilde{x}_n)$$

defines a holomorphic coordinate chart on \mathbb{F} . Thus, $(\tilde{s}_0, \dots, \tilde{s}_n, \tilde{x}_0, \dots, \tilde{x}_n, t)$ define holomorphic coordinates on $\mathbb{F} \times \mathbb{C}$ in which \mathcal{X} is given by $\tilde{s}_0 \tilde{x}_0 + \dots + \tilde{s}_n \tilde{x}_n = t$. Therefore $(\tilde{s}_0, \dots, \tilde{s}_n, \tilde{x}_0, \dots, \tilde{x}_n)$ define holomorphic coordinates on \mathcal{X} . Let

$$(3.5) \quad \begin{aligned} z_1 &= \frac{1}{2}(\tilde{s}_0 + \tilde{x}_0), \dots, z_{n+1} = \frac{1}{2}(\tilde{s}_n + \tilde{x}_n), \\ z_{n+2} &= \frac{1}{2i}(\tilde{s}_0 - \tilde{x}_0), \dots, z_{2n+2} = \frac{1}{2i}(\tilde{s}_n - \tilde{x}_n). \end{aligned}$$

Then (z_1, \dots, z_{2n+2}) define holomorphic coordinates on \mathcal{X} in which π has the form $(z_1, \dots, z_{2n+2}) \mapsto \sum_{j=1}^{2n+2} z_j^2$.

Let \mathcal{X}_t be a non-singular fiber of $\pi : \mathcal{X} \rightarrow \mathbb{C}\mathbb{P}^1$, i.e., fix $t \neq 0$. Then X_t is a complex submanifold of \mathbb{F} in which q is determined by the other coordinates of \mathbb{F} as $q = \frac{s_0 x_0 + \dots + s_n x_n}{t}$. Since $\mathbb{F} = (\mathbb{C}^{n+2} \setminus \{0\} \times \mathbb{C}^{n+2} \setminus \{0\}) / (\mathbb{C}^*)^2$, an element $[[\underline{s}, q, \underline{x}, x_{n+1}]] \in X_t$ cannot have $\underline{s} = 0$ or $\underline{x} = x_{n+1} = 0$. It follows that the map $\Phi_t : \mathcal{X}_t \rightarrow \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ given by

$$(3.6) \quad \Phi_t : ([[\underline{s}, q, \underline{x}, x_{n+1}]], t) \mapsto ([s_0 : \dots : s_n], [x_0 : \dots : x_{n+1}])$$

is well-defined. In fact, it is biholomorphic. Now define

$$\Phi : \mathcal{X} \setminus \mathcal{X}_0 \rightarrow (\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}) \times \mathbb{C}$$

by

$$(3.7) \quad \Phi([[\underline{s}, q, \underline{x}, x_{n+1}]], t) = (\Phi_{\frac{1}{t}}([[\underline{s}, q, \underline{x}, x_{n+1}]], \frac{1}{t}), \frac{1}{t}) \quad \text{for } t \in \mathbb{C}\mathbb{P}^1 \setminus \{0\}.$$

□

Remark 3.5. In the notation of Remark 3.1,

$$z_{\text{crit}} = ([\underline{0}, \kappa_1, \underline{0}, \kappa_2 - \kappa_1], 0).$$

In the polytope $\Delta^\kappa \subset \mathbb{R}^{2n+4}$ representing $(\mathbb{F}, \omega_{\mathbb{F}}^\kappa)$, $[\underline{0}, \kappa_1, \underline{0}, \kappa_2 - \kappa_1] \in \mathbb{F}$ corresponds to the vertex $v = (\underline{0}, \kappa_1, \underline{0}, \kappa_2 - \kappa_1)$. (See Figure 5 for the case $n = 1$). □

For each $\mu > 1$ we now introduce a 2-form Ω^μ on \mathcal{X} that makes (\mathcal{X}, π, J) into a Lefschetz fibration. Here J is the complex structure on \mathcal{X} coming from $\mathbb{F} \times \mathbb{C}\mathbb{P}^1$. Recall that if $\kappa = (\kappa_1, \kappa_2)$ lies in the cone C_1 , then we have a toric Kähler structure $\omega_{\mathbb{F}}^\kappa$ on \mathbb{F} . If $\mu > 1$, then $(1, \mu) \in C_1$ and we define

$$(3.8) \quad \Omega^\mu = (\pi_{\mathbb{F}}|_{\mathcal{X}})^*(\omega_{\mathbb{F}}^{(1, \mu)}).$$

Proposition 3.6. *For any $\mu > 1$, $(\mathcal{X}, \pi, J, \Omega^\mu)$ is a Lefschetz fibration whose non-singular fibers are symplectically isotopic to $(\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$. Moreover, the map*

$$\Phi_\infty : (\mathcal{X}_\infty, \Omega^\mu|_{\mathcal{X}_\infty}) \rightarrow (\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$$

described by (3.7) is a symplectomorphism.

Proof. Let $\mu > 1$. First we prove that $(\mathcal{X}, \pi, J, \Omega^\mu)$ is a Lefschetz fibration. Clearly, Ω^μ is closed. Its restriction to each non-singular fiber is

$$\Omega^\mu|_{\mathcal{X}_t} = \omega_{\mathbb{F}}^{(1, \mu)}|_{\mathcal{X}_t},$$

which is symplectic since all the X_t 's are complex submanifolds of \mathbb{F} for $t \neq 0$. We have already shown in Lemma 3.4 that π has the form $(z_1, \dots, z_{2n+2}) \mapsto \sum_{j=1}^{2n+2} z_j^2$ in a neighborhood of the critical point z_{crit} . By the definition of Ω^μ , the map $\pi_{\mathbb{F}}|_{\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1)}$ described in (3.4) is a biholomorphic map that identifies $(\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1), \Omega^\mu)$ and $(\mathbb{F} \setminus \mathbb{S}, \omega_{\mathbb{F}}^{(1, \mu)})$. Hence Ω^μ is J -Kähler in a neighborhood of the critical point. We conclude that $(\mathcal{X}, \pi, J, \Omega^\mu)$ is a Lefschetz fibration.

It remains to show that the non-singular fibers of $(\mathcal{X}, \pi, J, \Omega^\mu)$ are symplectically isotopic to $(\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$, i.e., that the forms

$$\alpha_t = (\Phi_t^{-1})^*(\omega_{\mathbb{F}}^{(1, \mu)}|_{X_t}) \quad \text{for } t \neq 0$$

are symplectically isotopic to σ_μ for all $t \neq 0$. Since all the α_t are symplectically isotopic, it suffices to prove it for $t = \infty$. But we have already seen in Lemma 3.2 that $\alpha_\infty = \sigma_\mu$ by construction. \square

3.3. The vanishing cycle of $(\mathcal{X}, \pi, J, \Omega^\mu)$. In order to compute the vanishing cycle of $(\mathcal{X}, \pi, J, \Omega^\mu)$, we need to understand the connection coming from the 2-form Ω^μ in a neighborhood of the node z_{crit} . By (3.8), the map $\pi_{\mathbb{F}}|_{\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1)}$ described in (3.4) identifies $(\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1), \Omega^\mu)$ with $(\mathbb{F} \setminus \mathbb{S}, \omega_{\mathbb{F}}^{(1, \mu)})$. Hence Ω^μ is symplectic on $\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1)$ and we can use the toric structure of \mathbb{F} to describe the following Darboux chart:

Lemma 3.7. *Let*

$$W^\mu = \{(\underline{s}, \underline{x}) \in \mathbb{C}^{2n+2} \mid |\underline{s}|^2 < 1, -|\underline{s}|^2 + |\underline{x}|^2 < \mu - 1\},$$

where $(\underline{s}, \underline{x}) = (s_0, \dots, s_n, x_0, \dots, x_n)$, $|\underline{s}|^2 = \sum_j |s_j|^2$ and $|\underline{x}|^2 = \sum_j |x_j|^2$.

The map

$$\psi_{\mathcal{X}}^\mu : W^\mu \rightarrow \mathcal{X} \cap (q \neq 0) \cap (x_{n+1} \neq 0)$$

given by

$$(\underline{s}, \underline{x}) \mapsto \left([\underline{s}, \sqrt{1 - |\underline{s}|^2}, \underline{x}, \sqrt{\mu - 1 + |\underline{s}|^2 - |\underline{x}|^2}], \frac{s_0 x_0 + \dots + s_n x_n}{\sqrt{1 - |\underline{s}|^2}} \right)$$

defines a Darboux chart on $(\mathcal{X} \setminus (\mathbb{S} \times \mathbb{C}\mathbb{P}^1), \Omega^\mu)$ such that $\psi_{\mathcal{X}}^\mu(\underline{0}, \underline{0}) = z_{\text{crit}}$.

Proof. As noted in the Remark 3.5, $z_{\text{crit}} = (z, 0) \in \mathbb{F} \times \mathbb{C}\mathbb{P}^1$ where

$$z = [[\underline{0}, 1, \underline{0}, 1]] = [\underline{0}, 1, \underline{0}, \sqrt{\mu - 1}].$$

By the comments preceding this lemma, it suffices to see that the map $\psi_{\mathbb{F}}^\mu : W^\mu \rightarrow \mathbb{F}$, where

$$\psi_{\mathbb{F}}^\mu(\underline{s}, \underline{x}) = [\underline{s}, \sqrt{1 - |\underline{s}|^2}, \underline{x}, \sqrt{\mu - 1 + |\underline{s}|^2 - |\underline{x}|^2}],$$

is a Darboux chart on $(\mathbb{F} \cap (q \neq 0) \cap (x_{n+1} \neq 0), \omega_{\mathbb{F}}^{(1, \mu)})$ centered at z .

We have the following commutative diagram

$$\begin{array}{ccc} W^\mu & \xrightarrow{\tilde{\psi}^\mu} & \Psi^{-1}((1, \mu)) \subset \mathbb{C}^{2n+4} \\ & \searrow \psi_{\mathbb{F}}^\mu & \downarrow \\ & & \mathbb{F} = \Psi^{-1}((1, \mu))/T^2, \end{array}$$

where $\tilde{\psi}^\mu$ is given by $(\underline{s}, \underline{x}) \mapsto (\underline{s}, \sqrt{1 - |\underline{s}|^2}, \underline{x}, \sqrt{\mu - 1 + |\underline{s}|^2 - |\underline{x}|^2})$. Hence it suffices to see that $\tilde{\psi}^\mu$ pulls back $\omega_{\mathbb{C}^{2n+4}}$ to $\omega_{\mathbb{C}^{2n+2}}$. But this is clear since $q, x_{n+1} \in \mathbb{R}$ on $\text{Im}(\tilde{\psi}^\mu)$, and therefore $dq \wedge d\bar{q}$ and $dx_{n+1} \wedge d\bar{x}_{n+1}$ pull back to 0 under $\tilde{\psi}^\mu$. \square

Remark 3.8. Each vertex of the polytope $\Delta^{(1, \mu)}$ gives a natural Darboux chart on $(\mathbb{F}, \omega_{\mathbb{R}}^{(1, \mu)})$, and the chart described in Lemma 3.7 is the one corresponding to $v = (\underline{0}, 1, \underline{0}, \mu - 1)$. By (3.2), an element $(\underline{\xi}, \nu, \underline{\eta}, \eta_{n+1})$ of $\Delta^{(1, \mu)}$ satisfies the equations

$$\nu = 1 - \sum_{j=0}^n \xi_j, \quad \eta_{n+1} = \mu - 1 + \sum_{j=0}^n (\xi_j - \eta_j).$$

Hence the projection $(\underline{\xi}, \nu, \underline{\eta}, \eta_{n+1}) \mapsto (\underline{\xi}, \underline{\eta})$ maps $\Delta^{(1, \mu)}$ isomorphically to a polytope $\tilde{\Delta}^{(1, \mu)}$ in \mathbb{R}^{2n+2} . Here the vertex v is mapped to 0. Let $F^{q, x_{n+1}}$ denote the union of the facets $\nu = 0$ and $\eta_{n+1} = 0$ in $\Delta^{(1, \mu)}$ and let $\tilde{F}^{q, x_{n+1}}$ be the corresponding facets in $\tilde{\Delta}^{(1, \mu)}$. If $\tilde{A} : \mathbb{C}^{2n+2} \rightarrow \mathbb{R}^{2n+2}$ is the map $(\underline{s}, \underline{x}) \mapsto (|s_0|^2, \dots, |s_n|^2, |x_0|^2, \dots, |x_n|^2)$, then $W^\mu = \tilde{A}^{-1}(\tilde{\Delta}^{(1, \mu)} \setminus \tilde{F}^{q, x_{n+1}})$ and we have the following commutative diagram:

$$\begin{array}{ccccc} \mathbb{C}^{2n+2} \supset & & W^\mu \xrightarrow{\tilde{\psi}^\mu} & A^{-1}(\Delta^{(1, \mu)} \setminus F^{q, x_{n+1}}) & \subset \mathbb{C}^{2n+4} \\ & & \downarrow \tilde{A} & \downarrow A & \\ \mathbb{R}^{2n+2} \supset & & \tilde{\Delta}^{(1, \mu)} \setminus \tilde{F}^{q, x_{n+1}} \xrightarrow{\cong} & \Delta^{(1, \mu)} \setminus F^{q, x_{n+1}} & \subset \mathbb{R}^{2n+4}, \end{array}$$

where $\tilde{\psi}^\mu$ is the map described in the proof of Lemma 3.7. \square

By Lemma 3.7, we have a commutative diagram of singular fibrations:

$$\begin{array}{ccc} (W^\mu, \omega_{\mathbb{C}^{2n+2}}) & \xrightarrow[\cong]{\psi_{\mathcal{X}}^\mu} & (\mathcal{X} \cap (q \neq 0) \cap (x_{n+1} \neq 0), \Omega^\mu) \\ & \searrow \pi_{\text{loc}} & \swarrow \pi \\ & \mathbb{C} & \end{array}$$

where

$$\pi_{\text{loc}}(\underline{s}, \underline{x}) = \frac{s_0 x_0 + \dots + s_n x_n}{\sqrt{1 - |\underline{s}|^2}}.$$

The fiber $W_t^\mu = \pi_{\text{loc}}^{-1}(t)$ in W^μ is given by

$$W_t^\mu = \{(\underline{s}, \underline{x}) \in W^\mu \mid f_1(\underline{s}, \underline{x}) = f_2(\underline{s}, \underline{x}) = 0\},$$

where

$$\begin{aligned} f_1(\underline{s}, \underline{x}) &= \sum_{j=0}^n (\Re(s_j) \Re(x_j) - \Im(s_j) \Im(x_j)) - \Re(t) \sqrt{1 - |\underline{s}|^2}, \\ f_2(\underline{s}, \underline{x}) &= \sum_{j=0}^n (\Im(s_j) \Re(x_j) + \Re(s_j) \Im(x_j)) - \Im(t) \sqrt{1 - |\underline{s}|^2}. \end{aligned}$$

Lemma 3.9. Consider the fibration $\pi_{\text{loc}} = \pi \circ \psi_{\mathcal{X}}^\mu : W^\mu \rightarrow \mathbb{C}$ with connection coming from the standard symplectic form $\omega_{\mathbb{C}^{2n+2}}$ on W^μ . Let $t > 0$. If $(\underline{s}, \underline{x}) \in \pi^{-1}(t)$, then $(\underline{x}, \underline{s})$ lies in the horizontal space $(T_{(\underline{s}, \underline{x})} W_t^\mu)^{\omega_{\mathbb{R}^{4n+4}}}$.

Proof. The horizontal space at $(\underline{s}, \underline{x}) \in W_t^\mu$ is the symplectic complement of $T_{(\underline{s}, \underline{x})} W_t^\mu$ in $(\mathbb{C}^{2n+2}, \omega_{\mathbb{C}^{2n+2}}) \cong (\mathbb{R}^{4n+4}, \omega_{\mathbb{R}^{4n+4}})$. Note that

$$T_{(\underline{s}, \underline{x})} W_t^\mu = (\text{span } \nabla f_1(\underline{s}, \underline{x}))^\perp \cap (\text{span } \nabla f_2(\underline{s}, \underline{x}))^\perp$$

which has symplectic complement

$$(T_{(\underline{s}, \underline{x})} W_t^\mu)^{\omega_{\mathbb{R}^{4n+4}}} = \text{span}\{J_0 \nabla f_1(\underline{s}, \underline{x}), J_0 \nabla f_2(\underline{s}, \underline{x})\}.$$

Hence the horizontal space is spanned by the vectors

$$\begin{pmatrix} \Im(x_0) - \Re(t) \frac{\Im(s_0)}{\sqrt{1-|\underline{s}|^2}} \\ \Re(x_0) + \Re(t) \frac{\Re(s_0)}{\sqrt{1-|\underline{s}|^2}} \\ \vdots \\ \Im(x_n) - \Re(t) \frac{\Im(s_n)}{\sqrt{1-|\underline{s}|^2}} \\ \Re(x_n) + \Re(t) \frac{\Re(s_n)}{\sqrt{1-|\underline{s}|^2}} \\ \Im(s_0) \\ \Re(s_0) \\ \vdots \\ \Im(s_n) \\ \Re(s_n) \end{pmatrix}, \begin{pmatrix} -\Re(x_0) - \Im(t) \frac{\Re(s_0)}{\sqrt{1-|\underline{s}|^2}} \\ \Im(x_0) + \Im(t) \frac{\Im(s_0)}{\sqrt{1-|\underline{s}|^2}} \\ \vdots \\ -\Re(x_n) - \Im(t) \frac{\Re(s_n)}{\sqrt{1-|\underline{s}|^2}} \\ \Im(x_n) + \Im(t) \frac{\Im(s_n)}{\sqrt{1-|\underline{s}|^2}} \\ -\Re(s_0) \\ \Im(s_0) \\ \vdots \\ -\Re(s_n) \\ \Im(s_n) \end{pmatrix}.$$

□

Using this, we are able to compute the vanishing cycle of $(\mathcal{X}, \pi, J, \Omega^\mu)$.

Proposition 3.10. *Let $t > 0$. The vanishing cycle of $(\mathcal{X}, \pi, J, \Omega^\mu)$ in the fiber \mathcal{X}_t is isotopic to*

$$\mathcal{V}_t^\mu = \left\{ ([\underline{x}, \sqrt{1-g(t)}, \underline{x}, \sqrt{\mu-1}], t) \in \mathcal{X} \mid |\underline{x}|^2 = g(t) \right\}$$

where $\underline{x} = (\bar{x}_0, \dots, \bar{x}_n)$ and

$$(3.9) \quad g(t) = \frac{-t^2 + \sqrt{t^4 + 4t^2}}{2}.$$

The vanishing cycle in \mathcal{X}_∞ is isotopic to

$$\mathcal{V}_\infty^\mu = \left\{ ([\underline{x}, 0, \underline{x}, \sqrt{\mu-1}], \infty) \in \mathcal{X} \mid |\underline{x}|^2 = 1 \right\}.$$

Proof. For each $t > 0$, define

$$L_t^\mu = \{(\underline{x}, \underline{x}) \in W^\mu \mid |\underline{x}|^2 = g(t)\}.$$

The condition on $|\underline{x}|^2$ ensures that $L_t^\mu \subset W_t^\mu$. We claim that parallel transport along the path $\gamma(r) = r$ for $0 < r \leq t$ takes all of L_t^μ to the node $(\underline{0}, \underline{0}) \in W_0^\mu$. Since L_t^μ is diffeomorphic to S^{2n+1} and

$$\psi_{\mathcal{X}}^\mu(L_t^\mu) = \mathcal{V}_t^\mu,$$

this shows that \mathcal{V}_t^μ is the vanishing cycle in \mathcal{X}_t .

Fix $t > 0$ and let $\gamma_t(r) = r$ for $0 < r \leq t$. Given $(\underline{x}, \underline{x}) \in L_t^\mu$, define the path

$$\tilde{\gamma}_t(r) = h_t(r)(\underline{x}, \underline{x}), \quad 0 < r \leq t$$

where

$$h_t(r) = \sqrt{\frac{g(r)}{g(t)}}.$$

This is a lift of γ_t with endpoints $(\underline{x}, \underline{x})$ and $(\underline{0}, \underline{0})$ and tangents

$$\frac{d}{dr} \tilde{\gamma}_t(r) = h_t'(r)(\underline{x}, \underline{x}).$$

For all r , $\tilde{\gamma}_t(r) \in L_r^\mu$, and by Lemma 3.9, the horizontal space at $\tilde{\gamma}_t(r)$ contains $\frac{d}{dr}\tilde{\gamma}_t(r)$. Hence parallel transport along γ_t takes L_t^μ to the node. Since $\lim_{t \rightarrow \infty} g(t) = 1$,

$$\mathcal{V}_\infty^\mu = \mathcal{V}_{\gamma_\infty}^\mu = \lim_{t \rightarrow \infty} \mathcal{V}_{\gamma_t}^\mu = \left\{ ([\underline{x}, 0, \underline{x}, \sqrt{\mu-1}], \infty) \in \mathcal{X} \mid |\underline{x}|^2 = 1 \right\}.$$

□

4. PROOFS OF THE MAIN RESULTS

We are now in a position to prove the main results described in Section 1.

Proof of Theorem 1.3. Given $\mu > 1$, Proposition 3.6 shows that $(\mathcal{X}, \pi, J, \Omega^\mu)$ is a Lefschetz fibration with one critical point

$$z_{\text{crit}} = ([\underline{0}, 1, \underline{0}, \sqrt{\mu-1}], 0) \in \mathcal{X}_0.$$

The biholomorphic trivialization $\Phi : \mathcal{X} \setminus \mathcal{X}_0 \rightarrow (\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}) \times \mathbb{C}$ defined by (3.7) induces a symplectomorphism

$$\Phi_\infty : (\mathcal{X}_\infty, \Omega^\mu |_{\mathcal{X}_\infty}) \cong (\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}, \sigma_\mu)$$

and by Proposition 3.10, the vanishing cycle in \mathcal{X}_∞ is isotopic to

$$\mathcal{V}_\infty^\mu = \left\{ ([\underline{x}, 0, \underline{x}, \sqrt{\mu-1}], \infty) \in \mathcal{X} \mid |\underline{x}|^2 = 1 \right\}.$$

Under the symplectomorphism Φ_∞ this corresponds to

$$\Phi_\infty(\mathcal{V}_\infty^\mu) = \{([\bar{x}_0 : \cdots : \bar{x}_n], [x_0 : \cdots : x_n : \sqrt{\mu-1}]) \in \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1} \mid |\underline{x}|^2 = 1\},$$

which is exactly the Lagrangian L^μ in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$. □

Proof of Theorem 1.1. Given $\mu > 1$, we consider the Lefschetz fibration $(\mathcal{X}, \pi, J, \Omega^\mu)$ from Theorem 1.3. By Proposition 2.7, the monodromy $\rho_\ell : \mathcal{X}_\infty \rightarrow \mathcal{X}_\infty$ along a positively oriented loop ℓ in $\mathbb{C}\mathbb{P}^1$ based at $\infty \in \mathbb{C}\mathbb{P}^1$ and circling 0 exactly once is symplectically isotopic to the Dehn twist $\tau_{\mathcal{V}_\infty^\mu}$ along the vanishing cycle \mathcal{V}_∞^μ . Since the fibration has only one singular fiber, ρ_ℓ is symplectically isotopic to the identity through an isotopy that arises from deforming the loop ℓ to the constant loop based at infinity in $\mathbb{C}\mathbb{P}^1$. Hence we have an isotopy $(\varphi_\lambda)_{0 \leq \lambda \leq 1}$ of \mathcal{X}_∞ such that φ_1 equals $\tau_{\mathcal{V}_\infty^\mu}$ and φ_0 is the identity. Recall that each fiber of \mathcal{X} contains the hypersurface $\mathbb{S} \times \{t\}$, with \mathbb{S} defined in (3.3). Parallel transport in $(\mathcal{X}, \pi, J, \Omega^\mu)$ along a path in $\mathbb{C}\mathbb{P}^1$ from t' to t takes any point $(z, t) \in \mathbb{S} \times \{t\} \subset \mathcal{X}_t$ to $(z, t') \in \mathbb{S} \times \{t'\} \subset \mathcal{X}_{t'}$. Hence φ_λ restricts to the identity on $\mathbb{S} \times \{\infty\}$ for all λ . Observe that $S = \Phi_\infty(\mathbb{S} \times \{\infty\})$ and that by property 3 of Theorem 1.3

$$\tau_{L^\mu} = \Phi_\infty \circ \tau_{\mathcal{V}_\infty^\mu} \circ \Phi_\infty^{-1}.$$

Hence $\Phi_\infty \circ \varphi_\lambda \circ \Phi_\infty^{-1}$, $0 \leq \lambda \leq 1$, is the desired isotopy of τ_{L^μ} . □

Under the projection to the first factor, the complex hypersurface S in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ defined in Theorem 1.1 is a $\mathbb{C}\mathbb{P}^n$ -bundle over $\mathbb{C}\mathbb{P}^1$. The fibers are linearly embedded copies of $\mathbb{C}\mathbb{P}^n$ in $\mathbb{C}\mathbb{P}^{n+1}$. As before, let

$$\begin{aligned} S_0 &= \mathbb{C}\mathbb{P}^n \times \{[0 : \cdots : 0 : 1]\} \\ S_\infty &= \{([s_0 : \cdots : s_n], [x_0 : x_1 : 0 : \cdots : 0]) \mid s_0 x_0 + s_1 x_1 = 0\}; \end{aligned}$$

these are sections of S . If $n = 1$, S is a Hirzebruch surface, S_0 is the small section (with negative self-intersection), and S_∞ is the large section (with positive self-intersection). As stated in Corollary 1.2, if we symplectically blow up $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ along S_0 and S_∞ , then as long as the blow-ups are small, the isotopy of the Dehn twist τ_{L^μ} remains. On the other hand, Proposition 1.4 (which is a consequence of the

proof of [CS05, Proposition 2]) shows that an isotopy cannot fix the submanifolds S_0 and S'_∞ . Figure 6 illustrates how the submanifolds S_0 , S_∞ and S'_∞ lie inside the polytope representing $(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^2, \sigma_\mu)$.




FIGURE 6. Polytope representing $(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^2, \sigma_\mu)$

Before proving Corollary 1.2 and Proposition 1.4, we review the symplectic blow-up construction as described in [MS98, Chapter 7]. First we consider a blow-up at the origin in \mathbb{C}^N . Let \mathcal{L} be the tautological line bundle over $\mathbb{C}\mathbb{P}^{N-1}$, i.e., the total space of \mathcal{L} is given by

$$\mathcal{L} = \{(z_1, \dots, z_N), [w_1, \dots, w_N]\} \in \mathbb{C}^N \times \mathbb{C}\mathbb{P}^{N-1} \mid w_j z_k = w_k z_j \text{ for all } j, k\}.$$

Let $\text{pr}_{\mathbb{C}^N} : \mathcal{L} \rightarrow \mathbb{C}^N$ and $\text{pr}_{\mathbb{C}\mathbb{P}^{N-1}} : \mathcal{L} \rightarrow \mathbb{C}\mathbb{P}^{N-1}$ be the natural projections. For each $\lambda > 0$ the 2-form

$$\omega_{\mathcal{L}}^\lambda = \text{pr}_{\mathbb{C}^N}^*(\omega_{\mathbb{C}^N}) + \lambda^2 \text{pr}_{\mathbb{C}\mathbb{P}^{N-1}}^*(\sigma_{\mathbb{C}\mathbb{P}^{N-1}})$$

is symplectic. For all $\delta \geq 0$, let $\mathcal{L}(\delta) = \text{pr}_{\mathbb{C}^N}^{-1}(B(\delta))$, where $B(\delta)$ is the ball of radius δ in \mathbb{C}^N . By [MS98, Lemma 7.11], for all $\lambda, \delta > 0$ $(\mathcal{L}(\delta) - \mathcal{L}(0), \omega_{\mathcal{L}}^\lambda)$ is symplectomorphic to the spherical shell $(B(\sqrt{\lambda^2 + \delta^2}) - B(\lambda), \omega_{\mathbb{C}^N})$. The blow-up $\tilde{\mathbb{C}}^N$ of 0 in \mathbb{C}^N of size λ is obtained by replacing the interior of the ball $B(\sqrt{\lambda^2 + \delta^2})$ in \mathbb{C}^N with $\mathcal{L}(\delta)$. Thus, we have replaced the ball $B(\lambda)$ in \mathbb{C}^N with the zero section $\mathcal{L}(0) \cong \mathbb{C}\mathbb{P}^{N-1}$, the exceptional divisor in $\tilde{\mathbb{C}}^N$. The manifold $\tilde{\mathbb{C}}^N$ has a natural symplectic structure that is independent of the choice of $\delta > 0$.

Let Q be a compact symplectic submanifold of codimension $2N$ in a symplectic manifold (M, ω) . Then ω can be written in a standard form in a neighborhood of Q . Details can be found in [MS98, p. 249-251]. The normal bundle $\nu_{Q|M}$ of Q in M has the structure of a $2N$ -dimensional symplectic vector bundle with a compatible complex structure J . It is associated to a principal $U(N)$ -bundle P , i.e., $\nu_{Q|M} = P \times_{U(N)} \mathbb{C}^N$. This bundle has a symplectic form

$$(4.1) \quad \omega' = \alpha + \pi_Q^*(\omega|_Q),$$

where $\pi_Q : \nu_{Q|M} \rightarrow Q$ is the natural projection and α is a closed 2-form on $\nu_{Q|M}$ that restricts to the standard symplectic form on the fibers \mathbb{C}^N . A small neighborhood

of Q in (M, ω) is symplectomorphic to the disk bundle $\nu_{Q|M}^{<\varepsilon} = P \times_{U(N)} B(\varepsilon)$ with symplectic form ω' for some $\varepsilon > 0$. Using the standard form of the symplectic structure near Q , we can define small blow-ups along Q in M . If $\lambda < \varepsilon$, then the size λ blow-up of Q in M is simply the result of blowing up the origin in the fiber $B(\varepsilon)$, i.e. if $\tilde{B}(\varepsilon)$ is the size λ blow-up of the origin in $B(\varepsilon)$, then the size λ blow-up of Q in M is obtained by removing a neighborhood of Q and glueing in the manifold $P \times_{U(N)} \tilde{B}(\varepsilon)$ in the obvious way.

Proof of Corollary 1.2. Under the trivialization Φ of $(\mathcal{X}, \pi, J, \Omega^\mu)$ defined in (3.7), the section S_0 in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ corresponds to

$$S_0 = \{[\underline{s}, q, \underline{x}, x_{n+1}] \in \mathbb{F} \mid q = 0, \underline{x} = \underline{0}\}.$$

Let $Q = S_0 \times \mathbb{C}\mathbb{P}^1$.

For all $t \in \mathbb{C}\mathbb{P}^1$, the fiber \mathcal{X}_t contains $S_0 \times \{t\}$ as a symplectic submanifold and the diagram

$$\begin{array}{ccccc} \mathcal{X}_t & \longrightarrow & \mathcal{X} & \longrightarrow & \mathbb{C}\mathbb{P}^1 \\ \uparrow & & \uparrow & & \parallel \\ S_0 \times \{t\} & \longrightarrow & Q & \longrightarrow & \mathbb{C}\mathbb{P}^1 \end{array}$$

commutes. It follows that the normal bundle of Q in \mathcal{X} can be obtained by glueing the normal bundles of $S_0 \cong S_0 \times \{t\}$ in $X_t \cong \mathcal{X}_t$ for all $t \in \mathbb{C}\mathbb{P}^1$. Now let $\tilde{\mathcal{X}}$ denote a small symplectic blow-up along Q in \mathcal{X} in the neighborhood of Q on which $(\Omega^\mu)'$ has the standard form (4.1). Then each fiber in $\tilde{\mathcal{X}}$ is of the form $\tilde{X}_t \times \{t\}$, where \tilde{X}_t is the blow-up of S_0 in X_t . Hence the isotopy of τ_{L^μ} lifts to the blow-up of S_0 in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$.

To see that we can blow up along S_∞ in $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$, we repeat this construction with S_0 replaced by

$$S_\infty = \{[\underline{s}, 0, \underline{x}, 0] \in \mathbb{F} \mid x_j = 0 \text{ for } j \in \{2, \dots, n\}, s_0 x_0 + s_1 x_1 = 0\}$$

and Q by the complex submanifold $S_\infty \times \mathbb{C}\mathbb{P}^1$ in \mathcal{X} . \square

Proof of Proposition 1.4. Assume we have an isotopy $(\varphi_\lambda)_{0 \leq \lambda \leq 1}$ such that $\varphi_1 = \tau_{L^\mu}$ and φ_λ restricts to the identity on $S_0 \sqcup S'_\infty$ for all λ . We can then deform $(\varphi_\lambda)_{0 \leq \lambda \leq 1}$ to a homotopy $(\tilde{\varphi}_\lambda)_{0 \leq \lambda \leq 1}$ from τ_{L^μ} to the identity such that each $\tilde{\varphi}_\lambda$ fixes the neighborhoods

$$\begin{aligned} \mathcal{N}(S_0) &= \mathbb{C}\mathbb{P}^n \times B(\varepsilon, p_0) \supset S_0, \\ \mathcal{N}(S'_\infty) &= \mathbb{C}\mathbb{P}^n \times B(\varepsilon, p_\infty) \supset S'_\infty, \end{aligned}$$

where $B(\varepsilon, p)$ denotes an embedded size ε (real) $(2n+2)$ -ball in $\mathbb{C}\mathbb{P}^{n+1}$ centered at $p = p_0, p_\infty$. We show that this leads to a contradiction as follows. Following [Gom95, p. 534-535], we create a new smooth manifold M by performing a smooth surgery on $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{n+1}$ that takes place in the neighborhoods $\mathcal{N}(S_0)$ and $\mathcal{N}(S_\infty)$. In the manifold M , the Dehn twist corresponds to a diffeomorphism that acts non-trivially on homology but is also homotopic to the identity. This is a contradiction.

Choose a diffeomorphism $\phi : B(\frac{\varepsilon}{2}) \setminus \{0\} \rightarrow B(\frac{\varepsilon}{2}) \setminus \{0\}$ of the punctured $(2n+2)$ -ball that turns it smoothly inside out, e.g.,

$$\phi : z \mapsto z \sqrt{\frac{\varepsilon^2}{4|z|^2} - 1}.$$

Let $\widehat{\mathbb{C}\mathbb{P}}^{n+1}$ be the manifold obtained by removing the points p_0 and p_∞ from $\mathbb{C}\mathbb{P}^{n+1}$ and identifying the open sets $B(\frac{\varepsilon}{2}, p_0) \setminus \{p_0\}$ and $B(\frac{\varepsilon}{2}, p_\infty) \setminus \{p_\infty\}$ via the map ϕ .

Let

$$M = \mathbb{C}\mathbb{P}^n \times \widehat{\mathbb{C}\mathbb{P}}^{n+1}.$$

Since the surgery took place on neighborhoods on which the Dehn twist τ_L and the homotopy $(\tilde{\varphi}_\lambda)_{0 \leq \lambda \leq 1}$ restrict to the identity, τ_L induces a diffeomorphism $\widehat{\tau}_L$ of M homotopic to the identity. The manifold M contains a $(2n+1)$ -cycle C with non-trivial intersection with L . Indeed, let

$$\Gamma = \{[x_0 : 0 : \cdots : 0 : \sqrt{\mu - x_0}] \in \mathbb{C}\mathbb{P}^{n+1} \mid 0 \leq x_0 \leq \mu\}$$

be the image of a path from p_0 to p_∞ in $\mathbb{C}\mathbb{P}^{n+1}$. Then $\mathbb{C}\mathbb{P}^n \times \Gamma$ gives rise to a $(2n+1)$ -cycle C in M that intersects the Lagrangian $(2n+1)$ -sphere L^μ exactly once, namely at the point

$$([1 : 0 : \cdots : 0], [1 : 0 : \cdots : 0 : \sqrt{\mu - 1}]) \in M,$$

and the intersection is transverse. In particular, L represents a non-trivial element in the homology of M . By [ST01, (1.5)], the Dehn twist $\widehat{\tau}_L$ acts on the homology class of C by adding the homology class of L to it. See also Figure 1. \square

REFERENCES

- [Aud04] Michèle Audin. *Torus actions on symplectic manifolds*, volume 93 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, revised edition, 2004.
- [CS05] Alessio Corti and Ivan Smith. Conifold transitions and Mori theory. *Math. Res. Lett.*, 12(5-6):767–778, 2005.
- [Dup07] Emiko Dupont. *A symplectic isotopy of a Dehn twist on a product of projective spaces*. PhD thesis, Stony Brook University, (2007).
- [Gom95] Robert E. Gompf. A new construction of symplectic manifolds. *Ann. of Math. (2)*, 142(3):527–595, 1995.
- [MS98] Dusa McDuff and Dietmar Salamon. *Introduction to symplectic topology*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1998.
- [MS04] Dusa McDuff and Dietmar Salamon. *J-holomorphic curves and symplectic topology*, volume 52 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.
- [Sei97] Paul Seidel. *Floer homology and the symplectic isotopy problem*. PhD thesis, Oxford University, (1997).
- [Sei03] Paul Seidel. A long exact sequence for symplectic Floer cohomology. *Topology*, 42(5):1003–1063, 2003.
- [ST01] Paul Seidel and Richard Thomas. Braid group actions on derived categories of coherent sheaves. *Duke Math. J.*, 108(1):37–108, 2001.
- [Tha94] Michael Thaddeus. Toric quotients and flips. In *Topology, geometry and field theory*, pages 193–213. World Sci. Publ., River Edge, NJ, 1994.

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794, USA