## A quantum solution to the arrow-of-time dilemma

Lorenzo Maccone\*

QUIT, Dip. A. Volta, 27100 Pavia, and Institute for Scientific Interchange, 10133 Torino, Italy.

The arrow of time dilemma: the laws of physics are invariant for time inversion, whereas the familiar phenomena we see everyday are not (i.e. entropy increases). I show that, within a quantum mechanical framework, all phenomena which leave a trail of information behind (and hence can be studied by physics) are those where entropy necessarily increases or remains constant. All phenomena where the entropy decreases must not leave any information of their having happened. This situation is completely indistinguishable from their not having happened at all. In the light of this observation, the second law of thermodynamics is reduced to a mere tautology: physics cannot study those processes where entropy has decreased, even if they were commonplace.

PACS numbers: 03.67.-a,03.65.Ud

Paradoxes have always been very fruitful in stimulating advances in physics. One which still lacks a satisfactory explanation is the Loschmidt paradox [1]. Namely, how can we obtain irreversible phenomena from reversible time-symmetric physical laws [2]? The irreversibility in Physics is summarized by the second law of thermodynamics: entropy, which measures the degradation of the usable energy in a system, never decreases in isolated systems. Many approaches have been proposed to solve this conundrum, but most ultimately resort to postulating low entropy initial states (see e.g. [3, 4]), which is clearly an *ad hoc* assumption [5]. Others suggest that the thermodynamic arrow of time is in some way connected to the cosmological one [6], that physical laws must be modified to embed irreversibility [7], that irreversibility arises from decoherence [8], or from some time-symmetric mechanism embedded in quantum mechanics [9], etc. Recent reviews on this problem are given in Ref. [10].

Here I propose a different approach, based on existing laws of physics (quantum mechanics). I show that entropy in a system can both increase and decrease (as time reversal dictates), but that all entropy-decreasing transformations cannot leave *any* trace of their having happened. Since no information on them exists, this is indistinguishable from the situation in which such transformations do not happen at all: "The past exists only insofar as it is recorded in the present" [11]. Then the second law is forcefully valid: the only physical evolutions we see in our past, and which can then be studied, are those where entropy has not decreased.

I start by briefly relating the thermodynamic entropy with the von Neumann entropy, and introducing the second law. I then present two thought experiments, where entropy is deleted together with all records of the entropy increasing processes: even though at some time the entropy of the system had definitely increased, afterward it is decreased again, but none of the observers can be aware of it. I conclude with a general derivation through the analysis of the entropy transfers that take place in physical transformations.

Entropy and the second law. Thermodynamic entropy is a quantity that measures how the usable energy in a physical process is degraded into heat. It can be introduced in many ways from different axiomatizations of thermodynamics. The von Neumann entropy of a quantum system in the state  $\rho$  is defined as  $S(\rho) \equiv -\text{Tr}[\rho \log_2 \rho]$ . When applicable, these two entropies coincide (except for an inconsequential multiplicative factor). This derives from an argument introduced by Einstein [12] and extended by Peres [13] (e.g. both the canonical and the microcanonical ensemble can be derived from quantum mechanical considerations [14, 15]). For our purposes, however, it is sufficient to observe that thermodynamic and von Neumann entropies can be inter-converted, employing Maxwelldemons [16, 17] or Szilard-engines [18, 19]: useful work can be extracted from a single thermal reservoir by increasing the von Neumann entropy of a memory space.

There are many different formulations of the second law, but we can summarize them by stating that, in any process in which an isolated system goes from one state to another, its thermodynamic entropy does not decrease [20]. There is a hidden assumption in this statement. Whenever an isolated system is obtained by joining two previously isolated systems, then the second law is valid only if the two systems are initially uncorrelated, i.e. if their initial joint entropy is the sum of their individual entropies. It is generally impossible to exclude that two systems might be correlated in some unknown way and there is no operative method to determine whether a system is uncorrelated from all others (e.g. given a box containing some gas it is impossible to exclude that the gas particles might be correlated with other systems). Thus, in thermodynamics all systems are considered uncorrelated, unless it is known otherwise. Without this assumption, it would be impossible to assign an entropy to any system unless the state of the whole universe is known: a normal observer is limited in the information she can acquire and on the control she can apply. This implies that thermodynamic entropy is a subjective quantity [21], even though for all practical situations this is completely irrelevant: the eventual correlations in all macroscopic systems are practically impossible to control and exploit. Even though they are ignored by the normal observer, correlations between herself and other systems do exist. Until they are eliminated, the other systems cannot decrease their entropy. A physical process may either reduce or increase these correlations. When they are reduced, this may seem to entail a diminishing of the entropy, but the observer will not be aware of it as her memories are correlations and will have been erased by necessity (each bit of memory is one bit of correlation and, until her memory has been erased, the correlations are not eliminated). Instead, when the physical process increases these correlations, she will see it as an increase in entropy. The observer will then only be aware of entropy non-decreasing processes. [Not even a super-observer that can keep track of all the correlations would ever see any entropy decrease. In fact, since he can discover and take advantage of all correlations between microscopic degrees of freedom, all processes are always zero-entropy processes from his super-observer point of view.]

The above analysis is limited to systems that are somehow correlated with the observer. One might then expect that she could witness entropy decreasing processes in systems that are completely factorized from her. That is indeed the case: statistical microscopic fluctuations can occasionally decrease the entropy of a system (the second law has only a statistical valence). However, an observer is macroscopic by definition, and all remotely interacting macroscopic systems become correlated very rapidly (e.g. Borel famously calculated that moving a gram of material on the star Sirius by one meter can influence the trajectories of the particles in a gas on earth on a time-scale of  $\mu s$  [22]). This is the same mechanism at the basis of quantum decoherence [8], and it entails that in practice the above analysis applies to all situations: no entropy decrease in macroscopic systems is ever observed.

In what follows I will make these ideas rigorous.

Since the two above definitions of entropy are equivalent, the von Neumann entropy also obeys the second law. In fact, isolated systems evolve with unitary evolutions, which leave the von Neumann entropy invariant. There may be an increase if the evolution is not exactly known or if it creates unknown correlations among subsystems. In the first case, the coarse-grained evolution is of the form  $\rho' = \sum_n p_n U_n \rho U_n^{\dagger}$ , where  $p_n$  is a probability and  $U_n$  are unitary operators. Then the final entropy  $S(\rho')$  may be larger than the initial entropy  $S(\rho)$ :

$$S(\rho') = S(\sum_{n} p_n U_n \rho U_n^{\dagger}) \ge \sum_{n} p_n S(U_n \rho U_n^{\dagger}) = S(\rho)(1)$$

(the inequality follows from the concavity of the entropy). In the second case, the entropy of subsystems can increase, as some unknown correlations between them may build up:

$$S(\rho_1') + S(\rho_2') \ge S(U(\rho_1 \otimes \rho_2)U^{\dagger}) = S(\rho_1) + S(\rho_2) , (2)$$

where  $\rho'_i$  and  $\rho_i$  are the final and initial states of the subsystems, U is the evolution coupling them, and the last equality holds if they are initially uncorrelated (the inequality follows from the subadditivity of the entropy).

Thought experiments. The quantum information theory mantra "Information is physical" [23] implies that any record [24] of an occurred event can be decorrelated from such event by an appropriate physical interaction. If all the records of an event are decorrelated from it, then by definition there is no way to know whether this event has ever happened. This situation is indistinguishable from its not having happened. If this event has increased the entropy, the subsequent erasing of all records can (will) produce an entropy decrease without violation of any physical law. We now analyze two such situations, an imperfect transmission of energy and a quantum measurement.

Alice's lab is perfectly isolated, so that to an outside observer (Bob), its quantum evolution is unitary. Analyze the situation in which Bob sends Alice some energy in the form of light, a multimode electromagnetic field in a zero-entropy pure state. We suppose that, to secure the energy Bob is sending her, she uses many detectors which are not matched to his modes. Given a system in almost any possible pure state, all its subsystems which are small enough are approximately in the canonical state [14]. This implies that, if each of Alice's detectors is sensitive to only a small part of Bob's modes, the detectors mostly see thermal radiation, and she feels them warming up. She will then be justified in assigning a nonzero thermodynamic entropy to her detectors, as she sees them basically as thermal-equilibrium systems. One might object that she is mistaken, since the states of the detectors are not uncorrelated. However, since she ignores the correlations, she cannot use such correlations to extract energy from the detectors. Alice concludes that most of the energy Bob sent her has been wasted as heat, raising the thermodynamic entropy of her lab. Suppose now that Bob has complete control of all the degrees of freedom in her lab. He knows and can exploit the correlations to recover all the energy he had initially given Alice. Of course, although possible in principle, he needs a dauntingly complex transformation, which requires him to be able to control a huge number of her lab's degrees of freedom (including the brain cells where her memories are, and the notepads where she wrote the temperatures!). To extract the energy, since it was initially locked in a pure state of the field, he must return it to a system in a zero-entropy pure state, i.e. factorized from all the other degrees of freedom of Alice's lab. Then he must erase *all* the correlations between them: at the end of Bob's recovery, Alice cannot remember feeling her detectors warm up, they are cool again, her notepads contain no temperature information, and all the energy initially in the electromagnetic field is again available, even though (from Alice's point of view) most of it was definitely locked into thermal energy at one time.

The second though-experiment [25] is a prototypical quantum measurement. Bob prepares a spin-1/2 particle oriented along the x axis, e.g. in a spin  $| \rightarrow \rangle$  state and hands it to Alice. She sends it through a Stern-Gerlach apparatus oriented along the z axis [13]. The



FIG. 1: a) Alice in her isolated lab performs a Stern-Gerlach measurement on a spin 1/2 particle initially oriented parallel to the x axis, i.e. in a state  $| \rightarrow \rangle$ . Since the apparatus is oriented along the z axis, this measurement creates one bit of entropy for Alice (not for Bob, who is isolated from her lab). b) Bob flips a switch that "cancels" Alice measurement by decorrelating from the spin all those degrees of freedom of her lab that have recorded the measurement outcome. Now the spin is returned to its initial state  $| \rightarrow \rangle$  and Alice cannot have any memory of what her measurement result was. Her entropy has decreased, but she cannot remember it ever having increased.

measurement consists in coupling the quantum system with some macroscopic degrees of freedom (a reservoir), not all of which are under the control of the experimenter [26], whence the irreversibility. Notice that,  $| \rightarrow \rangle = (| \uparrow \rangle + | \downarrow \rangle) / \sqrt{2}$ , where  $| \uparrow \rangle$  and  $| \downarrow \rangle$  are the eigenstates of a z measurement operator. Hence, this apparatus will increase the entropy of the spin system by one bit [16]: Before the readout, the spin state will be in the maximally mixed state  $(|\uparrow\rangle\langle\uparrow|+|\downarrow\rangle\langle\downarrow|)/2$ . After Alice has looked at the result, she has transferred this one bit of entropy, created by the measurement, to her memory. From the point of view of Bob, outside her isolated lab, Alice's measurement is simply a (quantum) correlation of her measurement apparatus to the spin. [A thorough analysis of the microscopic details and of the thermodynamics of this type of measurement is given in Ref. [16].] The initial state of the spin  $| \rightarrow \rangle = (| \uparrow \rangle + | \downarrow \rangle)/\sqrt{2}$ evolves into the correlated (entangled) state

$$(|\uparrow\rangle|$$
Alice sees "up" $\rangle + |\downarrow\rangle|$ Alice sees "down" $\rangle)/\sqrt{2}$ ,(3)

where the first ket in the two terms refers to the spin state, whereas the second ket refers to the rest of Alice's lab. Thus, from the point of view of Bob, Alice's measurement is an evolution similar to a controlled-NOT unitary transformation of the type  $U_{cnot}(|0\rangle + |1\rangle)|0\rangle =$  $|0\rangle|0\rangle + |1\rangle|1\rangle$ . Such a transformation can be easily inverted, as it is its own inverse. Analogously, Bob can flip a switch and invert Alice's measurement. At the end of his operation, all records of the measurement result (Alice's notepad, her brain cells, the apparatus gauges) will have been decorrelated from the spin state. She will remember having performed the measurement, but she will be (*must* be) unable to recall what the measurement result was. In addition, the spin has become uncorrelated from the measurement apparatus, so it is returned to a pure state. I emphasize that Bob's transformation is not necessarily a reversion of the dynamics of Alice's lab. [Notice that a "quantum eraser" [27] only permits to decide *a posteriori* which of two complementary measurements to perform using previously collected data: the measurement process is not actually erased, and the entropy does not decrease.]

In both the above experiments, from Alice's point of view, entropy definitely has been created after she has interacted with Bob's light or his spin. However, this entropy is subsequently coherently erased by Bob. At the end of the process, looking back at the evolution in her lab, she cannot see any violation of the second law: she has no (cannot have any) record of the fact that entropy at one point had increased.

Entropic considerations. The above thought experiments exemplify a general situation: entropy can decrease, but its decrease is accompanied by an erasure of any memory that the entropy-decreasing transformation has occurred. In fact, any interaction between an observer A and a system C which decreases their entropy by a certain quantity, must also reduce their quantum mutual information by the same amount (unless, of course, the entropy is dumped into a reservoir R). The quantum mutual information  $S(A:C) \equiv S(\rho_A) + S(\rho_C) - S(\rho_{AC})$ measures the amount of shared quantum correlations between the two systems A and C ( $\rho_{AC}$  being the state of the system AC, and  $\rho_A$  and  $\rho_C$  its partial traces, i.e. the states of A and C).

Taking the cue from [28], I now prove the above assertion, namely I show that

$$\Delta S(A) + \Delta S(C) - \Delta S(R) - \Delta S(A:C) = 0, \quad (4)$$

where  $\Delta S(X) \equiv S_t(\rho_X) - S_0(\rho_X)$  is the entropy difference between the final state at time t and the initial state of the system X, and where  $\Delta S(A:C) = S_t(A:C) - S_0(A:C)$  is the quantum mutual information difference. Choose the reservoir R so that the joint state of the systems ACR is pure and so that the evolution maintains the purity (R is a purification space). Then the initial and final entropies are  $S_0(AC) = S_0(R)$  and  $S_t(AC) = S_t(R)$ , respectively. Thus we find  $S_0(AB) = S_t(AB) - \Delta S(R)$ which, when substituted into the left-hand-side term of (4), shows that this term is null. [This proof is valid also if the evolution is not perfectly known, i.e. if it is given by a random unitary map, see Eq. (1).]

Now, to prove that the above reduction of entropy entails a memory erasure, I show that this erasure must follow from the elimination of quantum mutual information. A memory of an event is a physical system A which has nonzero classical mutual information on a system Cthat bears the consequences of that event. Then, the erasure of the memory follows from an elimination of the quantum mutual information S(A:C) if this last quantity is an upper bound to the classical mutual information I(A:C). Thus, we must show that for any POVM measurement  $\{\Pi_i^{(a)} \otimes \Pi_j^{(c)}\}$  extracting information separately from the two systems ( $\Pi_i^{(a)}$  acting on A and  $\Pi_i^{(c)}$  on C),

$$S(A:C) \ge I(A:C) , \qquad (5)$$

where I(A:C) is the mutual information of the POVM's measurement results. A simple proof of this statement exists (e.g. see [29, 30]): use the equality S(A:C) = $S(\rho_{AC}||\rho_A \otimes \rho_C)$ , where  $S(\rho||\sigma) \equiv \text{Tr}[\rho \log_2 \rho - \rho \log_2 \sigma]$  is the quantum relative entropy. This quantity is monotone for application of CP-maps [29], i.e.  $S(\rho||\sigma) \ge$  $S(\mathcal{N}[\rho]||\mathcal{N}[\sigma])$  for any transformation  $\mathcal{N}$  that can be written as  $\mathcal{N}[\rho] = \sum_k A_k \rho A_k^{\dagger}$ , with the Kraus operators  $A_k$  satisfying  $\sum_k A_k^{\dagger} A_k = \mathbb{1}$ . Consider the "measure and reprepare" channel, i.e. the transformation  $\mathcal{N}[\rho] = \sum_n \text{Tr}[\Pi_n \rho] |n\rangle \langle n|$  where  $\{|n\rangle\}$  is a basis, and  $\Pi_n$ is a POVM element (i.e. a positive operator such that  $\sum_n \Pi_n = \mathbb{1}$ ). It is a CP-map, since it has a Kraus form

$$A_{nm} = |n\rangle \langle v_m^{(n)}| \sqrt{p_m^{(n)}}, \text{ with } \Pi_n = \sum_m p_m^{(n)} |v_m^{(n)}\rangle \langle v_m^{(n)}|.(6)$$

Using the monotonicity of the relative entropy under the action of the map  $\mathcal{N}$ , we find

$$S(A:C) = S(\rho_{AC} || \rho_A \otimes \rho_C) \ge S(\mathcal{N}[\rho_{AC}] || \mathcal{N}[\rho_A \otimes \rho_C])$$
  
=  $\sum_{ij} p_{ij} \log_2 p_{ij} - \sum_{ij} p_{ij} \log_2 (q_i r_j) = I(A:C) ,$ 

where  $p_{ij} \equiv \text{Tr}[\Pi_i^{(a)} \otimes \Pi_j^{(c)} \rho_{AC}], q_i \equiv \text{Tr}[\Pi_i^{(a)} \rho_A], \text{ and } r_j \equiv \text{Tr}[\Pi_i^{(c)} \rho_C].$ 

The interpretation of Eq. (4) is that, if we want to decrease the entropy of the system C (somehow correlated with the observer A) without increasing the entropy of a reservoir R, we need to reduce the quantum mutual information between C and the observer A (e.g. in the Stern-Gerlach thought experiment, the system A is Alice's lab and C is the spin-1/2 particle: their final entropies are reduced by one bit at the expense of erasing two bits of quantum mutual information  $S_0(A : C)$ ). The fact that mutual information can be used to decrease entropy was already pointed out by Lloyd [28] and Zurek [31].

The implications of the above analysis can be seen explicitly by employing Eq. (4) twice, by considering an intermediate time when S(C) is higher than at the initial and final times. The entropy S(C) of the system is high at the intermediate time after an entropy-increasing transformation, and then (if no entropy-absorbing reservoir R is used) it can be reduced by a successive entropydecreasing transformation at the cost of reducing the mutual information between the observer and C. Even though the entropy S(C) (as measured from the point of view of the observer A) does decrease, the observer is not aware of it, as the entropy-decreasing transformation must factorize her from the system C containing information on the prior entropy-increasing event: her memories of such event must be part of the destroyed correlations. The deep reason for this is that, from her own point of view, the Born rule kicks in when the observer becomes entangled with another system. (An external super-observer may, instead, just see her becoming entangled with the other system, but then he cannot know the measurement result.) The Born rule is the only place where quantum mechanics allows irreversibility, but the correlations that stem from such rule *can* be undone, at least if one treats both the observer and the apparatus quantum mechanically. This means that the measurement can be undone, at the price that all the observer's memories must be erased.

What we have seen up to now is that any decrease in entropy of a system that is correlated with an observer entails a memory erasure of said observer, in the absence of reservoirs (or is a zero-entropy process for a superobserver that keeps track of all the correlations). That might seem to imply that an observer should be able to see entropy-decreasing processes when considering systems that are uncorrelated from her. In fact, at microscopic level, statistical fluctuations do decrease occasionally the entropy. However, the correlations between any two macroscopic systems build up continuously, and at amazing rates [22]: this is how decoherence arises [8]. Then no observer is really factorized with respect to any macroscopic system she observes. This implies that entropy decreases of a macroscopic system becomes unobservable (unless extreme care is taken to shield the system under analysis). Only microscopic systems can be considered factorized from an observer for a period of time long enough to see entropy decrease from fluctuations.

Conclusions. In this paper I gave a quantum solution to the Loschmidt paradox, showing that all physical transformations where entropy is decreased cannot relinquish any memory of their having happened from the point of view of any observer: both normal observers that interact with the studied systems and external superobservers that keep track of all the correlations. Thus they are irrelevant to physics. Quantum mechanics is necessary to this argument. In the above derivation, we have used the property that the entropy of a joint system can be smaller than that of each of its subsystems. This is true of von Neumann entropy, but not true if entropy is calculated using classical probability theory: then the entropy of a joint system is always larger than that of its subsystem with largest entropy. By how much must any system be extended until we can take advantage of this quantum reduction of the global entropy? It is clear from Borel's famous arguments [22] that the time scale in which a macroscopic system can be really considered as isolated is very small. As such, the arguments presented in this paper are of theoretical interest only, and have little or no practical consequence for any normal macroscopic system: the effects presented here become relevant only at a scale that approaches the whole universe very rapidly.

In closing, I indulge in a couple of more philosophical considerations. In a quantum cosmological setting, the above approach easily fits in the hypothesis that the quantum state of the whole universe is a pure (i.e. zeroentropy) state evolving unitarily (e.g. see [14, 32, 33]). One of the most puzzling aspects of our universe is the fact that its initial state had entropy so much lower than we see today, making the initial state highly unlikely [4]. Joining the above hypothesis of a zero-entropy pure state of the universe with the second law considerations analyzed in this paper, it is clear that such puzzle can be resolved. The universe may be in a zero entropy state. even though it appears (to us, internal observers) to possess a higher entropy: our situation is similar to the one of Alice, who, just after the measurement sees her lab in a nonzero entropy state, whereas to the super-observer Bob her lab maintains a zero-entropy state all along. However, it is clear that this approach does not require to deal with the quantum state of the whole universe, but it applies also to arbitrary physical systems.

In a quantum cosmological framework, Boltzmann's initial condition translates in the equivalent question of why the initial state of the universe is such that its subsystems are mostly unentangled. Such a state is highly improbable, as all states in a sufficiently large Hilbert space are almost completely entangled [34]. The most compelling answer to this question derives from Davies' argument that, as space expands because of cosmological expansion and inflation, new degrees of freedom are created, giving the potential for accommodating new entropy [35]: one may think that, as these new degrees of freedom are created, they are initialized in a factorized (possibly pure) state. However, this assumption is un5

warranted, and moreover, our current understanding of quantum mechanics does not allow the description of a situation where the number of degrees of freedom of a system (and hence its Hilbert space) changes dynamically [36]. The alternative solution presented in this paper sidesteps the problem: whatever the state of the universe, an internal observer would still only see the processes where entropy increases.

In addition, I recall that there is a substantial problem in rigorously defining past and future without resorting to the second law (which would then be reduced to a mere definition). In fact, the laws of physics are timereversal invariant. Hence, there is no preferred direction of time according to which we may establish a substantial difference between the two temporal directions past-tofuture and future-to-past [37]. Anthropocentrically, we could define the past as that of which we have memories of, and the future as that of which we do not have any memories. Of course, such definition cannot be made rigorous, since it resorts to a observers and their memories. However, even using this ambiguous, intuitive definition of past, it is clear that any event, which cannot have any correlation with us, does not pertain to our past just as if it had never happened.

I thank S. Lloyd, G.M. D'Ariano, V. Giovannetti, G.P. Beretta, S. Wolf, and A. Winter for useful discussions and feedback.

\* Current address: Massachusetts Institute of Technology, RLE,50 Vassar Street Bldg. 36-472B Cambridge, MA 02139 USA.

- J. Loschmidt, Sitzungsber. Kais. Akad. Wiss. Wien, Math. Naturwiss. Classe, II. Abteilung 73, 128 (1876).
- [2] More precisely: from laws symmetric for CPT-inversion.
- [3] R. Penrose, in *General Relativity, an Einstein Centennary Survey*, S. W. Hawking, W. Israel eds., (Cambridge Univ. Press, Cambridge, 1979).
- [4] R. M. Wald Eprint arXiv:gr-qc/0507094v1 (2005).
- [5] E. G. D. Cohen, T. H. Berlin, Physica 26, 717 (1960).
- [6] T. Gold, The nature of time (Cornell Univ. Press, Ithaca, 1967); P. C. W. Davies, The physics of time asymmetry, (Surrey univ. press, London, 1974); S. M. Carroll and J. Chen, arXiv:gr-qc/0505037 (2005).
- [7] G.C. Ghirardi, A. Rimini, T. Weber, Phys. Rev. D 34, 470 (1986); D. Z. Albert, *Quantum mechanics and experience* (Harvard Univ. press, Cambridge, 1994); E.P. Gyftopoulos, G.P. Beretta, *Thermodynamics: Foundations and Applications* (Dover, London, 2005).
- [8] H. D. Zeh, The Physical Basis of The Direction of Time, (Springer-Verlag, Berlin, 1999).
- [9] L. S. Schulman, *Time's arrow and quantum measurement* (Cambridge Univ. Press, Cambridge, 1997).
- [10] J.J. Halliwell, J. Pérez-Mercader, W.H. Zurek (eds.), Physical Origins of Time Asymmetry (Cambridge Univ. Press, Cambridge, 1996); H. Price, Time's arrow and Archimedes' point (Oxford Univ. Press, Oxford, 1996); S.F. Savitt (ed.), Time's arrows today (Cambridge Univ. Press, Cambridge, 1997).

- [11] J. A. Wheeler, as quoted in W. H. Zurek, Rev. Mod. Phys. 75, 715 (2003).
- [12] A. Einstein, Verh. Deut. Phys. Gesell. 16, 820 (1914).
- [13] A. Peres, Quantum Theory: Concepts and Methods, (Kluwer ac. publ., Dordrecht, 1993).
- [14] S. Popescu, A. J. Short, A. Winter, Nature Physics 2, 754 (2006).
- [15] J. Gemmer, A. Otte, G. Mahler, Phys. Rev. Lett. 86, 1927 (2001); H. Tasaki, Phys. Rev. Lett. 80, 1373 (1998).
- [16] S. Lloyd, Phys. Rev. A 56, 3374 (1997).
- [17] H. S. Leff, A. F. Rex (editors), Maxwell's Demon: Entropy, Information, Computing (Princeton Univ. Press, Princeton, 1991); C. H. Bennett, Int. J. Theor. Phys. 21, 905 (1982); V. Serreli, C.-F. Lee, E. R. Kay, and D. A. Leigh, Nature 445, 523 (2007).
- [18] L. Szilard, in *Quantum theory and Measurement*, J. A. Wheeler and W. H. Zurek ed. (Princeton Univ. Press, Princeton, 1983), pg. 539.
- [19] M. O. Scully, Phys. Rev. Lett. 87, 220601 (2001).
- [20] F. Reif, Fundamentals of statistical and thermal physics (Mc Graw Hill, Boston, 1965).
- [21] S. Lloyd in *Meeting the Entropy Challenge*, G.P. Beretta, A.F. Ghoniem, G.N. Hatsopoulos (AIP Conference Proceedings 1033, Melville, 2008).
- [22] E. Borel, Le Hasard (Alcan, Paris, 1914).
- [23] R. Landauer, Proceedings of the Workshop on Physics and Computation PhysComp'92 (IEEE Press, Los Alami-

tos, 1992) pg. 1.

- [24] By record I consider any physical system which is correlated with the event, such as its description on the pages of a book, or the neural pattern in a witness's brain.
- [25] B.S. DeWitt, R. N. Graham eds. The many-worlds Interpretation of Quantum Mechanics, (Princeton Univ. Press, 1973).
- [26] A. Peres, Phys. Rev. D 22, 879 (1980).
- [27] M. O. Scully, B.-G. Englert, and H. Walther, Nature 351, 111 (1991).
- [28] S. Lloyd, Phys. Rev. A **39**, 5378 (1989).
- [29] H. P. Yuen, M. Ozawa, Phys. Rev. Lett. **70**, 363 (1993).
  [30] D. DiVincenzo *et al.* Phys. Rev. Lett. 92, 067902 (2004), arXiv:quant-ph/0303088.
- [31] W. H. Zurek in Frontiers of Nonequilibrium Statistical Physics, Moore, G. T.; Scully, M. O., Eds. (Plenum

Press, New York, 1984) pg. 151, quant-ph/0301076.

- [32] M. Tegmark, Found. Phys. Lett. 9, 25 (1996).
- [33] D. Z. Albert, in PSA: Proceedings of the Biennial Meeting of the Philosophy of Science Association 2, 127 (1988).
- [34] P. Hayden, D. W. Leung, A. Winter, Comm. Math. Phys. 265, 95 (2006).
- [35] P. C. W. Davies, The physics of time asymmetry, (Surrey univ. press, London, 1974); P. C. W. Davies, About time: Einstein's unfinished revolution (Simon & Schuster, New York, 1995).
- [36] W. Unruh in S. F. Savitt (Editor), Time's arrows today: recent physical and philosophical work on the direction of time, (Cambridge Univ. Press, Cambridge, 1997).
- [37] M. Castagnino, M. Gadella, and O. Lombardi, Found. Phys. 36, 407 (2006).