

Correlation-induced metal insulator transition in a two-channel fermion-boson model

G. Wellein,¹ H. Fehske,² A. Alvermann,² and D. M. Edwards³

¹ Regionales Rechenzentrum Erlangen, Universität Erlangen-Nürnberg, 91058 Erlangen, Germany

² Institut für Physik, Ernst-Moritz-Arndt-Universität Greifswald, 17489 Greifswald, Germany

³ Department of Mathematics, Imperial College London, London SW7 2AZ, United Kingdom

We investigate charge transport within some background medium by means of an effective lattice model with a novel form of fermion-boson coupling. The bosons describe both correlations and fluctuations of the background. By analyzing ground-state and spectral properties of this two-channel transport model, we show how a metal-insulator quantum phase transition can occur for the half-filled band case. We establish connections to the Mott and Peierls transition scenarios.

The way a material evolves from a metallic to an insulating state is one of the most fundamental problems in solid state physics. Besides band structure and disorder effects, electron-electron and electron-phonon interactions are the driving forces behind metal-insulator transitions (MITs) in the majority of cases. While the so-called Mott-Hubbard MIT [1] is caused by strong Coulomb correlations, the Peierls MIT [2] is triggered by the coupling to vibrational excitations of the crystal. Both scenarios are known to compete in subtle ways.

The MIT problem can be addressed by the investigation of generic Hamiltonians for interacting electrons and phonons like the Hubbard or Holstein model, [3] or combinations of both. [4] These models have been proved to describe MIT phenomena for the half-filled band case, in particular for one-dimensional (1D) systems known to be susceptible to the formation of insulating spin-density-wave (SDW) or charge-density-wave (CDW) broken-symmetry groundstates. [5]

On the metallic side of the MIT, charge transport then takes place within a “background medium” that exhibits strong correlations, which anticipate the SDW or CDW on the insulating side. In that case, a particle, as it moves, creates local distortions of substantial energy in the background. The distortions may be able to relax. The rate at which this occurs depends on the system properties but also on the proximity to the MIT point.

Since this picture clearly has wide applicability, e.g., to transport in 1D CDW MX chain compounds, [5] 2D high- T_c superconducting cuprates, [6] 3D colossal magnetoresistive manganites, [7] and even biological systems, [8] the authors recently proposed a rather generic transport Hamiltonian [9, 10]

$$H = H_b - \lambda \sum_i (b_i^\dagger + b_i) + \omega_0 \sum_i b_i^\dagger b_i, \quad (1)$$

which covers basic features of more complicated Hubbard, t - J , Holstein or SSH type models. In Eq. (1) $H_b = -t_b \sum_{\langle i,j \rangle} c_j^\dagger c_i (b_i^\dagger + b_j)$ describes the boson-affected nearest neighbor (NN) hopping of fermionic particles (c_i^\dagger). Since a particle creates a boson (b_i^\dagger) only on the site it leaves and destroys a boson on the site it enters, it generates a “string” of local bosonic fluctuations with energy

ω_0 , which approximate the distortions in the background medium (reminiscent of t - J -model-like hole motion in a 2D antiferromagnet [11]).

The λ term allows a boson to decay spontaneously, thereby cutting the string that otherwise binds a fermion to its origin. But even for $\lambda = 0$, when transport is fully boson-assisted, there exist coherent hopping processes that propagate the particle and restore the background. [10] The lowest order process of this kind is

$$|\circ \cdot \cdot \rangle \rightarrow |* \circ \cdot \rangle \rightarrow |* * \circ \rangle \rightarrow |* * * \rangle \rightarrow |* * * \rangle \rightarrow |* * * \rangle \rightarrow |\circ * * \rangle \rightarrow |\circ * * \rangle \rightarrow |\cdot \circ \cdot \rangle, \quad (2)$$

where in steps 1-3, three bosons ($*$) are excited, which are consumed in steps 4-6 afterwards. In this process the fermion (\circ) becomes correlated with the background fluctuations. In some sense the effective next-NN hopping (2) represents an 1D analogue of the 2D “Trugman path” of a hole in a Néel spin background. [12]

A unitary transformation $b_i \mapsto b_i + \lambda/\omega_0$ eliminates the boson relaxation term in (1) in favor of a free-particle hopping channel, $H_f = -t_f \sum_{\langle i,j \rangle} c_j^\dagger c_i$ with $t_f = 2\lambda t_b/\omega_0$, in addition to the original one. That is,

$$H \mapsto H = H_b + H_f + \omega_0 \sum_i b_i^\dagger b_i, \quad (3)$$

i.e. the physics of our model is governed by two ratios: The relative strengths of the two transport channels (t_f/t_b) and the rate of bosonic fluctuations $(\omega_0/t_b)^{-1}$.

The model (3) has been solved numerically in the one-particle sector and revealed—despite its seeming simplicity—a surprisingly rich “phase diagram” with regimes of quasi-free, correlation and fluctuation dominated transport. [10] Depending on the parameters the particle’s motion resembles that of a spin or lattice polaron, or an electron coupled to bath degrees of freedom.

Whether our two-channel transport model (3) exhibits a quantum phase transition from a metallic to an insulating phase at certain commensurate band fillings is an open question. A tendency towards CDW formation is expected from perturbative arguments at $t_f = 0$ and large ω_0/t_b , where a gain in kinetic energy of $\mathcal{O}(t_b^2/\omega_0)$ stabilizes the CDW. On the other hand the free hopping channel H_f acts against any correlation induced change

ordering that might open a gap at the Fermi energy E_F , and strong bosonic fluctuations, i.e. small ω_0 , will also tend to destroy CDW order.

Simplified cluster models related to (1) have been studied previously [9]: (i) an infinite model with only one bosonic oscillator for the whole system, (ii) a 1D model based on a 2-site cluster. The exact solution of (i) and an approximate solution of (ii) both exhibit quantum phase transitions at $\lambda = 0$ between two Fermi liquids, with different Fermi surfaces, non-Fermi-liquid behavior persisting down to zero temperature at the critical point in case (ii). Both models, however, omit important aspects of model (1), e.g. the 3-boson hopping process (2).

Since the MIT is a subtle quantum mechanical problem that requires non-perturbative investigation schemes, in this work we will use unbiased numerical techniques to study the competition between itinerancy, correlations and fluctuations for the 1D half-filled band case, using the full model (1) without any restrictions.

To this end we employ exact diagonalization in combination with kernel polynomial expansion techniques, adapted for coupled fermion-boson systems. [13] The computational requirements are determined by the total dimension $D = \binom{N}{N_e} \binom{N+N_b}{N_b}$ of the Hilbert space, where N is the number of lattice sites, N_e counts the fermions, and N_b is the maximum number of bosons retained. A typical production job for a system with $N = 12$ at a given parameter set t_f/t_b and ω_0/t_b , runs on 300 processor cores of the HLRB-II supercomputer at LRZ Munich and consumes more than 30000 CPU hours. To check for convergence concerning N and N_b we re-calculated selected parameter sets for $N = 16$, resulting in D of about 5×10^{10} , with runs on up to 1000 processor cores.

Let us start with the discussion of the photoemission (PE) spectra. The spectral density of single-particle excitations associated with the injection of an electron with wave-vector k , $A^+(k, \omega)$ (inverse PE), and the corresponding quantity for the emission of an electron, $A^-(k, \omega)$ (PE), is given by

$$A^\pm(k, \omega) = \sum_n |\langle \psi_n^\pm | c_k^\pm | \psi_0 \rangle|^2 \delta[\omega \mp \omega_n^\pm]. \quad (4)$$

Here $c_k^+ = c_k^\dagger$, $c_k^- = c_k$, and $|\psi_0\rangle$ is the ground state in the N_e -particle sector while $|\psi_n^\pm\rangle$ denote the n -th excited states in the $N_e \pm 1$ -particle sectors with excitation energies $\omega_n^\pm = E_n^\pm - E_0$.

Figure 1 displays the wave-vector resolved single-particle spectra in the regime where distortions of the background are energy-intensive, i.e. the boson frequency ω_0 is high. If the free transport channel is dominant ($t_f = 5$ - upper graph), the occupied (unoccupied) band states, probed by PE (inverse PE), give rise to an almost particle-hole symmetric absorption spectrum $A_K^+(\omega - E_F) \simeq A_{K-\pi}^-(E_F - \omega)$. Thereby the main spectral weight resides in the uppermost (lowest) peaks




FIG. 1: Photoemission (black) and inverse photoemission spectra (red) for the half-filled band case with $\omega_0 = 2$ at $t_f(\lambda, \omega_0) = 5$ (upper panels) and $t_f = 0.01$ (lower panels), where $N = 12$, $N_b = 15$. Dashed lines give the integrated spectral weights, e.g. $S_K^+(\omega - E_F) = \int_0^\omega d\omega' A_K^+(\omega' - E_F)$, where $S_K = S_K^-(\infty) + S_K^+(\infty) = 1$, and $\sum_K S_K = N$. Here and in what follows periodic boundary conditions were used, leading to discrete $K(N)$ wave numbers. All energies are measured in units of $t_b = 1$, and ω is rescaled with respect to E_F .

of A_K^- (A_K^+) in each K -sector. The corresponding “coherent” band structure roughly follows the $-2t_f \cos K$ tight-binding band. Phonon satellites with less spectral weight occur near the Brillouin zone boundary predominantly, as a result of mixed electron-phonon excitations with total wave vector K . At $T = 0$ the Fermi energy is obtained from $\sum_K \int_{-\infty}^{E_F} A_K^-(\omega) d\omega = N_e = N/2$ (half-filling, no spin). We see that there is no gap between $A_{\pm\pi/2}^+$ and $A_{\pm\pi/2}^-$ at E_F . Moreover the spectral weight of both peaks is almost one, i.e. a particle injected (removed) with $K_F \pm \pi/2$ propagates unaffected by bosonic fluctuations. The system is a perfect metal.

If we decrease λ (t_f/t_b ratio) at fixed ω_0 , we enter

λ	ω_0	t_f	$E_{kin,f}$	$E_{kin,b}$	D	$S_c(\pi)$
5.0	0.5	20.0	-149.190	-13.915	80.336	0.0417
0.1	0.5	0.04	-2.745	-14.878	4.054	0.0419
5.0	2.0	5.0	-37.040	-5.291	20.290	0.0417
1.0	2.0	1.0	-7.008	-6.214	5.146	0.0425
0.1	2.0	0.1	-0.568	-7.036	–	0.0561
0.01	2.0	0.01	-0.023	-7.125	–	0.1056

TABLE I: Groundstate properties of the transport Hamiltonian (3) at half-filling ($N = 12$, $N_b = 15$).

mobility of the carrier. Accordingly the boson density is enlarged (suppressed) at the NN sites (site) of the particle. Clearly $\chi_{eb}(j)$ is small $\forall |j|$ if $\omega_0 \gg t_b, t_f$ because of the high energy cost.

As expected the fluctuation dominated regime is characterized by a large number of boson quanta in the groundstate. The position of the maximum in the boson weight function $|c_m|^2$ is shifted to slightly larger values as λ increases, i.e. the correlations become weaker.

Further information on the groundstate properties can be obtained from the kinetic energy parts $E_{kin,f/b} = \langle 0|H_{f/b}|0\rangle$. Table I shows that boson-assisted hopping becomes the major transport mechanism at small λ . On the metallic side of the MIT the creation and annihilation of bosons opens a coherent transport channel in the regime where strong correlations persist in the background. The Drude weight D , obtained from the f -sum rule $-D = \frac{1}{2}(E_{kin,f} + E_{kin,b}) + \int_0^\infty \sigma_{reg}(\omega)d\omega$, serves as a measure for this coherent transport. Here $\sigma_{reg}(\omega) = \sum_{n>0} \frac{|\langle n|j|0\rangle|^2}{\omega_n} \delta(\omega - \omega_n)$ is the regular part of the optical conductivity (with current $j = j_f + j_b$).

At the MIT point D vanishes for the infinite system. At the same time the optical gap opens. In the insulating phase the optical response is dominated by multi-boson emission and absorption processes. Thus the spectral weight contained in the regular part of $\sigma(\omega)$ is enhanced.

A small boson frequency allows for large fluctuations in the background, i.e. many bosons in our model. This supports via (2) transport on the one hand but, as in the one-particle sector, [10] also limits the mobility of a particle by many scattering events. Nevertheless D is expected to stay finite even for $\omega_0/t_b \rightarrow 0$.

To summarize, the two-channel transport Hamiltonian, introduced for studying the dynamics of charge carriers in a correlated/fluctuating medium, has previously only been properly analysed for a single carrier. [9, 10] Here, for the first time, we have investigated properly the interesting physics at finite carrier density. In particular, in 1D at half-filling, the model exhibits a quantum phase transition from a metallic to an insulating state. The MIT is driven by correlations, like the Mott-Hubbard transition, but in our case true long-range order devel-

ops because a CDW state is formed. This might point towards a Peierls transition scenario. The Peierls instability, however, is most pronounced in the adiabatic limit of small phonon frequencies, with many phonons involved establishing the CDW (lattice dimerization). By contrast we find that the CDW groundstate is a few-boson state. Fixing the boson relaxation $\lambda = t_f\omega_0/2t_b$, the system is more susceptible to CDW-formation if ω_0 increases while t_f decreases. In some respects this is evocative of the quantum phase transition in the (half-filled) spinless fermion Holstein model, which for large phonon frequencies and strong couplings can be mapped on the XXZ model. [15] The XXZ model undergoes a Kosterlitz-Thouless transition at the spin isotropy point. Mapping our model (1) for $\omega_0 \gg t_f, t_b$ to a spin model, the $S_z \rightarrow -S_z$ symmetry is broken, reflecting broken particle-hole symmetry. Note that the mass-asymmetric band structure that evolves in the insulating phase, with a very narrow valence and rather broad conduction band, is correlation induced, i.e. different in nature from simple two-band models.

Acknowledgements. This work was supported by the Bavarian KONWIHR project HQS@HPC and by DFG through SFB 652, B5.

-
- [1] N. F. Mott, *Metal-Insulator Transitions* (Taylor & Francis, London, 1990); F. Gebhard, *The Mott Metal-Insulator Transition* (Springer, Berlin, 1997).
 - [2] R. Peierls, *Quantum theory of solids* (Oxford University Press, Oxford, 1955).
 - [3] J. Hubbard, Proc. Roy. Soc. London, Ser. A **276**, 238 (1963); T. Holstein, Ann. Phys. (N.Y.) **8**, 325 (1959).
 - [4] Y. Takada and A. Chatterjee, Phys. Rev. B **67**, 081102(R) (2003); H. Fehske, G. Wellein, G. Hager, A. Weiße, and A. R. Bishop, Phys. Rev. B **69**, 165115 (2004); R. T. Clay and R. P. Hardikar, Phys. Rev. Lett. **95**, 096401 (2005); M. Tezuka, R. Arita, and H. Aoki, Phys. Rev. Lett. **95**, 226401 (2005).
 - [5] A. R. Bishop and B. I. Swanson, Los Alamos Sciences **21**, 133 (1993).
 - [6] D. M. Newns and C. C. Tsuei, J. Phys. Conf. Ser. **92**, 012007 (2007).
 - [7] L. G. L. Wegener and P. B. Littlewood, Phys. Rev. B **66**, 061905 (2002).
 - [8] S. Komineas, G. Kalosaka, and A. R. Bishop, Phys. Rev. E **65**, 061905 (2007).
 - [9] D. M. Edwards, Physica B **378B-380B**, 133 (2006).
 - [10] A. Alvermann, D. M. Edwards, and H. Fehske, Phys. Rev. Lett. **98**, 056602 (2007).
 - [11] W. F. Brinkman and T. M. Rice, Phys. Rev. B **2** (1970); G. Martinez and P. Horsch, Phys. Rev. B **44**, 317 (1991).
 - [12] S. A. Trugman, Phys. Rev. B **37**, 1597 (1988).
 - [13] A. Weiße, G. Wellein, A. Alvermann, and H. Fehske, Rev. Mod. Phys. **78**, 275 (2006); E. Jeckelmann and H. Fehske, Rivista del Nuovo Cimento **30**, 259 (2007).
 - [14] We found dispersive (flat) absorption bands in the PE (inverse PE) spectra as well, but for higher excitation

energies and with less spectral weight.

- [15] R. J. Bursill, R. H. McKenzie, and C. J. Hamer, Phys. Rev. Lett. **80**, 5607 (1998); M. Hohenadler, G. Wellein,

A. R. Bishop, A. Alvermann, and H. Fehske, Phys. Rev. B **74**, 245120 (2006).