

Symmetric construction of reference-frame-free qudits

Jun Suzuki,¹ Gelo Noel Macuja Tabia,^{1,2} and Berthold-Georg Englert^{1,2}

¹Centre for Quantum Technologies, National University of Singapore, Singapore 117543, Singapore

²Department of Physics, National University of Singapore, Singapore 117542, Singapore

(Dated: 12 February 2008)

By exploiting a symmetric scheme for coupling N spin-1/2 constituents (the physical qubits) to states with total angular momentum $N/2 - 1$, we construct rotationally invariant logical qudits of dimension $d = N - 1$. One can encode all qudit states, and realize all qudit measurements, by this construction. The rotational invariance of all relevant objects enables one to transmit quantum information without having aligned reference frames between the parties that exchange the qudits. We illustrate the method by explicit constructions of reference-frame-free qubits and qutrits.

PACS numbers: 03.67.Pp, 03.67.Lx, 03.65.Fd

The raw experimental data about physical systems or events are almost always tied to reference frames, defined by the coordinate systems to which the data refer. The comparison of data acquired by different observers then requires that they know how their reference frames are related to each other, such as whether the axes of the coordinate systems are aligned or rotated.

In the context of quantum information theory, the role of the reference frame has been reconsidered recently; see [1] for a summary. There is, in particular, an intimate connection with the concept of decoherence free (DF) subsystems and subspaces, which are important for experimental implementations of schemes for processing quantum information. In quantum cryptography, for example, the presence of decoherence lowers the efficiency of the quantum channel involved. Moreover, the lack of a shared reference frame between two distant parties becomes a practical problem when establishing a secure channel between them.

It is, therefore, reasonable to ask whether or not one could be free, in general, from the problem of decoherence or sharing of the reference frame. Put differently, one may ask if it is possible to construct arbitrary quantum states which are DF or reference-frame-free (RFF). This question has been widely discussed [1], and general arguments ensure the existence of such DF subsystems, DF subspaces, and RFF quantum states in any finite dimension. When it comes to their explicit construction, however, one encounters a situation in which one has to work out details whose number increases exponentially with the dimension of the Hilbert space in question.

In this contribution, we narrow this gap between the in-principle possibility and the in-practice difficulty by an explicit construction of all states of a d -dimensional RFF quantum system, the *RFF qudit*, out of $N = d + 1$ spin-1/2 constituents. We are thus making *logical* RFF qudits out of *physical* qubits, and the construction also identifies the d -dimensional DF subsystem.

The standard construction is based on successive addition of the angular momenta of the spin-1/2 ingredients with Clebsch-Gordan coefficients for the probability am-

plitudes. With more than two constituents, the classification of the final states depends on the order in which the individual spins are added, and the complexity grows very rapidly with the number of constituents [2].

Alternatively, there is the symmetric coupling that we will exploit here [3, 4]. The general symmetric coupling is known only for three angular momenta so far, and it is presently unknown if there is a symmetric coupling scheme for more than three angular momenta. In the special case of the coupling of N identical angular momenta, our study supports the possibility of the symmetric coupling [5]. In this paper we report the N spin-1/2 case and, taking advantage of the symmetric coupling scheme, we construct the RFF qudit immediately without the need for evaluating the Clebsch-Gordan coefficients.

The paper is organized as follows. We first present the symmetric coupling of N spin-1/2 constituents for the states with second-largest total angular momentum. Thereafter, we construct the general RFF qudit, and illustrate the procedure for RFF qubits and RFF qutrits.

The direct product of N spin-1/2 systems is a direct sum of irreducible representations of angular momentum,

$$\mathcal{D}_{1/2}^{\otimes N} = \bigoplus_{j \in J} c_j \mathcal{D}_j, \quad (1)$$

where the index set $J = \{N/2, N/2 - 1, \dots\}$ has $(N+1)/2$ elements if N is odd, and has $N/2 + 1$ elements if N is even. Here,

$$c_j = \frac{N!(2j+1)}{(N/2+j+1)!(N/2-j)!} \quad (2)$$

is the multiplicity of \mathcal{D}_j , the irreducible representation of angular momentum j . Our main concern is the subsystem of the second-largest angular momentum states, where we identify the d -dimensional DF subsystem or RFF qudit.

The states with second-largest angular momentum, $j_2 = N/2 - 1$, have a multiplicity of $c_{j_2} = N - 1$. Therefore, these states can be labeled by the eigenvalues of the z -component of the total angular momentum

$\vec{J} = \sum_{\ell=1}^N \vec{\sigma}^{(\ell)}/2$ together with the degeneracy label λ . That is, the state kets are denoted by $|j_2, m_2; \lambda\rangle$ with

$$m_2 = j_2, j_2 - 1, \dots, -j_2 \text{ and } \lambda = 1, 2, \dots, N - 1. \quad (3)$$

There are $d = 2j_2 + 1 = N - 1 = c_{j_2}$ states for each value of λ , so that we have $d^2 = (N-1)^2$ states for $j_2 = N/2 - 1$ in total.

Upon denoting the kets for the single-spin-1/2 states with $m = 1/2$ and $m = -1/2$ by $|0\rangle$ and $|1\rangle$, respectively, we have $|0^{\otimes N}\rangle = |00\dots 0\rangle$ for the unique state with maximal values of both j and m , that is $j_1 = m_1 = N/2$, and multiple applications of the ladder operator $J_- = J_x - iJ_y$ yield all states of maximal total angular momentum in the familiar way, $|j_1, m_1\rangle \propto J_-^{j_1 - m_1} |0^{\otimes N}\rangle$. Supplementing J_- are its d orthogonal partners $\Omega_-(\lambda)$, defined by

$$\Omega_-(\lambda) = \frac{1}{\sqrt{N}} \sum_{\ell=1}^N \omega_N^{\lambda\ell} \sigma_-^{(\ell)} \quad \text{with } \omega_N = e^{2\pi i/N}, \quad (4)$$

where $\sigma_-^{(\ell)}$ is the lowering operator for the ℓ th constituent. The angular momentum states with $m_2 = j_2$ are then obtained as $|j_2, j_2; \lambda\rangle = \Omega_-(\lambda) |0^{\otimes N}\rangle$, and successive applications of J_- give the remaining $|j_2, m_2; \lambda\rangle$. Since $\Omega_-(\lambda)$ and J_- commute with each other, we have,

$$\begin{aligned} |j_2, m_2; \lambda\rangle &= \sqrt{\frac{2(2j_2+1)(j_2+m_2)!}{j_2!(j_2-m_2)!}} \Omega_-(\lambda) J_-^{j_2-m_2} |0^{\otimes N}\rangle \\ &\propto \Omega_-(\lambda) |j_1, m_2+1\rangle, \end{aligned} \quad (5)$$

for which $\langle j_2, m_2; \lambda | j_2, m'_2; \lambda' \rangle = \delta_{m_2 m'_2} \delta_{\lambda \lambda'}$ states their orthonormality.

The discrete Fourier transformation that we chose in (4) is just one possibility for defining the $\Omega_-(\lambda)$ s and thus the kets $|j_2, m_2; \lambda\rangle$. More generally, any unitary $d \times d$ matrix U , with N th-row matrix elements $U_{N\ell} = N^{-1/2}$, can serve in $\Omega_-(\lambda) = \sum_{\ell=1}^N U_{\lambda\ell} \sigma_-^{(\ell)}$. For the specific choice of the discrete Fourier matrix, the projectors $|j_2, m_2; \lambda\rangle \langle j_2, m_2; \lambda|$ are invariant under the cyclic permutation $\vec{\sigma}^{(1)} \rightarrow \vec{\sigma}^{(2)} \rightarrow \dots \rightarrow \vec{\sigma}^{(N)} \rightarrow \vec{\sigma}^{(1)}$ of the spin-1/2 constituents. An analogous construction works for systems of N constituents with spin other than 1/2.

Here is a remark about the relation to the permutation symmetry. As is well known, the state space of representations of N identical angular momenta is also the state space of unitary representations of the symmetric group S_N . The elements of the permutation group, the unitary permutation operators, are invariant when the same unitary transformation is applied to all constituents, and therefore the permutation operators can be composed to form invariant operators. By requiring that the invariant objects are hermitian and nonnegative, one can then construct a proper quantum state. This fact was utilized in constructing the Werner state for two parties [6]. Consult Refs. [7, 8] for recent generalizations to more parties.

The basic operators for the RFF qudit in $d = N/2 - 1 = 2j_2 + 1$ dimension are the d^2 operators

$$Q_{\lambda\lambda'} = \sum_{m_2=-j_2}^{j_2} |j_2, m_2, \lambda\rangle \langle j_2, m_2, \lambda'| = Q_{\lambda'\lambda}^\dagger. \quad (6)$$

They commute with the vector operator of total angular momentum, $\vec{J} Q_{\lambda\lambda'} = Q_{\lambda\lambda'} \vec{J}$, and are closed under multiplication, $Q_{\lambda\lambda'} Q_{\lambda''\lambda'''} = \delta_{\lambda'\lambda''} Q_{\lambda\lambda''}$. It follows that $Q_{\lambda\lambda'}$ can be written in the tensor product form

$$Q_{\lambda\lambda'} = |\lambda\rangle \langle \lambda'| \otimes I_d, \quad (7)$$

where the first factor refers to the *signal* qudit, and the d -dimensional identity I_d refers to the *idler* qudit.

A qudit quantum state, specified by a $d \times d$ density matrix $\rho = \rho^\dagger \geq 0$ with matrix elements $\rho_{\lambda\lambda'}$, is then implemented by

$$\rho^{(\text{RFF})} = \sum_{\lambda, \lambda'=1}^d |\lambda\rangle \rho_{\lambda\lambda'} \langle \lambda'| \otimes \frac{1}{d} I_d, \quad (8)$$

where the signal qudit is in the state ρ and the idler qudit is in the completely mixed state. Whereas $\rho^{(\text{RFF})}$ is a mixed state of the N spin-1/2 constituents, with a binary entropy of $S(\rho^{(\text{RFF})}) = S(\rho) + \log_2 d$, the state of the signal qudit can be pure or mixed, whatever is the nature of the given qudit state ρ .

Any qudit positive-operator-valued measure (POVM), $\sum_k \Pi_k = I_d$ with $\Pi_k = \Pi_k^\dagger \geq 0$, can be realized as a POVM for the signal qudit by the analogous construction

$$\Pi_k^{(\text{RFF})} = \sum_{\lambda, \lambda'=1}^d |\lambda\rangle (\Pi_k)_{\lambda\lambda'} \langle \lambda'| \otimes I_d, \quad (9)$$

so that $\text{Tr}\{\rho^{(\text{RFF})} \Pi_k^{(\text{RFF})}\} = \text{Tr}\{\rho \Pi_k\}$ for all outcomes Π_k of the POVM under consideration.

The ambiguity in defining $\Omega_-(\lambda)$, mentioned in the paragraph following Eq. (5), carries over to $\rho^{(\text{RFF})}$ and $\Pi_k^{(\text{RFF})}$, which are equally ambiguous. But once the $\Omega_-(\lambda)$ are chosen, the above construction gives a definite implementation of ρ and Π_k .

As a first illustration we consider the RFF qubit ($d = 2$, $j_2 = 1/2$) composed of $N = 3$ spin-1/2 constituents. It is both convenient and systematic to express all operators in terms of the unitary and hermitian swap operators $P_{jk} = P_{kj} = (1 + \vec{\sigma}^{(j)} \cdot \vec{\sigma}^{(k)})/2$ (for $j \neq k$) that permute the j th and k th constituents: $P_{jk} \vec{\sigma}^{(k)} = \vec{\sigma}^{(j)} P_{jk}$. Since these swap operators are obviously invariant under the collective rotations, so are

$$Q_{12} = Q_{21}^\dagger = \frac{1}{3} (P_{12} + \omega_3 P_{23} + \omega_3^2 P_{31}), \quad (10)$$

and

$$\left. \begin{aligned} Q_{11} &= Q_{11}^\dagger = Q_{12}Q_{21} \\ Q_{22} &= Q_{22}^\dagger = Q_{21}Q_{12} \end{aligned} \right\} = \frac{1}{2} - \frac{1}{6}(P_{12} + P_{23} + P_{31}) \\ \mp \frac{i}{\sqrt{12}}[P_{31}, P_{12}]. \quad (11)$$

The components of the hermitian Pauli vector $\vec{\sigma}^{(\text{RFF})}$ for the RFF qubit are then given by [9]

$$\sigma_x^{(\text{RFF})} + i\sigma_y^{(\text{RFF})} = 2Q_{12}, \quad \sigma_z^{(\text{RFF})} = Q_{11} - Q_{22}, \quad (12)$$

and $I_{j=1/2} = Q_{11} + Q_{22}$ is the projector onto the $j_2 = 1/2$ subspace of the signal and idler qubits. Explicit expression, such as $\sigma_z^{(\text{RFF})} = -\vec{\sigma}^{(1)} \cdot (\vec{\sigma}^{(2)} \times \vec{\sigma}^{(3)})/\sqrt{12}$, emphasize once more that these operators are rotationally invariant, indeed, and therefore any orientation in space of the x, y , and z axes of the Cartesian reference frame is as good as any other.

Here is an example of a state preparation. By preparing the third spin-1/2 constituent in a completely mixed state and the first and second in their singlet state, the experimenter puts the three physical qubits into the mixed state

$$\rho_3 = \frac{1}{4}(1 - \vec{\sigma}^{(1)} \cdot \vec{\sigma}^{(2)}) = \frac{1}{2}(1 - P_{12}) \\ = \frac{1}{2}(Q_{11} + Q_{22} - Q_{12} - Q_{21}) \hat{=} \frac{1}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \quad (13)$$

which is a pure state for the signal qubit. Likewise, $\rho_1 = (1 - P_{23})/2$ and $\rho_2 = (1 - P_{31})/2$ are pure states of the signal qubit, and together these states make up a trine.

When one considers the higher-dimensional general case, it is useful to employ a more systematic construction of the RFF qudit. One possible way is to expand the state in terms of the generators for a $SU(d)$ Lie group with real coefficients. The standard Gell-Mann matrices together with the positivity requirement provide proper d level quantum states [10, 11]. Alternatively, one can use the unitary Heisenberg-Weyl-Schwinger (HWS) operator basis [12]. The complete set of unitary operators are given by $U^j V^k$ ($j, k = 1, 2, \dots, d$), where the unitary operators U and V have period d , $U^d = V^d = 1$, and $U^j V^k = \omega_d^{-jk} V^k U^j$ with $\omega_d = \exp(2\pi i/d)$. In the following we will use this HWS operator basis.

Explicitly, the RFF U and V unitary operators are given by

$$U_d = \sum_{\lambda=1}^d \omega_d^\lambda Q_{\lambda\lambda}, \quad V_d = \sum_{\lambda=1}^{d-1} Q_{\lambda\lambda+1} + Q_{d1}. \quad (14)$$

For the RFF qubit of Eqs. (10)–(12), these are simply $U_2 = -\sigma_z^{(\text{RFF})}$ and $V_2 = \sigma_x^{(\text{RFF})}$.

The next example is the case of four spin-1/2 constituents. This system is known to provide the DF qubit

subspace spanned by two spin-0 states, which have been studied using the standard Clebsch-Gordan coefficients, and have been experimentally demonstrated [13]. Our emphasis here is the usefulness of the RFF qutrit construction. As a byproduct, the symmetric coupling yields an alternative choice for the basic spin-0 states, with properties different from the spin-0 states obtained by the Clebsch-Gordan coupling scheme.

For $N = 4$, the second-largest angular momentum states have $j_2 = 1$ and are triply degenerate. With $\omega_4 = i$, they are

$$\begin{aligned} |1, 1; \lambda\rangle &= \frac{1}{2}(\omega_4^\lambda |1000\rangle + \omega_4^{2\lambda} |0100\rangle \\ &\quad + \omega_4^{3\lambda} |0010\rangle + |0001\rangle), \\ |1, 0; \lambda\rangle &= \frac{1}{\sqrt{8}}[(\omega_4^\lambda + 1)(|1001\rangle - |0110\rangle) \\ &\quad + (\omega_4^{2\lambda} + 1)(|0101\rangle - |1010\rangle) \\ &\quad + (\omega_4^{3\lambda} + 1)(|0011\rangle - |1100\rangle)], \\ |1, -1; \lambda\rangle &= -\frac{1}{2}(\omega_4^\lambda |0111\rangle + \omega_4^{2\lambda} |1011\rangle \\ &\quad + \omega_4^{3\lambda} |1101\rangle + |1110\rangle), \end{aligned} \quad (15)$$

when expressed as superpositions of the basic product states of the four spin-1/2 constituents. And the construction (14) gives

$$\begin{aligned} U_3 &= \frac{\omega_3^2}{4}[2P_{13}P_{24} - P_{12}P_{34} - P_{14}P_{23} \\ &\quad - \frac{\sqrt{3}}{2}([P_{12}, P_{13}] + [P_{23}, P_{24}] \\ &\quad + [P_{34}, P_{31}] + [P_{41}, P_{42}])], \\ V_3 &= \frac{1}{4}[iP_{12}P_{34} - iP_{14}P_{23} + (1+i)(P_{12} - P_{34}) \\ &\quad + (1-i)(P_{23} - P_{14}) + P_{13} - P_{24}] \end{aligned} \quad (16)$$

for the basic HWS unitary operators.

Similarly, the $j_3 = 0$ states are given by

$$\begin{aligned} |0, 0; \lambda\rangle &= \frac{1}{\sqrt{6}}[\omega_3^\lambda (|1001\rangle + |0110\rangle) \\ &\quad + \omega_3^{2\lambda} (|0101\rangle + |1010\rangle) \\ &\quad + (|0011\rangle + |1100\rangle)]. \end{aligned} \quad (17)$$

Quite obviously, the two respective projectors ($\lambda = 1, 2$)

$$\begin{aligned} |0, 0; \lambda\rangle\langle 0, 0; \lambda| &= \frac{1}{3}(S_{12}S_{34} + S_{13}S_{24} + S_{14}S_{23}) \\ &\quad + \frac{i(-1)^\lambda}{\sqrt{12}}([S_{12}, S_{13}] - [S_{23}, S_{24}] \\ &\quad + [S_{34}, S_{31}] - [S_{41}, S_{42}]) \end{aligned} \quad (18)$$

are unchanged or interchanged when the spin-1/2 constituents are permuted. Here, $S_{jk} = (1 - \vec{\sigma}^{(j)} \cdot \vec{\sigma}^{(k)})/4$ is the singlet state between j th and k th constituents. This

invariance is in marked contrast to the lack of permutation invariance in the projectors

$$\begin{aligned} |S_1\rangle\langle S_1| &= \frac{1}{3}(-S_{12}S_{34} + 2S_{13}S_{24} + 2S_{14}S_{23}), \\ |S_2\rangle\langle S_2| &= S_{12}S_{34} \end{aligned} \quad (19)$$

onto the $j_3 = 0$ states of the standard successive coupling [14],

$$\begin{aligned} |S_1\rangle &= \frac{1}{2}(|0101\rangle + |1010\rangle - |1001\rangle - |0110\rangle), \\ |S_2\rangle &= \frac{1}{\sqrt{12}}(2|0011\rangle + 2|1100\rangle - |0101\rangle \\ &\quad - |1010\rangle - |0110\rangle - |1001\rangle). \end{aligned} \quad (20)$$

We note that the singlet states (17) have been introduced in Ref. [15]. A detailed comparison of the properties of the states (17) and the states (20) establishes that they should be regarded as quite different singlet bases [15, 16]. Put differently, the total angular momentum and its third component together with the degeneracy do not specify the state uniquely; one needs other quantum numbers in addition. In the case of N qubits, for example, the product states of the computational basis are uniquely labeled by N binary quantum numbers. Although this ‘‘missing label problem’’ is well known [17], it seems that it has not received due attention in quantum information theory as yet.

Lastly, for completeness we report the RFF qubit Pauli operators for the $j_3 = 0$ qubit. They are

$$\begin{aligned} \sigma_x^{(\text{RFF})} &= \frac{-2}{3}(2S_{12}S_{34} - S_{14}S_{23} - S_{13}S_{24}), \\ \sigma_y^{(\text{RFF})} &= \frac{-2}{\sqrt{3}}(S_{13}S_{24} - S_{14}S_{23}), \\ \sigma_z^{(\text{RFF})} &= \frac{-i}{\sqrt{3}}([S_{12}, S_{13}] - [S_{23}, S_{24}] \\ &\quad + [S_{34}, S_{31}] - [S_{41}, S_{42}]), \end{aligned} \quad (21)$$

and the projector onto the singlet states is

$$I_{j=0} = \frac{2}{3}(S_{12}S_{34} + S_{13}S_{24} + S_{14}S_{23}). \quad (22)$$

All RFF qubit states and operators are now constructed from $I_{j=0}$ and $\vec{\sigma}^{(\text{RFF})}$ in the usual manner.

We note that tracing out either one of the spin-1/2 constituents from the four-qubit RFF Pauli operators in (21) leads to the three-qubit RFF Pauli operators of (12). Accordingly, they are different realizations of the same $\text{su}(2)$ Lie algebra by representations with different dimensions. In other words, the amount of information carried by the logical RFF qubit is exactly the same irrespective of the construction in terms of three or four physical qubits.

Further, since the Hilbert spaces for the RFF qutrit and RFF qubit are two terms in the direct sum of (1), one can utilize the two RFF quantum systems jointly.

The simultaneous laboratory implementation of a RFF qutrit and a RFF qubit, in rotationally invariant states of four physical qubits, would be very interesting indeed.

In summary, we have given the systematic and rather simple construction of RFF quantum states in any arbitrary dimension. The symmetric coupling of N spin-1/2 constituents distinguishes our scheme from others. This novel coupling scheme immediately leads to the general explicit expression for the RFF qubits, which also provides alternative representations for a DF subsystem in any arbitrary dimension. We remark that the new expression for the DF subsystem possesses a higher symmetry under permutations than the subsystems obtained from other constructions, which is possibly advantageous for experimental implementations. Future studies should address this issue.

JS and BGE wish to thank Hans Briegel for the hospitality they enjoyed at the Institute for Quantum Optics and Quantum Information in Innsbruck. This work is supported by the National Research Foundation and Ministry of Education, Singapore. Support from A*STAR Temasek Grant No. 012-104-0040, NUS Grant WBS R-144-000-116-101, and SERC Grant 052 101 0043 is gratefully acknowledged.

-
- [1] S. D. Bartlett, T. Rudolph, and R. W. Spekkens, *Rev. Mod. Phys.* **79**, 555 (2007).
 - [2] L. C. Biedenharn and J. D. Louck, *Encyclopedia of Mathematics and Its Applications*, Vols. **8** and **9** (Addison-Wesley Publishing Company, 1981).
 - [3] A. Chakrabarti, *Ann. Inst. Henri Poincaré* **1**, 301 (1964).
 - [4] J.-M. Lévy-Leblond and M. Lévy-Nahas, *J. Math. Phys.* **6**, 1372 (1965).
 - [5] J. Suzuki, in preparation.
 - [6] R. F. Werner, *Phys. Rev. A* **40**, 4277 (1989).
 - [7] T. Eggeling and R. F. Werner, *Phys. Rev. A* **63**, 042111 (2001).
 - [8] D. Chruściński and A. Kossakowski, *Open Sys. & Info. Dyn.* **14**, 25 (2007).
 - [9] Equation (38) in S. D. Filippo, *Phys. Rev. A* **62**, 052307 (2000), states the RFF qubit operators (12) in the language of C^* algebras.
 - [10] G. Kimura, *Phys. Lett. A* **314**, 339 (2003).
 - [11] M. S. Byrd and N. Khaneja, *Phys. Rev. A* **68**, 062322 (2003).
 - [12] See, e.g., J. Schwinger, *Quantum Mechanics. Symbolism of Atomic Measurements* (Springer-Verlag, 2001).
 - [13] M. Bourennane, M. Eibl, S. Gaertner, C. Kurtsiefer, A. Cabello, and H. Weinfurter, *Phys. Rev. Lett.* **92**, 107901 (2004).
 - [14] There exist other expressions for the singlets with Clebsch-Gordan coefficients [2].
 - [15] A. Higuchi and A. Sudbery, *Phys. Lett. A* **273**, 213 (2001).
 - [16] S. Brierley and A. Higuchi, *J. Phys. A* **40**, 8455 (2007).
 - [17] A. Peccia and R. T. Sharp, *J. Math. Phys.* **17**, 1313 (1976).