

# Quantum mechanical sum rules for two model systems

M. Belloni\*

*Physics Department*

*Davidson College*

*Davidson, NC 28035 USA*

R. W. Robinett<sup>†</sup>

*Department of Physics*

*The Pennsylvania State University*

*University Park, PA 16802 USA*

(Dated: February 15, 2008)

## Abstract

Sum rules have played an important role in the development of many branches of physics since the earliest days of quantum mechanics. We present examples of one-dimensional quantum mechanical sum rules and apply them in two familiar systems, the infinite well and the single  $\delta$ -function potential. These cases illustrate the different ways in which such sum rules can be realized, and the varying mathematical techniques by which they can be confirmed. Using the same methods, we also evaluate the second-order energy shifts arising from the introduction of a constant external field, namely the Stark effect.

PACS numbers: 03.65.-w, 03.65.Db, 11.55.Hx

## I. INTRODUCTION

Quantum mechanical identities which relate time-dependent expectation values, influential in the early days of quantum theory, continue to act as useful pedagogical tools in the modern curriculum. For example, the results often known as Ehrenfest's theorem(s),<sup>1</sup>

$$\langle \hat{p} \rangle_t = m \frac{d\langle x \rangle_t}{dt} \quad \text{and} \quad m \frac{d^2\langle x \rangle_t}{dt^2} = - \left\langle \frac{dV(x)}{dx} \right\rangle_t \quad (1)$$

can be used to show that time-dependent quantum expectation values are related to their corresponding classical equations of motion.<sup>2</sup> Identities restricted to time-independent expectation values evaluated using energy eigenstates,  $|n\rangle$ , such as the quantum virial theorem

$$\langle n | \hat{T} | n \rangle = \left\langle n \left| \frac{\hat{p}^2}{2m} \right| n \right\rangle = \frac{1}{2} \left\langle n \left| x \frac{dV(x)}{dx} \right| n \right\rangle \quad (2)$$

and related hypervirial theorems,<sup>3</sup> are historically and pedagogically valuable as they too have clear classical analogs and can often be evaluated without resorting to direct integration.

Similar relationships involving off-diagonal matrix elements, especially quantum mechanical sum rules, were also used to dramatic effect in the early days of quantum theory. For example, the Thomas-Reiche-Kuhn (TRK) energy-weighted sum rule,<sup>4</sup> again for energy eigenstates,

$$\sum_k (E_k - E_n) |\langle n | x | k \rangle|^2 = \frac{\hbar^2}{2m} \quad (\text{TRK sum rule}) \quad (3)$$

was used to describe the physics of electric-dipole interactions with atoms. It was originally obtained by requiring that the Kramers-Heisenberg dispersion relation reduce to the Thomas scattering formula at high energies. Written in the form

$$\sum_k \frac{2m(E_k - E_n)}{\hbar^2} |\langle n | x | k \rangle|^2 = \sum_k f_{n,k} = 1 \quad (4)$$

this was an important experimental check of the oscillator strengths ( $f_{n,k}$ ) and an early confirmation of quantum results. Kramer was able to derive this relation in the context of matrix mechanics, reproducing the matrix version of the famous commutation relation  $[x, p] = i\hbar$ .<sup>5</sup> Other early uses of sum rules included Bethe's study of energy loss mechanisms for charged particles in matter,<sup>6</sup> which made use of the relation

$$\sum_k (E_k - E_n) |\langle n | e^{iqx} | k \rangle|^2 = \frac{\hbar^2 q^2}{2m} \quad (\text{Bethe sum rule}), \quad (5)$$

eventually leading to the Bethe-Bloch formula.

Since then, sum rules have been used in many areas of physics, including in atomic,<sup>7</sup> molecular,<sup>8</sup> solid state,<sup>9,10,11,12</sup> nuclear,<sup>13,14,15,16</sup> and especially in particle physics.<sup>17,18,19,20,21</sup> One well-known paper applying sum rule methods to QCD<sup>22</sup> is the 10<sup>th</sup> most highly cited paper in the particle physics literature and over 2,000 papers on QCD sum rules have been published, with 60 appearing in 2007 alone.<sup>23</sup>

The power of such sum rule identities is that they encode a large amount of information about both the energy spectrum and energy eigenfunctions of the system in a compact form, often in a way which is amenable to experimental confirmation. This in turn can probe assumptions about the fundamental interactions assumed or the calculation methods used to approximate physical systems. For example, QCD sum rules have been used to extract values of both the light and heavy quark masses, which are not otherwise directly measurable quantities.

Despite their historical and contemporary importance, sum rules are not often treated in the context of standard quantum mechanics courses. The TRK sum rule is sometimes included in undergraduate quantum mechanics books,<sup>24</sup> but often only as a problem, and typically only using the harmonic oscillator. This lack of coverage may well be due to the paucity of tractable examples in familiar model systems to which students typically have exposure, or the level of mathematical analysis required to verify even the simplest cases.

The purpose of this paper is to provide a suite of one-dimensional sum rules and to demonstrate the mathematical techniques required for their confirmation in two model quantum mechanical systems, the infinite well and the single (attractive)  $\delta$ -function potential. In each case, the sum rules saturate differently and rely on different mathematical methods (summation techniques and contour integration methods) illustrating the diverse ways in which such sum rules are realized. The level of mathematical detail required, however, is low enough to be easily accessible to advanced undergraduate students.

Explicitly confirming that these identities are indeed satisfied is not an empty exercise since it is possible to obtain surprising results, even from relatively simple systems such as the rigid rotator.<sup>25</sup> In addition, energy-weighted sum rule calculations are actually not exotic, since perturbation theory is discussed in standard textbooks in quantum mechanics. The expression for the second-order shift in the energy due to a perturbation  $V'(x)$  is given

by

$$E_n^{(2)} = \sum_{k \neq n} \frac{|\langle n | V'(x) | k \rangle|^2}{(E_n^{(0)} - E_k^{(0)})} \quad (6)$$

which is a form of energy-weighted sum rule. Using this connection, we will find that we can make use of the exactly same techniques derived for confirming sum rules to evaluate the shift due to the addition of a constant external field,  $V'(x) = Fx$ , namely the Stark shift, in each of the model systems we consider.

Introducing the concept of sum rules, using the tractable examples considered here, can certainly help students appreciate their use in later research applications. It can also help put the mathematical methods used in the same context as more familiar second-order perturbation theory calculations, one of the most important applications of undergraduate quantum mechanics, showing how the related required sums over intermediate states can be sometimes done in closed-form, and compared with explicit results.

## II. SUM RULE EXAMPLES

The derivation of many energy weighted sum rules has been succinctly described<sup>11</sup> as making use of a ‘...well-known technique which involves closure and evaluating a double commutator in two different ways.’ Such calculations rely on the fact that the solutions of the system under study form a complete set of states. For example, consider a system with energy eigenstates satisfying  $\hat{H}|n\rangle = E_n|n\rangle$ . Then, for an arbitrary operator,  $\hat{O}$ , we have the sum over off-diagonal matrix elements

$$\begin{aligned} \sum_{all\ k} |\langle n | \hat{O} | k \rangle|^2 &= \sum_{all\ k} \langle n | \hat{O} | k \rangle \langle k | \hat{O} | n \rangle \\ &= \langle n | \hat{O} \left\{ \sum_{all\ k} |k\rangle \langle k| \right\} \hat{O} | n \rangle \\ &= \langle n | \hat{O}^2 | n \rangle. \end{aligned} \quad (7)$$

We note that the sum over the complete set of intermediate states,  $|k\rangle$ , can include both an infinite sum (for discrete levels), an integral (for continuum states), or both.

For the special case of  $\hat{O} = x$ , we obtain the simplest dipole matrix element sum rule listed in Bethe and Jackiw,<sup>26,27</sup> namely

$$\sum_k |\langle n | x | k \rangle|^2 = \langle n | x^2 | n \rangle \quad (x\text{-closure sum rule}) \quad (8)$$

with an identical sum rule relation for the off-diagonal matrix elements of the momentum operator.

To derive the Thomas-Reiche-Kuhn (TRK) sum rule, we start with two commutation relations,

$$[\hat{p}, x] = \frac{\hbar}{i} \quad \text{and} \quad [\hat{H}, x] = \frac{1}{2m}[\hat{p}^2, x] = \frac{\hbar}{mi} \hat{p} \quad (9)$$

where we assume a standard 1D Hamiltonian of the form

$$\hat{H} = \frac{\hat{p}^2}{2m} + V(x). \quad (10)$$

The first of these relations can be written in the form

$$\frac{\hbar}{i} = \langle n | \hat{p}x - x\hat{p} | n \rangle = \sum_{\text{all } k} \{ \langle n | \hat{p} | k \rangle \langle k | x | n \rangle - \langle n | x | k \rangle \langle k | \hat{p} | n \rangle \} \quad (11)$$

where we have inserted a complete set of states. The second relation in Eqn. (9) can be written as

$$\langle n | \hat{p} | k \rangle = \frac{im}{\hbar} \langle n | [\hat{H}, x] | k \rangle = \frac{im(E_n - E_k)}{\hbar} \langle n | x | k \rangle \quad (12)$$

with a similar expression for  $\langle k | \hat{p} | n \rangle$ . When used in Eqn. (11), this gives the desired result,

$$\frac{\hbar^2}{2m} = \sum_k (E_k - E_n) |\langle n | x | k \rangle|^2. \quad (13)$$

Wang<sup>28</sup> has derived a very general expression for the energy-difference weighted sum rules for the matrix elements of a well-behaved function of  $x$ ,  $F(x)$ , namely

$$\sum_k (E_k - E_n) |\langle n | F(x) | k \rangle|^2 = \frac{\hbar^2}{2m} \left\langle n \left| \frac{dF(x)}{dx} \frac{dF^\dagger(x)}{dx} \right| n \right\rangle \quad (14)$$

which simplifies if the function is Hermitian so that  $F(x) = F^\dagger(x)$ . This general result can be used to immediately reproduce the TRK sum rule by using  $F(x) = x$ . We can then also derive the Bethe sum rule<sup>6</sup> by using  $\hat{O} = e^{iqx}$  in which case we find

$$\sum_k (E_k - E_n) |\langle n | e^{iqx} | k \rangle|^2 = \frac{\hbar^2 q^2}{2m}. \quad (15)$$

If we use  $F(x) = x^2$ , we obtain the so-called ‘monopole sum rule,’ which has been used in applications to nuclear collective excitations,<sup>14</sup>

$$\sum_k (E_k - E_n) |\langle n | x^2 | k \rangle|^2 = \frac{2\hbar^2}{m} \langle n | x^2 | n \rangle. \quad (16)$$

Wang<sup>28</sup> also discussed sum rules involving functions of the momentum operator, and ‘mixed’  $x, \hat{p}$  relations.

Bethe and Jackiw<sup>26,27</sup> derive several other sum rules for dipole moment matrix elements by using multiple commutation relations with the Hamiltonian, thus generalizing Eqn. (9), and yielding higher powers of the energy difference:

$$\begin{aligned} \sum_k (E_k - E_n)^2 |\langle n|x|k \rangle|^2 &= \frac{\hbar^2}{m^2} \langle n|\hat{p}^2|n \rangle \\ &= \frac{2\hbar^2}{m} \{E_n - \langle n|V(x)|n \rangle\} . \end{aligned} \quad (17)$$

$$\sum_k (E_k - E_n)^3 |\langle n|x|k \rangle|^2 = \frac{\hbar^4}{2m^2} \left\langle n \left| \frac{d^2V(x)}{dx^2} \right| n \right\rangle \quad (18)$$

and

$$\sum_k (E_k - E_n)^4 |\langle n|x|k \rangle|^2 = \frac{\hbar^4}{m^2} \left\langle n \left| \left( \frac{dV(x)}{dx} \right)^2 \right| n \right\rangle \quad (19)$$

where Eqns. (18) and (19) are described as the “*force times momentum*” and “*force squared*” sum rules, respectively.

We note that not all of these sum rules are guaranteed to converge<sup>26</sup> and in our case, because of the singular nature of the potentials used here as idealized models (the infinite well and the single- $\delta$  cases) several of these sum rules will not be applicable.

### III. THE INFINITE SQUARE WELL

The infinite square well (ISW) potential is the most frequently presented of all textbook examples of bound state systems and is frequently used as a model system to introduce students to tractable examples of research level physics, such as wave packet revivals.<sup>29</sup> We can confirm many of the sum rules discussed above for this case, making use of relatively straightforward mathematical techniques to evaluate the infinite sums which appear. (The only example we can find in the literature of the evaluation of sum rules in the context of the infinite well is a short discussion in Ref. [14].)

We consider the standard ISW potential, defined by

$$V(x) = \begin{cases} 0 & \text{for } 0 < x < a \\ \infty & \text{for } x < 0 \text{ and } x > a \end{cases} . \quad (20)$$

The energy eigenstates and corresponding eigenvalues are

$$\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right) \quad \text{and} \quad E_n = \frac{\hbar^2 n^2 \pi^2}{2ma^2} \quad (21)$$

where  $n = 1, 2, \dots$  and the expectation value of  $x^2$  required for the closure sum rule in Eqn. (8) is easily calculated to be

$$\langle n|x^2|n\rangle = a^2 \left( \frac{1}{3} - \frac{1}{2n^2\pi^2} \right). \quad (22)$$

The energy differences needed for the various sum rule calculations are given by

$$E_k - E_n = \frac{\hbar^2 \pi^2}{2ma^2} (k^2 - n^2), \quad (23)$$

while the off-diagonal matrix elements are given by

$$\begin{aligned} \langle n|x|k\rangle &= \frac{2}{a} \int_0^a \sin\left(\frac{n\pi x}{a}\right) x \sin\left(\frac{k\pi x}{a}\right) dx \\ &= \begin{cases} 0 & k+n \text{ even} \\ -(8na/\pi^2)[k/(k^2 - n^2)^2] & k+n \text{ odd} \end{cases} \end{aligned} \quad (24)$$

so that for  $n$  even (odd) only odd (even) values of  $k$  will contribute. This result is due to the energy eigenfunctions' generalized parity property relative to the center of the well at  $x = a/2$ . For the closure identity in Eqn. (8), we also need to include the diagonal matrix element,

$$\langle n|x|n\rangle = \frac{a}{2}. \quad (25)$$

This term does not contribute to the other sum rules, since the  $k = n$  term is suppressed by the  $(E_k - E_n)$  energy difference factor. In contrast to potential energy functions that are symmetric about the origin, such as the harmonic oscillator potential and the single  $\delta$ -function potential, the ISW potential as defined above is not symmetric and one must consider the  $k = n$  case for the closure identity.

The position closure sum rule in Eqn. (8) then reads

$$\sum_{\text{all } k} |\langle n|x|k\rangle|^2 = \left(\frac{a}{2}\right)^2 + \left(\frac{8na}{\pi^2}\right)^2 \sum_k \frac{k^2}{(k^2 - n^2)^4} \quad (26)$$

where the summation is over even (odd) values  $k$  if  $n$  is odd (even). This is the first of many examples we will encounter where we require infinite summations of the form

$$S_p^{(\pm)}(z) = \sum_k \frac{1}{(k^2 - z^2)^p} \quad (27)$$

where  $z$  takes on integral values, and where the summation is over odd,  $S^{(-)}$ , or even,  $S^{(+)}$ , values of  $k$ . For example, the required summation in Eqn. (26) can be written in the form

$$\sum_k \frac{k^2}{(k^2 - n^2)^4} = \sum_k \frac{(k^2 - n^2 + n^2)}{(k^2 - n^2)^4} = S_3^{(\pm)}(n) + n^2 S_4^{(\pm)}(n). \quad (28)$$

We provide a brief, but complete, review of how all of the sums required in this section can be evaluated using standard series expansions in Appendix A. We note, however, that modern computer algebra systems (such as Mathematica) can easily handle such sums. Students may be allowed on first pass to use such tools and then asked to delve more deeply into the methods used to obtain the general mathematical results for this class of problems.

For example, in modified Mathematica syntax, the summation over even integers  $k$  (relevant for  $n$  odd), yields:

```
Sum[k^2/(k^2-z^2)^4,{k,2,Infinity,2}]
= (-12 Pi Cot[Pi z/2] - 6 Pi^2 z Csc[Pi z/2]^2
+ 2 Pi^4 z^3 Cot[Pi z/2]^2 Csc[Pi z/2]^2
+ Pi^4 z^3 Csc[Pi z/2]^4)/768z^5
```

so that for odd integer values of  $z = n$ , we have (by hand or by using `Assuming-> z ∈ Integers` in Mathematica)

$$\sum_{k \text{ even}} \frac{k^2}{(k^2 - n^2)^4} = \frac{\pi^4 n^3 - 6\pi^2 n}{768n^5} = \frac{\pi^4}{768n^2} - \frac{\pi^2}{128n^4}. \quad (29)$$

We obtain the same result (same function of  $n$ ) for the summation over odd values of  $k$  (relevant for even  $n$ ). A trivial modification (one character in fact) of the Mathematica code is all that is required. Using this result in Eqn. (26), we then find that

$$\begin{aligned} \sum_{\text{all } k} |\langle n|x|k \rangle|^2 &= |\langle n|x|n \rangle|^2 + \sum_{k \neq n} |\langle n|x|k \rangle|^2 \\ &= \frac{a^2}{4} + \frac{64a^2 n^2}{\pi^4} \left( \frac{\pi^4}{768n^2} - \frac{\pi^2}{128n^4} \right) \\ &= a^2 \left( \frac{1}{3} - \frac{1}{2n^2 \pi^2} \right) = \langle n|x^2|n \rangle \end{aligned} \quad (30)$$

as expected.

The TRK sum rule is then given by

$$\sum_k (E_k - E_n) |\langle n|x|k \rangle|^2 = \left( \frac{\hbar^2}{2m} \right) \left( \frac{64n^2}{\pi^2} \right) \sum_k \frac{k^2}{(k^2 - n^2)^3} \quad (31)$$



where the summation over  $k$  is only for even (odd) values for  $n$  odd (even). These sums can also be done in closed form and one finds

$$\begin{aligned}\mathcal{I}_n^{(+)}(z) &\equiv \sum_{k \text{ even}} \frac{k^2}{(k^2 - z^2)^3} = S_2^{(+)}(z) + z^2 S_3^{(+)}(z) \\ &= \frac{1}{64z^3} \left[ \pi^2 z \csc^2\left(\frac{\pi z}{2}\right) + 2\pi \cot\left(\frac{\pi z}{2}\right) - \pi^3 z^2 \cot\left(\frac{\pi z}{2}\right) \cos^2\left(\frac{\pi z}{2}\right) \right]\end{aligned}\quad (32)$$

$$\begin{aligned}\mathcal{I}_n^{(-)}(z) &\equiv \sum_{k \text{ odd}} \frac{k^2}{(k^2 - z^2)^3} = S_2^{(-)}(z) + z^2 S_3^{(-)}(z) \\ &= \frac{1}{64z^3} \left[ \pi^2 z \sec^2\left(\frac{\pi z}{2}\right) - 2\pi \tan\left(\frac{\pi z}{2}\right) + \pi^3 z^2 \tan\left(\frac{\pi z}{2}\right) \sec^2\left(\frac{\pi z}{2}\right) \right]\end{aligned}\quad (33)$$

and we note the similarities in form. Inserting the appropriate odd and even values of  $n$ , in each case we find that

$$\mathcal{I}_n^{(+)}(n) = \mathcal{I}_n^{(-)}(n) = \frac{\pi^2}{64n^2}\quad (34)$$

for all integral values of  $n$ . This result, when substituted into Eqn. (31), directly confirms the TRK sum rule.

Verification of the monopole sum rule in Eqn. (16) requires a small, but important modification of the summation methods. The off-diagonal matrix elements required for  $k \neq n$  are

$$\langle n|x^2|k\rangle = \frac{(-1)^{k-n} 8a^2 n}{\pi^2} \left( \frac{k}{(k^2 - n^2)^2} \right),\quad (35)$$

while for  $k = n$ , one uses the result in Eqn. (22). Since the  $k = n$  term does not contribute to the sum (because of the associated energy difference factor) the left-hand side of Eqn. (16) reduces to

$$\sum_k (E_k - E_n) |\langle n|x^2|k\rangle|^2 = \left( \frac{\hbar^2 \pi^2}{2ma^2} \right) \left( \frac{64n^2 a^4}{\pi^4} \right) \sum_{k \neq n} \frac{k^2}{(k^2 - n^2)^3}\quad (36)$$

and we must sum over all values of  $k \neq n$  since the even/odd pattern seen in the dipole matrix elements is not present in this case.

In order to evaluate this sum, just as discussed in Appendix A, we can first generalize the sum to non-integer values of  $n$ , and then rewrite the sum as

$$T(z; n) \equiv \sum_{k \neq n} \frac{k^2}{(k^2 - z^2)^3} = \left[ \sum_{\text{all } k} \frac{k^2}{(k^2 - z^2)^3} \right] - \frac{n^2}{(n^2 - z^2)^3}\quad (37)$$

where the second term corresponds to the ‘missing’ term in the  $k \neq n$  summation. The first sum can be evaluated for arbitrary  $z$ , giving the result

$$T(z; n) = \left( \frac{\pi \cot(\pi z) + \pi^2 z \csc^2(\pi z) - 2\pi^3 z^2 \cot(\pi z) \csc^2(\pi z)}{16z^3} \right) - \frac{n^2}{(n^2 - z^2)^3}.\quad (38)$$

Since we will be taking the limit where  $z \rightarrow n$  (and integral), we write  $z = n + \epsilon$  for general  $n$ , and we find that both terms have factors which diverge as  $1/\epsilon^3$ ,  $1/\epsilon^2$ , and  $1/\epsilon$ . If, however, we expand both terms about  $z = n$  (*i.e.* in small values of  $\epsilon$ ) we find that these divergences cancel, leaving the finite result

$$\lim_{z \rightarrow n} T(z; n) = \lim_{\epsilon \rightarrow 0} T(n + \epsilon; n) = T(n) = \frac{\pi^2}{16n^2} \left( \frac{1}{3} - \frac{1}{2n^2\pi^2} \right) \quad (39)$$

which when inserted into Eqn. (36) reproduces the right-hand side of Eqn. (16).

Many of the other sum rules discussed in Sec. II, such as those that require derivatives of the potential energy function, Eqns. (18) and (19), are not well-defined for the infinite square well (or the single  $\delta$ -function in Sec. IV) due to the singular nature of the potential energy function. While the matrix elements  $\langle n|e^{iqx}|k\rangle$  required for the Bethe sum rule in Eqn. (5) are easily obtained in closed form, the summation methods discussed here are not immediately applicable.

We can now use identical methods to evaluate the second-order shift of the energy levels of the infinite square well due to the addition of a linear potential,  $V'(x) = Fx$ , namely the Stark effect. In this geometry, where the ISW potential is not symmetric, the first-order energy shift is non-vanishing and is given by

$$E_n^{(1)} = \langle n|Fx|n\rangle = \frac{aF}{2}. \quad (40)$$

The second-order shift has been evaluated by Mavromatis for the ground state<sup>30</sup> and then extended to a general state<sup>31,32</sup> by using variations on the Dalgarno-Lewis method.<sup>33</sup> If we explicitly write the standard expression for the second-order energy shift, we have

$$E_n^{(2)} = \sum_{k \neq n} \frac{|\langle n|Fx|k\rangle|^2}{(E_n^{(0)} - E_k^{(0)})} = - \left( \frac{F^2 2ma^2}{\hbar^2} \right) \left( \frac{8na}{\pi^2} \right)^2 \sum_k \frac{k^2}{(k^2 - n^2)^5} \quad (41)$$

which is formally identical to the class of summations discussed here. Using either Mathematica or the results of Appendix A we find that the required sum (for either  $n$  even or odd) is given by

$$\sum_k \frac{k^2}{(k^2 - n^2)^5} = \frac{15\pi^2 n - \pi^4 n^3}{3072n^7} \quad (42)$$

so that

$$E_n^{(2)} = -F^2 \left( \frac{ma^4}{\hbar^2} \right) \left( \frac{15 - (n\pi)^2}{24\pi^2 n^4} \right). \quad (43)$$

The overall  $n$ -dependent form agrees with the results of Mavromatis,<sup>30,31,32</sup> who considered the related problem of the symmetric infinite well, for which the first-order correction vanishes. This result is interesting in itself as the second order shift for the ground state is negative (as it always should be, since all states contributing to Eqn. (41) are higher in energy) but for  $n = 2$  and higher, the shift changes sign. This is in contrast to the behavior of the harmonic oscillator, where the second-order shift is always negative, independent of quantum number.

#### IV. THE SINGLE $\delta$ -FUNCTION POTENTIAL

Another popular model system in which to investigate sum rule and perturbation theory results is the single (attractive)  $\delta$ -function potential, defined here by

$$V_\delta(x) = -g\delta(x). \quad (44)$$

The use of  $\delta$ -function potentials as simply soluble models of potential barriers or wells has a long history in quantum mechanics, going back at least to Kronig and Penney<sup>34</sup> who considered a ‘*series of equidistant rectangular barriers*’ and then took the limit where the ‘*...the breadth  $b$  of these barriers is made infinitely small and their height  $V_0$  infinitely large...*’ while not actually using the  $\delta$ -function notation.

Morse and Feshbach<sup>35</sup> explicitly considered the form in Eqn. (44), make note of the correct (dis)continuity condition on the energy eigenfunction at the origin, namely

$$\psi'(0^+) - \psi'(0^-) = -\frac{2mg}{\hbar^2}\psi(0), \quad (45)$$

cite it as being ‘*useful in the study of nuclear forces,*’ and go on to discuss the single bound state as well as scattering solutions. Frost<sup>36</sup> considered single and multiple attractive  $\delta$ -function potentials as models of ‘*hydrogen-like atoms,*’ the hydrogen molecule-ion and more complex systems. He was perhaps the first to explicitly comment on the similarities of the energy eigenvalue and eigenfunction for the single bound state of this system to the ground state of the Coulomb problem. Since then, single and multiple  $\delta$ -function potentials have been widely used in model calculations in both the pedagogical and research literature.<sup>37</sup>

We note that compared to the two other most widely used simple 1D models, the infinite well and harmonic oscillator, the  $\delta$ -function potential has the advantage that it admits both

bound and continuum solutions, as does the Coulomb potential, and so it presents new features compared to purely discrete spectra.

The single bound ( $E < 0$ ) state for the potential in Eqn. (44) is given by

$$\psi_0(x) = \sqrt{K_0} e^{-K_0|x|} \quad (46)$$

where  $K_0 = \frac{mg}{\hbar^2}$  with the corresponding bound state energy eigenvalue

$$E_0 = -\frac{mg^2}{2\hbar^2} = -\frac{\hbar^2 K_0^2}{2m}. \quad (47)$$

One can then note many comparisons to the ground state of the hydrogen atom, if one defines the Coulomb potential as  $V_c(r) = -\frac{g}{r}$  and one defines and substitutes  $a_0 \equiv \frac{1}{K_0}$  in Eqns. (46) and (47). Not only does the form of the ground state energy in Eqn. (47) match that of the Coulomb potential, but the form of the energy eigenfunction in Eqn. (46) does as well.

For use in confirming the closure relations in Eqn. (8), we find that for the ground state energy eigenfunction we have

$$\langle 0|x^2|0\rangle = \frac{1}{2K_0^2}. \quad (48)$$

The  $E > 0$  continuum states can be classified by their parity and are given by

$$\psi_k^{(-)}(x) = \frac{1}{\sqrt{\pi}} \sin(kx) \quad (49)$$

$$\psi_k^{(+)}(x) = \frac{1}{\sqrt{\pi(k^2 + K_0^2)}} (K_0 \sin(k|x|) - k \cos(kx)) \quad (50)$$

both of which have the same free-particle energy  $E_k = \hbar^2 k^2 / 2m$ . The combination of the single bound state in Eqn. (46) and the continuum states in Eqns. (49) and (50) have been explicitly shown<sup>38</sup> to form a complete set of states. The effect of the continuum states on a simple perturbation theory calculation has also been demonstrated by Kiang.<sup>39</sup>

For the various sum rules, we will consider here the  $|n\rangle = |0\rangle$  case only, as others using purely continuum states do not converge. Because of the symmetry of the system, parity arguments dictate that the only non-zero dipole matrix elements connecting the single ground state to the continuum will arise from the  $\psi_k^{(-)}(x)$  states, and we find that

$$\langle 0|x|k^{(-)}\rangle = \int_{-\infty}^{+\infty} \left( \sqrt{K_0} e^{-K_0|x|} \right) x \left( \frac{1}{\sqrt{\pi}} \sin(kx) \right) dx = 4\sqrt{\frac{K_0^3}{\pi}} \frac{k}{(K_0^2 + k^2)^2}. \quad (51)$$

The energy differences are then given by

$$E_k - E_0 = \frac{\hbar^2}{2m}(k^2 + K_0^2). \quad (52)$$

and we note the similarities in form between these two expressions and the corresponding results for the ISW in Eqns. (23) and (24).

The dipole matrix element closure relation in Eqn. (8) then becomes

$$\sum_k |\langle 0|x|k^{(-)}\rangle|^2 = \left(\frac{16K_0^3}{\pi}\right) \int_0^\infty \frac{k^2}{(k^2 + K_0^2)^4} dk = \frac{1}{2K_0^2} = \langle 0|x^2|0\rangle \quad (53)$$

where the integral can be done by standard methods, and agrees with the value in Eqn. (48).

The left hand-side of the TRK sum rule in Eqn. (3) gives

$$\int_0^\infty (E_k - E_0) |\langle 0|x|k^{(-)}\rangle|^2 dk = \frac{\hbar^2}{2m} \left(\frac{16K_0^3}{\pi}\right) \int_0^{+\infty} \frac{k^2}{(k^2 + K_0^2)^3} dk = \frac{\hbar^2}{2m} \quad (54)$$

also as expected. We note the similarity in form of these integral expressions to the summation results for the infinite well in Eqns. (31) and (26).

In order to confirm the monopole sum rule in Eqn. (16), we require the off-diagonal matrix elements of  $x^2$  for which only the even continuum states in Eqn. (50) contribute, giving

$$\langle 0|x^2|k^{(+)}\rangle = 8\sqrt{\frac{K_0^3}{\pi(k^2 + K_0^2)}} \frac{k}{(k^2 + K_0^2)^2}. \quad (55)$$

We then find that

$$\sum_k (E_k - E_0) |\langle 0|x^2|k^{(+)}\rangle|^2 = \left(\frac{\hbar^2}{2m}\right) \left(\frac{64K_0^3}{\pi}\right) \int_0^\infty \frac{k^2}{(k^2 + K_0^2)^4} dk = \frac{\hbar^2}{mK_0^2} \quad (56)$$

and we note that factors of  $(k^2 + K_0^2)$  from the energy difference and energy eigenfunction normalization in the numerator and denominator respectively cancel.

The Stark effect for the single  $\delta$ -function potential has been analyzed using exact results from the Airy function solutions,<sup>40,41</sup> as well as using the Dalgarno-Lewis method.<sup>42</sup> Using the dipole matrix elements derived above, we can evaluate the second-order energy shift directly, using the same kinds of straightforward integrals encountered so far. We find that

$$\begin{aligned} E_0^{(2)} &= \int_k \frac{|\langle 0|Fx|k^{(-)}\rangle|^2}{(E_0^{(0)} - E_k^{(0)})} \\ &= -\frac{2mF^2}{\hbar^2} \left(\frac{16K_0^3}{\pi}\right) \int_0^\infty \left\{ \frac{1}{(k^2 + K_0^2)} \right\} \frac{k^2}{(k^2 + K_0^2)^4} dk \end{aligned} \quad (57)$$

which agrees with the results of Refs. [40]-[42], when put into this notation. We note that the entire contribution to the Stark shift in the ground state energy in this case comes from the continuum states and this result is one of the few examples of the explicit evaluation of the contribution of the continuum terms in such a calculation.

We recall that for the hydrogen atom ground state, the total second-order shift<sup>43</sup> can be written in the form

$$E_0^{(2)}(\text{H-atom}) = -\frac{9}{4} \left( \frac{F^2 a_0^3}{g} \right) \quad (58)$$

using the form of the Coulomb potential and perturbation theory, this result comes from summing over the contributions of both the bound states and continuum states. Ruffa<sup>44</sup> has evaluated the continuum contribution to the expression in Eqn. (58) in terms of a single integral and finds a net contribution of 0.4184 to the total  $9/4 = 2.25$  value of the prefactor. It is perhaps then fairer to compare the second-order Stark result in Eqn. (57), namely  $5/8 = 0.625$ , to that partial contribution.

Finally, the Bethe sum rule is given by

$$\mathcal{B} = \sum_k (E_k - E_0) |\langle 0 | e^{iqx} | k \rangle|^2 = \frac{\hbar^2 q^2}{2m} \quad (59)$$

and in this case we will have two contributions to the left-hand-side, coming from the even (*e*) or odd (*o*) continuum states, namely

$$\begin{aligned} \mathcal{B} &= \mathcal{B}_e + \mathcal{B}_o \\ &\equiv \int_0^\infty (E_k - E_0) |\langle 0 | \cos(qx) | k^{(+)} \rangle|^2 dk \\ &\quad + \int_0^\infty (E_k - E_0) |\langle 0 | \sin(qx) | k^{(-)} \rangle|^2 dk \end{aligned} \quad (60)$$

and we consider each term separately. The first matrix element of interest is

$$\begin{aligned} \mathcal{I}_o &= \langle 0 | \sin(qx) | k^{(-)} \rangle \\ &= \sqrt{\frac{4K_0}{\pi}} \int_0^\infty e^{-K_0 x} \sin(qx) \sin(kx) dx \\ &= \sqrt{\frac{4K_0}{\pi}} \left[ \frac{2kqK_0}{[(k+q)^2 + K_0^2][(k-q)^2 + K_0^2]} \right] \end{aligned} \quad (61)$$

where we use the symmetry of the energy eigenfunctions to evaluate the integral over positive values of  $x$  only. Recalling that  $E_k - E_0 = \hbar^2(k^2 + K_0^2)/2m$ , we find

$$\mathcal{B}_o = \frac{\hbar^2 q^2}{2m} \left( \frac{16K_0^3}{\pi} \right) \int_0^\infty \frac{k^2(k^2 + K_0^2)}{[(k+q)^2 + K_0^2]^2 [(k-q)^2 + K_0^2]^2} dk. \quad (62)$$

Use of an integrated mathematics package (again Mathematica) returns the correct value for the integral, provided one correctly interprets the many cautionary restrictions on the values of  $K_0$  and  $q$ . Given the relatively complicated nature of the intermediate results coming from such programs, however, it is again important to be able to check the expressions ‘by hand.’ In this case, it simply involves extending the integral over the entire real line (since the integrand is an even function of  $k$ ) and then using contour integration methods (see Appendix B for details), giving

$$\mathcal{B}_o = \left( \frac{\hbar^2 q^2}{2m} \right) \left\{ \frac{K_0^2 + q^2/2}{K_0^2 + q^2} \right\}. \quad (63)$$

For the even case, we require the matrix element

$$\begin{aligned} \mathcal{I}_e &= \langle 0 | \cos(qx) | k^{(+)} \rangle \\ &= \sqrt{\frac{4K_0}{\pi(K_0^2 + k^2)}} \int_0^\infty e^{-K_0 x} \cos(qx) [K_0 \sin(kx) - k \cos(kx)] dx \\ &= \sqrt{\frac{4K_0}{\pi(K_0^2 + k^2)}} \left[ \frac{-2kK_0 q^2}{[(k+q)^2 + K_0^2][(k-q)^2 + K_0^2]} \right] \end{aligned} \quad (64)$$

and the even contribution to the sum rule becomes

$$\begin{aligned} \mathcal{B}_e &= \int_0^\infty (E_k - E_0) |\langle 0 | \cos(qx) | k^{(+)} \rangle|^2 dk \\ &= \left( \frac{\hbar^2 q^2}{2m} \right) \left( \frac{8K_0^3 q^2}{\pi} \right) \int_{-\infty}^{+\infty} \frac{k^2}{[(k+q)^2 + K_0^2]^2 [(k-q)^2 + K_0^2]^2} dk. \end{aligned} \quad (65)$$

The integral can again be done with similar contour methods giving the result

$$\mathcal{B}_e = \left( \frac{\hbar^2 q^2}{2m} \right) \left\{ \frac{q^2/2}{K_0^2 + q^2} \right\} \quad (66)$$

which can be combined with Eqn. (63) to give

$$\mathcal{B} = \mathcal{B}_o + \mathcal{B}_e = \frac{\hbar^2 q^2}{2m} \quad (67)$$

as expected.

## V. CONCLUSIONS AND DISCUSSION

We have presented an array of familiar (and not-so-familiar) one-dimensional sum rules, a number of which have proved useful in the development of many fields of physics. Using two

standard model systems as testbeds, we have illustrated the diverse ways in which such sum rules are confirmed, emphasizing the different mathematical techniques (infinite summation tricks and contour integration methods) used in each case. While the evaluation of the necessary summations or integrals can be simplified by the use of integrated mathematics programs, we have also provided the details necessary to demonstrate the same results from first principles.

We have also noted the striking similarities of some of the expressions which arise for the same sum rules in the infinite square well and single  $\delta$ -potential cases. Despite the qualitatively very different physical behavior of the two systems, they both begin with free-particle solutions. The infinite wall boundaries of the ISW force quantized eigenstates with  $E_n = \hbar^2 k_n^2 / 2m$ , while the attractive  $\delta$ -function gives the identical dispersion relation for the  $E > 0$  states, but with continuous  $k$ -values. The  $\delta$ -function case also includes one  $E < 0$  state for which the sinusoidal solution is analytically continued to the localized exponential form in Eqn. (46). The connections between these two model systems are seldom if ever stressed, but appear very naturally in these sum rule calculations.

We hope that the suite of exemplary problems discussed here can be useful to instructors in lectures as well as for homework problems, in both the advanced undergraduate and graduate quantum mechanics curriculum, especially by putting this important tool of theoretical physics into a historical and research context.

## APPENDIX A: INFINITE SUMS FOR THE SQUARE WELL PROBLEM

Many of the sum rule and second-order perturbation theory results in Sec. III for the infinite square well involve the evaluation of infinite sums of the forms

$$S_p^{(+)}(z) = \sum_{\text{even } k} \frac{1}{(k^2 - z^2)^p} \quad \text{or} \quad S_p^{(-)}(z) = \sum_{\text{odd } k} \frac{1}{(k^2 - z^2)^p} \quad (\text{A1})$$

where the both expressions are eventually evaluated using integral values of  $z = n$ , with  $n$  odd and even respectively so no divergences occur. While multi-purpose computer programs such as Mathematica can recognize and correctly evaluate such sums, it can be important for some students (and many instructors) to also be able to derive them ‘*from scratch*.’ To that end, in this Appendix we provide a very brief, but self-contained and complete, review of the mathematical tools necessary for their derivation from more basic results with which



students at this level should be quite familiar.

We begin by considering the general expression

$$S_p(z) \equiv \sum_{k=1}^{\infty} \frac{1}{(k^2 - z^2)^p} \quad (\text{A2})$$

where the summation is over all positive integer values of  $k$ . The basic result we require is for the  $p = 1$  case, namely

$$S_1(z) = \sum_{k=1}^{\infty} \frac{1}{(k^2 - z^2)} = \frac{1}{2z^2} - \frac{\pi \cot(\pi z)}{2z} \quad (\text{A3})$$

which appears, for example, in Gradshteyn and Ryzhik.<sup>45</sup> This standard ‘handbook’ result can, in turn, be derived at a more fundamental level from a Fourier series expansion<sup>46</sup> by evaluating the Fourier components of the expansion

$$\cos(zx) = \frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos(nx) + b_n \sin(nx)] \quad (\text{A4})$$

over the interval  $(-\pi, +\pi)$ ; note that here  $z$  is considered a constant. The Fourier coefficients can be evaluated using standard integrals and we obtain

$$\cos(zx) = \frac{\sin(z\pi)}{z\pi} - \sum_{n=1}^{\infty} \left[ \frac{2z \sin(\pi z) \cos(n\pi)}{\pi(n^2 - z^2)} \right] \cos(nx) \quad (\text{A5})$$

since the  $b_n = 0$  by symmetry. If we then specialize to  $x = \pi$ , and use the fact that  $\cos^2(n\pi) = 1$ , we find

$$\pi \cot(\pi z) = \frac{1}{z} - 2z \sum_{n=1}^{\infty} \frac{1}{(n^2 - z^2)} \quad (\text{A6})$$

and we note that this partial fraction expansion of  $\cot(\pi z)$  correctly encodes the information on the divergences of the function at all integral (positive, negative, and zero) values of  $z$ . Rewriting this expression, we find that

$$S_1(z) \equiv \sum_{n=1}^{\infty} \frac{1}{(n^2 - z^2)} = \frac{1}{2z} \left( \frac{1}{z} - \pi \cot(\pi z) \right) = \frac{1}{2z^2} - \frac{\pi \cot(\pi z)}{2z} \quad (\text{A7})$$

confirming the handbook result of Eqn. (A3). Such sums are already useful in that they can be used to evaluate quantities such as the Riemann zeta function, defined by

$$\zeta(s) \equiv \sum_{n=1}^{\infty} \frac{1}{n^s} \quad \text{giving} \quad \zeta(2) = S_1(z=0) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \quad (\text{A8})$$

as the  $z \rightarrow 0$  limit of Eqn. (A7). (Such results can be directly connected to integrals which appear frequently in the evaluation of quantities related to blackbody radiation which students encounter in standard textbooks<sup>47</sup> on statistical mechanics and can therefore be reinforced through such examples.)

If we differentiate the result in Eqn. (A7) with respect to  $z$ , we find that

$$\frac{d}{dz}S_1(z) = 2z \sum_{k=1}^{\infty} \frac{1}{(k^2 - z^2)^2} = 2zS_2(z) \quad (\text{A9})$$

so that in general

$$S_{p+1}(z) = \frac{1}{2z} \frac{dS_p(z)}{dz} \quad (\text{A10})$$

thereby generating sums of arbitrarily high power. For example, this gives

$$S_2(z) = \frac{\csc^2(\pi z)[-2 + 2\pi^2 z^2 + 2 \cos(2\pi z) + \pi z \sin(2\pi z)]}{8z^4} \quad (\text{A11})$$

implying that  $\zeta(4) = S_2(z=0) = \pi^4/90$ .

Since our interest is often in summations restricted to the even or odd integers, we write

$$S_1(z) = \sum_{k=1}^{\infty} \frac{1}{(k^2 - z^2)} = \sum_{k \text{ even}}^{\infty} \frac{1}{(k^2 - z^2)} + \sum_{k \text{ odd}}^{\infty} \frac{1}{(k^2 - z^2)} \equiv S_1^{(+)}(z) + S_1^{(-)}(z). \quad (\text{A12})$$

We then note that

$$\begin{aligned} S_1^{(+)}(z) &= \sum_{k \text{ even}}^{\infty} \frac{1}{(k^2 - z^2)} = \sum_{l=1}^{\infty} \frac{1}{((2l)^2 - z^2)} \\ &= \frac{1}{4} \sum_{l=1}^{\infty} \frac{1}{(l^2 - (z/2)^2)} = \frac{1}{4} S_1\left(\frac{z}{2}\right) \\ &= \frac{1}{2z^2} - \frac{\pi \cot(\pi z/2)}{4z} \end{aligned} \quad (\text{A13})$$

which then gives

$$S_1^{(-)}(z) = S_1(z) - S_1^{(+)}(z) = \frac{\pi}{4z} \left[ \cot\left(\frac{\pi z}{2}\right) - 2 \cot(\pi z) \right] = \frac{\pi \tan(\pi z/2)}{4z} \quad (\text{A14})$$

where we use half-angle formulae in the last step. Both of the expressions in Eqns. (A13) and (A14) can, of course, be confirmed using Mathematica.

The sums over higher powers of even/odd values of  $n$  required to evaluate  $S_p^{(+)}(x)$  and  $S_p^{(-)}(x)$  in Eqn. (A1) are then obtained by repeated use of the differentiation trick in Eqn. (A10). For example, we obtain results such as

$$S_2^{(+)}(z) = \frac{\csc^2(\pi z/2)[-4 + \pi^2 z^2 + 4 \cos(\pi z) + \pi z \sin(\pi z)]}{16z^4} \quad (\text{A15})$$

and

$$S_2^{(-)}(z) = \frac{\pi \sec^2(\pi z/2)[\pi z - \sin(\pi z)]}{16z^3}. \quad (\text{A16})$$

## APPENDIX B: CONTOUR INTEGRALS

The explicit evaluation of the integrals in Eqns. (62) and (65) by contour integration techniques can be done by extending the region of integration over the entire real-axis. A contour consisting of a semi-circle of radius  $R$  can be then used as the integrands both have simple (double) poles at  $z_0^{(\pm)} = \pm q + iK_0$  in the upper-half plane. The contribution to the contour integral over the circular arc vanishes as  $R \rightarrow \infty$ , leaving

$$\int_{-\infty}^{+\infty} F(k) dk = 2\pi i \sum_i \mathcal{R}_i \quad (\text{B1})$$

where the residues are given by

$$\mathcal{R}_i = \frac{1}{(n-1)!} \left\{ \left( \frac{d}{dz} \right)^{n-1} [(z - z_0^{(i)})^n F(z)] \right\}_{z \rightarrow z_0^{(i)}} \quad (\text{B2})$$

for  $z_0^{(i)} = z_0^{(\pm)}$  and where in this case  $n = 2$ .

---

\* Electronic address: mabelloni@davidson.edu

† Electronic address: rick@phys.psu.edu

<sup>1</sup> P. Ehrenfest, “Bemerkung über die angenäherte Gültigkeit der klassischen Mechanik innerhalb der Quantenmechanik,” *Z. Phys.* **45**, 455-457 (1927).

<sup>2</sup> D. F. Styer, “The motion of wave packets through their expectation values and uncertainties,” *Am. J. Phys.* **58**, 742-744 (1990).

<sup>3</sup> J. Hirschfelder, “Classical and quantum mechanical hypervirial theorems,” *J. Chem. Phys.* **33**, 1462-1466 (1960); J. H. Epstein and S. T. Epstein, “Some applications of hypervirial theorems to the calculation of average values,” *Am. J. Phys.* **30**, 266-268 (1961).

<sup>4</sup> W. Thomas, “Über die Zahl der Dispersionselektronen, die einem stationären Zustände zugeordnet sind (Vorläufige Mitteilung),” *Naturwissenschaften* **13** 627 (1925); W. Kuhn, “Über die Gesamtstärke der von einem Zustände ausgehenden Absorptionslinien,” *Z. Phys.* **33**, 408-412;

- F. Reiche and W. Thomas, "Über die Zahl der dispersionselektronen, die einem stationären Zustand zugeordnet sind," *Z. Phys.* **34**, 510-525 (1925).
- <sup>5</sup> J. Mehra and H. Rechenberg, *The Historical Development of Quantum Theory: The Formulation of Matrix Mechanics and its Modifications 1925-1926*, Vol. 3 (Springer-Verlag, New York, 1982), Chap. IV.
- <sup>6</sup> H. Bethe, "Zur Theorie des Durchgangs schneller Korpuskularstrahlen durch Materie," *Ann. Phys. (Leipzig)* **5**, 325-400 (1930); translated as "Theory of the passage of fast corpuscular rays through matter," in *Selected Works of Hans A. Bethe with Commentary*, H. Bethe, World Scientific Series in 20th Century Physics, Vol. 18, (World Scientific, Singapore, 1997), pp. 77-154.
- <sup>7</sup> A. Guiner, G. Fournat, C. Walker, and K. Yudowitch, *Small-angle Scattering of X-rays* (Wiley, New York, 1955).
- <sup>8</sup> I. Shimamura, "Sum rules for the rotational structure in the molecular transition spectrum," *J. Phys. B. At. Mol. Phys.* **15**, 93-100 (1982).
- <sup>9</sup> L. Van Hove, "Correlations in space and time and Born approximation scattering in systems of interacting particles," *Phys. Rev.* **95**, 249-262 (1954).
- <sup>10</sup> R. A. Ferrell and R. E. Glover III, "Conductivity of superconducting films: A sum rule," *Phys. Rev.* **109**, 1398-1399 (1958).
- <sup>11</sup> M. Tinkham and R. A. Ferrell, "Determination of the superconducting skin depth from the energy gap and sum rule," *Phys. Rev. Lett.* **8**, 331-333 (1959).
- <sup>12</sup> D. N. Basov *et al.*, "Sum rules and interlayer conductivity of high- $T_c$  cuprates," *Science* **283**, 49-52 (1999).
- <sup>13</sup> R. Leonardi and M. Rosa-Clot, "Isospin sum rules in nuclear physics," *Phys. Rev. Lett.* **23**, 874-877 (1969).
- <sup>14</sup> O. Bohigas, A. M. Lane, and J. Martorell, "Sum rules for nuclear collective excitations," *Phys. Rep.* **5**, 267-316 (1979).
- <sup>15</sup> G. Orlandini and M. Traini, "Sum rules for electron-nuclear scattering," *Rep. Prog. Phys.* **54**, 257-338 (1991).
- <sup>16</sup> T. D. Cohen, R. J. Furnstahl, D. K. Griegel, and X. Jin, "QCD sum rules and applications to nuclear physics," *Prog. Part. Nucl. Phys.* **35**, 221-298 (1995).
- <sup>17</sup> V. de Alfaro, S. Fubini, C. Rossetti, and G. Furlan, "Sum rules for strong interactions," *Phys.*

- Lett. **21**, 576-579 (1966); S. L. Adler and R. F. Dashen, *Current Algebras and Applications to Particle Physics* (Benjamin, New York, 1968).
- <sup>18</sup> J. D. Bjorken, “Asymptotic sum rules at infinite momentum,” Phys. Rev. **179**, 1547-1553 (1969).
- <sup>19</sup> S. Narison, *QCD Spectral Sum Rules* (World Scientific, New York, 1989).
- <sup>20</sup> L. J. Reinders, H. Rubinstein, and S. Yazaki, “Hadron properties from QCD sum rules,” Phys. Rep. **127**, 1-97 (1985).
- <sup>21</sup> N. Isgur, M. Wise and M. Yousefmir, “Excited charm baryons in semileptonic  $\Lambda_b$  decay and their contributions to the Bjorken sum rule,” Phys. Lett. B **254**, 215-219 (1991); M. B. Voloshin, “Optical sum rule for form factors of heavy mesons,” Phys. Rev. D **46**, 3062-3592 (1992); C.-K. Chow, “Conservation laws and sum rules in the heavy quark limit,” Phys. Rev. D **51**, 3587-3592 (1995).
- <sup>22</sup> M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, “QCD and resonance physics: Sum rules,” Nucl. Phys. B **147**, 385-447 (1979).
- <sup>23</sup> Citation data from <http://www.slac.stanford.edu/spires/> where one finds, for example, that Ref. [22] has been cited over 3,200 times.
- <sup>24</sup> E. Merzbacher, *Quantum Mechanics* (Wiley, New York, 1961), pp. 445-447; D. S. Saxon, *Elementary Quantum Mechanics* (Holden-Day, San Francisco, 1968), pp. 225-226; R. Shankar, *Principles of Quantum Mechanics* (Plenum, New York, 1980), p. 466; S. Gasiorowicz, *Quantum Physics*, Second edition (Wiley, New York, 1996), p. 277.
- <sup>25</sup> E. Hadjimichael, W. Currie, and S. Fallieros, “The Thomas-Reiche-Kuhn sum rule and the rigid rotator,” Am. J. Phys. **65**, 335-341 (1997); F. M. Fernandez, “The Thomas-Reiche-Kuhn sum rule for the rigid rotator,” Int. J. Math. Ed. Sci. Tech. **33**, 636-640 (2002).
- <sup>26</sup> H. Bethe and R. Jackiw, *Intermediate Quantum Mechanics*, Second edition (Benjamin, New York, 1968), Chapter 11.
- <sup>27</sup> R. Jackiw, “Quantum mechanical sum rules,” Phys. Rev. **157**, 1220-1225 (1967).
- <sup>28</sup> S. Wang, “Generalization of the Thomas-Reiche-Kuhn and the Bethe sum rules,” Phys. Rev. A **60**, 262-266 (1999).
- <sup>29</sup> R. Bluhm, V. A. Kostelecky, and J. A. Porter, “The evolution and revival structure of localized quantum wave packets,” Am. J. Phys. **64**, 944-953 (1996); D. F. Styer, “Quantum revivals versus classical periodicity in the infinite square well,” Am. J. Phys. **69**, 56-62 (2001).
- <sup>30</sup> H. A. Mavromatis, “The Dalgarno-Lewis summation technique: Some comments and examples,”

- Am. J. Phys. **59**, 738-744 (1991).
- <sup>31</sup> H. A. Mavromatis, “New summation expressions obtained by combining perturbation theory formalisms,” *Int. J. Comp. Math.* **50**, 119-123 (1993).
- <sup>32</sup> H. A. Mavromatis, “Sum rules, as a tool for obtaining mathematical series,” *Int. J. Math. Educ. Sci. Tech.* **26**, 267-313 (1995).
- <sup>33</sup> A. Dalgarno and J. T. Lewis, “The exact calculation of long range forces between atoms by perturbation theory,” *Proc. R. Soc. London, Ser. A* **70**, 70-74 (1955).
- <sup>34</sup> R. de L. Kronig and W. G. Penney, “Quantum mechanics of electrons in crystal lattices,” *Proc. R. Soc. London, Ser. A* **130**, 499-513 (1931).
- <sup>35</sup> P. M. Morse and H. Feshbach, *Methods of Theoretical Physics* (McGraw-Hill, New York, 1953), pp. 1644-1645.
- <sup>36</sup> A. A. Frost, “Delta potential function model for electronic energies in molecules,” *J. Chem. Phys.* **22**, 1613 (1954).
- <sup>37</sup> For a long list of references on the use of  $\delta$ -function potentials in the pedagogical and research literature, see I. R. Lapidus, “Bound states and scattering from the one-dimensional hydrogen atom at a boundary,” *Am. J. Phys.* **51**, 1137-1139 (1983).
- <sup>38</sup> K. R. Brownstein, “Calculation of a bound state wavefunction using free state wavefunctions only,” *Am. J. Phys.* **43**, 173-176 (1974); W. C. Damert, “Completeness of the energy eigenstates for a delta function potential,” *Am. J. Phys.* **43**, 531-534 (1974); S. H. Patil, “Completeness of the energy eigenfunctions for the one-dimensional  $\delta$ -function potential,” *Am. J. Phys.* **68**, 712-714 (2000).
- <sup>39</sup> D. Kiang, “Contribution of the continuum in perturbation theory,” *Am. J. Phys.* **45**, 308-309 (1977).
- <sup>40</sup> V. S. Popov, V. L. Eletsii, and A. V. Turbiner, “Higher orders of perturbation theory and summation of series in quantum mechanics and field theory,” *Zh. Eksp. Teor. Fiz.* **74**, 445-465 (1978). [*Sov. Phys. JETP* **47**, 232-242 (1978).]
- <sup>41</sup> F. M. Fernandez and E. A. Castro, “Stark effect in a one-dimensional model atom,” *Am. J. Phys.* **53**, 757-760 (1985).
- <sup>42</sup> M. A. Maize and C. A. Burkholder, “Electric polarizability and the solution of an inhomogeneous differential equation,” *Am. J. Phys.* **63**, 244-247 (1995).
- <sup>43</sup> S. Borowitz, *Fundamentals of Quantum Mechanics* (W. A. Benjamin, New York, 1967), pp. 327-

330; E. Merzbacher, *Quantum Mechanics*, Second edition (Wiley, New York, 1970), pp. 420-425; A. Z. Capri, *Nonrelativistic Quantum Mechanics*, Third edition (World Scientific, Singapore, 2002), pp. 278-282.

<sup>44</sup> A. R. Ruffa, "Continuum wave functions in the calculation of sums involving off-diagonal matrix elements," *Am. J. Phys.* **41**, 234-241 (1973).

<sup>45</sup> I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Corrected and Enlarged Edition prepared by A. Jeffrey, (Academic Press, New York, 1980), p. 36.

<sup>46</sup> For example, see J. Mathews and R. L. Walker, *Mathematical Methods of Physics*, Second edition (Benjamin, Menlo Park, 1970), pp. 50-55. It is noted there that such sums can be evaluated using the Mittag-Leffler theorem of complex variable theory, and one often sees that name associated with sums of this type.

<sup>47</sup> F. Reif, *Fundamentals of Statistical and Thermal Physics* (McGraw-Hill, New York, 1965).