

# EQUIVARIANT CLASSIFICATION OF 2-TORUS MANIFOLDS

ZHI LÜ AND MIKIYA MASUDA

## 1. INTRODUCTION

In this paper, we consider the equivariant classification of locally standard 2-torus manifolds. A 2-torus manifold is a closed smooth manifold of dimension  $n$  with an effective action of a 2-torus group  $(\mathbb{Z}_2)^n$  of rank  $n$ , and it is said to be locally standard if it is locally isomorphic to a faithful representation of  $(\mathbb{Z}_2)^n$  on  $\mathbb{R}^n$ . The orbit space  $Q$  of a locally standard 2-torus  $M$  by the action is a nice manifold with corners. When  $Q$  is a simple convex polytope,  $M$  is called a small cover and studied in [4]. A typical example of a small cover is a real projective space  $\mathbb{R}P^n$  with a standard action of  $(\mathbb{Z}_2)^n$ . Its orbit space is an  $n$ -simplex. On the other hand, a typical example of a compact non-singular toric variety is a complex projective space  $\mathbb{C}P^n$  with a standard action of  $(\mathbb{C}^*)^n$  where  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ .  $\mathbb{C}P^n$  has complex conjugation and its fixed point set is  $\mathbb{R}P^n$ . More generally, any compact non-singular toric variety admits complex conjugation and its fixed point set often provides an example of a small cover. Similarly to the theory of toric varieties, an interesting connection among topology, geometry and combinatorics is discussed for small covers in [4], [5] and [7]. Although locally standard 2-torus manifolds form a much wider class than small covers, one can still expect such a connection. See [9] for the study of 2-torus manifolds from the viewpoint of cobordism.

The orbit space  $Q$  of a locally standard 2-torus manifold  $M$  contains a lot of topological information on  $M$ . For instance, when  $Q$  is a simple convex polytope (in other words, when  $M$  is a small cover), the betti numbers of  $M$  (with  $\mathbb{Z}_2$  coefficient) are described in terms of face numbers of  $Q$  ([4]). This is not the case for a general  $Q$ , but the euler characteristic of  $M$  can be described in terms of  $Q$  (Theorem 4.1). Although  $Q$  contains a lot of topological information on  $M$ ,  $Q$  is not sufficient to reproduce  $M$ , i.e., there are many locally standard 2-torus manifolds with the same orbit space in general. We need two data to reproduce  $M$  from  $Q$ . One is a characteristic function on  $Q$  introduced in [4]. It is a map from the set of codimension-one faces of  $Q$  to  $(\mathbb{Z}_2)^n$  satisfying a certain linearly independence condition. Roughly speaking, a characteristic function provides information on the set of non-free orbits in  $M$ . The other data is a principal  $(\mathbb{Z}_2)^n$ -bundle over  $Q$  which provides information on the set of free orbits in  $M$ . It turns out that the orbit space  $Q$  together with these two data uniquely determines a locally standard 2-torus manifold up to equivariant homeomorphism (Lemma 3.1).

---

**Keywords.** 2-torus manifold, equivariant classification.

**2000AMS Classification:** 57S10, 14M25, 52B70.

The first author is supported by grants from NSFC (No. 10371020 and No. 10671034) and JSPS (No. P02299).

When  $Q$  is a simple convex polytope, any principal  $(\mathbb{Z}_2)^n$ -bundle over it is trivial; so only a characteristic function matters in this case ([4]).

The set of isomorphism classes in all principal  $(\mathbb{Z}_2)^n$ -bundles over  $Q$  can be identified with  $H^1(Q; (\mathbb{Z}_2)^n)$ . Let  $\Lambda(Q)$  be the set of all characteristic functions on  $Q$ . Then each element in  $H^1(Q; (\mathbb{Z}_2)^n) \times \Lambda(Q)$  determines a locally standard 2-torus manifold with orbit space  $Q$ . However, different elements in the product may produce equivariantly homeomorphic locally standard 2-torus manifolds. Let  $\text{Aut}(Q)$  be the group of self-homeomorphisms of  $Q$  as a manifold with corners. It naturally acts on  $H^1(Q; (\mathbb{Z}_2)^n) \times \Lambda(Q)$  and one can see that equivariant homeomorphism classes in locally standard 2-torus manifolds with orbit space  $Q$  can be identified with the coset  $(H^1(Q; (\mathbb{Z}_2)^n) \times \Lambda(Q)) / \text{Aut}(Q)$ , see Proposition 5.5.

It is not easy in general to count elements in the coset above, but we can manage when  $Q$  is a compact surface with only one boundary. In this case, codimension-one faces sit in the boundary circle, so a characteristic function on  $Q$  is nothing but a coloring on a circle (with vertices) with three colors.

The paper is organized as follows. In section 2, we introduce the notion of locally standard 2-torus manifold and give several examples. Following Davis and Januszkiewicz [4], we define a characteristic function and construct a locally standard 2-torus manifold from a characteristic function and a principal bundle in section 3. In section 4 we describe the euler characteristic of a locally standard 2-torus manifold in terms of its orbit space. Section 5 discusses three equivalence relations among locally standard 2-torus manifolds and identify them with some cosets. We count the number of colorings on a circle in section 6. Applying this result, we find in section 7 the number of equivariant homeomorphism classes in locally standard 2-torus manifolds when the orbit space is a compact surface with only one boundary.

## 2. 2-TORUS MANIFOLDS

We denote the quotient additive group  $\mathbb{Z}/2\mathbb{Z}$  by  $\mathbb{Z}_2$  throughout this paper. The natural action of a 2-torus  $(\mathbb{Z}_2)^n$  of rank  $n$  on  $\mathbb{R}^n$  defined by

$$(x_1, \dots, x_n) \longmapsto ((-1)^{g_1} x_1, \dots, (-1)^{g_n} x_n), \quad (g_1, \dots, g_n) \in (\mathbb{Z}_2)^n$$

is called *the standard representation* of  $(\mathbb{Z}_2)^n$ . The orbit space is a positive cone  $\mathbb{R}_{\geq 0}^n$ . Any real  $n$ -dimensional faithful representation of  $(\mathbb{Z}_2)^n$  is obtained from the standard representation by composing a group automorphism of  $(\mathbb{Z}_2)^n$ , up to isomorphism. Therefore the orbit space of the faithful representation space can also be identified with  $\mathbb{R}_{\geq 0}^n$ .

A 2-torus manifold  $M$  is a closed smooth manifold of dimension  $n$  with an effective smooth action of  $(\mathbb{Z}_2)^n$ . We say that  $M$  is *locally standard* if for each point  $x$  in  $M$ , there is a  $(\mathbb{Z}_2)^n$ -invariant neighborhood  $V_x$  of  $x$  such that  $V_x$  is equivariantly homeomorphic to an invariant open subset of a real  $n$ -dimensional faithful representation space of  $(\mathbb{Z}_2)^n$ .

**Remark.** The notion of a torus manifold is introduced in [8]. It is a closed smooth manifold of dimension  $2n$  with an effective smooth action of a compact  $n$ -dimensional

torus  $(S^1)^n$  having a fixed point. (More precisely speaking, an orientation data on  $M$  called an omniorientation in [2] is incorporated in the definition.) There is also a notion of local standardness in this setting ([4]). Although many interesting examples of torus manifolds are locally standard (e.g. this is the case for compact non-singular toric varieties with restricted action of the compact torus, more generally for torus manifolds with vanishing odd degree cohomology, [11]), the local standardness is not assumed in the study of [8] and [10] because a combinatorial object called a multi-fan can be defined without assuming it. However, the existence of a fixed point is not assumed for a 2-torus manifold unlike a torus manifold.

For a locally standard 2-torus manifold  $M$ , the orbit space  $Q$  of  $M$  naturally becomes a manifold with corners (see [3] for the details of a manifold with corners). Therefore the notion of a face can be defined for  $Q$ . In this paper we assume that a face is connected. We call a face of dimension 0 a *vertex*, a face of dimension one an *edge* and a codimension-one face a *facet*.

An  $n$ -dimensional convex polytope  $P$  is said to be *simple* if exactly  $n$  facets meet at each of its vertices. Each point of a simple convex polytope  $P$  has a neighborhood which is affine isomorphic to an open subset of the positive cone  $\mathbb{R}_{\geq 0}^n$ , so  $P$  is an  $n$ -dimensional manifold with corners. A locally standard 2-torus manifold  $M$  is said to be a *small cover* when its orbit space is a simple convex polytope, see [4].

We call a closed, connected, codimension-one submanifold of  $M$  *characteristic* if it is a connected component of the set fixed pointwise by some  $\mathbb{Z}_2$  subgroup. Since  $M$  is compact,  $M$  has only finitely many characteristic submanifolds. The action of  $(\mathbb{Z}_2)^n$  is free outside the union of all characteristic submanifolds, in other words, a point of  $M$  with non-trivial isotropy subgroup is contained in some characteristic submanifold of  $M$ .

Through the quotient map  $M \rightarrow Q$ , a fixed point in  $M$  corresponds to a vertex of  $Q$  and a characteristic submanifold of  $M$  corresponds to a facet of  $Q$ . A connected component of the intersection of  $k$  characteristic submanifolds of  $M$  corresponds to a codimension- $k$  face of  $Q$ , so a codimension- $k$  face of  $Q$  is a connected component of the intersection of  $k$  facets. In particular, any codimension-two face of  $Q$  is a connected component of the intersection of two facets of  $Q$ , which means that  $Q$  is *nice*, see [3].

We shall give examples of locally standard 2-torus manifolds.

**Example 2.1.** A real projective space  $\mathbb{R}P^n$  with the standard  $(\mathbb{Z}_2)^n$ -action defined by

$$[x_0, x_1, \dots, x_n] \longmapsto [x_0, (-1)^{g_1}x_1, \dots, (-1)^{g_n}x_n], \quad (g_1, \dots, g_n) \in (\mathbb{Z}_2)^n$$

is a locally standard 2-torus manifold. It has  $n+1$  isolated points and  $n+1$  characteristic submanifolds. The orbit space of  $\mathbb{R}P^n$  by this action is an  $n$ -simplex, so this locally standard 2-torus manifold is actually a small cover.

**Example 2.2.** Let  $S^1$  denote the unit circle in the complex plane  $\mathbb{C}$  and consider two involutions on  $S^1 \times S^1$  defined by

$$t_1 : (z, w) \longmapsto (-z, w), \quad t_2 : (z, w) \longmapsto (z, \bar{w}).$$

Since  $t_1$  and  $t_2$  are commutative, they define a  $(\mathbb{Z}_2)^2$ -action on  $S^1 \times S^1$ , and it is easy to see that  $S^1 \times S^1$  with this action is a locally standard 2-torus manifold. It has no fixed point and the orbit space is  $\mathbb{R}P^1 \times I = S^1 \times I$  where  $I$  is a closed interval.

**Example 2.3.** If  $M_1$  and  $M_2$  are both locally standard 2-torus manifolds of the same dimension, then the equivariant connected sum of them along their free orbits produces a new locally standard 2-torus manifold. For example, we take  $\mathbb{R}P^2$  in Example 2.1 and  $S^1 \times S^1$  in Example 2.2 and do the equivariant connected sum of them along their free orbits. The orbit space of the resulting locally standard 2-torus manifold  $M$  is the connected sum of a 2-simplex with  $S^1 \times I$  at their interior points.  $M$  has five characteristic submanifolds and three of them have a fixed point but the other two have no fixed point.

If  $M$  is a locally standard 2-torus manifold of dimension  $n$  and a subgroup of  $(\mathbb{Z}_2)^n$  has an isolated fixed point, then the isolated point must be fixed by the entire group  $(\mathbb{Z}_2)^n$ . This follows from the local standardness of  $M$ . The following is an example of a closed  $n$ -manifold with an effective  $(\mathbb{Z}_2)^n$ -action which is not a locally standard 2-torus manifold.

**Example 2.4.** Consider two involutions on the unit sphere  $S^2$  of  $\mathbb{R} \times \mathbb{C}$  defined by

$$t_1 : (x, z) \mapsto (-x, -z), \quad t_2 : (x, z) \mapsto (x, \bar{z}).$$

Since  $t_1$  and  $t_2$  are commutative, they define a  $(\mathbb{Z}_2)^2$ -action on  $S^2$ . But  $S^2$  with this action is not a locally standard 2-torus manifold because the fixed point set of  $t_1 t_2$  consists of two isolated points  $(0, \pm\sqrt{-1})$  but they are not fixed by the entire group  $(\mathbb{Z}_2)^2$ .

### 3. CHARACTERISTIC FUNCTIONS AND PRINCIPAL BUNDLES

Let  $Q$  be an  $n$ -dimensional nice manifold with corners. We denote by  $\mathcal{F}(Q)$  the set of facets of  $Q$ . A codimension- $k$  face of  $Q$  is a connected component of the intersection of  $k$  facets. We call a map

$$\lambda : \mathcal{F}(Q) \longrightarrow (\mathbb{Z}_2)^n$$

a *characteristic function* on  $Q$  if it satisfies the following linearly independent condition:

if a codimension- $k$  face  $F$  of  $Q$  is a connected component of the intersection of  $k$  facets  $F_1, \dots, F_k$ , then  $\lambda(F_1), \dots, \lambda(F_k)$  are linearly independent when viewed as vectors of the vector space  $(\mathbb{Z}_2)^n$  over the field  $\mathbb{Z}_2$ .

We denote by  $G_F$  the subgroup of  $(\mathbb{Z}_2)^n$  generated by  $\lambda(F_1), \dots, \lambda(F_k)$ .

**Remark.** When  $n \leq 2$ , it is easy to see that any  $Q$  admits a characteristic function. When  $n = 3$ ,  $Q$  admits a characteristic function if the boundary of  $Q$  is a union of 2-spheres, which follows from the Four Color Theorem, but  $Q$  may not admit a characteristic function otherwise. When  $n \geq 4$ , there is a simple convex polytope which admits no characteristic function, see [4, Nonexamples 1.22].

A characteristic function arises naturally from a locally standard 2-torus manifold  $M$  of dimension  $n$  with orbit space  $Q$ . A facet of  $Q$  is the image of a characteristic

submanifold of  $M$  by the quotient map  $\pi: M \rightarrow Q$ . To each element  $F \in \mathcal{F}(Q)$  we assign the nonzero element of  $(\mathbb{Z}_2)^n$  which fixes pointwise the characteristic submanifold  $\pi^{-1}(F)$ . The local standardness of  $M$  implies that this assignment satisfies the linearly independent condition above required for a characteristic function.

Besides the characteristic function, a principal  $(\mathbb{Z}_2)^n$ -bundle over  $Q$  will be associated with  $M$  as follows. We take a small invariant open tubular neighborhood for each characteristic submanifold of  $M$  and remove their union from  $M$ . Then the  $(\mathbb{Z}_2)^n$ -action on the resulting space is free and its orbit space can naturally be identified with  $Q$ , so it gives a principal  $(\mathbb{Z}_2)^n$ -bundle over  $Q$ .

We have associated a characteristic function and a principal  $(\mathbb{Z}_2)^n$ -bundle with a locally standard 2-torus manifold. Conversely, one can reproduce the locally standard 2-torus manifold from these two data. This is done by Davis-Januszkiewicz [4] when  $Q$  is a simple convex polytope, but their construction still works in our setting. Let  $\xi = (E, \kappa, Q)$ , where  $\kappa: E \rightarrow Q$ , be a principal  $(\mathbb{Z}_2)^n$ -bundle over  $Q$  and let  $\lambda: \mathcal{F}(Q) \rightarrow (\mathbb{Z}_2)^n$  be a characteristic function on  $Q$ . We define an equivalence relation  $\sim$  on  $E$  as follows: for  $u_1, u_2 \in E$

$$u_1 \sim u_2 \iff \kappa(u_1) = \kappa(u_2) \text{ and } u_1 = u_2 g \text{ for some } g \in G_F$$

where  $F$  is the face of  $Q$  containing  $\kappa(u_1) = \kappa(u_2)$  in its relative interior and  $G_F$  is the subgroup of  $(\mathbb{Z}_2)^n$  defined at the beginning of this section. Then the quotient space  $E/\sim$ , denoted by  $M(\xi, \lambda)$ , naturally inherits the  $(\mathbb{Z}_2)^n$ -action from  $E$ .

The following is proved in [4] when  $Q$  is a simple convex polytope, but the same proof works in our setting.

**Lemma 3.1.** *If a locally standard 2-torus manifold  $M$  over  $Q$  has  $\xi$  as the associated principal  $(\mathbb{Z}_2)^n$ -principal bundle and  $\lambda$  as the characteristic function, then there is an equivariant homeomorphism from  $M(\xi, \lambda)$  to  $M$  which covers the identity on  $Q$ .*

#### 4. EULER CHARACTERISTIC OF A LOCALLY STANDARD 2-TORUS MANIFOLD

The following formula describes the euler characteristic  $\chi(M)$  of a locally standard 2-torus manifold  $M$  in terms of its orbit space.

**Theorem 4.1.** *If  $M$  is a locally standard 2-torus manifold over  $Q$ , then*

$$\chi(M) = \sum_F 2^{\dim F} \chi(F, \partial F) = \sum_F 2^{\dim F} (\chi(F) - \chi(\partial F))$$

where  $F$  runs over all faces of  $Q$ .

*Proof.* As observed in Section 3,  $M$  is the disjoint union of  $2^{\dim F}$  copies of  $F \setminus \partial F$  over all faces  $F$  of  $Q$ . This implies the former identity in the theorem. The latter identity is well-known. In fact, it follows from the homology exact sequence for a pair  $(F, \partial F)$ .  $\square$

When  $\dim M = 2$ ,  $Q$  is a surface with boundary and each boundary component is a circle with at least two vertices if it has a vertex.

**Corollary 4.2.** *If  $\dim M = 2$  and  $Q$  has  $m$  vertices, then  $\chi(M) = 4\chi(Q) - m$ .*

*Proof.* Since  $\partial Q$  is a union of circles,  $\chi(Q, \partial Q) = \chi(Q)$ . If a boundary circle has no vertex, then it is an edge without boundary and its euler characteristic is zero. So we may neglect it. If  $F$  is an edge with a vertex, then it has two endpoints and  $\chi(F, \partial F) = \chi(F) - \chi(\partial F) = -1$ , and if  $F$  is a vertex, then  $\chi(F, \partial F) = \chi(F) = 1$ . Since the number of edges with a vertex and the number of vertices are both  $m$ , it follows from Theorem 4.1 that

$$\chi(M) = 2^2\chi(Q) - 2m + m = 4\chi(Q) - m.$$

□

**Remark.** When  $\dim M = 2$ , it is not difficult to see that  $M$  is orientable if and only if  $Q$  is orientable and the characteristic function  $\lambda: \mathcal{F}(Q) \rightarrow (\mathbb{Z}_2)^2$  associated with  $M$  assigns exactly two elements to each boundary component of  $Q$  with a vertex, cf. [12]. Therefore one can find the homeomorphism type of  $M$  from the corollary above and the characteristic function  $\lambda$ .

## 5. CLASSIFICATION OF LOCALLY STANDARD 2-TORUS MANIFOLDS

In this section we introduce three notions of equivalence in locally standard 2-torus manifolds over  $Q$  and identify each set of equivalence classes with a coset of  $H^1(Q; (\mathbb{Z}_2)^n) \times \Lambda(Q)$  by some action.

Following Davis and Januszkiewicz [4] we say that two locally standard 2-torus manifolds  $M$  and  $M'$  over  $Q$  are *equivalent* if there is a homeomorphism  $f: M \rightarrow M'$  together with an element  $\sigma \in \mathrm{GL}(n, \mathbb{Z}_2)$  such that

- (1)  $f(gx) = \sigma(g)f(x)$  for all  $g \in (\mathbb{Z}_2)^n$  and  $x \in M$ , and
- (2)  $f$  induces the identity on the orbit space  $Q$ .

When we classify locally standard 2-torus manifolds up to the above equivalence, it suffices to consider locally standard 2-torus manifolds of the form  $M(\xi, \lambda)$  by Lemma 3.1. We denote by  $\xi^\sigma$  the principal  $(\mathbb{Z}_2)^n$ -bundle  $\xi$  with  $(\mathbb{Z}_2)^n$ -action through  $\sigma \in \mathrm{GL}(n, \mathbb{Z}_2)$ . Then it would be obvious that  $M(\xi', \lambda')$  is equivalent to  $M(\xi, \lambda)$  if and only if there exists  $\sigma \in \mathrm{GL}(n, \mathbb{Z}_2)$  such that  $\xi' = \xi^\sigma$  and  $\lambda' = \sigma \circ \lambda$ .

We denote by  $\mathcal{P}(Q)$  the set of all principal  $(\mathbb{Z}_2)^n$ -bundles over  $Q$ . Since the classifying space of  $(\mathbb{Z}_2)^n$  is an Eilenberg-MacLane space  $K((\mathbb{Z}_2)^n, 1)$ ,  $\mathcal{P}(Q)$  can naturally be identified with  $H^1(Q; (\mathbb{Z}_2)^n)$  and the action of  $\sigma$  sending  $\xi$  to  $\xi^\sigma$  is nothing but the action on  $H^1(Q; (\mathbb{Z}_2)^n)$  induced from the automorphism  $\sigma$  on the coefficient  $(\mathbb{Z}_2)^n$ . With this understood, the above fact implies the following.

**Proposition 5.1.** *The set of equivalence classes in locally standard 2-torus manifolds over  $Q$  bijectively corresponds to the coset*

$$\mathrm{GL}(n, \mathbb{Z}_2) \backslash (H^1(Q; (\mathbb{Z}_2)^n) \times \Lambda(Q))$$

*by the diagonal action.*

The action of  $\mathrm{GL}(n, \mathbb{Z}_2)$  on  $H^1(Q; (\mathbb{Z}_2)^n) \times \Lambda(Q)$  is free when  $Q$  has a vertex by the following lemma.

**Lemma 5.2.** *If  $Q$  has a vertex, then the action of  $\mathrm{GL}(n, \mathbb{Z}_2)$  on  $\Lambda(Q)$  is free and  $|\Lambda(Q)| = |\mathrm{GL}(n, \mathbb{Z}_2) \backslash \Lambda(Q)| \prod_{k=1}^n (2^n - 2^{k-1})$ .*

*Proof.* Suppose that  $\lambda = \sigma \circ \lambda$  for some  $\lambda \in \Lambda(Q)$  and  $\sigma \in \mathrm{GL}(n, \mathbb{Z}_2)$ . Take a vertex of  $Q$  and let  $F_1, \dots, F_n$  be the facets of  $Q$  meeting at the vertex. Then

$$(\lambda(F_1), \dots, \lambda(F_n)) = \sigma(\lambda(F_1), \dots, \lambda(F_n)).$$

Since the matrix  $(\lambda(F_1), \dots, \lambda(F_n))$  is non-singular,  $\sigma$  is the identity matrix. This proves the former statement in the lemma. Then the latter statement follows from the well-known fact that  $|\mathrm{GL}(n, \mathbb{Z}_2)| = \prod_{k=1}^n (2^n - 2^{k-1})$ , see [1].  $\square$

Lemma 5.2 is also helpful to count the number of elements in  $\Lambda(Q)$ . Here is an example.

**Example 5.3.** (The number of characteristic functions on a prism.) There exist seven combinatorially inequivalent 3-polytopes with six vertices (see [6, Theorem 6.7]) and only one of them is simple, which is a prism  $P^3$ .

Let us count the number of characteristic functions on  $P^3$ .  $P^3$  has five facets, consisting of three square facets and two triangular facets. We denote the three square facets by  $F_1, F_2, F_4$ , and the two triangular facets by  $F_3, F_5$ . The facets  $F_1, F_2, F_3$  intersect at a vertex and we may assume that a characteristic function  $\lambda$  on  $P^3$  takes the standard basis  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$  of  $(\mathbb{Z}_2)^3$  on  $F_1, F_2, F_3$  respectively through the action of  $\mathrm{GL}(3, \mathbb{Z}_2)$  on  $(\mathbb{Z}_2)^3$ . The characteristic function  $\lambda$  must satisfy the linearly independent condition at each vertex of  $P^3$ . This requires that the values of  $\lambda$  on the remaining facets  $F_4, F_5$  must be as follows:

$$(\lambda(F_4), \lambda(F_5)) = (\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3, \mathbf{e}_3) \quad \text{or} \quad (\mathbf{e}_1 + \mathbf{e}_2, a\mathbf{e}_1 + b\mathbf{e}_2 + \mathbf{e}_3)$$

where  $a, b \in \mathbb{Z}_2$ . Therefore

$$|\mathrm{GL}(3, \mathbb{Z}_2) \backslash \Lambda(P^3)| = 5 \quad \text{and} \quad |\Lambda(P^3)| = 5 |\mathrm{GL}(3, \mathbb{Z}_2)| = 840$$

by Lemma 5.2.

Another natural equivalence relation among locally standard 2-torus manifolds is equivariant homeomorphism. An *automorphism* of  $Q$  is a self-homeomorphism of  $Q$  as a manifold with corners, and we denote the group of automorphisms of  $Q$  by  $\mathrm{Aut}(Q)$ . Similarly, an *automorphism* of  $\mathcal{F}(Q)$  is a bijection from  $\mathcal{F}(Q)$  to itself which preserves the poset structure of  $\mathcal{F}(Q)$  defined by inclusions of faces, and we denote the group of automorphisms of  $\mathcal{F}(Q)$  by  $\mathrm{Aut}(\mathcal{F}(Q))$ . An automorphism of  $Q$  induces an automorphism of  $\mathcal{F}(Q)$ , so we have a natural homomorphism

$$(5.1) \quad \Phi: \mathrm{Aut}(Q) \rightarrow \mathrm{Aut}(\mathcal{F}(Q)).$$

We note that  $\mathrm{Aut}(\mathcal{F}(Q))$  acts on  $\Lambda(Q)$  by sending  $\lambda \in \Lambda(Q)$  to  $\lambda \circ h$  for  $h \in \mathrm{Aut}(\mathcal{F}(Q))$ .

**Lemma 5.4.**  *$M(\xi, \lambda)$  is equivariantly homeomorphic to  $M(\xi', \lambda')$  if and only if there is an  $h \in \mathrm{Aut}(Q)$  such that  $\lambda' = \lambda \circ \Phi(h)$  and  $h^*(\xi') = \xi$  in  $\mathcal{P}(Q)$ , where  $h^*(\xi')$  denotes the bundle induced from  $\xi'$  by  $h$ .*

*Proof.* If  $M(\xi, \lambda)$  is equivariantly homeomorphic to  $M(\xi', \lambda')$ , then there is an equivariant homeomorphism  $H: M(\xi', \lambda') \rightarrow M(\xi, \lambda)$  and it is easy to see that the automorphism of  $Q$  induced from  $H$  is the desired  $h$  in the theorem.

Conversely, suppose that there is an  $h \in \Lambda(Q)$  such that  $\lambda' = \lambda \circ \Phi(h)$  and  $\xi' = h^*(\xi)$  in  $\mathcal{P}(Q)$ . Then there is a bundle map  $\hat{h}: \xi' \rightarrow \xi$  which covers  $h$ , and  $\hat{h}$  descends to a map  $H$  from  $M(\xi', \lambda')$  to  $M(\xi, \lambda)$  because  $\lambda' = \lambda \circ \Phi(h)$ . It is not difficult to see that  $H$  is an equivariant homeomorphism.  $\square$

$\text{Aut}(Q)$  naturally acts on  $H^1(Q; (\mathbb{Z}_2)^n)$  and the canonical bijection between  $\mathcal{P}(Q)$  and  $H^1(Q; (\mathbb{Z}_2)^n)$  is equivariant with respect to the actions of  $\text{Aut}(Q)$ .

**Proposition 5.5.** *The set of equivariant homeomorphism classes in all locally standard 2-torus manifolds over  $Q$  bijectively corresponds to the coset*

$$(H^1(Q, (\mathbb{Z}_2)^n) \times \Lambda(Q)) / \text{Aut}(Q)$$

*by the diagonal action of  $\text{Aut}(Q)$ . If  $Q$  is a simple convex polytope, then the set of equivariant homeomorphism classes in all small covers over  $Q$  bijectively corresponds to the coset  $\Lambda(Q) / \text{Aut}(\mathcal{F}(Q))$ .*

*Proof.* The former statement in the proposition follows from Lemma 5.4. If  $Q$  is a simple polytope, then  $H^1(Q; (\mathbb{Z}_2)^n) = 0$ . Therefore, the latter statement in the proposition follows if we prove that the map  $\Phi$  in (5.1) is surjective when  $Q$  is a simple convex polytope.

A simple polytope  $Q$  has a simplicial polytope  $Q^*$  as its dual and the face poset  $\mathcal{F}(Q)$  is same as  $\mathcal{F}(Q^*)$  with reversed inclusion relation. Therefore  $\text{Aut}(\mathcal{F}(Q)) = \text{Aut}(\mathcal{F}(Q^*))$ . Since  $Q^*$  is simplicial, an element  $\varphi$  of  $\text{Aut}(\mathcal{F}(Q^*))$  is realized by a simplicial automorphism on the boundary of  $Q^*$ , so it extends to an automorphism of  $Q^*$ . Since  $Q$  is dual to  $Q^*$ , the automorphism of  $Q^*$  determines a bijection on the vertex set of  $Q$  and hence an automorphism of  $Q$  which induces the chosen  $\varphi$ .  $\square$

Our last equivalence relation is a combination of the previous two relations. We say that two locally standard 2-torus manifolds  $M$  and  $M'$  over  $Q$  are *weakly equivariantly homeomorphic* if there is a homeomorphism  $f: M \rightarrow M'$  together with  $\sigma \in \text{GL}(n, \mathbb{Z}_2)$  such that  $f(gx) = \sigma(g)f(x)$  for any  $g \in (\mathbb{Z}_2)^n$  and  $x \in M$ . We note that  $f$  induces an automorphism of  $Q$  but it may not be the identity on  $Q$ . The observation above shows that  $M(\xi, \lambda)$  and  $M(\xi', \lambda')$  are weakly equivariantly homeomorphic if and only if there are  $h \in \text{Aut}(Q)$  and  $\sigma \in \text{GL}(n, \mathbb{Z}_2)$  such that  $\xi' = h^*(\xi^\sigma)$  and  $\lambda' = \sigma \circ \lambda \circ h$ . It follows that

**Proposition 5.6.** *The set of weakly equivariant homeomorphism classes in locally standard 2-torus manifolds over  $Q$  bijectively corresponds to the double coset*

$$\text{GL}(n, \mathbb{Z}_2) \backslash (H^1(Q; (\mathbb{Z}_2)^n) \times \Lambda(Q)) / \text{Aut}(Q)$$

*by the diagonal actions of  $\text{Aut}(Q)$  and  $\text{GL}(n, \mathbb{Z}_2)$ . If  $Q$  is a simple convex polytope, then the set of weakly equivariant homeomorphism classes in small covers over  $Q$  bijectively corresponds to the double coset*

$$(5.2) \quad \text{GL}(n, \mathbb{Z}_2) \backslash \Lambda(Q) / \text{Aut}(\mathcal{F}(Q)).$$



**Remark.** When  $Q$  is a right-angled regular hyperbolic polytope (such  $Q$  is the dodecahedron, the 120-cell or an  $m$ -gon with  $m \geq 5$ ), it is shown in [7, Theorem 3.3] that the double coset (5.2) agrees with the set of hyperbolic structures in small covers over  $Q$ . This together with Mostow rigidity implies that when  $\dim Q \geq 3$ , that is, when  $Q$  is the dodecahedron or the 120-cell, the double coset (5.2) agrees with the set of homeomorphism classes in small covers over  $Q$  ([7, Corollary 3.4]), i.e., the natural surjective map from the double coset to the set of homeomorphism classes in small covers over  $Q$  is bijective for such  $Q$ . However, this last statement does not hold for an  $m$ -gon  $Q$  with  $m \geq 6$  although it holds for  $m = 3, 4, 5$ , see the remark following Example 6.5 in the next section.

## 6. ENUMERATION OF COLORINGS ON A CIRCLE

When  $\dim Q = 2$ , each boundary component is a circle with at least two vertices if it has a vertex, and any two non-zero elements in  $(\mathbb{Z}_2)^2$  form a basis of  $(\mathbb{Z}_2)^2$ ; so a characteristic function on  $Q$  is equivalent to coloring arcs on the boundary circles with three colors in such a way that any two adjacent arcs have different colors.

Let  $S(m)$  be a circle with  $m$  ( $\geq 2$ ) vertices. A coloring on  $S(m)$  (with three colors) means to color arcs of  $S(m)$  in such a way that any adjacent arcs have different colors. We denote by  $\Lambda(m)$  the set of all colorings on  $S(m)$  and set

$$A(m) := |\Lambda(m)|.$$

**Lemma 6.1.**  $A(m) = 2^m + (-1)^m 2$ .

*Proof.* Let  $L(m)$  be a segment with  $m+1$  vertices including the endpoints, so  $L(m)$  has  $m$  segments. The number of coloring segments of  $L(m)$  with three colors in such a way that any adjacent segments have different colors is  $3 \cdot 2^{m-1}$ . If the two end segments have different colors, then it produces a coloring on  $S(m)$  by gluing the end points of  $L(m)$ . If the two end segments have the same color, then it produces a coloring on  $S(m-1)$  by gluing the end segments of  $L(m)$ . Thus, we have that

$$(6.1) \quad A(m) + A(m-1) = 3 \cdot 2^{m-1}.$$

It follows that

$$A(m) - 2A(m-1) = -(A(m-1) - 2A(m-2)) = \dots = (-1)^{m-3}(A(3) - 2A(2))$$

and a simple observation shows that  $A(3) = A(2) = 6$ , so

$$(6.2) \quad A(m) - 2A(m-1) = (-1)^m 6.$$

The lemma then follows from (6.1) and (6.2).  $\square$

We think of  $S(m)$  as the unit circle of  $\mathbb{C}$  with  $m$  vertices  $e^{2\pi k/m}$  ( $k = 0, 1, \dots, m-1$ ). Let  $\mathfrak{D}_m$  be the dihedral group of order  $2m$  consisting of  $m$  rotations of  $\mathbb{C}$  by angles  $2\pi k/m$  ( $k = 0, 1, \dots, m-1$ ) and  $m$  reflections with respect to lines in  $\mathbb{C}$  obtained by rotating the real axis by angles  $\pi k/m$  ( $k = 0, 1, \dots, m-1$ ). Then the action of  $\mathfrak{D}_m$  on  $S(m)$  preserves the vertices so that  $\mathfrak{D}_m$  acts on the set  $\Lambda(m)$ . With this understood we have

**Theorem 6.2.** *Let  $\varphi$  denote the Euler's totient function, that is,  $\varphi(1) = 1$  and  $\varphi(N)$  for a positive integer  $N(\geq 2)$  is the number of positive integers both less than  $N$  and coprime to  $N$ . Then*

$$|\Lambda(m)/\mathfrak{D}_m| = \frac{1}{2m} \left( \sum_{2 \leq d|m} \varphi(m/d)A(d) + \frac{1 + (-1)^m}{2} \cdot 3 \cdot 2^{m/2} \cdot \frac{m}{2} \right).$$

*Proof.* The famous Burnside Lemma or Cauchy-Frobenius Lemma (see [1]) says that if  $G$  is a finite group and  $X$  is a finite  $G$ -set, then

$$|X/G| = \frac{1}{|G|} \sum_{g \in G} |X^g|$$

where  $X^g$  denotes the set of  $g$ -fixed points in  $X$ . We apply this formula to our  $\mathfrak{D}_m$ -set  $\Lambda(m)$ . Let  $a \in \mathfrak{D}_m$  be the rotation by angle  $2\pi/m$  and  $b \in \mathfrak{D}_m$  be the reflection with respect to the real axis. Then we have

$$(6.3) \quad |\Lambda(m)/\mathfrak{D}_m| = \frac{1}{2m} \sum_{k=0}^{m-1} (|\Lambda(m)^{a^k}| + |\Lambda(m)^{a^k b}|).$$

Here, if  $d$  is the greatest common divisor of  $k$  and  $m$ , then  $\Lambda(m)^{a^k} = \Lambda(m)^{a^d}$  because the subgroup generated by  $a^k$  is same as that by  $a^d$ . Since  $\Lambda(m)^{a^d} = \Lambda(d)$  and  $\Lambda(1)$  is empty, we have

$$(6.4) \quad \sum_{k=0}^{m-1} |\Lambda(m)^{a^k}| = \sum_{2 \leq d|m} \varphi(m/d)A(d).$$

On the other hand, since  $a^k b$  is a reflection with respect to the line in  $\mathbb{C}$  obtained by rotating the real axis by angle  $\pi k/m$ , we have

$$(6.5) \quad |\Lambda(m)^{a^k b}| = \begin{cases} 3 \cdot 2^{m/2} & \text{when } m \text{ is even and } k \text{ is odd,} \\ 0 & \text{otherwise.} \end{cases}$$

Putting (6.4) and (6.5) into (6.3), we obtain the formula in the theorem.  $\square$

**Example 6.3.** As is well known,  $\varphi(p^n) = p^{n-1}(p-1)$  for any prime number  $p$  and positive integer  $n$ , and  $\varphi(ab) = \varphi(a)\varphi(b)$  for relatively prime positive integers  $a$  and  $b$ . We set

$$B(m) := |\Lambda(m)/\mathfrak{D}_m|.$$

Using the formula in Theorem 6.2 together with Lemma 6.1, one finds that

$$\begin{aligned} B(2) &= 3, & B(3) &= 1, & B(4) &= 6, & B(5) &= 3, & B(6) &= 13, \\ B(7) &= 9, & B(8) &= 30, & B(9) &= 29, & B(10) &= 78, \end{aligned}$$

$$B(2^k) = 2^{2^k - k - 1} + 3 \cdot 2^{2^{k-1} - 2} + \sum_{i=1}^k 2^{2^{i-1} - i - 1}$$

$$B(p^k) = \sum_{i=1}^k \frac{1}{2p^i} (2^{p^i} - 2^{p^{i-1}})$$

$$B(2p) = \frac{1}{4p} (4^p + (3p+1)2^p + 6p - 6)$$

$$B(pq) = \frac{1}{2pq} (2^{pq} - 2^p - 2^q + 2) + \frac{1}{2p} (2^p - 2) + \frac{1}{2q} (2^q - 2)$$

where  $p$  is an odd prime number and  $q$  is another odd prime number.

**Remark.** The same argument as above works for coloring  $S(m)$  with  $s$  colors. In this case the identity in Lemma 6.1 turns into

$$A_s(m) = (s-1)^m + (-1)^m (s-1)$$

and if we denote by  $\Lambda_s(m)$  the set of all coloring on  $S(m)$  with  $s$  colors, then the formula in Theorem 6.2 turns into

$$|\Lambda_s(m)/\mathfrak{D}_m| = \frac{1}{2m} \left( \sum_{2 \leq d|m} \varphi(m/d) A_s(d) + \frac{1 + (-1)^m}{2} \cdot s \cdot (s-1)^{m/2} \cdot \frac{m}{2} \right).$$

The computation of  $|\mathrm{GL}(2, \mathbb{Z}_2) \backslash \Lambda(m)/\mathfrak{D}_m|$  can be done in a similar fashion to the above but is rather complicated. We note that the action of  $\mathrm{GL}(2, \mathbb{Z}_2)$  on  $\Lambda(m)$  is permutation of the 3 colors used to color  $S(m)$ .  $\mathrm{GL}(2, \mathbb{Z}_2)$  consists of 6 elements and three of them are of order 2 and two of them is of order 3.

**Theorem 6.4.** *Let  $\alpha$  and  $\beta$  be the functions defined as follows:*

$$\begin{aligned} \alpha(1) &= 1, & \alpha(2) &= 3, & \alpha(3) &= 2, & \alpha(6) &= 4, \\ \beta(1) &= 0, & \beta(2) &= 2, & \beta(3) &= 2, & \beta(6) &= 4. \end{aligned}$$

Then  $|\mathrm{GL}(2, \mathbb{Z}_2) \backslash \Lambda(m)/\mathfrak{D}_m|$  is given by

$$\frac{1}{2m} \left[ \sum_{d|m} \left\{ \varphi(m/d) \cdot \frac{1}{6} \left( \alpha((m/d, 6)) A(d) + \beta((m/d, 6)) A(d-1) \right) \right\} + E(m) \right]$$

where  $(m/d, 6)$  denotes the greatest common divisor of  $m/d$  and 6,  $A(q) = 2^q + (-1)^q 2$  as before, and

$$E(m) = \begin{cases} \frac{m}{6} A\left(\frac{m+1}{2}\right) & \text{if } m \text{ is odd,} \\ m \cdot 2^{m/2-1} & \text{if } m \text{ is even.} \end{cases}$$

*Proof.* Applying the Burnside Lemma to our  $\mathfrak{D}_m$ -set  $\Gamma(m) := \mathrm{GL}(2, \mathbb{Z}_2) \backslash \Lambda(m)$ , we have

$$(6.6) \quad \begin{aligned} |\mathrm{GL}(2, \mathbb{Z}_2) \backslash \Lambda(m) / \mathfrak{D}_m| &= \frac{1}{2m} \sum_{g \in \mathfrak{D}_m} |\Gamma(m)^g| \\ &= \frac{1}{2m} \sum_{k=0}^{m-1} (|\Gamma(m)^{a^k}| + |\Gamma(m)^{a^{kb}}|) = \frac{1}{2m} \left[ \sum_{d|m} \varphi(m/d) |\Gamma(m)^{a^d}| + \sum_{k=0}^{m-1} |\Gamma(m)^{a^{kb}}| \right]. \end{aligned}$$

We need to analyze  $|\Gamma(m)^{a^d}|$  with  $d|m$  and  $|\Gamma(m)^{a^{kb}}|$ .

First we shall treat  $|\Gamma(m)^{a^d}|$  with  $d|m$ . Note that  $\lambda \in \Lambda(m)$  is a representative of  $\Gamma(m)^{a^d}$  if and only if there is  $\sigma \in \mathrm{GL}(2, \mathbb{Z}_2)$  such that

$$(6.7) \quad \sigma \circ \lambda = \lambda \circ a^d.$$

Since  $a^d$  is of order  $m/d$ , the repeated use of (6.7) shows that

$$(6.8) \quad \sigma^{m/d} = 1.$$

The identity (6.7) implies that the  $\lambda$  satisfying (6.7) can be determined by the coloring restricted to the union of a consecutive  $d$  arcs, say  $T$ , and it also tells us how to recover  $\lambda$  from the coloring on  $T$ .

Let  $\mu$  be a coloring on  $T$ . We shall count colorings  $\lambda$  on  $S(m)$  which are extensions of  $\mu$  and satisfy (6.7) for some  $\sigma \in \mathrm{GL}(2, \mathbb{Z}_2)$ . To each  $\sigma$  satisfying (6.8), there is a unique extension to  $S(m)$  which satisfies (6.7). However, the extended one may not be a coloring, i.e., two arcs meeting at a junction of  $T$  and its translations by rotations  $(a^d)^r$  ( $r = 1, \dots, m/d - 1$ ) may have the same color. Let  $t$  and  $t'$  be the end arcs of  $T$  such that the rotation of  $t$  by  $a^{d-1}$  is  $t'$ . (Note: When  $d = 1$ , we understand  $t = t'$  and then the subsequent argument works.) The extended one is a coloring if and only if

$$(6.9) \quad \sigma(\mu(t)) \neq \mu(t').$$

As is easily checked, the number of  $\sigma$  satisfying conditions (6.8) and (6.9) is  $\alpha((m/d, 6))$  if  $\mu(t) \neq \mu(t')$  and is  $\beta((m/d, 6))$  if  $\mu(t) = \mu(t')$ . On the other hand, the number of  $\mu$  with  $\mu(t) \neq \mu(t')$  is  $A(d)$  and that with  $\mu(t) = \mu(t')$  is  $A(d-1)$ . It follows that the number of  $\lambda$  satisfying (6.7) for some  $\sigma$  is  $\alpha((m/d, 6))A(d) + \beta((m/d, 6))A(d-1)$ . This proves that

$$(6.10) \quad |\Gamma(m)^{a^d}| = \frac{1}{6} \left( \alpha((m/d, 6))A(d) + \beta((m/d, 6))A(d-1) \right)$$

since the action of  $\mathrm{GL}(2, \mathbb{Z}_2)$  on  $\Lambda(m)$  is free by Lemma 5.2 and the order of  $\mathrm{GL}(2, \mathbb{Z}_2)$  is 6.

Next we shall treat  $|\Gamma(m)^{a^{kb}}|$ . The argument is similar to the above. As before,  $\lambda \in \Lambda(m)$  is a representative of  $\Gamma(m)^{a^{kb}}$  if and only if there is  $\sigma \in \mathrm{GL}(2, \mathbb{Z}_2)$  such that

$$(6.11) \quad \sigma \circ \lambda = \lambda \circ a^{kb}.$$

Since  $a^{kb}$  is of order two, the repeated use of (6.11) shows that

$$(6.12) \quad \sigma^2 = 1.$$

Suppose that  $m$  is odd. Then the line fixed by  $a^{kb}$  goes through a vertex, say  $v$ , of  $S(m)$  and the midpoint of the arc, say  $e'$ , of  $S(m)$  opposite to the vertex  $v$ . Let  $H$  be

the union of  $(m+1)/2$  consecutive arcs starting from  $v$  and ending at  $e'$ . Let  $e$  be the other end arc of  $H$  different from  $e'$ . The arc  $e$  has  $v$  as a vertex. Let  $\nu$  be a coloring on  $H$  and let  $\sigma \in \text{GL}(2, \mathbb{Z}_2)$  satisfy (6.12). Then  $\nu$  has an extension to a coloring of  $S(m)$  satisfying (6.11) if and only if

$$\sigma(\nu(e)) \neq \nu(e) \quad \text{and} \quad \sigma(\nu(e')) = \nu(e').$$

It follows that  $\nu(e)$  must be different from  $\nu(e')$  and there is only one  $\sigma$  satisfying the two identities above for each such  $\nu$ . Since the number of  $\nu$  with  $\nu(e) \neq \nu(e')$  is  $A((m+1)/2)$ , so is the number of  $\lambda \in \Lambda(m)$  satisfying (6.11) for some  $\sigma$ . It follows that  $|\Gamma(m)^{a^k b}| = \frac{1}{6}A((m+1)/2)$  and hence

$$(6.13) \quad \sum_{k=0}^{m-1} |\Gamma(m)^{a^k b}| = \frac{m}{6}A((m+1)/2).$$

Suppose that  $m$  is even and  $k$  is odd. Then the line fixed by  $a^k b$  goes through the midpoints of two opposite arcs, say  $e$  and  $e'$ , of  $S(m)$ . Let  $H$  be the union of consecutive  $m/2 + 1$  arcs starting from  $e$  and ending at  $e'$ . Let  $\nu$  be a coloring on  $H$  and let  $\sigma \in \text{GL}(2, \mathbb{Z}_2)$  satisfy (6.12). Then  $\nu$  has an extension to a coloring of  $S(m)$  satisfying (6.11) if and only if

$$\sigma(\nu(e)) = \nu(e) \quad \text{and} \quad \sigma(\nu(e')) = \nu(e').$$

If  $\nu(e) \neq \nu(e')$  then such  $\sigma$  must be the identity, and if  $\nu(e) = \nu(e')$  then there are two such  $\sigma$  one of which is the identity. Since the number of  $\nu$  with  $\nu(e) \neq \nu(e')$  is  $A(m/2 + 1)$  and that with  $\nu(e) = \nu(e')$  is  $A(m/2)$ , the number of  $\lambda \in \Lambda(m)$  satisfying (6.11) for some  $\sigma$  is  $A(m/2 + 1) + 2A(m/2)$ . It follows that

$$(6.14) \quad \sum_{k=0, k:\text{odd}}^{m-1} |\Gamma(m)^{a^k b}| = \frac{m}{12}(A(m/2 + 1) + 2A(m/2)).$$

Suppose that  $m$  is even and  $k$  is even. Then the line fixed by  $a^k b$  goes through two opposite vertices, say  $v$  and  $v'$ , of  $S(m)$ . Let  $H$  be the union of consecutive  $m/2$  arcs starting from  $v$  and ending at  $v'$ . Let  $e$  and  $e'$  be the end arcs of  $H$  which respectively have  $v$  and  $v'$  as a vertex. Let  $\nu$  be a coloring on  $H$  and let  $\sigma \in \text{GL}(2, \mathbb{Z}_2)$  satisfy (6.12). Then  $\nu$  has an extension to a coloring of  $S(m)$  satisfying (6.11) if and only if

$$\sigma(\nu(e)) \neq \nu(e) \quad \text{and} \quad \sigma(\nu(e')) \neq \nu(e').$$

If  $\nu(e) \neq \nu(e')$  then there is only one such  $\sigma$ , and if  $\nu(e) = \nu(e')$  then there are two such  $\sigma$ . Since the number of  $\nu$  with  $\nu(e) \neq \nu(e')$  is  $A(m/2)$  and that with  $\nu(e) = \nu(e')$  is  $A(m/2 - 1)$ , the number of  $\lambda \in \Lambda(m)$  satisfying (6.11) for some  $\sigma$  is  $A(m/2) + 2A(m/2 - 1)$ . It follows that

$$(6.15) \quad \sum_{k=0, k:\text{even}}^{m-1} |\Gamma(m)^{a^k b}| = \frac{m}{12}(A(m/2) + 2A(m/2 - 1)).$$

Thus, when  $m$  is even, it follows from (6.14) and (6.15) that

$$(6.16) \quad \sum_{k=0}^{m-1} |\Gamma(m)^{a^{kb}}| = \frac{m}{12} (A(m/2 + 1) + 3A(m/2) + 2A(m/2 - 1)) = m \cdot 2^{m/2-1}$$

where we used  $A(q) = 2^q + (-1)^q 2$  at the latter identity.

The theorem now follows from (6.6), (6.10), (6.13) and (6.16).  $\square$

**Remark.** When  $m$  is even,  $\Lambda(m)$  contains exactly three colorings with two colors and it defines the unique element in the double coset  $\mathrm{GL}(2, \mathbb{Z}_2) \backslash \Lambda(m) / \mathfrak{D}_m$ .

**Example 6.5.** We set

$$C(m) := |\mathrm{GL}(2, \mathbb{Z}_2) \backslash \Lambda(m) / \mathfrak{D}_m|.$$

Using the formula in Theorem 6.4, one finds that

$$\begin{aligned} C(2) = 1, \quad C(3) = 1, \quad C(4) = 2, \quad C(5) = 1, \quad C(6) = 4, \quad C(7) = 3 \\ C(8) = 8, \quad C(9) = 8, \quad C(10) = 18, \quad C(11) = 21, \quad C(12) = 48. \end{aligned}$$

We conclude this section with a remark. When  $Q$  is an  $m$ -gon ( $m \geq 3$ ), a small cover over  $Q$  is a closed surface with euler characteristic  $4 - m$  and the cardinality of the set of homeomorphism classes in small covers over  $Q$  is one (resp. two) when  $m$  is odd (resp. even). On the other hand, the double coset (5.2) agrees with  $\mathrm{GL}(2, \mathbb{Z}_2) \backslash \Lambda(m) / \mathfrak{D}_m$  and we see from Theorem 6.4 that its cardinality is strictly larger than 2 when  $m \geq 6$ . So, the natural surjective map from the double coset (5.2) to the set of homeomorphism classes in small covers over  $Q$  is not injective when  $Q$  is an  $m$ -gon with  $m \geq 6$ . However, it is bijective when  $m = 3, 4, 5$ , see Example 6.5.

## 7. LOCALLY STANDARD 2-TORUS MANIFOLDS OF DIMENSION TWO

We shall enumerate the number of equivariant homeomorphism classes in locally standard 2-torus manifolds with orbit space  $Q$  when  $Q$  is a compact surface with only one boundary.

**Theorem 7.1.** *Suppose that  $Q$  is a compact surface with only one boundary component with  $m$  ( $\geq 2$ ) vertices and set*

$$h(Q) := |H^1(Q; (\mathbb{Z}_2)^2) / \mathrm{Aut}(Q)|.$$

*Then the number of equivariant homeomorphism classes in locally standard 2-torus manifolds over  $Q$  is  $h(Q)B(m)$ , where  $B(m) = |\Lambda(m) / \mathfrak{D}_m|$  is the number discussed in the previous section.*

*Proof.* By Corollary 5.5 it suffices to count the number of orbits in  $H^1(Q; (\mathbb{Z}_2)^2) \times \Lambda(Q)$  under the diagonal action of  $\mathrm{Aut}(Q)$ . Since  $Q$  has only one boundary component and  $m$  vertices,  $\Lambda(Q)$  can be identified with  $\Lambda(m)$  in Section 6 and  $\mathrm{Aut}(\mathcal{F}(Q))$  is isomorphic to the dihedral group  $\mathfrak{D}_m$ .

Let  $H$  be the normal subgroup of  $\mathrm{Aut}(Q)$  which acts on  $H^1(Q; (\mathbb{Z}_2)^2)$  trivially. We claim that the restriction of the natural homomorphism

$$(7.1) \quad \mathrm{Aut}(Q) \rightarrow \mathrm{Aut}(\mathcal{F}(Q)) \cong \mathfrak{D}_m$$

to  $H$  is still surjective. An automorphism of  $Q$  (as a manifold with corners) which rotates the boundary circle and fixes the exterior of its neighborhood is an element of  $H$ . Therefore  $H$  contains all rotations in  $\mathfrak{D}_m$ . It is not difficult to see that any closed surface admits an involution which has one-dimensional fixed point component and acts trivially on the cohomology with  $\mathbb{Z}_2$  coefficient. Since  $Q$  is obtained from a closed surface by removing an invariant open disk centered at a point in the one-dimensional fixed point set,  $Q$  admits an involution which reflects the boundary circle and lies in  $H$ . This implies the claim.

Let  $K$  be the kernel of the homomorphism  $\text{Aut}(Q) \rightarrow \text{Aut}(\mathcal{F}(Q))$ . Then

$$(7.2) \quad |(H^1(Q; (\mathbb{Z}_2)^2) \times \Lambda(Q)) / \text{Aut}(Q)| = |(H^1(Q; (\mathbb{Z}_2)^2) / K \times \Lambda(Q)) / \text{Aut}(Q)|.$$

For any element  $g$  in  $\text{Aut}(Q)$ , there is an element  $h$  in  $H$  such that  $gh$  lies in  $K$  because the map (7.1) restricted to  $H$  is surjective. Since  $H$  acts trivially on  $H^1(Q; (\mathbb{Z}_2)^2)$ , this shows that an  $\text{Aut}(Q)$ -orbit in  $H^1(Q; (\mathbb{Z}_2)^2)$  is same as an  $K$ -orbit. This means that the induced action of  $\text{Aut}(Q)$  on  $H^1(Q; (\mathbb{Z}_2)^2) / K$  is trivial. Therefore the right hand side at (7.2) reduces to

$$|H^1(Q; (\mathbb{Z}_2)^2) / \text{Aut}(Q)| |\Lambda(Q) / \text{Aut}(Q)|.$$

Here the first factor is  $h(Q)$  by definition and the second one agrees with  $|\Lambda(m) / \mathfrak{D}_m| = B(m)$  because of the surjectivity of the map (7.1), proving the theorem.  $\square$

**Example 7.2.**  $H^1(Q; (\mathbb{Z}_2)^2)$  is isomorphic to  $H^1(Q; \mathbb{Z}_2) \oplus H^1(Q; \mathbb{Z}_2)$  and the action of  $\text{Aut}(Q)$  on the direct sum is diagonal. When  $Q$  is a disk,  $h(Q) = 1$ . When  $Q$  is a real projective plane with an open disk removed,  $H^1(Q; \mathbb{Z}_2)$  is isomorphic to  $\mathbb{Z}_2$  and the action of  $\text{Aut}(Q)$  on it is trivial. Therefore,  $h(Q) = 4$  in this case. When  $Q$  is a torus with an open disk removed,  $H^1(Q; \mathbb{Z}_2)$  is isomorphic to  $(\mathbb{Z}_2)^2$ . The action of  $\text{Aut}(Q)$  on it is non-trivial and it is not difficult to see that  $h(Q) = 5$  in this case.

## REFERENCES

- [1] J. L. Alperin and R. B. Bell, *Groups and representations*, Graduate Texts in Mathematics **162** (1995), Springer-verlag.
- [2] V. M. Buchstaber and T.E. Panov, *Torus actions and their applications in topology and combinatorics*, University Lecture Series, 24. American Mathematical Society, Providence, RI, 2002.
- [3] M. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. **117** (1983), 293-324.
- [4] M. Davis and T. Januszkiewicz, *Convex polytopes, Coxeter orbifolds and torus actions*, Duke Math. J. **61** (1991), 417-451.
- [5] M. Davis, T. Januszkiewicz and R. Scott, *Nonpositive curvature of blow-ups*, Sel. math. New ser. **4** (1998), 491-547.
- [6] G. Ewald, *Combinatorial Convexity and Algebraic Geometry*, Graduate Texts in Math., Springer, 1996.
- [7] A. Garrison and R. Scott, *Small covers of the dodecahedron and the 120-cell*, Proc. Amer. Math. Soc. **131** (2002), 963-971.
- [8] A. Hattori and M. Masuda, *Theory of multi-fans*, Osaka J. Math., **40** (2003), 1-68.
- [9] Z. Lü, *2-torus manifolds, cobordism and small covers*, arXiv:math/0701928.
- [10] M. Masuda, *Unitary toric manifolds, multi-fans and equivariant index*, Tohoku Math. J. **51** (1999), 237-265.

- [11] M. Masuda and T. Panov, *On the cohomology of torus manifolds*, Osaka J. Math. **43** (2006), 711-746.
- [12] H. Nakayama and Y. Nishimura, *The orientability of small covers and coloring simple polytopes*, Osaka J. Math. **42** (2005), 243-256.

INSTITUTE OF MATHEMATICS, SCHOOL OF MATHEMATICAL SCIENCE, FUDAN UNIVERSITY, SHANGHAI, 200433, P.R. CHINA.

*E-mail address:* zlu@fudan.edu.cn

DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, SUMIYOSHI-KU, OSAKA 558-8585, JAPAN.

*E-mail address:* masuda@sci.osaka-cu.ac.jp