Optimal Controlled teleportation via several kinds of three-qubit states

Ting Gao^{1,2}, Feng-Li Yan^{2,3}, and You-Cheng Li³

¹ College of Mathematics and Information Science,

Hebei Normal University, Shijiazhuang 050016, China

² Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-Str. 1, D-85748 Garching, Germany

³ College of Physics and Information Engineering,

Hebei Normal University, Shijiazhuang 050016, China

(Dated: October 30, 2018)

The probability of successfully controlled teleportating an unknown qubit using a general threeparticle state is investigated. We give the analytic expressions of maximal probabilities of successfully controlled teleportating an unknown qubit via several kinds of tripartite states including a tripartite GHZ state and a tripartite W-state.

PACS numbers: 03.67.Hk, 89.70.+c

I. INTRODUCTION

Bennett *et al.* [1] showed that an arbitrary unknown state of a qubit could be teleported from a sender to a spatially distant receiver with the aid of longrange Einstein-Podolsky-Rosen (EPR) correlations and the transmission of two bits of classical information. Since then, quantum teleportation has been developed by many authors [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18] due to its important applications in quantum communication [19] and quantum computation. In past several years quantum teleportation has been also experimentally demonstrated by several groups [20, 22].

The controlled quantum teleportation scheme was presented by Karlsson and Bourennane [12]. In the scheme, an unknown state can be perfectly transported from one place to another place via previously shared GHZ state by means of local operations and classical communications (LOCC) under the permission of the third party. The signal state can not be transmitted unless all three sides agree to cooperate. The controlled quantum teleportation is useful in networked quantum information processing and cryptographic conferencing [24, 25, 26, 27], and controlled quantum secure direct communication [28] and has other interesting applications, such as in opening account on the agreement of managers in a network. Recently, a number of works on controlled quantum teleportation have also been proposed [13, 14, 17, 18], where they restrict themselves to the special quantum channels, such as GHZ state or W state. If a nonmaximally entangled state is taken as quantum channel, then one can not teleport a qubit with unit probability and unit fidelity. However, it is possible to teleport a qubit with a probability p < 1, which is called probabilistic quantum teleportation [15, 16]. More recently, the probabilistic scheme has been generalized to teleport N qubits [10, 11].

The entanglement property lies at the very heart of quantum information theory. The reason is that entanglement is the physical resource to perform some of the most important quantum information tasks, such as quantum teleportation, quantum computation etc. In [29], Verstraete, Popp, and Cirac introduced a new concept which they called localizable entanglement (LE). This quantity not only has a very well defined physical meaning that treats entanglement as a truly physical resource, but also establishes a very close connection between entanglement and correlation functions. The LE E_{ij} is defined as the maximum of the average entanglement between the spins *i* and *j* over all possible outcomes

$$E_{ij} = \max_{\varepsilon} \sum_{s} p_s E(|\phi_s\rangle), \qquad (1)$$

where p_s denotes the probability to obtain the two-spin state $|\phi_s\rangle$ after performing the measurement $|s\rangle$ in the rest of the system, $E(|\phi_s\rangle)$ is the chosen measure of entanglement of $|\phi_s\rangle$. The determination of the LE is a formidable task since it involves optimization over all possible local measurement strategies, and thus can not be determined in general. However, Verstaete, Popp, and Cirac gave tight upper bound and lower pound in case of $E(|\phi_s\rangle)$ being the concurrence of $|\phi_s\rangle$. We determined the exact value of this kind of LE of the general tripartite state, and obtained the analytic expression of another kind of LE, the maximal successful probability of controlled teleporting a qubit of unknown information from a sender to a remote receiver via the control of a third agent by the use of a general three-qubit state [30].

In this paper, we give the exact values of the maximal probabilities of successfully controlled teleportating an unknown qubit via many kinds of tripartite states including a tripartite GHZ state and a tripartite W-state.

The paper is outlined as follows. In Section II, we present a scheme for controlled probabilistic quantum teleportation of an arbitrary unknown qubit with a general three-qubit state. Furthermore, the successful probability of this teleportation is also obtained. In Section III, we show how to select measuring basis to reach the maximal successful probability of controlled teleporting the complete information about an arbitrary unknown state of a qubit using some kinds of three-qubit states. A brief summary is given in Section IV.

II. THE CONTROLLED QUANTUM TELEPORTATION USING A GENERAL THREE-PARTICLE STATE

Acín *et al.* [31] gave the minimal decomposition of any pure three-qubit state in terms of orthogonal product states built from local bases—a generalization of the two-quantum-bit Schmidt decomposition. They proved that for any pure three-quantum-bit state the existence of local bases which allow one to build a set of five orthogonal product states in terms of which the state can be written in a unique form. That is, for every pure state of a composite system, 123, there exist orthonormal states $|0\rangle_1$, $|1\rangle_1$ for system 1, orthonormal states $|0\rangle_2$, $|1\rangle_2$ for system 2, and orthonormal states $|0\rangle_3$, $|1\rangle_3$ for system 3 such that

$$\begin{split} |\Psi\rangle_{123} &= a_0 |000\rangle_{123} + a_1 e^{i\mu} |100\rangle_{123} + a_2 |101\rangle_{123} \\ &+ a_3 |110\rangle_{123} + a_4 |111\rangle_{123}, \\ a_i &\ge 0, \ 0 \le \mu \le \pi, \ \Sigma_{i=0}^4 a_i^2 = 1. \end{split}$$
(2)

It is uniquely characterized by the five entanglement parameters.

Suppose that Alice is to deliver an unknown state to a distant receiver Bob supervised by the controller Charlie via a quantum channel of a normalized general pure three-qubit state in (2), where particle 1 belongs to Charlie, particle 2 is in Alice's side, while Bob has particle 3. Let $a_0 \neq 0$ through out the paper. Since if $a_0 = 0$, then $|\Psi\rangle_{123}$ is a tensor product state of a pure state of particle 1 and a pure state of particles 2 and 3, but not a true tripartite entangled state. Bob can get the qubit of quantum information carried by the unknown state only if he obtains the permission of Charlie (i.e., Charlie is trustworthy and cooperative).

After getting the approval of Charlie, Alice and Bob begin their teleportation under the control of Charlie.

The controller Charlie measures his particle in the basis

$$\begin{aligned} |x\rangle &= \cos\frac{\theta}{2}|0\rangle + e^{\mathbf{i}\varphi}\sin\frac{\theta}{2}|1\rangle, \\ |x\rangle^{\perp} &= \sin\frac{\theta}{2}|0\rangle - e^{\mathbf{i}\varphi}\cos\frac{\theta}{2}|1\rangle, \end{aligned}$$
(3)

and broadcasts his measurement result. Here $\theta \in [0, \pi], \varphi \in [0, 2\pi]$.

The tripartite state $|\Psi\rangle_{123}$ can be reexpressed as

$$|\Psi\rangle_{123} = \sqrt{p_1} |x\rangle_1 |\Phi_1\rangle_{23} + \sqrt{p_2} |x\rangle_1^{\perp} |\Phi_2\rangle_{23}.$$
 (4)

Here

$$p_{1} = \sin^{2} \frac{\theta}{2} + a_{0}^{2} \cos \theta + a_{0}a_{1} \cos(\mu - \varphi) \sin \theta, \quad (5)$$

$$p_{2} = \cos^{2} \frac{\theta}{2} - a_{0}^{2} \cos \theta - a_{0}a_{1} \cos(\mu - \varphi) \sin \theta, \quad (6)$$

$$|\Phi_{1}\rangle_{23} = \frac{1}{\sqrt{p_{1}}} [(a_{0} \cos \frac{\theta}{2} + a_{1}e^{i(\mu - \varphi)} \sin \frac{\theta}{2})|00\rangle_{23}$$

$$+ a_{2}e^{-i\varphi} \sin \frac{\theta}{2}|01\rangle_{23} + a_{3}e^{-i\varphi} \sin \frac{\theta}{2}|10\rangle_{23}$$

$$+ a_{4}e^{-i\varphi} \sin \frac{\theta}{2}|11\rangle_{23}], \quad (7)$$

$$|\Phi_{2}\rangle_{23} = \frac{1}{\sqrt{p_{2}}} [(a_{0} \sin \frac{\theta}{2} - a_{1}e^{i(\mu - \varphi)} \cos \frac{\theta}{2})|00\rangle_{23}$$

$$- a_{2}e^{-i\varphi} \cos \frac{\theta}{2}|01\rangle_{23} - a_{3}e^{-i\varphi} \cos \frac{\theta}{2}|10\rangle_{23}$$

$$-a_4 e^{-i\varphi} \cos\frac{\tilde{\theta}}{2} |11\rangle_{23}]. \tag{8}$$

After Charlie's measurement, the quantum channel is collapsed to $|\Phi_1\rangle_{23}$ and $|\Phi_2\rangle_{23}$ with probability p_1 and p_2 , respectively.

By Schmidt decomposition,

$$|\Phi_1\rangle_{23} = \sqrt{\lambda_{10}}|0_2'0_3'\rangle + \sqrt{\lambda_{11}}|1_2'1_3'\rangle,$$
 (9)

$$|\Phi_2\rangle_{23} = \sqrt{\lambda_{20}}|\bar{0}_2\bar{0}_3\rangle + \sqrt{\lambda_{21}}|\bar{1}_2\bar{1}_3\rangle, \qquad (10)$$

where $\{0'_2, 1'_2\}$ and $\{\overline{0}_2, \overline{1}_2\}$ ($\{0'_3, 1'_3\}$, and $\{\overline{0}_3, \overline{1}_3\}$) are orthonormal bases of system 2 (system 3), and Schmidt coefficients

$$\lambda_{10} = \frac{1 - \sqrt{1 - C_1^2}}{2}, \ \lambda_{11} = \frac{1 + \sqrt{1 - C_1^2}}{2}, \\ \lambda_{20} = \frac{1 - \sqrt{1 - C_2^2}}{2}, \ \lambda_{21} = \frac{1 + \sqrt{1 - C_2^2}}{2}.$$
(11)

Here $C_1 = \frac{|a_0a_4e^{-i\varphi}\sin\theta+2(a_1a_4e^{i\mu}-a_2a_3)e^{-2i\varphi}\sin^2\frac{\theta}{2}|}{p_1}$ and $C_2 = \frac{|a_0a_4e^{-i\varphi}\sin\theta-2(a_1a_4e^{i\mu}-a_2a_3)e^{-2i\varphi}\cos^2\frac{\theta}{2}|}{p_2}$ are the concurrence of $|\Phi_1\rangle_{23}$ and $|\Phi_2\rangle_{23}$, respectively.

For simplicity, we write (9) and (10) as

$$|\Phi_{1}\rangle_{23} = \sqrt{\lambda_{10}}|0_{2}0_{3}\rangle + \sqrt{\lambda_{11}}|1_{2}1_{3}\rangle, \qquad (12)$$

$$|\Phi_2\rangle_{23} = \sqrt{\lambda_{20}}|0_20_3\rangle + \sqrt{\lambda_{21}}|1_21_3\rangle.$$
 (13)

Suppose that the unknown quantum state the sender Alice wants to teleport to Bob is

$$|\psi\rangle_4 = \alpha |0\rangle_4 + \beta |1\rangle_4, |\alpha|^2 + |\beta|^2 = 1.$$
 (14)

If the measurement outcome of Charlie is $|x\rangle_1$, then the collect state of particles 2, 3 and 4 is

$$\begin{aligned} &|\psi\rangle_{4}|\Phi_{1}\rangle_{23} \\ &= \sqrt{\lambda_{10}|\alpha|^{2} + \lambda_{11}|\beta|^{2}} [\frac{1}{\sqrt{2}} (|\phi^{+}\rangle_{24} \frac{\alpha\sqrt{\lambda_{10}}|0\rangle_{3} + \beta\sqrt{\lambda_{11}}|1\rangle_{3}}{\sqrt{\lambda_{10}|\alpha|^{2} + \lambda_{11}|\beta|^{2}}} \\ &+ |\phi^{-}\rangle_{24} \frac{\alpha\sqrt{\lambda_{10}}|0\rangle_{3} - \beta\sqrt{\lambda_{11}}|1\rangle_{3}}{\sqrt{\lambda_{10}|\alpha|^{2} + \lambda_{11}|\beta|^{2}}})] \\ &+ \sqrt{\lambda_{11}|\alpha|^{2} + \lambda_{10}|\beta|^{2}} [\frac{1}{\sqrt{2}} (|\psi^{+}\rangle_{24} \frac{\beta\sqrt{\lambda_{10}}|0\rangle_{3} + \alpha\sqrt{\lambda_{11}}|1\rangle_{3}}{\sqrt{\lambda_{11}|\alpha|^{2} + \lambda_{10}|\beta|^{2}}} \\ &+ |\psi^{-}\rangle_{24} \frac{\beta\sqrt{\lambda_{10}}|0\rangle_{3} - \alpha\sqrt{\lambda_{11}}|1\rangle_{3}}{\sqrt{\lambda_{11}|\alpha|^{2} + \lambda_{10}|\beta|^{2}}})], \end{aligned}$$
(15)

where $|\phi^{\pm}\rangle = \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle), |\psi^{\pm}\rangle = \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle).$ Alice makes a Bell measurement on her particles 2 and 4. She obtains $|\phi^{+}\rangle_{24}, |\phi^{-}\rangle_{24}, |\psi^{+}\rangle_{24}, \text{ and } |\psi^{-}\rangle_{24}$ with probability $\left(\frac{\sqrt{\lambda_{10}|\alpha|^{2}+\lambda_{11}|\beta|^{2}}}{\sqrt{2}}\right)^{2}, \left(\frac{\sqrt{\lambda_{10}|\alpha|^{2}+\lambda_{11}|\beta|^{2}}}{\sqrt{2}}\right)^{2}, \left(\frac{\sqrt{\lambda_{11}|\alpha|^{2}+\lambda_{10}|\beta|^{2}}}{\sqrt{2}}\right)^{2}, \text{ respectively. Then she conveys her measurement outcome to Bob over a classical communication channel.}$

In order to achieve teleportation, Bob needs to introduce an auxiliary particle b with the initial state $|0\rangle_b$ and performs a collective unitary transformation

$$U_{3b} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \frac{\sqrt{\lambda_{10}}}{\sqrt{\lambda_{11}}} & \sqrt{1 - \frac{\lambda_{10}}{\lambda_{11}}} \\ 0 & 0 & -\sqrt{1 - \frac{\lambda_{10}}{\lambda_{11}}} & \frac{\sqrt{\lambda_{10}}}{\sqrt{\lambda_{11}}} \end{pmatrix} \text{ on the state}$$

of particles 3 and b. Then the measurement on his auxiliary particle b follows. If his measurement result is $|0\rangle_b$, Bob can fix up the state of his particle 3, recovering $|\psi\rangle$, by applying an appropriate local unitary operation. The achievable successful probability of teleporting the unknown state in (14)

via
$$|\Phi_1\rangle_{23}$$
 is $2\left(\frac{\sqrt{\lambda_{10}|\alpha|^2 + \lambda_{11}|\beta|^2}}{\sqrt{2}}\right)^2 \left(\frac{\sqrt{\lambda_{10}}}{\sqrt{\lambda_{10}|\alpha|^2 + \lambda_{11}|\beta|^2}}\right)^2 + 2\left(\frac{\sqrt{\lambda_{11}|\alpha|^2 + \lambda_{10}|\beta|^2}}{\sqrt{2}}\right)^2 \left(\frac{\sqrt{\lambda_{10}}}{\sqrt{\lambda_{11}|\alpha|^2 + \lambda_{10}|\beta|^2}}\right)^2 = 2\lambda_{10}$. Similarly, if the measurement result of Charlie is $|x\rangle_{\perp}^+$, the

harly, if the measurement result of Charlie is $|x\rangle_1^+$, the achievable successful probability of teleporting the state in (14) via $|\Phi_2\rangle_{23}$ is $2\lambda_{20}$.

Therefore, probability p of successfully controlled teleporting an unknown qubit (14) using a general threeparticle state in (2) is

$$p = 2p_1\lambda_{10} + 2p_2\lambda_{20}$$

= $1 - \left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\right).$ (16)

Here

$$P(\theta,\varphi) = p_1^2(1-C_1^2)$$

$$= \frac{1}{4}a_0^2a_1^2\cos 2(\varphi-\mu) + 3a_1a_2a_3a_4\cos\mu + \frac{1}{8}(3-4a_0^2) + 4a_0^4 + 2a_0^2a_1^2 - 12a_2^2a_3^2 - 4a_0^2a_4^2 - 12a_1^2a_4^2) + \frac{1}{8}\cos 2\theta(1-4a_0^2+4a_0^4-2a_0^2a_1^2-2a_0^2a_1^2\cos 2(\varphi-\mu)) - 4a_2^2a_3^2 + 8a_1a_2a_3a_4\cos\mu + 4a_0^2a_4^2 - 4a_1^2a_4^2) - \cos\theta(\frac{1}{2}-a_0^2-2a_2^2a_3^2 + 4a_1a_2a_3a_4\cos\mu - 2a_1^2a_4^2) + a_0\sin\theta[2a_2a_3a_4\cos\varphi + a_1(1-2a_4^2)\cos(\varphi-\mu)] - \frac{1}{2}a_0\sin 2\theta[2a_2a_3a_4\cos\varphi + a_1(1-2a_0^2-2a_4^2)\cos(\varphi-\mu)],$$
(17)

$$Q(\theta,\varphi) = p_1^2(1-C_2^2)$$

$$= \frac{1}{4}a_0^2a_1^2\cos 2(\varphi-\mu) + 3a_1a_2a_3a_4\cos\mu + \frac{1}{8}(3-4a_0^2) + 4a_0^4 + 2a_0^2a_1^2 - 12a_2^2a_3^2 - 4a_0^2a_4^2 - 12a_1^2a_4^2) + \frac{1}{8}\cos 2\theta(1-4a_0^2+4a_0^4-2a_0^2a_1^2-2a_0^2a_1^2\cos 2(\varphi-\mu)) - 4a_2^2a_3^2 + 8a_1a_2a_3a_4\cos\mu + 4a_0^2a_4^2 - 4a_1^2a_4^2) + \cos \theta(\frac{1}{2}-a_0^2-2a_2^2a_3^2 + 4a_1a_2a_3a_4\cos\mu - 2a_1^2a_4^2) - a_0\sin \theta[2a_2a_3a_4\cos\varphi + a_1(1-2a_4^2)\cos(\varphi-\mu)] - \frac{1}{2}a_0\sin 2\theta(2a_2a_3a_4\cos\varphi + a_1(1-2a_0^2-2a_4^2)\cos(\varphi-\mu)))$$
(18)

Obviously,

$$Q(\theta,\varphi) = P(\pi - \theta,\varphi + \pi) \text{ if } \varphi \in [0,\pi], Q(\theta,\varphi) = P(\pi - \theta,\varphi - \pi) \text{ if } \varphi \in [\pi,2\pi],$$
(19)

$$P(\theta,\varphi) = Q(\pi - \theta,\varphi + \pi) \text{ if } \varphi \in [0,\pi], P(\theta,\varphi) = Q(\pi - \theta,\varphi - \pi) \text{ if } \varphi \in [\pi,2\pi],$$
(20)

and

$$\frac{\sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}}{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}} = \frac{\sqrt{P(\pi,\varphi)} + \sqrt{Q(\pi,\varphi)}}{\sqrt{P(\theta,2\pi)} + \sqrt{Q(\theta,2\pi)}}.$$
(21)

It is clear that the successful probability is the same as the above (16) if the operation order is changed. That is, if Alice makes a Bell state measurement on her particles 2 and 4 first, the third party Charlie's measurement on his particle 1 follows, after that Bob operates his particle to acquire a qubit of quantum information Alice sends, instead of the above operation order, then they can also achieve the successful probability (16) of controlled teleportation.

III. THE MAXIMAL SUCCESSFUL PROBABILITY OF CONTROLLED QUANTUM TELEPORTATION USING THREE-PARTICLE STATES WITH $a_1a_2a_3a_4 \sin \mu = 0$

In [30], we determined the analytic expression of the maximal successful probability of controlled teleportation by using the general tripartite state (2) with $a_0a_1a_2a_3a_4 \sin \mu \neq 0$. In this section, we give the analytic expression of the localizable entanglement (LE), the maximum of probability of successfully controlled teleporting an unknown qubit state (14) via every three-qubit state (2) satisfying $a_1a_2a_3a_4 \sin \mu = 0$ and investigate how to achieve it (that is, Charlie finds optimal measurement basis).

Obviously, the maximum of (16) is

$$p_{\max} = \max\{p\}$$

$$= \max\{2p_1\lambda_{10} + 2p_2\lambda_{20}\}$$

$$= 1 - \min\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\}.$$
(22)

In order to get the maximum p_{max} of p in (22), we need only obtain the minimum of $\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)}$

$$\min\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\}.$$
 (23)

In other words, to reach the maximal probability of exact controlled teleportation through an arbitrary partially entangled quantum channel (2), the supervisor Charlie needs only to choose optimal measurement basis, i.e. he selects θ_0 and φ_0 such that $\min\{\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)}\} = \sqrt{P(\theta_0, \varphi_0)} + \sqrt{Q(\theta_0, \varphi_0)}$.

Note that the minimum of $\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)}$ should occur at the points such that $P(\theta, \varphi) = 0$, $Q(\theta, \varphi) = 0$,

$$\frac{\partial \left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\right)}{\partial \theta} = 0, \qquad (24)$$

$$\frac{\partial \left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\right)}{\partial \varphi} = 0, \qquad (25)$$

or, the boundary of θ and φ . From Eq.(24) and Eq.(25), there are

$$\frac{\partial P(\theta,\varphi)}{\partial \theta} \frac{\partial Q(\theta,\varphi)}{\partial \varphi} - \frac{\partial Q(\theta,\varphi)}{\partial \theta} \frac{\partial P(\theta,\varphi)}{\partial \varphi} = 0, (26)$$

$$P(\theta,\varphi) \left(\frac{\partial Q(\theta,\varphi)}{\partial\varphi}\right)^2 - Q(\theta,\varphi) \left(\frac{\partial P(\theta,\varphi)}{\partial\varphi}\right)^2 = 0, (27)$$

$$P(\theta,\varphi) \left(\frac{\partial Q(\theta,\varphi)}{\partial \theta}\right)^2 - Q(\theta,\varphi) \left(\frac{\partial P(\theta,\varphi)}{\partial \theta}\right)^2 = 0. (28)$$

Let $y = \cot \theta$, $t = \cot \frac{\theta}{2}$, $\theta \in (0, \pi)$, then $\sin \theta = \frac{1}{\sqrt{y^2+1}} = \frac{2t}{1+t^2}, \cos \theta = \frac{y}{\sqrt{1+y^2}} = \frac{t^2-1}{t^2+1}, t = y + \sqrt{y^2+1}, t \in (0, +\infty)$. These will be useful throughout the paper. Next we give controller's optimal measurement basis (3) for every kind of given quantum channel (2) satisfying $a_1a_2a_3a_4\sin\mu = 0$. That is, we determine the two parameters θ and φ in measurement basis (3). To state clearly, we classify the quantum channel (2) into the following cases. Next we examine quantum channels (2) with the following different characterizations, give the maximal probability of exact teleportation via any three-qubit state (2) with $a_1a_2a_3a_4\sin\mu = 0$, and characterizes the tripartite states that can collapse to an EPR pair after Charlie's measurement.

A.
$$a_1 = a_2 = a_3 = 0$$
, and $a_0 a_4 \neq 0$

The quantum channel (2) with three coefficients being 0 is the only one satisfying $a_1 = a_2 = a_3 = 0$ and $a_0a_4 \neq 0$, since others are biseparable (one party is not entangled with the other two parties) and can not be used as quantum channel of controlled teleportation.

It can be seen that

$$\min\left\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\right\} = \min\left\{\sqrt{\frac{1}{4}[\cos\theta - (1 - 2a_0^2)]^2} + \sqrt{\frac{1}{4}[\cos\theta + (1 - 2a_0^2)]^2}\right\} = |1 - 2a_0^2|,$$
(29)

for each $\varphi \in [0, 2\pi]$, and all θ satisfying $|\cos \theta| \le |1-2a_0^2|$. It implies that

$$p_{\max} = 1 - |1 - 2a_0^2|. \tag{30}$$

Note that if $\cos \theta = 1 - 2a_0^2 \neq 0$ or $\cos \theta = -1 + 2a_0^2 \neq 0$, then $P(\theta, \varphi) = 0$ or $Q(\theta, \varphi) = 0$, which means that after Charlie's measurement the state of Alice's particles 2 and 3 can be an EPR pair with probability $2a_0^2(1-a_0^2)$. Moreover, if $\cos \theta = 1 - 2a_0^2 = 0$ or $\cos \theta = -1 + 2a_0^2 = 0$, i.e. $\theta = \frac{\pi}{2}$, $a_0 = a_4 = \frac{1}{\sqrt{2}}$, then $P(\theta, \varphi) = Q(\theta, \varphi) = 0$, that is, after Charlie measures quantum channel in basis $\{\frac{|0\rangle + e^{i\varphi}|1\rangle}{\sqrt{2}}, \frac{|0\rangle - e^{i\varphi}|1\rangle}{\sqrt{2}}\}$, particles 2 and 3 are collapsed to a Bell state with probability 1. It follows that perfect quantum teleportation can be achieved if $a_0 = \frac{1}{\sqrt{2}}$ (i.e quantum channel is in GHZ state) and $\theta = \frac{\pi}{2}$. That is, one can send perfect unknown state to another using GHZ state as a quantum channel via controller's measurement in the basis $\{\frac{|0\rangle + e^{i\varphi}|1\rangle}{\sqrt{2}}, \frac{|0\rangle - e^{i\varphi}|1\rangle}{\sqrt{2}}\}$, where $\varphi \in [0, 2\pi]$.

B. $a_1 = a_4 = 0$, and $a_0 a_2 a_3 \neq 0$

Here we consider the quantum channel with coefficients satisfying $a_1 = a_4 = 0$ and $a_0 a_2 a_3 \neq 0$. These states are called tri-Bell states in [31]. Note that W-state

$$|W\rangle = \frac{1}{\sqrt{3}}(|001\rangle + |010\rangle + |100\rangle)$$
 (31)

is contained here, since the quantum channel (2) in case of $a_1 = a_4 = 0$ and $a_0 = a_2 = a_3 = \frac{1}{\sqrt{3}}$ is LOCC equivalent to W-state.

If $P(\theta, \varphi) = 0$, then $\cos \theta = \frac{1+2a_2a_3}{1-2a_0^2+2a_2a_3}$, or $\cos \theta = \frac{-1+2a_2a_3}{-1+2a_0^2+2a_2a_3}$. It is not difficult to prove that $|\frac{1+2a_2a_3}{1-2a_0^2+2a_2a_3}| > 1$, $|\frac{-1+2a_2a_3}{-1+2a_0^2+2a_2a_3}| > 1$ if $a_2 \neq a_3$, and $\frac{-1+2a_2a_3}{-1+2a_0^2+2a_2a_3} = -1$ if $a_2 = a_3$. Thus, $P(\theta, \varphi) = 0$ if and only if $a_2 = a_3$, and $\theta = \pi$. From (19), there is $Q(\theta, \varphi) = 0$ if and only if $a_2 = a_3$, and $\theta = 0$. Combining (19) and (20), there is

$$= \begin{pmatrix} \sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} \\ \sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} \end{pmatrix} \Big|_{P(\theta,\varphi)=0}$$
(32)
$$= a_0^2.$$

Therefore, for the quantum channel (2) such that $a_1 = a_4 = 0$, $a_2 = a_3$ and $a_0 a_2 a_3 \neq 0$, can be collapsed to a Bell state with probability $p_1 = p_2 = 1 - a_0^2$ by Charlie measuring his particle in the basis $\{|0\rangle, |1\rangle\}$.

Next we suppose that $P(\theta, \varphi) \neq 0$ and $Q(\theta, \varphi) \neq 0$. From (28), we derive that $a_0^4 a_2^2 a_3^2 \left(-1+2a_0^2+4a_2^2 a_3^2\right) \cos \theta \sin^2 \theta = 0$. Thus, $\theta = 0, \frac{\pi}{2}, \pi$. By checking, $\theta = 0, \frac{\pi}{2}, \pi$ are roots of (24). Note that if $\theta = 0, \pi$, then $a_2 \neq a_3$ by $P(\theta, \varphi) \neq 0$ and $Q(\theta, \varphi) \neq 0$. Obviously, $\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)} = a_0^2 + |a_2^2 - a_3^2| > 0$ if $\theta = 0, \pi$ and $a_2 \neq a_3$; $\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)} = \sqrt{1 - 4a_2^2 a_3^2} > 0$ if $\theta = \frac{\pi}{2}$. Therefore,

$$\min\left\{ \sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} \right\} = \min\left\{ a_0^2 + |a_2^2 - a_3^2|, \sqrt{1 - 4a_2^2 a_3^2} \right\} > 0.$$
(33)

That is, for quantum channel (2) with $a_1 = a_4 = 0$, $a_0a_2a_3 \neq 0$ and $a_0^2 + |a_2^2 - a_3^2| < \sqrt{1 - 4a_2^2a_3^2}$, the controller should choose measurement basis $\{|0\rangle, |1\rangle\}$; otherwise, he selects measurement basis (3) with $\theta = \frac{\pi}{2}$ (i.e. he measures in the basis $\{\frac{|0\rangle+e^{i\varphi}|1\rangle}{\sqrt{2}}, \frac{|0\rangle-e^{i\varphi}|1\rangle}{\sqrt{2}}\}$, where $\varphi \in [0, 2\pi]$), thus, the controlled teleportation in Section II can achieve maximal successful probability $p_{\text{max}} < 1$.

C. one is $a_1 = a_2 = 0$ and $a_0 a_3 a_4 \neq 0$, the other is $a_1 = a_3 = 0$ and $a_0 a_2 a_4 \neq 0$

Now we discuss the quantum channel (2) with the characterization $a_1 = a_2 = 0$ and $a_0 a_3 a_4 \neq 0$.

First, $P(\theta, \varphi) \neq 0$ and $Q(\theta, \varphi) \neq 0$. If $P(\theta, \varphi) = 0$, then $\cos \theta = \frac{1-2a_0^2-4a_0^2a_4\sqrt{-a_3^2}}{1-4a_0^2+4a_0^2+4a_0^2a_4^2}$, or $\cos \theta = \frac{1-2a_0^2+4a$

$$\min\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\} = \sqrt{\frac{P(\frac{\pi}{2},\varphi)}{1 - 4a_0^2a_4^2}} + \sqrt{\frac{Q(\frac{\pi}{2},\varphi)}{1 - 4a_0^2a_4^2}} = \sqrt{1 - 4a_0^2a_4^2} > 0.$$
(34)

$$p_{\max} = 1 - \sqrt{1 - 4a_0^2 a_4^2} < 1, \tag{35}$$

where $\theta = \frac{\pi}{2}$. Similarly, when $a_1 = a_3 = 0$, and $a_0 a_2 a_4 \neq 0$, we have

$$p_{\max} = 1 - \sqrt{1 - 4a_0^2 a_4^2} < 1, \tag{36}$$

where $\theta = \frac{\pi}{2}$. That is, the controlled teleportation via quantum channel (2) satisfying $a_1 = a_2 = 0$ and $a_0a_3a_4 \neq 0$, or $a_1 = a_3 = 0$ and $a_0a_2a_4 \neq 0$ can only succeed with optimal probability $p_{\text{max}} = 1 - \sqrt{1 - 4a_0^2a_4^2} < 1$.

D. one is $a_2 = a_4 = 0$ and $a_0a_1a_3 \neq 0$, the other is $a_3 = a_4 = 0$ and $a_0a_1a_2 \neq 0$

For the quantum channel with characterization either $a_2 = a_4 = 0$ and $a_0a_1a_3 \neq 0$, or $a_3 = a_4 = 0$ and $a_0a_1a_2 \neq 0$, since $P(\theta, \varphi) = p_1^2$, $Q(\theta, \varphi) = p_2^2$, so $\min\{\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)}\} = 1$, and $p_{\max} = 0$ for both cases. It can also be seen directly from the quantum channel (2) with these two characterizations being biseparable states.

E. $a_2 = a_3 = 0$ and $a_0 a_1 a_4 \neq 0$

The quantum channel (2) with coefficients satisfying $a_2 = a_3 = 0$, but $a_0 a_1 a_4 \neq 0$, are extended GHZ states according to the classification in [31].

For this kind of quantum channel, we can derive that

$$P(\theta,\varphi) = \frac{1}{4} [1 - 2a_4^2 + 2a_0a_1\cos(\varphi - \mu)\sin\theta - (1 - 2a_0^2 - 2a_4^2)\cos\theta]^2, \quad (37)$$
$$Q(\theta,\varphi) = \frac{1}{4} [-1 + 2a_4^2 + 2a_0a_1\cos(\varphi - \mu)\sin\theta]^2$$

$$-(1 - 2a_0^2 - 2a_4^2)\cos\theta]^2, \qquad (38)$$

and

$$p_{\max} = 1 - \min\{\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)}\} = 1 - |1 - 2a_4^2|$$
(39)
$$(1 - 2a_4^2) = 1 - |1 - 2a_4^2|$$
(39)

where θ, φ satisfy $|2a_0a_1\cos(\varphi - \mu)\sin\theta - (1 - 2a_0^2 - 2a_4^2)\cos\theta| \le |1 - 2a_4^2|$. Note that the set S of (θ, φ) such that $|2a_0a_1\cos(\varphi - \mu)\sin\theta - (1 - 2a_0^2 - 2a_4^2)\cos\theta| \le |1 - 2a_4^2|$ is a region, as we can see easily that $(\frac{\pi}{2}, \frac{\pi}{2} + \mu)$, $(\frac{\pi}{2}, \frac{3\pi}{2} + \mu) \in S$ in case of $0 \le \mu \le \frac{\pi}{2}$, and $(\frac{\pi}{2}, \frac{\pi}{2} + \mu)$, $(\frac{\pi}{2}, -\frac{\pi}{2} + \mu) \in S$ in case of $\frac{\pi}{2} \le \mu \le \pi$. From (37) and (38), we have that $P(\theta, \varphi) = Q(\theta, \varphi) = Q(\theta, \varphi)$

From (37) and (38), we have that $P(\theta, \varphi) = Q(\theta, \varphi) = 0$ if and only if $a_4 = \frac{1}{\sqrt{2}}$ and $a_1 \cos(\varphi - \mu) \sin \theta + a_0 \cos \theta = 0$, which means that as long as Charlie measures the quantum channel

$$a_0|000\rangle + a_1 e^{i\mu}|100\rangle + a_4|111\rangle, \ a_0^2 + a_1^2 = a_4^2 = \frac{1}{2}, \ (40)$$

in the basis (3) satisfying $a_1 \cos(\varphi - \mu) \sin \theta + a_0 \cos \theta = 0$, where $\theta \in [0, \pi]$, $\varphi \in [0, 2\pi]$, Alice and Bob can obtain an EPR pair with certainty (i.e. with probability p = 1), and Alice teleports her one qubit information to Bob with probability 1 and with unit fidelity. Note that the state

$$a_0|000\rangle + a_1|100\rangle + a_4|111\rangle, \ a_0^2 + a_1^2 = a_4^2 = \frac{1}{2},$$
 (41)

is LOCC equivalent to the state in (40). That is, the states in (41) or (40) can be used for perfect controlled quantum teleportation. Clearly, one can achieve perfect teleportation via these states by the controller making a measurement on his particle using the basis (3) satisfying $a_1 \cos(\varphi - \mu) \sin\theta + a_0 \cos\theta = 0$. More important, this kind of states in (41) or (40) are different from GHZ state according to the classification in [31].

F. $a_1 = 0$ and $a_0 a_2 a_3 a_4 \neq 0$

In this section, we investigate the quantum channel with coefficients having $a_1 = 0$ and $a_0 a_2 a_3 a_4 \neq 0$. Evidently,

$$\frac{\sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}}{\sqrt{P(\pi,\varphi)} + \sqrt{Q(\pi,\varphi)}} = a_0^2 + \sqrt{(1-a_0^2)^2 - 4a_2^2a_3^2} = a_0^2 + \sqrt{[a_4^2 + (a_2 - a_3)^2][a_4^2 + (a_2 + a_3)^2]},$$
(42)

and $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$ in case of $\theta = 0, \pi$.

We first examine the condition of $P(\theta, \varphi) = 0$. If $P(\theta, \varphi) = 0$, then

$$\cos\varphi = \left[2\sin\theta(1-\cos\theta)a_0a_2a_3a_4\right]^{-1}\left[-\frac{1}{4}+a_2^2a_3^2+a_0^2a_4^2+\cos\theta(\frac{1}{2}-a_0^2-2a_2^2a_3^2)-\frac{1}{4}\cos^2\theta(1-4a_0^2+4a_0^4-4a_2^2a_3^2+4a_0^2a_4^2)\right].$$
(43)

Note that

$$z = [2\sin\theta(1-\cos\theta)a_0a_2a_3a_4]^{-1}[-\frac{1}{4}+a_2^2a_3^2 + a_0^2a_4^2 + \cos\theta(\frac{1}{2}-a_0^2-2a_2^2a_3^2) - \frac{1}{4}\cos^2\theta(1-4a_0^2+4a_0^4-4a_2^2a_3^2+4a_0^2a_4^2)] \quad (44)$$
$$= \frac{-a_0^4t^4 + (-2a_0^2+2a_0^4+4a_0^2a_4^2)t^2 - (1-a_0^2)^2 + 4a_2^2a_3^2}{8ta_0a_2a_3a_4}$$

tends to $-\infty$ when t tends to 0 and $+\infty$, and z = -1holds only if $a_2 = a_3$ and $t = \frac{a_4}{a_0}$. It follows that $z \le -1$, where the equality occurs if and only if $a_2 = a_3$ and $\theta = \theta_0$, where $\cot \frac{\theta_0}{2} = \frac{a_4}{a_0}$ and $0 < \theta_0 < \pi$. Therefore, only if $a_2 = a_3$, $\varphi = \pi$, $\theta = \theta_0$, there is $P(\theta, \varphi) = 0$.

By (19) and (20), we have $Q(\theta, \varphi) = 0$ iff $a_2 = a_3$, $\varphi = 0$, and $\theta = \pi - \theta_0$, and

$$\begin{aligned} \left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\right)\Big|_{P(\theta,\varphi)=0} \\ &= \sqrt{Q(\theta_0,\pi)}\Big|_{a_2=a_3} \\ &= \left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\right)\Big|_{Q(\theta,\varphi)=0} \\ &= \sqrt{P(\pi-\theta_0,0)}\Big|_{a_2=a_3} \\ &= \frac{\sqrt{a_0^8 + 2a_0^4a_4^2 - 2a_0^6a_4^2 + a_4^4 - 2a_0^2a_4^4 - 3a_0^4a_4^4 + 8a_0^2a_3^2a_4^4 - 4a_3^4a_4^4}{a_0^2 + a_4^2} \\ &= \sqrt{1 - 4a_3^2 + 4a_3^4 - 4a_4^2 + 12a_3^2a_4^2 + 4a_4^4}. \end{aligned}$$
(45)

That is, $\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} = \sqrt{1 - 4a_3^2 + 4a_4^3 - 4a_4^2 + 12a_3^2a_4^2 + 4a_4^4}$ if $P(\theta,\varphi) = 0$ or $Q(\theta,\varphi) = 0$.

It is shown that an EPR pair can be obtained with probability $p_1 = p_2 = \frac{a_0^2(1-a_0^2+a_4^2)}{a_0^2+a_4^2}$ if $a_2 = a_3$ and Charlie using measurement basis (3) with $(\theta, \varphi) = (\theta_0, \pi)$, or $(\theta, \varphi) = (\pi - \theta_0, 0)$, where $\cot \frac{\theta_0}{2} = \frac{a_4}{a_0}$ and $0 < \theta_0 < \pi$.

Next we assume that $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$. From (26) and (27), there are

 $\sin\varphi[2a_2a_3a_4\cos\varphi\sin\theta - 2a_0^3\cos\theta + a_0\cos\theta(1 - 2a_4^2)] = 0$

and

$$\frac{\sin\varphi[a_0(1-2a_4^2)\cos\theta\sin^2\theta}{+2a_0^3\cos^3\theta+2a_2a_3a_4\cos\varphi\sin^3\theta]} = 0$$

respectively. By $2a_2a_3a_4\cos\varphi\sin\theta - 2a_0^3\cos\theta + a_0\cos\theta(1-2a_4^2) = 0$, and $a_0(1-2a_4^2)\cos\theta\sin^2\theta + 2a_0^3\cos^3\theta + 2a_2a_3a_4\cos\varphi\sin^3\theta = 0$, there are $\theta = \frac{\pi}{2}, \varphi = \frac{\pi}{2}, \frac{3\pi}{2}$. Evidently,

$$\sqrt{P(\frac{\pi}{2}, \frac{\pi}{2})} + \sqrt{Q(\frac{\pi}{2}, \frac{\pi}{2})} = \sqrt{P(\frac{\pi}{2}, \frac{3\pi}{2})} + \sqrt{Q(\frac{\pi}{2}, \frac{3\pi}{2})} = \sqrt{1 - 4a_2^2a_3^2 - 4a_0^2a_4^2}.$$
(46)

Next we need only to examine the case $\sin \varphi = 0$, i.e. $\varphi = 0, \pi, 2\pi$, while $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$ and $\theta \in (0, \pi)$.

Now let us consider the case $\varphi = 0$. From (28), we obtain

$$-4a_{0}^{3}(2ya_{0}a_{2}a_{3}+a_{4})[4y^{2}a_{0}^{2}a_{2}a_{3}a_{4}^{2}+2ya_{0}(2a_{0}^{4}+2a_{0}^{2}a_{4}^{2}-3a_{0}^{2}-4a_{2}^{2}a_{3}^{2}-a_{4}^{2}+1)a_{4} +a_{2}a_{3}(2a_{0}^{2}-4a_{0}^{2}a_{4}^{2}+4a_{2}^{2}a_{3}^{2}+2a_{4}^{2}-1)]$$

$$=0,$$
(47)

which implies that

=

$$y = \cot \theta_1 = -\frac{a_4}{2a_0 a_2 a_3},\tag{48}$$

$$y = \cot \theta_2 = \begin{cases} \frac{a_3(1-2a_3^2-2a_4^2)}{2a_0a_2a_4}, & \text{if } a_2 > a_3, \\ \frac{a_2(1-2a_2^2-2a_4^2)}{2a_0a_3a_4}, & \text{if } a_2 < a_3, \end{cases}$$
(49)

$$y = \cot \theta_3 = \begin{cases} \frac{a_2(1-2a_2^2-2a_4^2)}{2a_0a_3a_4}, & \text{if } a_2 > a_3, \\ \frac{a_3(1-2a_3^2-2a_4^2)}{2a_0a_2a_4}, & \text{if } a_2 < a_3. \end{cases}$$
(50)

Note that when $a_2 = a_3$, then there is $\cot \theta_2 = \cot \theta_3 = \cot(\pi - \theta_0) = \frac{1 - 2a_3^2 - 2a_4^2}{2a_0 a_4} = \frac{a_0^2 - a_4^2}{2a_0 a_4}$, which implies that $\theta_2 = \theta_3 = \pi - \theta_0$ (i.e. $Q(\theta_2, 0) = 0$) since $\theta_2, \theta_3, \pi - \theta_0 \in (0, \pi)$. Hence $a_2 \neq a_3$ in the expression $y = \cot \theta_2$ and $y = \cot \theta_3$ because of the hypothesis $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$. By checking, it can be proved that θ_1, θ_2 are the roots of $\frac{\partial(\sqrt{P(\theta, 0)} + \sqrt{Q(\theta, 0)})}{2\theta} = 0$. Thus

$$\min\left\{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\right\} = \begin{cases} \min\left\{\sqrt{P(\theta_1,0)} + \sqrt{Q(\theta_1,0)}, \sqrt{P(\theta_2,0)} + \sqrt{Q(\theta_2,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\right\}, & \text{if } a_2 \neq a_3, \\ \min\left\{\sqrt{P(\pi - \theta_0,0)}\right|_{a_2 = a_3}, \sqrt{P(\theta_1,0)} + \sqrt{Q(\theta_1,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\right\}, & \text{if } a_2 = a_3. \end{cases}$$
(51)

For the case $\varphi = \pi$ and the case $\varphi = 2\pi$, from (19), (20),

and (21), we have

$$\min\{\sqrt{P(\theta,\pi)} + \sqrt{Q(\theta,\pi)}\}\$$

$$= \min\{\sqrt{P(\pi-\theta,0)} + \sqrt{Q(\pi-\theta,0)}\}\$$

$$= \min\{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\}\$$

$$= \min\{\sqrt{P(\theta,2\pi)} + \sqrt{Q(\theta,2\pi)}\}.$$
(52)

Therefore, if $a_2 \neq a_3$, then

$$= \min\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\} = \min\{\sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}, \sqrt{P(\theta_1,0)} + \sqrt{Q(\theta_1,0)}, \sqrt{P(\theta_2,0)} + \sqrt{Q(\theta_2,0)}, \sqrt{P(\frac{\pi}{2},\frac{\pi}{2})} + \sqrt{Q(\frac{\pi}{2},\frac{\pi}{2})}\};$$
(53)

if
$$a_2 = a_3$$
, then

$$= \min\left\{ \sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} \right\}$$

=
$$\min\left\{ \sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}, \sqrt{P(\pi-\theta_0,0)} \Big|_{a_2=a_3}, \sqrt{P(\theta_1,0)} + \sqrt{Q(\theta_1,0)}, \sqrt{P(\frac{\pi}{2},\frac{\pi}{2})} + \sqrt{Q(\frac{\pi}{2},\frac{\pi}{2})} \right\}$$
(54)



In this section, we consider the quantum channel (2) with coefficients satisfying $a_4 = 0$ and $a_0 a_1 a_2 a_3 \neq 0$.

Note that

$$\begin{split} & \sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)} \\ &= \sqrt{P(\pi,\varphi)} + \sqrt{Q(\pi,\varphi)} \\ &= a_0^2 + \sqrt{(1-a_0^2)^2 - 4a_2^2 a_3^2} \\ &= a_0^2 + \sqrt{[a_1^2 + (a_2 - a_3)^2][a_1^2 + (a_2 + a_3)^2]}, \end{split}$$

and $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$ in case of $\theta = 0, \pi$. Next we suppose that $\sin \theta \neq 0$, that is $\theta \in (0, \pi)$.

We begin with the discussion of the condition such that $P(\theta, \varphi) = 0$. We can see that

$$P(\theta,\varphi) = \frac{1}{4} [1 + 2a_0 a_1 \cos(\varphi - \mu) \sin \theta - 2a_2 a_3 - \cos \theta (1 - 2a_0^2 - 2a_2 a_3)] [1 + 2a_0 a_1 \cos(\varphi - \mu) \sin \theta + 2a_2 a_3 - \cos \theta (1 - 2a_0^2 + 2a_2 a_3)] = \frac{1}{(1+t^2)^2} [1 + (t^2 - 1)a_0^2 + 2ta_0 a_1 \cos(\varphi - \mu) - 2a_2 a_3] \times [1 + (t^2 - 1)a_0^2 + 2ta_0 a_1 \cos(\varphi - \mu) + 2a_2 a_3] = 0$$

$$(55)$$

implies that

$$\begin{aligned}
\cos(\varphi - \mu) &= z_1 = \frac{-1 + a_0^2 - t^2 a_0^2 - 2a_2 a_3}{2t a_0 a_1}, \\
\cos(\varphi - \mu) &= z_2 = \frac{-1 + a_0^2 - t^2 a_0^2 + 2a_2 a_3}{2t a_0 a_1}.
\end{aligned}$$
(56)

Note that both $z_1 = \frac{-1+a_0^2-t^2a_0^2-2a_2a_3}{2ta_0a_1}$ and $z_2 = \frac{-1+a_0^2-t^2a_0^2+2a_2a_3}{2ta_0a_1}$ tend to $-\infty$ when $t \to 0$ and $t \to +\infty$. Since $z_1 \leq -2\sqrt{\frac{1-a_0^2+2a_2a_3}{2ta_0a_1}}\sqrt{\frac{t^2a_0^2}{2ta_0a_1}} = -\frac{\sqrt{a_1^2+(a_2+a_3)^2}}{a_1} < -1$, there is no φ such that $\cos(\varphi - \mu) = z_1 = \frac{-1+a_0^2-t^2a_0^2-2a_2a_3}{2ta_0a_1}$. Note that $z_2 \leq -2\sqrt{\frac{1-a_0^2-2a_2a_3}{2ta_0a_1}}\sqrt{\frac{t^2a_0^2}{2ta_0a_1}} = -\frac{\sqrt{a_1^2+(a_2-a_3)^2}}{a_1} \leq -1$, where the equality $z_2 = -1$ holds iff $t = \frac{a_1}{a_0}$ and $a_2 = a_3$. It follows that $\cos(\varphi - \mu) = z_2 = \frac{-1+a_0^2-t^2a_0^2+2a_2a_3}{2ta_0a_1}$ iff $a_2 = a_3, \ \varphi = \pi + \mu$, and $\theta = \theta_0$, where $\cot \frac{\theta_0}{2} = \frac{a_1}{a_0}$.

Thus, $P(\theta, \varphi) = 0$ if and only if $a_2 = a_3$, $\varphi = \pi + \mu$, and $\theta = \theta_0$. Therefore,

$$\left. \begin{array}{l} \left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} \right) \Big|_{P(\theta,\varphi)=0} \\ = \left. \sqrt{Q(\theta_0,\mu+\pi)} \Big|_{a_2=a_3} \\ = \left. \frac{\sqrt{\left(a_0^4 + a_1^2 + a_0^2 a_1^2 - 2 a_1^2 a_3^2\right) \left(a_0^4 + a_1^2 + a_0^2 a_1^2 + 2 a_1^2 a_3^2\right)}}{a_0^2 + a_1^2} \right.$$
(57)

By (19), we know that $Q(\theta, \varphi) = 0$ if and only if $a_2 = a_3, \varphi = \mu$, and $\theta = \pi - \theta_0$ and

$$\begin{cases} \left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\right)\Big|_{Q(\theta,\varphi)=0} \\ = \left. \sqrt{P(\pi-\theta_0,\mu)} \right|_{a_2=a_3} \\ = \left. \left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\right) \right|_{P(\theta,\varphi)=0}. \end{cases}$$
(58)

From above, we can see that the quantum channel with $a_4 = 0$, $a_2 = a_3$, and $a_0a_1a_2a_3 \neq 0$ can be collapsed to an EPR pair with probability $p_1 = p_2 = \frac{a_0^2(1-a_0^2-a_1^2)}{a_0^2+a_1^2}$ via Charlie's appropriate measurement (Charlie measures his particle in measurement basis (3) with $(\theta, \varphi) = (\theta_0, \mu + \pi)$ or $(\theta, \varphi) = (\pi - \theta_0, \mu)$).

In the following we suppose that $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$. From (26) and (27), we get

$$\begin{aligned} \sin(\varphi - \mu)[a_0\cos\theta + a_1\cos(\varphi - \mu)\sin\theta] &= 0,\\ \sin(\varphi - \mu)[a_0\cos\theta + a_1\cos(\varphi - \mu)\sin\theta] \\ \times [\sin^2\theta + 2a_0^2\cos^2\theta + 2a_0a_1\cos\theta\cos(\varphi - \mu)\sin\theta] &= 0. \end{aligned}$$
It follows that the minimum of $\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}$ must occur at either the hyperplane $\sin(\varphi - \mu) = 0$ or the hyperplane $\sin(\varphi - \mu) = 0$ or the hyperplane $\sin(\varphi - \mu) = 0$ or

the hyperplane $a_0 \cos \theta + a_1 \cos(\varphi - \mu) \sin \theta = 0$ when $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$ and $\theta \in (0, \pi)$. Now we consider the two hyperplanes.

If $a_0 \cos \theta + a_1 \cos(\varphi - \mu) \sin \theta = 0$, then $\cos(\varphi - \mu) = -\frac{a_0 \cot \theta}{a_1}$,

$$\left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} \right) \Big|_{\cos(\varphi-\mu) = -\frac{a_0 \cot \theta}{a_1}}$$

$$= \sqrt{1 - 4a_2^2 a_3^2}.$$
(60)

For $\sin(\varphi - \mu) = 0$, i.e. $\varphi = \mu, \mu + \pi$ in case of $\mu \in (0, \pi]$, or $\varphi = 0, \pi, 2\pi$ in case of $\mu = 0$, we first investigate the case $\varphi = \mu$. From (28), we have

$$\begin{aligned} &(a_0\cos\theta + a_1\sin\theta)[2a_0a_1(a_0^2 - a_1^2)\sin 2\theta - 2a_0^2(\sin^2\theta \\ &+ 2a_1^2\cos 2\theta) + \sin^2\theta(1 - 2a_1^2 - 4a_2^2a_3^2)] = 0, \end{aligned}$$

that is, $a_0 \cos \theta + a_1 \sin \theta = 0$, or $2a_0a_1(a_0^2 - a_1^2)\sin 2\theta - 2a_0^2(\sin^2 \theta + 2a_1^2\cos 2\theta) + \sin^2 \theta(1 - 2a_1^2 - 4a_2^2a_3^2) = 0$. From $a_0 \cos \theta + a_1 \sin \theta = 0$, there is

$$\cot \theta_1 = -\frac{a_1}{a_0}$$

Clearly, θ_1 is a root of (24). From

=

$$2a_0a_1(a_0^2 - a_1^2)\sin 2\theta - 2a_0^2(\sin^2\theta + 2a_1^2\cos 2\theta) +\sin^2\theta(1 - 2a_1^2 - 4a_2^2a_3^2) = 0,$$
(62)

there is

$$\cot \theta_2 = \frac{a_0^2 - a_1^2 - |a_2^2 - a_3^2|}{2a_0a_1},$$

$$\cot \theta_3 = \frac{a_0^2 - a_1^2 + |a_2^2 - a_3^2|}{2a_0a_1},$$

where $a_2 \neq a_3$. (Since when $a_2 = a_3$ there is $\theta_2 = \theta_3 = \pi - \theta_0$, i.e. $Q(\pi - \theta_0, \mu) = 0$, which contradict with the hypothesis $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$). By checking, we see θ_3 is a root of $\frac{\partial (\sqrt{P(\theta,\mu)} + \sqrt{Q(\theta,\mu)})}{\partial \theta} = 0$, while θ_2 is not. Hence,

$$= \begin{cases} \min\{\sqrt{P(\theta,\mu)} + \sqrt{Q(\theta,\mu)}\} \\ \min\{\sqrt{P(\theta_1,\mu)} + \sqrt{Q(\theta_1,\mu)}, \sqrt{P(\theta_3,\mu)} + \sqrt{Q(\theta_3,\mu)}, \sqrt{P(0,\mu)} + \sqrt{Q(0,\mu)}\} \\ \min\{\sqrt{P(\theta_1,\mu)} + \sqrt{Q(\theta_1,\mu)}, \sqrt{P(\pi-\theta_0,\mu)}|_{a_2=a_3}, \sqrt{P(0,\mu)} + \sqrt{Q(0,\mu)}\} \\ , & \text{if } a_2 \neq a_3. \end{cases}$$
(63)

For $\varphi = \mu + \pi, \mu + 2\pi$, there is

$$\min\{\sqrt{P(\theta,\mu+\pi)} + \sqrt{Q(\theta,\mu+\pi)}\} = \min\{\sqrt{P(\pi-\theta,\mu)} + \sqrt{Q(\pi-\theta,\mu)}\} = \min\{\sqrt{P(\theta,\mu)} + \sqrt{Q(\theta,\mu)}\} = \min\{\sqrt{P(\theta,\mu+2\pi)} + \sqrt{Q(\theta,\mu+2\pi)}\}.$$
(64)

from (20) and (19).

Thus, if $a_2 = a_3$, then

$$\min \left\{ \sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)} \right\} \\ = \min \left\{ a_0^2 + \sqrt{(1 - a_0^2)^2 - 4a_2^2 a_3^2}, \sqrt{1 - 4a_2^2 a_3^2}, \sqrt{P(\theta_1, \mu)} + \sqrt{Q(\theta_1, \mu)}, \sqrt{Q(\theta_0, \pi + \mu)} \right|_{a_2 = a_3} \right\};$$
(65)

if $a_2 \neq a_3$, then

$$\min \left\{ \sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)} \right\}$$

= $\min \left\{ a_0^2 + \sqrt{(1 - a_0^2)^2 - 4a_2^2 a_3^2}, \sqrt{1 - 4a_2^2 a_3^2}, \sqrt{P(\theta_1, \mu)} + \sqrt{Q(\theta_1, \mu)}, \sqrt{P(\theta_3, \mu)} + \sqrt{Q(\theta_3, \mu)} \right\}$
(66)
Here $\cot \frac{\theta_0}{2} = \frac{a_1}{a_0}, \cot \theta_1 = -\frac{a_1}{a_0}, \cot \theta_3 = \frac{a_0^2 - a_1^2 + |a_2^2 - a_3^2|}{2a_0 a_1}.$

H. One is $a_2 = 0$ and $a_0a_1a_3a_4 \neq 0$, the other is $a_3 = 0$ and $a_0a_1a_2a_4 \neq 0$

We now discuss the quantum channel (2), the coefficients of which satisfying $a_2 = 0$, but $a_0 a_1 a_3 a_4 \neq 0$. When $\sin \theta = 0$, i.e. $\theta = 0, \pi$, then

$$\sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}$$

$$= \sqrt{P(\pi,\varphi)} + \sqrt{Q(\pi,\varphi)}$$

= $a_0^2 + \sqrt{(1-a_0^2)^2 - 4a_1^2 a_4^2}$
= $a_0^2 + \sqrt{[a_3^2 + (a_1 - a_4)^2][a_3^2 + (a_1 + a_4)^2]},$ (67)

and $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$ in case of $\theta = 0, \pi$. Next we suppose that $\sin \theta \neq 0$.

First, we prove that $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$. If $P(\theta, \varphi) =$

0, then

$$\begin{aligned} &\cos(\varphi - \mu) \\ &= \frac{1}{2a_0^2 a_1^2 \sin^2 \theta} \Big[-a_0^3 a_1 \sin 2\theta + 8\sqrt{-a_0^2 a_1^2 a_3^2 a_4^2 \cos^2 \frac{\theta}{2} \sin^6 \frac{\theta}{2}} \\ &-4a_0 a_1 (1 - 2a_4^2) \cos \frac{\theta}{2} \sin^3 \frac{\theta}{2} \Big], \\ &\cos(\varphi - \mu) \\ &= -\frac{1}{2a_0^2 a_1^2 \sin^2 \theta} \Big[a_0^3 a_1 \sin 2\theta + 8\sqrt{-a_0^2 a_1^2 a_3^2 a_4^2 \cos^2 \frac{\theta}{2} \sin^6 \frac{\theta}{2}} \\ &+4a_0 a_1 (1 - 2a_4^2) \cos \frac{\theta}{2} \sin^3 \frac{\theta}{2} \Big]. \end{aligned}$$

But $8\sqrt{-a_0^2a_1^2a_3^2a_4^2\cos^2(\frac{\theta}{2})\sin^6(\frac{\theta}{2})}$ is an imaginary number, so $P(\theta,\varphi) \neq 0$. From (19), there is also $Q(\theta,\varphi) \neq 0$. That is, no matter what kind of measurement basis Charlie choose, Alice and Bob can never share an EPR pair after his measurement.

Second, the minimum of $\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)}$ should occur at the point satisfying (24) and (25), or $\theta = 0$. From (26) and (27), there are

$$\sin(\varphi - \mu)[a_0\cos\theta + a_1\cos(\varphi - \mu)\sin\theta] = 0, \quad (68)$$

and

$$\sin(\varphi - \mu)[a_0\cos\theta + a_1\cos(\varphi - \mu)\sin\theta][2a_0^2\cos^2\theta + a_0a_1\cos(\varphi - \mu)\sin2\theta + (1 - 2a_4^2)\sin^2\theta] = 0.$$
(69)

It follows that the minimum of $\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)}$ should occur at the point such that $\sin(\varphi - \mu) = 0$, or $a_0 \cos \theta + a_1 \cos(\varphi - \mu) \sin \theta = 0$, or $\theta = 0$.

From $a_0 \cos \theta + a_1 \cos(\varphi - \mu) \sin \theta = 0$, we know that $\cos(\varphi - \mu) = -\frac{a_0 \cot \theta}{a_1}$, and

$$\left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} \right) \Big|_{\{\theta \in (0,\pi), \cos(\varphi-\mu) = -\frac{a_0 \cot \theta}{a_1}\}}$$

$$= \sqrt{1 - 4a_0^2 a_4^2 - 4a_1^2 a_4^2}.$$

$$(70)$$

For the case $\sin(\varphi - \mu) = 0$, since equality (64) holds, we need only to consider the case $\varphi = \mu$. From (28),we derive

$$\begin{aligned} \sin 3\theta \left(2 a_0^2 a_1 - a_1 + 4 a_0^4 a_1 + 2 a_1^3 - 12 a_0^2 a_1^3\right) \\ + \sin \theta \left(3 a_1 - 6 a_0^2 a_1 + 4 a_0^4 a_1 - 6 a_1^3 + 4 a_0^2 a_1^3\right) \\ + \cos \theta \left(a_0 - 2 a_0^3 - 2 a_0 a_1^2 - 4 a_0^3 a_1^2 - 4 a_0 a_1^4\right) \\ + \cos 3\theta \left(2 a_0^3 - a_0 + 2 a_0 a_1^2 - 12 a_0^3 a_1^2 + 4 a_0 a_1^4\right) \\ = \left\{-16 \cot^3 \theta a_0^3 a_1^2 + 16 \cot^2 \theta a_0^2 a_1 \left(a_0^2 - 2 a_1^2\right) \\ + 4 \cot \theta a_0 \left[1 - 2 a_1^2 - 4 a_1^4 + 2 a_0^2 (4 a_1^2 - 1)\right] \\ + 4 \left(1 - 2 a_0^2\right) a_1 \left(1 - 2 a_1^2\right)\right\} \sin^3 \theta \\ = 0, \end{aligned} \tag{71}$$

which follows that

$$\cot \theta = \cot \theta_1 = -\frac{a_1}{a_0},$$

$$\cot \theta = \cot \theta_2 = \frac{1 - 2a_1^2}{2a_0a_1},$$

$$\cot \theta = \cot \theta_3 = \frac{-1 + 2a_0^2}{2a_0a_1},$$

where θ_1 and θ_2 are the roots of equation $\frac{\partial(\sqrt{P(\theta,\mu)}+\sqrt{Q(\theta,\mu)})}{\partial\theta} = 0$, while θ_3 is not. It is not difficult to obtain

$$= \min \left\{ \sqrt{P(\theta, \mu)} + \sqrt{Q(\theta, \mu)} \right\}$$

=
$$\min \left\{ \sqrt{P(\theta_1, \mu)} + \sqrt{Q(\theta_1, \mu)}, \sqrt{P(\theta_2, \mu)} + \sqrt{Q(\theta_2, \mu)} \right\}$$

$$\sqrt{P(0, \mu)} + \sqrt{Q(0, \mu)} \right\}.$$

(72)

Therefore, for quantum channel (2) with $a_2 = 0$ and $a_0 a_1 a_3 a_4 \neq 0$, there is

$$\min\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\} = \min\{\sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}, \sqrt{1 - 4a_0^2a_4^2 - 4a_1^2a_4^2}, \sqrt{P(\theta_1,\mu)} + \sqrt{Q(\theta_1,\mu)}, \sqrt{P(\theta_2,\mu)} + \sqrt{Q(\theta_2,\mu)}\}$$
(73)

Similarly, for quantum channel (2) with $a_3 = 0$ and $a_0 a_1 a_2 a_4 \neq 0$ we also obtain

$$\min\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\} = \min\{a_0^2 + \sqrt{(1-a_0^2)^2 - 4a_1^2a_4^2}, \sqrt{1-4a_0^2a_4^2 - 4a_1^2a_4^2}, \sqrt{P(\theta_1,\mu)} + \sqrt{Q(\theta_1,\mu)}, \sqrt{P(\theta_2,\mu)} + \sqrt{Q(\theta_2,\mu)}\}\}$$
(74)
Here $\cot \theta_1 = -\frac{a_1}{a_0}$.

I. $\mu = 0$ and $a_0 a_1 a_2 a_3 a_4 \neq 0$

In this section, we investigate the quantum channel (2) with $\mu = 0$ and $a_0 a_1 a_2 a_3 a_4 \neq 0$. Evidently,

$$\sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}
 = \sqrt{P(\pi,\varphi)} + \sqrt{Q(\pi,\varphi)}
 = a_0^2 + \sqrt{(1-a_0^2)^2 - 4(a_2a_3 - a_1a_4)^2}
 = a_0^2 + \sqrt{[(a_2 - a_3)^2 + (a_1 + a_4)^2][(a_2 + a_3)^2 + (a_1 - a_4)^2]}
 (75)$$

and $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$ in case of $\theta = 0, \pi$. In the following, we suppose that $\theta \neq 0, \pi$, i.e. $\theta \in (0, \pi)$.

We first consider the case $P(\theta, \varphi) = 0$. If $P(\theta, \varphi) = 0$, then

$$\begin{aligned} \cos\varphi &= z_1 \\ &= \frac{1}{2 t a_0 a_1^2} \{ -t^2 a_0^2 a_1 - (1 - a_0^2) a_1 \\ &- 2(a_2 a_3 - a_1 a_4) a_4 - 2[t^2 a_0^2 a_1 a_2 a_3 a_4 + \\ &(a_2 a_3 - a_1 a_4)(a_1 a_3 + a_2 a_4)(a_1 a_2 + a_3 a_4)]^{\frac{1}{2}} \}, \\ &\cos\varphi &= z_2 \end{aligned}$$

$$= \frac{1}{2 t a_0 a_1^2} \{ -t^2 a_0^2 a_1 - (1 - a_0^2) a_1 \\ -2(a_2 a_3 - a_1 a_4) a_4 + 2[t^2 a_0^2 a_1 a_2 a_3 a_4 + (a_2 a_3 - a_1 a_4)(a_1 a_3 + a_2 a_4)(a_1 a_2 + a_3 a_4)]^{\frac{1}{2}} \}.$$

Note that if $a_1 a_4 - a_2 a_3 > 0$, then $t \in [t_0, +\infty)$; if $a_1 a_4 - a_2 a_3 \leq 0$, then $t \in (0, +\infty)$. Here, $t_0 = \sqrt{\frac{(a_1 a_4 - a_2 a_3)(a_1 a_3 + a_2 a_4)(a_1 a_2 + a_3 a_4)}{a_0^2 a_1 a_2 a_3 a_4}}$. Obviously, both z_1 and z_2 go to $-\infty$ when $t \to +\infty$. We can prove that $z = z_1 = \frac{1}{2t a_0 a_1^2} \{-t^2 a_0^2 a_1 - t^2 a_0^2 a_1 -$

We can prove that $z = z_1 = \frac{1}{2ta_0a_1^2} \{-t^2a_0^2a_1 - (1-a_0^2)a_1 - 2(a_2a_3 - a_1a_4)a_4 - 2[t^2a_0^2a_1a_2a_3a_4 + (a_2a_3 - a_1a_4)(a_1a_3 + a_2a_4)(a_1a_2 + a_3a_4)]^{\frac{1}{2}}\}$ has no intersection point with the straight line z = -1. Since z_1 is a continuous function of t and tends to $-\infty$ when t tends to $+\infty$, so $z_1 < -1$. Thus, there is no φ satisfying $\cos \varphi = z_1$. Let us look at $z = z_2$. $z_2 = -1$ implies that

$$\begin{bmatrix} -t^{2} a_{0}^{2} a_{1} - (1 - a_{0}^{2})a_{1} - 2(a_{2}a_{3} - a_{1}a_{4})a_{4} + 2ta_{0}a_{1}^{2} \end{bmatrix}^{2} -4\{t^{2} a_{0}^{2} a_{1} a_{2} a_{3} a_{4} + (a_{2} a_{3} - a_{1} a_{4}) \times (a_{1}a_{3} + a_{2}a_{4})(a_{1}a_{2} + a_{3}a_{4})\} = a_{1}^{2} \begin{bmatrix} -t^{2} a_{0}^{2} + 2t a_{0}(a_{1} + a_{4}) - 1 + a_{0}^{2} + 2a_{2} a_{3} - 2a_{1} a_{4} \end{bmatrix} \times \begin{bmatrix} -t^{2} a_{0}^{2} + 2t a_{0}(a_{1} - a_{4}) - 1 + a_{0}^{2} - 2a_{2} a_{3} + 2a_{1} a_{4} \end{bmatrix} = 0,$$
(76)

which follows that

$$t = t_1 = \frac{a_1 - a_4 - \sqrt{-(a_2 + a_3)^2}}{a_0},$$

$$t = t_2 = \frac{a_1 - a_4 + \sqrt{-(a_2 + a_3)^2}}{a_0},$$

$$t = t_3 = \frac{a_1 + a_4 - \sqrt{-(a_2 - a_3)^2}}{a_0},$$

$$t = t_4 = \frac{a_1 + a_4 + \sqrt{-(a_2 - a_3)^2}}{a_0}.$$

Here t_1 and t_2 are imaginary numbers, while t_3 and t_4 are real numbers only if $a_2 = a_3$. By checking, when $a_2 = a_3$, $t_3 = t_4 = \frac{a_1+a_4}{a_0}$ is a root of equation $z_2 = -1$, and the maximum point of function $z = z_2$. Since $z = z_2$ is a continuous function of t, tends to $-\infty$ when t tends to $+\infty$, and has only one intersection point with straight line z = -1, there must be $z_2 \leq -1$, where the equality holds iff $a_2 = a_3$ and $t = t_3 = t_4 = \frac{a_1 + a_4}{a_0}$. It means that $P(\theta, \varphi) = 0$ iff $a_2 = a_3$, $\varphi = \pi$, and $\theta = \theta_0$, where $\cot \frac{\theta_0}{2} = \frac{a_1 + a_4}{a_0}$. From (19), we can show that $Q(\theta, \varphi) = 0$ iff $a_2 = a_3$, $\varphi = 0$, and $\theta = \pi - \theta_0$. Therefore

$$\begin{split} & (\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}) \big|_{P(\theta,\varphi)=0} \\ &= \sqrt{Q(\theta_0,\pi)} \big|_{a_2=a_3} \\ &= \sqrt{P(\pi-\theta_0,0)} \big|_{a_2=a_3} \\ &= (\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}) \big|_{Q(\theta,\varphi)=0} \\ &= \sqrt{(1-2a_4^2)^2 + 4a_3^2(-1+a_1^2+a_3^2+2a_1a_4+3a_4^2)} \end{split}$$

It is shown that the quantum channel (2) with $\mu = 0$, $a_0a_1a_2a_3a_4 \neq 0$, and $a_2 = a_3$, collapses to an EPR pair with probability $p_1 = p_2 = \frac{a_0^2(1-a_0^2+3a_1^2+4a_1a_4+a_4^2)}{a_0^2+(a_1+a_4)^2}$ after Charlie's measurement in the basis (3) with either $\varphi = \pi$ and $\theta = \theta_0$, or $\varphi = 0$ and $\theta = \pi - \theta_0$ in case of $a_2 = a_3$.

In the following we suppose that $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$. By (26), there is

$$\sin \varphi \{ 2 \cos \varphi (a_2 a_3 - a_1 a_4) (a_1 a_3 + a_2 a_4) (a_1 a_2 + a_3 a_4) -y a_0 [a_2 a_3 a_4 (1 + 2a_1^2 - 2a_2^2 - 2a_3^2) -2 a_1 (a_2^2 a_3^2 - a_2^2 a_4^2 - a_3^2 a_4^2)] \} = 0.$$

(77) Note that $a_2a_3a_4(1+2a_1^2-2a_2^2-2a_3^2)-2a_1(a_2^2a_3^2-a_2^2a_4^2-a_3^2a_4^2) = a_1a_4^2 \neq 0$ if $a_2a_3 = a_1a_4$. Therefore,

$$a_2a_3a_4(1+2a_1^2-2a_2^2-2a_3^2)-2a_1(a_2^2a_3^2-a_2^2a_4^2-a_3^2a_4^2)=0$$

implies that $\cos \varphi = 0$. Thus, (77) implies that $\sin \varphi = 0$, $\cos \varphi = 0$, or

$$y = \frac{2\cos\varphi(a_2a_3 - a_1a_4)[(a_1^2 + a_4^2)a_2a_3 + a_1(a_2^2 + a_3^2)a_4]}{a_0[a_2a_3a_4(1 + 2a_1^2 - 2a_2^2 - 2a_3^2) - 2a_1(a_2^2a_3^2 - a_2^2a_4^2 - a_3^2a_4^2)]}$$

in case of sin $2\varphi \neq 0$.

From (27), there is

$$\begin{aligned} 4y^{3}a_{0}^{3}(a_{2}a_{3}-a_{1}a_{4})(a_{1}a_{3}+a_{2}a_{4})(a_{1}a_{2}+a_{3}a_{4}) \\ +8y^{2}a_{0}^{2}a_{1}\cos\varphi(a_{2}a_{3}-a_{1}a_{4})(a_{1}a_{3}+a_{2}a_{4})(a_{1}a_{2}+a_{3}a_{4}) \\ +ya_{0}\{4a_{1}^{2}\cos^{2}\varphi(a_{2}a_{3}-a_{1}a_{4})(a_{1}a_{3}+a_{2}a_{4})(a_{1}a_{2}+a_{3}a_{4}) \\ +(a_{1}+2a_{2}a_{3}a_{4}-2a_{1}a_{4}^{2})[a_{2}a_{3}(-2a_{4}^{3}-4a_{1}^{2}a_{4}\\ +a_{4}+2a_{1}a_{2}a_{3})-2a_{1}a_{4}^{2}(a_{2}^{2}+a_{3}^{2})]\} \\ +2\cos\varphi(a_{1}+2a_{2}a_{3}a_{4}-2a_{1}a_{4}^{2})[a_{2}^{2}a_{3}^{2}(a_{1}^{2}+a_{4}^{2})\\ +a_{1}a_{2}a_{3}a_{4}(1-2a_{1}^{2}-2a_{4}^{2})-a_{1}^{2}a_{4}^{2}(a_{2}^{2}+a_{3}^{2})] \\ =0. \end{aligned}$$

$$\tag{79}$$

Combining (78) and (79), we obtain that

$$\cos \varphi(a_1 + 2a_2a_3a_4 - 2a_1a_4^2) \{a_0^2 [2a_1(a_2^2a_3^2 - a_2^2a_4^2 - a_3^2a_4^2) - a_2a_3a_4(1 + 2a_1^2 - 2a_2^2 - 2a_3^2)]^2
+ 4\cos^2 \varphi(a_2a_3 - a_1a_4)^2(a_1a_3 + a_2a_4)^2(a_1a_2 + a_3a_4)^2 \}
= 0,$$
(80)

which implies that $\cos \varphi = 0$.

Next we only need to examine the two cases $\sin \varphi = 0$, and $\cos \varphi = 0$, that is, $\varphi = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi$.

Now let look at the hyperplane $\varphi = 0$. From (28), we have

$$\begin{bmatrix} 2ya_0(a_2a_3 - a_1a_4) + 2a_1a_2a_3 + a_4 - 2a_1^2a_4 \\ \times \{a_2a_3 - 2a_0^2a_2a_3 - 2a_1^2a_2a_3 + 4a_0^2a_1^2a_2a_3 \\ -4a_2^3a_3^3 - 2a_1a_4 + 6a_0^2a_1a_4 - 4a_0^4a_1a_4 + 2a_1^3a_4 \\ -4a_0^2a_1^3a_4 + 8a_1a_2^2a_3^2a_4 - 2a_2a_3a_4^2 + 4a_0^2a_2a_3a_4^2 \\ -4a_1^2a_2a_3a_4^2 + 2a_1a_4^3 - 4a_0^2a_1a_4^3 \\ -4y^2a_0^2(a_1a_3 + a_2a_4)(a_1a_2 + a_3a_4) \\ -2ya_0(-2a_0^2a_1a_2a_3 + 2a_1^3a_2a_3 + a_4 - 3a_0^2a_4 \\ +2a_0^4a_4 + a_1^2a_4 - 2a_1^4a_4 - 4a_2^2a_3^2a_4 + 6a_1a_2a_3a_4^2 \\ -a_4^3 + 2a_0^2a_4^3 - 2a_1^2a_4^3) \} = 0.$$

$$\tag{81}$$

First, we consider the case $a_1a_4 - a_2a_3 \neq 0$. In this case, (81) has the following solution

$$\begin{split} y &= \cot \theta_1 \; = \; \frac{2a_1a_2a_3 + a_4 - 2a_1^2a_4}{2a_0(a_1a_4 - a_2a_3)}, \\ & \text{where if } a_2 = a_3, \text{then } a_1a_4 - a_2a_3 \neq \frac{1}{2}; \\ y &= \cot \theta_2 \; = \; \frac{-2a_1a_2a_4 + a_3(1 - 2a_1^2 - 2a_3^2 - 2a_4^2)}{2a_0(a_1a_3 + a_2a_4)}, \\ & \text{where } a_2 \neq a_3; \\ y &= \cot \theta_3 \; = \; \frac{-2a_1a_3a_4 + a_2(1 - 2a_1^2 - 2a_2^2 - 2a_4^2)}{2a_0(a_1a_2 + a_3a_4)}, \\ & \text{where } a_2 \neq a_3. \end{split}$$

By checking, we show that θ_1 is a root of $\frac{\partial(\sqrt{P(\theta,0)}+\sqrt{Q(\theta,0)})}{\partial\theta}\Big|_{a_1a_4-a_2a_3\neq 0} = 0$, while θ_2 is a root only if $a_2 > a_3$, and θ_3 is a root only if $a_2 < a_3$.

Second, we consider the case $a_1a_4 - a_2a_3 = 0$. From (81), we get

$$\begin{split} y_4 &= \cot \theta_4 = \ \frac{(1-2a_0^2)(-1+a_0^2-a_1^2+a_4^2+|a_2^2-a_3^2|)}{4a_0(1-a_0^2)a_1}, \\ y_5 &= \cot \theta_5 = \ \frac{(1-2a_0^2)(-1+a_0^2-a_1^2+a_4^2-|a_2^2-a_3^2|)}{4a_0(1-a_0^2)a_1}, \\ y_6 &= \ \frac{(1-2a_0^2)(-1+a_0^2-a_1^2+a_4^2-|a_2^2-a_3^2|)}{4a_0(1-a_0^2)a_1}, \\ y_6 &= \ \frac{(1-2a_0^2)(-1+a_0^2-a_1^2+a_4^2-|a_2^2-a_3^2|)}{4a_0(1-a_0^2)a_1}, \end{split}$$

By checking, we know that only θ_4 is a root of $\frac{\partial(\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)})}{\partial \theta}|_{a_1a_4 - a_2a_3 = 0} = 0.$

Therefore,

$$\min\{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\}$$

$$= \begin{cases} \min\{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\}, \sqrt{P(\theta_2,0)} + \sqrt{Q(\theta_2,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, \text{ if } a_2 > a_3 \text{ and } a_1a_4 - a_2a_3 \neq 0, \\ \min\{\sqrt{P(\theta_1,0)} + \sqrt{Q(\theta_1,0)}, \sqrt{P(\theta_3,0)} + \sqrt{Q(\theta_3,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, \text{ if } a_2 < a_3 \text{ and } a_1a_4 - a_2a_3 \neq 0, \\ \min\{\sqrt{P(\theta_4,0)} + \sqrt{Q(\theta_4,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, \text{ if } a_2 \neq a_3 \text{ and } a_1a_4 - a_2a_3 \neq 0, \\ \min\{\sqrt{P(\pi - \theta_0,0)}|_{a_2 = a_3}, \sqrt{P(\theta_1,0)} + \sqrt{Q(\theta_1,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, \text{ if } a_2 \neq a_3 \text{ and } a_1a_4 - a_2a_3 = 0, \\ \min\{\sqrt{P(\pi - \theta_0,0)}|_{a_2 = a_3}, \sqrt{P(\theta_1,0)} + \sqrt{Q(\theta_1,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, \text{ if } a_2 = a_3. \end{cases}$$

$$\tag{82}$$

For $\varphi = \pi, 2\pi$, by (52), there is

$$\min\{\sqrt{P(\theta,\pi)} + \sqrt{Q(\theta,\pi)}\} = \min\{\sqrt{P(\theta,2\pi)} + \sqrt{Q(\theta,2\pi)}\} = \min\{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\}.$$
(83)

Now we investigate the hyperplanes $\varphi = \frac{\pi}{2}$ and $\varphi = \frac{3\pi}{2}$. From $\frac{d(\sqrt{P(\theta, \frac{\pi}{2})} + \sqrt{Q(\theta, \frac{\pi}{2})})}{d\theta} = 0$, there is $\cos \theta = 0$, i.e. $\theta = \frac{\pi}{2}$. Similarly, by $\frac{d(\sqrt{P(\theta, \frac{3\pi}{2})} + \sqrt{Q(\theta, \frac{3\pi}{2})})}{d\theta} = 0$, we also get $\theta = \frac{\pi}{2}$. It is direct from (19) and (20) that

Therefore,

$$= \min\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\} \\ = \min\{\sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}, \sqrt{P(\frac{\pi}{2},\frac{\pi}{2})} + \sqrt{Q(\frac{\pi}{2},\frac{\pi}{2})}, \\ \min\{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\}\}.$$
(85)



In this section, we examine the quantum channel (2) with coefficients satisfying $\mu = \pi$ and $a_0 a_1 a_2 a_3 a_4 \neq 0$.

Let us start with the special case $\sin \theta = 0$. In this case, there is

First, we look for the condition of $P(\theta, \varphi) = 0$, $Q(\theta, \varphi) = 0$. If $P(\theta, \varphi) = 0$, then

$$\cos \varphi = z_{1} \\
= \frac{1}{2a_{0}a_{1}^{2}t} [t^{2}a_{0}^{2}a_{1} + (1 - a_{0}^{2})a_{1} - 2a_{4}(a_{2}a_{3} + a_{1}a_{4}) - 2\sqrt{-t^{2}a_{0}^{2}a_{1}a_{2}a_{3}a_{4} + (a_{2}a_{3} + a_{1}a_{4})(a_{1}a_{3} - a_{2}a_{4})(a_{1}a_{2} - a_{3}a_{4})], \\
\cos \varphi = z_{2} \\
= \frac{1}{2a_{0}a_{1}^{2}t} [t^{2}a_{0}^{2}a_{1} + (1 - a_{0}^{2})a_{1} - 2a_{4}(a_{2}a_{3} + a_{1}a_{4}) + 2\sqrt{-t^{2}a_{0}^{2}a_{1}a_{2}a_{3}a_{4} + (a_{2}a_{3} + a_{1}a_{4})(a_{1}a_{3} - a_{2}a_{4})(a_{1}a_{2} - a_{3}a_{4})]. \\$$
(87)

Here $(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) > 0$, and $t \in (0, \sqrt{\frac{(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4)}{a_0^2a_1a_2a_3a_4}}]$, since $(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) \leq 0$ implies that $P(\theta, \varphi) \neq 0$. Directly, $(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) > 0$ if and only if $a_2^2 + a_3^2 < \frac{a_2a_3(a_1^2 + a_4^2)}{a_1a_4}$.

Note that

$$(1 - a_0^2)a_1 - 2a_4(a_2a_3 + a_1a_4) = a_1(a_1^2 - a_4^2) + a_1(a_2^2 + a_3^2) - 2a_2a_3a_4 > 0$$

in case of $a_1 > a_4$,

$$\begin{array}{l} (1-a_0^2)a_1 - 2a_4(a_2a_3 + a_1a_4) \\ = a_1(a_1^2 - a_4^2) + a_1(a_2^2 + a_3^2) - 2a_2a_3a_4 \\ < \frac{(a_1^2 - a_4^2)(a_2a_3 + a_1a_4)}{a_4} \\ < 0 \end{array}$$

in case of $a_1 < a_4$ and $(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) > 0$,

and

$$[(1 - a_0^2)a_1 - 2a_4(a_2a_3 + a_1a_4)]^2 - \{2\sqrt{(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4)}\}^2 = a_1^2[(a_1 - a_4)^2 + (a_2 - a_3)^2][(a_1 + a_4)^2 + (a_2 + a_3)^2] > 0.$$
(88)

It follows that

$$\begin{array}{l} (1 - a_0^2)a_1 - 2a_4(a_2a_3 + a_1a_4) \\ + 2\sqrt{(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4)} \\ = (a_1 - a_4)(a_1^2 + 2a_3^2 + a_1a_4) + 2a_3\sqrt{(a_1 - a_4)^2(a_3^2 + a_1a_4)} \\ > 0 \end{array}$$

$$\begin{array}{l} (89) \end{array}$$

and

in case of $a_1 > a_4$;

$$\begin{array}{l} (1-a_0^2)a_1 - 2a_4(a_2a_3 + a_1a_4) \\ + 2\sqrt{(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4)} \\ = (a_1 - a_4)(a_1^2 + 2a_3^2 + a_1a_4) + 2a_3\sqrt{(a_1 - a_4)^2(a_3^2 + a_1a_4)} \\ < 0 \end{array}$$

$$\begin{array}{l} (91) \end{array}$$

and

$$\begin{array}{l} {}^{44}) & (1-a_0^2)a_1 - 2a_4(a_2a_3 + a_1a_4) \\ & -2\sqrt{(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4)} \\ = & (a_1 - a_4)(a_1^2 + 2a_3^2 + a_1a_4) - 2a_3\sqrt{(a_1 - a_4)^2(a_3^2 + a_1a_4)} \\ < & 0 \end{array}$$

 $\begin{array}{l} (1-a_0^2)a_1 - 2a_4(a_2a_3 + a_1a_4) \\ -2\sqrt{(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4)} \\ = (a_1 - a_4)(a_1^2 + 2a_3^2 + a_1a_4) - 2a_3\sqrt{(a_1 - a_4)^2(a_3^2 + a_1a_4)} \\ = 0 \end{array}$ in case of $a_1 < a_4$. Thus, if $a_1 > a_4$, then both z_1 and z_2 go to $+\infty$ when $t \to 0$; if $a_1 < a_4$, then both z_1 and z_2 go to $-\infty$ when $t \to 0$; if $a_1 < a_4$, then both z_1 and z_2 go to $-\infty$ when $t \to 0$.

(90)

Now suppose that $a_1 > a_4$. From $z_1 = 1$, there is

$$\frac{\left[t^2 a_0^2 a_1 - 2t a_0 a_1^2 + (1 - a_0^2) a_1 - 2a_4 (a_2 a_3 + a_1 a_4)\right]^2}{-\left[2\sqrt{-t^2 a_0^2 a_1 a_2 a_3 a_4 + (a_2 a_3 + a_1 a_4)(a_1 a_3 - a_2 a_4)(a_1 a_2 - a_3 a_4)}\right]^2} = 0,$$
(93)

the solutions of which are

$$t = t_{11} = \frac{a_1 + a_4 - \sqrt{-(a_2 + a_3)^2}}{a_0},$$

$$t = t_{12} = \frac{a_1 + a_4 + \sqrt{-(a_2 + a_3)^2}}{a_0},$$

$$t = t_{13} = \frac{a_1 - a_4 - \sqrt{-(a_2 - a_3)^2}}{a_0},$$

$$t = t_{14} = \frac{a_1 - a_4 + \sqrt{-(a_2 - a_3)^2}}{a_0}.$$
(94)

Here, t_{11} and t_{12} are imaginary numbers, and t_{13} and t_{14} are positive real numbers only if $a_2 = a_3$ and $a_1 > a_4$. It is not difficult to check that θ_0 is a root of $z_1 = 1$ and the minimum point of z_1 in case of $a_1 > a_4$ and $a_2 = a_3$. Here $t = \cot \frac{\theta_0}{2} = \frac{a_1 - a_4}{a_0}$. $z_1 \ge 1$ comes directly since $z = z_1$ is a continuous function of t, goes to $+\infty$ when $t \to 0$, and has only one intersection point with straight line z = 1, where the equality $z_1 = 1$ holds iff $a_2 = a_3, a_1 > a_4$ and $\cot \frac{\theta_0}{2} = \frac{a_1 - a_4}{a_0}$. We can show that $z_2 > 1$ in case of $a_1 > a_4$ in the same way.

Similarly, we can prove that if $a_1 < a_4$, then $z_1 < -1$, and $z_2 \leq -1$, where $z_2 = -1$ iff $a_2 = a_3$, $a_1 < a_4$, and $t = \cot \frac{\overline{\theta}_0}{2} = \frac{-a_1+a_4}{a_0}$. Therefore, $P(\theta, \varphi) = 0$ iff $\varphi = 0$ and $\theta = \theta_0$ in case of $a_2 = a_3$ and $a_1 > a_4$, or $\varphi = \pi$ and $\theta = \overline{\theta}_0$ in case of $a_2 = a_3$ and $a_1 < a_4$. It can be derived that

$$\left(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)} \right) \Big|_{P(\theta,\varphi)=0, \ \theta \in (0,\pi)}$$

$$= \sqrt{Q(\theta_0,0)} \Big|_{\{a_2=a_3,a_1>a_4\}}$$

$$= \sqrt{Q(\overline{\theta}_0,\pi)} \Big|_{\{a_2=a_3,a_1

$$= \sqrt{(1-2a_4^2)^2 + 4a_3^2(-1+a_1^2+a_3^2-2a_1a_4+3a_4^2)}.$$

$$(95)$$$$

Immediately, from (19) and (20), $Q(\theta, \varphi) = 0$ iff $\varphi = \pi$ and $\theta = \pi - \theta_0$ in case of $a_2 = a_3$ and $a_1 > a_4$, or $\varphi = 0$ and $\theta = \pi - \overline{\theta}_0$ in case of $a_2 = a_3$ and $a_1 < a_4$, and

$$= \frac{(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)})|_{Q(\theta,\varphi)=0}}{(\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)})|_{P(\theta,\varphi)=0}}.$$
(96)

From above, it can be seen that the quantum channel (2), the coefficients of which satisfy $\mu = \pi$, $a_0a_1a_2a_3a_4 \neq 0$ and $a_2 = a_3$, can be collapsed to a Bell state with probability $p_1 = p_2 = \frac{a_0^2(1-a_0^2+3a_1^2-4a_1a_4+a_4^2)}{a_0^2+(a_1-a_4)^2}$ by Charlie's measurement in the basis (3) with $(\theta, \varphi) = (\theta_0, 0)$ or $(\theta, \varphi) = (\pi - \theta_0, \pi)$ in case of $a_1 > a_4$, or $(\theta, \varphi) = (\theta_0, \pi)$ or $(\theta, \varphi) = (\pi - \theta_0, 0)$ in case of $a_1 < a_4$. In particular, this quantum channel can also be purified to an EPR pair with probability $2a_1^2 + 2a_2^2$ via controller's measurement in the basis $|0\rangle, |1\rangle$ in case of $a_1 = a_4$.

In the following, we suppose that $P(\theta, \varphi)Q(\theta, \varphi) \neq 0$. In order to obtain the minimum of $\sqrt{P(\theta, \varphi)} + \sqrt{Q(\theta, \varphi)}$, we need to find the points such that equations (26) and (27) hold. From (26), we get $\sin \varphi = 0$, or

$$ya_0[2a_1a_2^2a_3^2 - a_2a_3a_4(-2a_0^2 - 4a_1^2 - 2a_4^2 + 1) -2a_1a_4^2(a_2^2 + a_3^2)] -2\cos\varphi(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) = 0.$$
(97)

By (27), we have $\sin \varphi = 0$, or

. . .

$$\begin{aligned} &4y^3a_0^3(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) \\ &-8y^2a_0^2a_1\cos\varphi(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) \\ &+ya_0[(a_1 - 2a_2a_3a_4 - 2a_1a_4^2)(2a_1a_2^2a_3^2 - a_2a_3a_4) \\ &+4a_1^2a_2a_3a_4 - 2a_1a_2^2a_4^2 - 2a_1a_3^2a_4^2 + 2a_2a_3a_4^3) \\ &+4a_1^2\cos^2\varphi(a_2a_3 + a_1a_4)(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4)] \\ &+2\cos\varphi(-a_1 + 2a_2a_3a_4 + 2a_1a_4^2)[a_2^2a_3^2(a_1^2 + a_4^2) \\ &-a_1a_2a_3a_4(-2a_1^2 - 2a_4^2 + 1) - a_1^2(a_2^2 + a_3^2)a_4^2] \\ &= 0. \end{aligned}$$
(98)

If $(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) = 0$, then (97) and (98) become

$$y = 0, \tag{99}$$

and

$$y[a_4^2 - (1 - 2a_0^2)a_1^2] - 2a_0a_1\cos\varphi(a_1^2 + a_4^2) = 0, \quad (100)$$

respectively. However, if $(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) \neq 0$, then (97) and (98) become

$$= \frac{ya_0\{-2a_1a_4^2(a_3^2+a_2^2)+a_2a_3[2a_1a_2a_3+a_4+2(a_1^2-a_2^2-a_3^2)a_4]\}}{2(a_2a_3+a_1a_4)(a_1a_3-a_2a_4)(a_1a_2-a_3a_4)}$$
(101)

and

$$\frac{y(1+y^2)a_0^3a_2^2a_3^2a_4^2(2a_2a_3a_4-a_1(1-2a_4^2))^2}{(a_2a_3+a_1a_4)(a_1a_3-a_2a_4)(a_1a_2-a_3a_4)} = 0, (102)$$

respectively. That is, y = 0, $\cos \varphi = 0$.

For getting the minimum point of $\sqrt{P(\theta,\varphi)}$ + $\sqrt{Q(\theta,\varphi)}$, it is enough for us to consider the hyperplanes $\sin \varphi = 0$, and $\cos \varphi = 0$. For hyperplane $\sin \varphi = 0$, we need only to consider the case $\varphi = 0$ by (19), (20), and (21). For hyperplane $\cos \varphi = 0$, we need only to consider the case $\varphi = \frac{\pi}{2}$ by (19) and (20).

Now, let us consider the case $\varphi = 0$. In this case (28) becomes

$$\begin{aligned} & \left[-2a_{1}a_{2}a_{3} + a_{4} - 2a_{1}^{2}a_{4} + 2ya_{0}(a_{2}a_{3} + a_{1}a_{4}) \right] \\ & \times \left[4y^{2}a_{0}^{2}(a_{1}a_{3} - a_{2}a_{4})(a_{1}a_{2} - a_{3}a_{4}) \right. \\ & \left. + 2ya_{0}(2a_{1}a_{2}a_{3} - 4a_{1}^{3}a_{2}a_{3} - 2a_{1}a_{2}^{3}a_{3} \right. \\ & \left. - 2a_{1}a_{2}a_{3}^{3} - a_{2}^{2}a_{4} + 4a_{1}^{2}a_{2}^{2}a_{4} + 2a_{2}^{4}a_{4} - a_{3}^{2}a_{4} \right. \\ & \left. + 4a_{1}^{2}a_{3}^{2}a_{4} + 2a_{3}^{4}a_{4} - 8a_{1}a_{2}a_{3}a_{4}^{2} + 2a_{2}^{2}a_{4}^{3} + 2a_{3}^{2}a_{4}^{3} \right. \\ & \left. + (a_{2} - 2a_{1}^{2}a_{2} - 2a_{2}^{3} + 2a_{1}a_{3}a_{4} - 2a_{2}a_{4}^{2}) \right. \\ & \left. \times (a_{3} - 2a_{1}^{2}a_{3} - 2a_{3}^{3} + 2a_{1}a_{2}a_{4} - 2a_{3}a_{4}^{2}) \right] \\ = 0. \end{aligned}$$

If $(a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) \neq 0$, then

$$y = \cot \theta_1 = \frac{2 a_1 a_2 a_3 - a_4 + 2a_1^2 a_4}{2 a_0 (a_2 a_3 + a_1 a_4)},$$

$$y = \cot \theta_2 = \frac{-a_2 + 2a_1^2 a_2 + 2a_3^2 - 2a_1 a_3 a_4 + 2a_2 a_4^2}{2 a_0 (a_1 a_2 - a_3 a_4)},$$
 (104)

$$y = \cot \theta_3 = \frac{-a_3 + 2a_1^2 a_3 + 2a_3^2 - 2a_1 a_2 a_4 + 2a_3 a_4^2}{2 a_0 (a_1 a_3 - a_2 a_4)},$$

where θ_1 is a root of equation (24), while θ_2 is a root of equation (24) only in case of $a_2 < a_3$, and θ_3 is a root of equation (24) only in case of $a_2 > a_3$. If $(a_1 a_2 - a_3 a_4) =$ 0, and $a_1 \neq a_4$, then $a_2 \neq a_3$, and

$$y = \cot \theta_4 = \frac{-a_1 + 2a_1^3 + 2a_1a_3^2}{2a_0(a_1^2 + a_3^2)},$$

$$y = \cot \theta_5 = \frac{-a_1 + 2a_1^3 + 2a_1a_3^2}{2a_0(a_1^2 - a_4^2)},$$
(105)

where θ_4 is a root of equation (24), while θ_5 is a root of equation (24) only if $a_2 > a_3$. If $(a_1 a_3 - a_2 a_4) = 0$, and $a_1 \neq a_4$, then $a_2 \neq a_3$, and

$$y = \cot \theta_6 = \frac{-a_1 + 2a_1^3 + 2a_1a_2^2}{2a_0(a_1^2 + a_2^2)}, y = \cot \theta_7 = \frac{-a_1 + 2a_1^3 + 2a_1a_2^2}{2a_0(a_1^2 - a_4^2)},$$
(106)

where θ_6 is a root of equation (24), while θ_7 is a root only if $a_2 < a_3$. If $a_1 = a_4$ and $a_2 = a_3$, then

$$y = \cot \theta_8 = -\frac{a_0 a_1}{2(a_1^2 + a_2^2)},\tag{107}$$

where θ_8 is a root of equation (24).

Therefore,

$$\min \left\{ \sqrt{P(\theta, 0)} + \sqrt{Q(\theta, 0)} \right\}$$

$$= \begin{cases} \min \left\{ \left(\sqrt{P(\theta, 0)} + \sqrt{Q(\theta, 0)} \right) |_{a_2 > a_3} \right\}, & \text{if } a_2 > a_3, \\ \min \left\{ \left(\sqrt{P(\theta, 0)} + \sqrt{Q(\theta, 0)} \right) |_{a_2 < a_3} \right\}, & \text{if } a_2 < a_3, \\ \min \left\{ \left(\sqrt{P(\theta, 0)} + \sqrt{Q(\theta, 0)} \right) |_{a_2 = a_3} \right\}, & \text{if } a_2 = a_3. \end{cases}$$

$$(108)$$

(103)

Here.

$$\min\{\left(\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\right)|_{a_2 > a_3}\} \\ = \begin{cases} \min\{\sqrt{P(\theta_1,0)} + \sqrt{Q(\theta_1,0)}, \sqrt{P(\theta_3,0)} + \sqrt{Q(\theta_3,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, & \text{if } (a_1a_3 - a_2a_4)(a_1a_2 - a_3a_4) \neq 0, \\ \min\{\sqrt{P(\theta_4,0)} + \sqrt{Q(\theta_4,0)}, \sqrt{P(\theta_5,0)} + \sqrt{Q(\theta_5,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, & \text{if } a_1a_2 - a_3a_4 = 0 \text{ and } a_1 \neq a_4, \\ \min\{\sqrt{P(\theta_6,0)} + \sqrt{Q(\theta_6,0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, & \text{if } a_1a_3 - a_2a_4 = 0 \text{ and } a_1 \neq a_4; \\ (109) \end{cases}$$

$$\min\{\left(\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\right)|_{a_{2} < a_{3}}\} = \begin{cases} \min\{\sqrt{P(\theta_{1},0)} + \sqrt{Q(\theta_{1},0)}, \sqrt{P(\theta_{2},0)} + \sqrt{Q(\theta_{2},0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, & \text{if } (a_{1}a_{3} - a_{2}a_{4})(a_{1}a_{2} - a_{3}a_{4}) \neq 0, \\ \min\{\sqrt{P(\theta_{4},0)} + \sqrt{Q(\theta_{4},0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, & \text{if } a_{1}a_{2} - a_{3}a_{4} = 0 \text{ and } a_{1} \neq a_{4}, \\ \min\{\sqrt{P(\theta_{6},0)} + \sqrt{Q(\theta_{6},0)}, \sqrt{P(\theta_{7},0)} + \sqrt{Q(\theta_{7},0)}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, & \text{if } a_{1}a_{3} - a_{2}a_{4} = 0 \text{ and } a_{1} \neq a_{4}; \\ (110) \end{cases}$$

$$\min\{\left(\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\right)|_{a_{2}=a_{3}}\} = \begin{cases} \min\{\sqrt{P(\theta_{1},0)} + \sqrt{Q(\theta_{1},0)}, \sqrt{Q(\theta_{0},0)}|_{a_{2}=a_{3},a_{1}>a_{4}}, \sqrt{P(0,0)} + \sqrt{Q(0,0)}\}, \text{ if } a_{1} > a_{4}, \\ \min\{\sqrt{P(\theta_{1},0)} + \sqrt{Q(\theta_{1},0)}, \sqrt{Q(\overline{\theta}_{0},\pi)}|_{a_{2}=a_{3},a_{1}(111)$$

By (19), (20) and (21), we have

$$\min\{\sqrt{P(\theta,\pi)} + \sqrt{Q(\theta,\pi)}\} = \min\{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\} = \min\{\sqrt{P(\theta,2\pi)} + \sqrt{Q(\theta,2\pi)}\}$$
(112)

Now let us examine the case $\varphi = \frac{\pi}{2}$. In this case, (28) goes to

$$4ya_0^4[1 - 2a_0^2 - 4(a_2a_3 + a_1a_4)^2][a_2^2a_3^2 + 2a_1a_2a_3a_4 - a_4^2(a_2^2 + a_3^2)](1 + y^2)^{-\frac{3}{2}} = 0, (113)$$

which implies that $\theta = \frac{\pi}{2}$. It is shown that $\theta = \frac{\pi}{2}$ is the only possible extreme point of both $\sqrt{P(\theta, \frac{\pi}{2})} + \sqrt{Q(\theta, \frac{\pi}{2})}$ and $\sqrt{P(\theta, \frac{3\pi}{2})} + \sqrt{Q(\theta, \frac{3\pi}{2})}$, and

$$\frac{\sqrt{P(\frac{\pi}{2},\frac{\pi}{2})} + \sqrt{Q(\frac{\pi}{2},\frac{\pi}{2})}}{\sqrt{P(\frac{\pi}{2},\frac{3\pi}{2})} + \sqrt{Q(\frac{\pi}{2},\frac{3\pi}{2})}} = \sqrt{1 - 4(a_2a_3 + a_1a_4)^2 - 4a_0^2a_4^2}.$$
(114)

To sum up,

$$\min\{\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)}\} = \min\{\sqrt{P(0,\varphi)} + \sqrt{Q(0,\varphi)}, \sqrt{P(\frac{\pi}{2},\frac{\pi}{2})} + \sqrt{Q(\frac{\pi}{2},\frac{\pi}{2})}, \\ \min\{\sqrt{P(\theta,0)} + \sqrt{Q(\theta,0)}\}, \\ (\sqrt{P(\theta,\varphi)} + \sqrt{Q(\theta,\varphi)})|_{P(\theta,\varphi)=0}\}.$$
(115)

In brief, we give the analytic expression of the localizable entanglement (LE), the maximum of probability of successfully controlled teleporting an unknown qubit state (14) via every three-qubit state (2) satisfying $a_1a_2a_3a_4 \sin \mu = 0$ and investigate how to achieve it (that is, Charlie finds optimal measurement basis, performs optimal measurements, and communicates the results).

The authors thank Prof. J. I. Cirac for his fruitful discussions and for his hospitality during their stay at Max-Planck-Institut für Quantenoptik. This work was supported by the National Natural Science Foundation of China under Grant No: 10671054, Hebei Natural Science Foundation of China under Grant Nos: A2005000140, 07M006, and the Key Project of Science and Technology Research of Education Ministry of China under Grant No: 207011.

 C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W. K. Wooters, Phys. Rev. Lett. 70, 1895 (1993).

IV. CONCLUSION

14

- [2] M. Fujii, Phys. Rev. A 68, 050302 (2003).
- [3] N. Ba An, Phys. Rev. A 68, 022321 (2003).
- [4] W. P. Bowen, N. Treps, B. C. Buchler, R. Schnabel, T. C. Ralph, Hans-A. Bachor, T. Symul, and P. K. Lam, Phys. Rev. A 67, 032302 (2003).
- [5] T. J. Johnson, S. D. Bartlett, and B. C. Sanders, Phys. Rev. A 66, 042326 (2002).
- [6] L. Vaidman, Phys. Rev. A 49, 1473 (1994).
- [7] S. L. Braunstein and H. J. Kimble, Phys. Rev. Lett. 80, 869 (1998).
- [8] W. Son, J. Lee, M. S. Kim, and Y.-J. Park, Phys. Rev. A 64, 064304 (2001).
- [9] D. Bruß, D. P. DiVincenzo, A. Ekert, C. A. Fuchs, C. Macchiavello, and J. A. Smolin, Phys. Rev. A, 57, 2368 (1998).
- [10] G. Gordon and G. Rigolin, Phys. Rev. A 73, 042309 (2006).
- [11] T. Gao, F. L. Yan, and Z. X. Wang, Quantum Information and Computation, 4, 186 (2004).
- [12] A. Karlsson and M. Bourennane, Phys. Rev. A 58, 4394 (1998).
- [13] C. P. Yang, S. I. Chu, and S. Han, Phys. Rev. A 70, 022329 (2004).
- [14] A. K. Pati, Phys. Rev. A **61**, 022308 (2000).
- [15] P. Agrawal and A. K. Pati, Phys. Lett. A **305**, 12 (2002).
- [16] A. K. Pati and P. Agrawal, J. Opt. B: Quantum Semiclassical Opt. 6, S844 (2004).
- [17] F. L. Yan and D. Wang, Phys. Lett. A **316**, 297 (2003).

- [18] F. G. Deng, C. Y. Li, Y. S. Li, H. Y. Zhou, and Y. Wang, Phys. Rev. A 72, 022338 (2005).
- [19] F. L. Yan, and X. Q. Zhang, The European Physical Journal B, 41 75 (2004).
- [20] D. Bouwmeester, J. W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, Nature (London) **390**, 575 (1997).
- [21] A. Furusawa, J. L. Srensen, S. L. Braunstein, C. A. Fuchs, H. J. Kimble, and E. S. Polzik, Science 282, 706 (1998).
- [22] M. A. Nielsen, E. Knill, and R. Laflamme, Nature (London) **396**, 52 (1998).
- [23] M. Hillery, V. Bužek, and A. Berthiaume, Phys. Rev. A 59, 1829 (1999).
- [24] B. Aoun, M.Tarifi, e-print, quantum-ph/0401076.
- [25] E. Biham, B. Huttner, and T. Mor, Phys. Rev. A 54, 2651 (1996).
- [26] P. D. Townsend, Nature **385**, 47 (1997).
- [27] S. Bose, V. Vedral, and P. L. Knight, Phys. Rev. A 57, 822 (1998).
- [28] T. Gao, Z. Naturforsch. **59a**, 597 (2004).
- [29] F. Verstraete, M. Popp, and J. I. Cirac, Phys. Rev. Lett. 92, 027901 (2004).
- [30] T. Gao, F. L. Yan, and Y. C. Li, e-print, quantumph/0710.1055.
- [31] A. Acín, A. Andrianov, L. Costa, E. Jané, J. I. Latorre, and R. Tarrach, Phys. Rev. Lett. 85, 1560 (2000).