

Labelling Algorithms for Paired-domination Problems in Block and Interval Graphs ^{*}

Lei Chen¹ Changhong Lu^{2,1,†} Zhenbing Zeng¹

¹ Shanghai Key Laboratory of Trustworthy Computing,
East China Normal University, Shanghai, 200062, China

² Department of Mathematics,
East China Normal University, Shanghai, 200062, China

Abstract Let $G = (V, E)$ be a graph without isolated vertices. A set $S \subseteq V$ is a paired-domination set if every vertex in $V - S$ is adjacent to a vertex in S and the subgraph induced by S contains a perfect matching. The paired-domination problem is to determine the paired-domination number, which is the minimum cardinality of a paired-dominating set. Motivated by a mistaken algorithm given by Chen, Kang and Ng [Paired domination on interval and circular-arc graphs, Disc. Appl. Math. 155(2007),2077-2086], we present two linear time algorithms to find a minimum cardinality paired-dominating set in block and interval graphs. In addition, we prove that paired-domination problem is *NP*-complete for bipartite graphs, chordal graphs, even split graphs.

Keywords: Algorithm, Block graph, Interval graph, Paired-domination, NP-complete
2000 Mathematics Subject Classification: 05C69; 05C85; 68R10

^{*}Supported in part by National Natural Science Foundation of China (Nos. 60673048 and 10471044) and National Basic Research Program of China (2003CB318003).

[†]Correspond author. E-mail: chlu@math.ecnu.edu.cn

1 Introduction

Domination and its variations in graphs have been extensively studied, cf. [1, 2]. A set of vertices S is a *dominating set* for a graph $G = (V, E)$ if every vertex in $V - S$ is adjacent to a vertex in S . The *domination problem* is to determine the *domination number* $\gamma(G)$, which is the minimum cardinality of a dominating set for G .

Let $G = (V, E)$ be a simple graph without isolated vertices. For a vertex $v \in V$, the *open neighborhood* of v is defined as $N(v) = \{u \in V \mid uv \in E\}$ and the *closed neighborhood* of v is defined as $N[v] = N(v) \cup \{v\}$. The distance between u and v , denoted by $d_G(u, v)$, is the minimum length of a path between u and v . For a subset S of V , the subgraph of G induced by the vertices in S is denoted by $G[S]$. A *matching* in a graph G is a set of pairwise nonadjacent edges in G . A *perfect matching* M in G is a matching such that every vertex of G is incident to an edge of M . Some other notations and terminology not introduced in here can be found in [9].

A set $S \subseteq V$ is a *paired-dominating set* of G if S is a dominating set of G and the induced subgraph $G[S]$ has a perfect matching. If $e = uv \in M$, where M is a perfect matching of $G[S]$, we say that u and v are *paired in S* . The *paired-domination problem* is to determine the *paired-domination number* $\gamma_p(G)$, which is the minimum cardinality of a paired-dominating set for a graph G . The paired-domination problem was introduced by Haynes and Slater [3]. If we think of each $s \in S \subseteq V$ as the location of a guard capable of protecting each vertex in $N[S]$, then “domination” requires every vertex to be protected, and for paired-domination, we will require the guards’ location to be selected as adjacent pairs of vertices so that each guard is assigned one other and they are designated as backups for each other.

Linear time algorithm for paired domination problem is available for trees [6]; Polynomial time algorithm for paired domination problem is available for circular-arc graphs [5]. Other results on this subject can be found in [4, 8]. Although a linear time algorithm for paired-domination problem on interval graphs was given in [5], it is incorrect. In this paper, we employ the labelling technique to give efficient algorithms for finding a minimum paired-dominating set in block graphs (which contains trees) and interval graphs. In section 2, we begin with presenting a linear time algorithm for paired-domination problem in block graphs and then prove the correctness of the algorithm. Our algorithm can deduce a quite simple algorithm for paired-domination problem in trees. In section 3, we first show that the algorithm in [5] for finding a minimum paired-dominating set in interval graphs is false. Then we give an intuitive algorithm for the paired-domination problem of interval graphs. In [3], authors proved that the paired domination problem is *NP*-complete for undirected graphs. In section 4, we show that it is still *NP*-complete for bipartite graphs, chordal graphs, even split graphs.

2 Algorithm for paired-domination problem in Block graphs

In a graph $G = (V, E)$ with $|V| = n$ and $|E| = m$, a vertex x is a *cut-vertex* if deleting x and all edges incident to it increases the number of connected components. A *block* of G is a maximal connected subgraph of G without a cut-vertex. If G itself is connected and has no cut-vertex, then G is a block. The intersection of two blocks contains at most one vertex, and a vertex is a cut-vertex if and only if it is the intersection of two or more blocks. A *block graph* is a connected graph whose blocks are complete graphs. If every block is K_2 , then it is a tree.

As we know, every block graph not isomorphic to complete graph has at least two *end blocks*, which are blocks with only one cut-vertex. Beginning with an end block and working recursively inward, we can find a vertex ordering v_1, v_2, \dots, v_n in $O(n+m)$ time such that $v_i v_j \in E$ and $v_i v_k \in E$ implies that $v_j v_k \in E$ for $i < j < k \leq n$. Note that if B is an end block with cut-vertex x of block graph G , then the vertices in B is following continually and x is the last vertex of B in the vertex ordering v_1, v_2, \dots, v_n .

Let v_1, v_2, \dots, v_n be the vertex ordering of block graph G such that $v_i v_j \in E$ and $v_i v_k \in E$ implies that $v_j v_k \in E$ for $i < j < k \leq n$. We define the following notations:

1. $F(v_i) = v_j$, $j = \max\{k \mid v_i v_k \in E, i < k\}$. v_j is called the *father* of v_i and v_i is a *child* of v_j . For technical reasons, we say complete graph has an end block and v_n is a cut-vertex. Obviously, v_j must be a cut-vertex in block graphs.
2. $C(v_i) = \{v_j \mid F(v_j) = v_i\}$.
3. $PD(G)$ is a minimum paired-dominating set of G .
4. For any block graph G , we define a rooted tree $T(G)$ about G , whose vertex set is $V(G)$, and uv is an edge of $T(G)$ if and only if $F(u) = v$. The root of $T(G)$ is v_n . Moreover let T_{v_i} is a subtree of $T(G)$ rooted at v_i and every vertex in T_{v_i} except v_i is a descendant of v_i . For a vertex $v_i \in V(G)$, $D_G(v_i)$ denotes the vertex set consisting of the descendants of v_i in $T(G)$ and $D_G[v_i] = D_G(v_i) \cup \{v_i\}$. That is, $D_G[v_i] = V(T_{v_i})$.

In our algorithm, we will use two labels on each vertex u , denoted by $(D(u), L(u))$:

$$D(u) = \begin{cases} 0 & \text{if } u \text{ is not dominated;} \\ 1 & \text{if } u \text{ is dominated.} \end{cases}$$

$$L(u) = \begin{cases} 0 & \text{if } u \text{ is not put into } PD(G); \\ 1 & \text{if } u \text{ is put into } PD(G), \text{ but it has no paired vertex in } PD(G); \\ 2 & \text{if } u \text{ is put into } PD(G), \text{ and it has a paired vertex in } PD(G). \end{cases}$$

Now, we give an algorithm to determine a minimum paired-dominating set in block graphs.

Algorithm MPDB. Find a minimum paired-dominating set of a block graph.

Input. A block graph $G = (V, E)$ with a vertex ordering v_1, v_2, \dots, v_n ($n \geq 2$) such that $v_i v_j \in E$ and $v_i v_k \in E$ implies that $v_j v_k \in E$ for $i < j < k \leq n$. Each vertex v_i has a label $(D(v_i), L(v_i)) = (0, 0)$. $F(v_i) = v_j$ with $j = \max\{k \mid v_k v_i \in E, k > i\}$, $C(v_i) = \{v_j \mid F(v_j) = v_i\}$.

Output. A minimum paired-dominating set PD of G .

Method.

```

For  $i = 1$  to  $n$  do
  If  $(D(v_i) = 0$  and  $i \neq n)$  then
     $L(F(v_i)) = 1$ ;
     $D(u) = 1$  for every vertex  $u \in N[F(v_i)]$ ;
  endif
  If  $(D(v_i) = 1)$  then
    Let  $C'(v_i) = \{w \mid w \in C(v_i) \text{ and } L(w) = 1\}$ ;
     $L(w) = 2$  for every vertex  $w \in C'(v_i)$ ;
    Let  $C''(v_i) = \{w \mid w \in C'(v_i) \text{ and } w \in V(M), M \text{ is a maximum matching in } G[C'(v_i)]\}$ .
    If  $(C'(v_i) - C''(v_i) \neq \emptyset)$ , then
       $L(v_i) = 2$ ;
       $D(u) = 1$  for every vertex  $u \in N[v_i]$ ;
      Take a vertex  $w \in C'(v_i) - C''(v_i)$ , for every vertex  $v \in C'(v_i) - C''(v_i) - \{w\}$ 
       $L(v') = 2$  for some vertex  $v' \in C(v)$  such that  $L(v') = 0$ ;
    endif
  endif
endifor
If  $(D(v_n) = 0$  or  $L(v_n) = 1)$  then
   $L(v_n) = 2$ ;
   $L(w) = 2$  for some vertex  $w \in C(v_n)$  such that  $L(w) = 0$ ;
   $D(v_n) = 1$ ;
endif
Output  $PD = \{v \mid L(v) = 2\}$ 
end

```

Next we will verify the validity of the algorithm MPDB. For a block graph G of order $n \geq 2$. when the algorithm MPDB terminates, any vertex $u \in V(G)$ has a label $D(u) = 1$ and any vertex $v \in PD$ has a label $L(v) = 2$. Hence, the output PD is a paired-dominating set of G . It suffices to prove that PD is a minimum paired-dominating set of G .

S_i is the set of vertices defined by $v \in S_i$ if and only if v has the label $(1, 2)$ when v_i is the considering vertex in the loop of the algorithm MPDB for $i = 1, 2, \dots, n$. In particular, we define S_{n+1} as the set of vertices with label $(1, 2)$ after v_n is considered in the loop of the algorithm MPDB. In order to prove that the output PD is a minimum paired-dominating set of G , we proceed by induction on i and show that, when v_i ($1 \leq i \leq n$) is the considering vertex in the loop of the algorithm MPDB, there is a minimum paired-dominating set S in G such that $S_i \subseteq S$. Obviously, $S_1 = \emptyset$. This is certainly true for $i = 1$. Assume that there is a minimum

paired-dominating set S in G such that $S_i \subseteq S$ ($1 \leq i \leq n$). We show that S_{i+1} holds for a given $i + 1$ by the following lemmas.

Lemma 1 *Let B be an end block of a block graph G . If there is a vertex $u \in V(B)$ such that u is not a cut-vertex with $D(u) = 0$ and $F(u) = v$ ($v \neq u$), then $L(v) = 1$ and $D(w) = 1$ for every vertex $w \in N[v]$. Moreover $PD(G) = PD(G')$, where $G' = G - (D_G(v) - \{u\})$.*

Proof G' has at least two vertices, hence $PD(G')$ is also a paired-dominating set of G . Then $|PD(G)| \leq |PD(G')|$.

Now we show that $|PD(G')| \leq |PD(G)|$. If $v \in PD(G)$ and v is paired with $x \in B$, then $(PD(G) - \{x\}) \cup \{u\}$ is also a paired-domination of G' . If $v \in PD(G)$ and v is paired with $x \notin B$, it is clear that $PD(G)$ is a paired-domination of G' . Assume that $v \notin PD(G)$, then there are two vertices $x, y \in B$ such that $x, y \in PD(G)$. Hence, $(PD(G) - \{x, y\}) \cup \{u, v\}$ is also a paired-domination of G' .

v must be in $PD(G')$ implies that v is also in $PD(G)$. Note that the paired vertex with v may be in B or not. So we set $L(v) = 1$ and $D(w) = 1$ for every vertex $w \in N[v]$. \square

From Lemma 1, when we consider a vertex v_i in an end block whose label is $(0, 0)$ and $i \neq n$, then we will put its father $F(v_i)$ into PD . Since its paired vertex can not be determined at this time, we temporarily label $L(F(v_i)) = 1$. Lemma 1 implies that we can consider a possible smaller block graph G' since $PD(G) = PD(G')$, where $G' = G - (D_G(v) - \{u\})$.

When we consider a vertex v_i such that $D(v_i) = 1$, we will take its child set to see if there is a child must be paired with v_i . After that we will also consider a possible smaller block graph.

Lemma 2 *Let G be a block graph and v_i be a considering vertex with $D(v_i) = 1$ in some step of the loop. Set $C'(v_i) = \{v_j \mid v_j \in C(v_i) \text{ and } L(v_j) = 1\}$, $C''(v_i) = \{w \mid w \in C'(v_i) \text{ and } w \in V(M), M \text{ is a maximum matching in } G[C'(v_i)]\}$ and $S_i(v_j) = S_i \cap D_G(v_j)$ for $j \leq i$. Then, $L(w) = 2$ for every vertex $w \in C'(v_i)$.*

- (1) *If $L(v_i) = 0$ and $C'(v_i) - C''(v_i) = \emptyset$, then $PD(G) = PD(G') \cup S_{i+1}(v_i)$, where $G' = G - D_G[v_i] \neq K_1$ or $G' = G - D_G(v_i) = K_2$.*
- (2) *If $L(v_i) = 1$ and $C'(v_i) - C''(v_i) = \emptyset$, then $PD(G) = PD(G') \cup S_{i+1}(v_i)$, where $G' = G - (D_G(v_i) - \{x\})$ and x is a vertex whose father is v_i and $L(x) = 0$.*
- (3) *If $C'(v_i) - C''(v_i) \neq \emptyset$, then $L(v_i) = 2$ and $D(u) = 1$ for every $u \in N[v_i]$, and for every vertex $v \in C'(v_i) - C''(v_i)$ except one vertex w , $L(v') = 2$ for some vertex $v' \in C(v)$ such that $L(v') = 0$. Moreover, $PD(G) = PD(G') \cup (S_{i+1}(v_i) - S_{i+1}(w) - \{w\})$, where $G' = G - (D_G(v_i) - D_G[w])$.*

Proof $L(w) = 1$ for $w \in C'(v_i)$ implies that w must be put into PD , so $L(w) = 2$ for every vertex $w \in C'(v_i)$ and we will determine their paired vertices in this step.

(1) If v_i is not a cut-vertex in graph G , then $S_{i+1} = S_i$ and we do nothing. Suppose that v_i is a cut-vertex. Obviously, $S_{i+1}(v_i) = S_i(v_i) \cup C'(v_i)$ and $PD(G') \cup S_{i+1}(v_i)$ is a paired-dominating set of G , where $G' = G - D_G[v_i] \neq K_2$ or $G' = G - D_G(v_i) = K_2$. Now we will show that $PD(G') \cup S_{i+1}(v_i)$ is a minimum paired-dominating set of G .

By the hypothesis on induction, there exists a minimum paired-dominating set S of G with $S_i(v_i) \subseteq S$. Since the label of every vertex in $C'(v_i)$ is $(1, 1)$, it must be in a minimum paired-dominating set of G . When v_i is the considering vertex, these vertices in S_i have no influence on the label of remained vertices in possible smaller graph. We can assume that S is a minimum paired-dominating set with $S_i(v_i) \cup C'(v_i) \subseteq S$. We first claim that every vertex in $C'(v_i)$ is paired with another vertex in $C'(v_i)$. Let $C'_p = \{x | x \notin C'(v_i) \text{ and } x \text{ is paired with a vertex } y \text{ in } S \text{ such that } y \in C'(v_i)\}$. Since vertices in S_i have been paired each other, $C'_p \cap S_i = \emptyset$. If $v_i \notin S$, since vertices in $C'(v_i)$ can be paired each other, $S - C'_p$ is a smaller paired-dominating set of G , a contradiction. Then, $v_i \in C'_p$ and we assume that v_i is paired with $x \in C'(v_i)$. Note that $|C'_p| \geq 2$. Let $z \in C'_p$ be paired with $y \in C'(v_i)$ and $xy \in E$. Obviously, $z \in D_G(v_i)$. If $t \in S$ for every $t \in N(v_i)$, then x can be paired with y and $S - \{v_i, z\}$ is a smaller paired-dominating set of G , a contradiction. If there exists some vertex $t \in N(v_i)$ and $t \notin S$, then $S \cup \{t\} - \{z\}$ is also a paired-dominating set of G . Anyway, we get that these vertices in $C'(v_i)$ are paired each other in S .

If $G' = G - D_G(v_i) = K_2$, it is easy to know that $S - S_{i+1}(v_i)$ is a paired-dominating set of $G' = K_2$. Hence, we know that $PD(G') \cup S_{i+1}(v_i) = PD(G)$, where $G' = G - D_G(v_i) = K_2$.

Suppose that $G' = G - D_G[v_i] \neq K_1$. We will show that $S - S_{i+1}(v_i)$ is a paired-dominating set of G' . Hence, $PD(G') \cup S_{i+1}(v_i) = PD(G)$, where $G' = G - D_G[v_i] \neq K_1$.

If $v_i \notin S$, then it is obvious that $S - S_{i+1}(v_i)$ is a paired-dominating set of G' . Assume that $v_i \in S$ and its paired vertex in S is x . If the father of v_i , denoted by $F(v_i)$, is in S , and its paired vertex is not v_i (that is $x \neq F(v_i)$), $S - \{x, v_i\}$ is a smaller paired domination set of G , a contradiction. If $F(v_i)$ is not in S , $S - \{x\} \cup \{F(v_i)\}$ is also a paired-dominating set of G . Without loss of generality, we assume that v_i is paired with $F(v_i)$ in S . If $N(F(v_i)) \subseteq S$, then $S - \{v_i, F(v_i)\}$ is a smaller paired-dominating set of G , a contradiction. Assume that $z \in N(F(v_i))$ is not in S . $S - \{v_i\} \cup \{z\}$ is also a paired-dominating set of G . Hence, we can find a minimum paired domination set S of G such that $S_{i+1}(v_i) \subseteq S$ and $v_i \notin S$. Note that $D(v_i) = 1$. i.e., v_i has been dominated by some vertex in $S_{i+1}(v_i)$. Hence, $S - S_{i+1}(v_i)$ is a paired-dominating set of G' , where $G' = G - D_G[v_i] \neq K_1$.

(2) When v_i is the considering vertex in the loop of the algorithm MPDB, v_i has been labelled by $(1, 1)$. There exists a vertex $x \in C(v_i)$ with label $(1, 0)$ dominated by v_i . In this situation, v_i must be put into PD after $F(v_i)$ is considered in the algorithm MPDB, and the paired vertex of v_i may be a vertex in $C(v_i)$. Similarly, we can prove that those vertices in $C'(v_i)$ are paired each other, hence, the paired vertex of v_i may be a vertex in $C(v_i)$ and its label is $(1, 0)$. Thus, let $G' = G - (D_G(v_i) - \{x\})$, where x is a vertex whose father is v_i and $L(x) = 0$. Using the

same argument, we get that $PD(G) = PD(G') \cup S_{i+1}(v_i)$. The detail is left to readers.

(3) Since $G[C''(v_i)]$ has a perfect matching and it is also a maximum matching in $G[C'(v_i)]$, the set $C'(v_i) - C''(v_i)$ is an independent set. For every vertex $v \in C'(v_i) - C''(v_i)$ except one w , let $v' \in C(v)$ with $L(v') = 0$. (It always can be found since the label of v is (1,1)) and the set of these vertices is denoted by CC , then $|CC| = |C'(v_i)| - |C''(v_i)| - 1$. After v_i was considered in the loop of the algorithm, every vertex in $C'(v_i) \cup CC \cup \{v_i\}$ has the label (1,2). Hence, $S_{i+1} = S_i \cup C'(v_i) \cup CC \cup \{v_i\}$.

Let $G' = G - (D_G(v_i) - D_G[w'])$. It is easy to check that $PD(G') \cup (S_{i+1}(v_i) - \{w\})$ is a paired-dominating set of G . Next we will show that $PD(G') \cup (S_{i+1}(v_i) - \{w\})$ is a minimum paired-dominating set of G . We can similarly find a minimum paired-dominating set S of G such that $S_i \cup C'(v_i) \subseteq S$.

Let $C'_p = \{x | x \in S, x \notin C'(v_i) \text{ and its paired vertex is in } C'(v_i)\}$. Then $|C'_p| \geq |C'(v_i)| - |C''(v_i)|$. If $v_i \notin S$, then the set $S - C'_p \cup CC \cup \{v_i\}$ is a minimum paired-dominating set of G . If $v_i \in S$ and $v_i \in C'_p$, then $S - (C'_p - \{v_i\}) \cup CC$ is also a minimum paired-dominating set of G . If $v_i \in S$ and $v_i \notin C'_p$, we also get that $S - C'_p \cup CC \cup \{w'\}$ is a minimum paired-dominating set of G , where $w' \in C(w)$ has the label (1,0). Anyway, we can assume that these vertices in $C''(v_i)$ are paired in S each other and $\{v_i\} \cup CC \subseteq S$. And for every $v \in C'(v_i) - C''(v_i) - \{w\}$, v is paired in S with $v' \in CC$.

Next we will say that v_i and w can be paired in S . Let $w'(v')$, respectively) is the paired vertex of $w(v_i$, respectively) in S . We assert that there is a vertex $v'' \in N_G(v')$ such that $v'' \notin S$, for, otherwise, $S - \{w', v'\}$ is a smaller paired-dominating set of G . Hence, $S - \{w'\} \cup \{v''\}$ is also a minimum paired-dominating set of G . Thus we can assume w is paired with v_i in S .

Let $S' = S - (S_{i+1}(v_i) - S_{i+1}(w) - \{w\})$. Then S' is a paired-dominating set of G' , where $G' = G - (D_G(v_i) - D_G[w])$. Hence, that is $PD(G') \cup (S_{i+1}(v_i) - S_{i+1}(w) - \{w\})$ is a minimum paired-dominating set of G . In this case, $PD(G) = PD(G') \cup (S_{i+1}(v_i) - S_{i+1}(w) - \{w\})$, where $G' = G - (D_G(v_i) - D_G[w'])$.

The discussion above implies that w, v_i are paired in S , so let $L(v_i) = 2$ and $D(u) = 1$ for every vertex in $N[v_i]$. Since every vertex $v \in C'(v_i) - C''(v_i)$ except one vertex w is paired with $v' \in C(v)$, which is labelled by (1,0), we label v' by (1,2) in this step. \square

Using Lemmas 1 and 2 recursively, we have that $PD(G) = PD(G') \cup S_{n+1}$ when v_n has been considered in the loop of the algorithm, where G' is empty or G' is K_2 . If G' is K_2 , it implies that $D(v_n) = 0$ or $L(v_n) = 1$ after the loop of the algorithm. For the former, v_n need to be dominated and no vertex in $N(v_n)$ is put in PD . For the latter, v_n must be put in to PD and it need a paired vertex. Hence, in the end of the algorithm MPDB, we change the labels of v_n and w to (1,2), where $w \in C(v_n)$ with the label (1,0).

Theorem 3 *Given a vertex ordering, described in the beginning of this section, of a block graph G , the algorithm MPDB can produce a minimum paired-dominating set of G in $O(m+n)$ time,*

where $m = |E(G)|$ and $n = |V(G)|$.

Proof From discussion above, the algorithm MPDB can produce a minimum paired-dominating set of a block graph G . Furthermore, every vertex in $V(G)$ and every edge in $E(G)$ are scanned in a constant number. Although we must find a maximum matching in $G[C'(v_i)]$, it can be done in linear time, because $G[C'(v_i)]$ is disjoint union of some clique. If every clique has even number of vertices, then $G[C'(v_i)]$ has a perfect matching, otherwise it has not a perfect matching. \square

In [6], a linear time algorithm was given to determine a minimum paired-dominating set in trees. Since block graphs contain trees, we can also use the algorithm MPDB to produce a paired-dominating set in trees. The only difference is that $G[C'(v_i)]$ can not have a perfect matching in tree. Here, we give a very simple algorithm for tree which can be deduced from MPDB at once.

Algorithm MPDT. Find a minimum paired-dominating set of a tree.

Input. A tree $T = (V, E)$ with a vertex ordering v_1, v_2, \dots, v_n such that $d_T(v_i, v_n) < d_T(v_j, v_n)$ implies that $i < j$. Each vertex v_i has a label $(D(v_i), L(v_i)) = (0, 0)$. $F(v_i) = v_j$ with $v_i v_j \in E$ and $j > i$; $C(v_i) = \{v_j \mid F(v_j) = v_i\}$.

Output. A minimum paired-dominating set PD of T .

Method.

```

For  $i = 1$  to  $n$  do
  If  $(D(v_i) = 0$  and  $i \neq n)$  then
     $L(F(v_i)) = 1$ ;
     $D(u) = 1$  for every vertex  $u \in N[F(v_i)]$ ;
  endif
  If  $(D(v_i) = 1)$  then
    Let  $C'(v_i) = \{w \mid w \in C(v_i) \text{ and } L(w) = 1\}$ ;
    If  $C'(v_i) \neq \emptyset$  then
       $L(v_i) = 2$ ;
       $D(u) = 1$  for every vertex  $u \in N[v_i]$ ;
       $L(w) = 2$  for every vertex  $w \in C'(v_i)$ ;
      Take a vertex  $w \in C'(v_i)$ , for every vertex  $v \in C'(v_i) - \{w\}$ 
       $L(v') = 2$  for some vertex  $v' \in C(v)$  such that  $L(v') = 0$ ;
    endif
  endif
endif
endfor
If  $(D(v_n) = 0$  or  $L(v_n) = 1)$  then
   $L(v_n) = 2$ ;
   $L(w) = 2$  for some vertex  $w \in C(v_n)$  such that  $L(w) = 0$ ;

```



```

     $D(v_n) = 1;$ 
endif
Output  $PD = \{v | L(v) = 2\}$ 
end

```

Corollary 4 *Algorithm MPDT can produce a minimum paired-dominating set of a tree T in $O(m + n)$, where $m = |E(T)|$ and $n = |V(T)|$.*

3 Algorithm for paired-domination problem in Interval graph

An *interval representation* of a graph is a family of intervals assigned to the vertices so that vertices are adjacent if and only if the corresponding intervals intersect. A graph having such a representation is an *interval graph*. Booth and Lueker [7] gave an $O(|V(G)| + |E(G)|)$ -time algorithm for recognizing an interval graph and constructing an interval representation using *PQ-tree*. In [5], a linear algorithm was given to produce a minimum paired-dominating set of an interval graph. But this algorithm is incorrect. Next, we will introduce this algorithm and given a counterexample.

In [5], it is assumed that the input graph is given by an interval representation I that is a set of n sorted intervals labelled by $1, 2, \dots, n$ in increasing order of their right endpoints. The left endpoint of interval i is denoted by a_i and the right endpoint by b_i ($1 < a_i \leq b_i \leq 2n$). Define the following notations.

(1) For a set S of intervals, the largest left(right) endpoint of the intervals in S is denoted by $\max a(S)$ ($\max b(S)$); the interval in S with the largest right endpoint is denoted by $last(S)$. Let $\max a(S) = 0$ ($\max b(S) = 0$) if S is empty. For endpoint e , use $IFB(e)$ to denote the set of all intervals whose right endpoint are less than e . For any interval j , let l_j be the interval such that intervals l_j and j have nonempty intersection and $a(l_j)$ is minimum.

(2) For $j \in \{1, 2, \dots, n\}$, define $V_j = \{i : i \in \{1, 2, \dots, n\} \text{ and } a_i \leq b_j\}$. Let $PD(j) = \{S : S \subseteq V_j, S \text{ is a paired-dominating set of } G[V_j]\}$. Let $PD(i, j) = \{S : S \subseteq V_j, S \text{ is a paired-dominating set of } G[V_j], i, j \in S \text{ and } i, j \text{ are paired in } S\}$. Let $MPD(j) = \text{Min}(PD(j))$, $MPD(i, j) = \text{Min}(PD(i, j))$. The left endpoint sets $A_i = \{a_j : b_{i-1} < a_j < b_j\}$ for $i \in I$, where $b_0 = 0$.

Introduce two intervals $n + 1$ and $n + 2$ with $a_{n+1} = 2n + 1$, $a_{n+2} = 2n + 2$, $b_{n+1} = 2n + 3$, and $b_{n+2} = 2n + 4$. Let I_p be the set of intervals obtained by augmenting I with the two intervals $n + 1$ and $n + 2$.

Algorithm MPD


Input: A set I_p of sorted intervals.

Output: A minimum cardinality paired-dominating set of G with interval representation I_p .

1. Find $\max a(IFB(a_j))$ for all $j \in I_p$.

2. Find l_j for all $j \in I_p$.
 3. Scan the endpoints of I_p to find the left endpoint sets $A_i = \{a_j : b_{i-1} < a_j < b_i\}$ for $i \in I$, where $b_0 = 0$.
 4. $MPD(0) = \emptyset$.
 5. for $j = 1$ to $n + 2$ do
 6. Find the left endpoint set A_k containing $\max a(IFB(\min(a_j, a_{l_j})))$.
 7. Let b_k be the right endpoint of the interval k associated with the left endpoint set A_k .
 8. $MPD(j) = \{l_j, j\} \cup MPD(k)$.
 9. end for
- Output $MPD(n + 2)$.

It is obvious that $MPD(n + 2) - \{n + 1, n + 2\}$ got from the algorithm MPD is a paired-dominating set of G . The following counterexample implies that $MPD(n + 2) - \{n + 1, n + 2\}$ may be not a minimum paired-dominating set of G .



The figure above is a counterexample. The left figure is an interval representation of the graph in the right figure. The number is ordered by the right endpoint of intervals. The parameters used in MPD are as follows:

i	a_i	b_i	$\max a(IFB(a_i))$	l_i	A_i
1	0	3	0	2	$\{1, 2\}$
2	1	4	0	1	\emptyset
3	2	6	0	1	$\{5\}$
4	7	9	2	5	$\{7, 8\}$
5	5	10	1	3	\emptyset
6	8	11	2	5	\emptyset
7	13	15	8	8	\emptyset
8	14	16	8	7	\emptyset

Execute algorithm MPD as follows:

- $j = 1$, $\max a(IFB(\min(a_1, a_2))) = 0$, $k = 0$, $MPD(1) = \{1, 2\} \cup MPD(0) = \{1, 2\}$;
 $j = 2$, $\max a(IFB(\min(a_1, a_2))) = 0$, $k = 0$, $MPD(2) = \{1, 2\} \cup MPD(0) = \{1, 2\}$;
 $j = 3$, $\max a(IFB(\min(a_3, a_1))) = 0$, $k = 0$, $MPD(3) = \{1, 3\} \cup MPD(0) = \{1, 3\}$;
 $j = 4$, $\max a(IFB(\min(a_4, a_5))) = 1$, $k = 1$, $MPD(4) = \{4, 5\} \cup MPD(1) = \{4, 5, 1, 2\}$;
 $j = 5$, $\max a(IFB(\min(a_5, a_3))) = 0$, $k = 0$, $MPD(5) = \{3, 5\} \cup MPD(0) = \{3, 5\}$;
 $j = 6$, $\max a(IFB(\min(a_6, a_5))) = 1$, $k = 1$, $MPD(6) = \{5, 6\} \cup MPD(1) = \{5, 6, 1, 2\}$;
 $j = 7$, $\max a(IFB(\min(a_7, a_8))) = 8$, $k = 4$, $MPD(7) = \{1, 2\} \cup MPD(4) = \{7, 8, 4, 5, 1, 2\}$;

$j = 8$, $\max a(IFB(\min(a_7, a_8))) = 8$, $k = 4$, $MPD(8) = \{1, 2\} \cup MPD(4) = \{7, 8, 4, 5, 1, 2\}$;

Hence the result set of the algorithm MPD is $\{4, 5, 1, 2\}$. But it is easy to see that $\{3, 5\}$ is a minimum paired-dominating set of this graph. Note that Lemma 2.3 and Lemma 2.5 in [5] are not right. The detail is left to readers.

Next, we employ the labelling technique to give a linear algorithms for finding a minimum paired-dominating set of an interval graph. Let $G = (V, E)$ be an interval graph and its interval representation is I . For every vertex $u_i \in V$, I_i is the corresponding interval, and let a_i (b_i , respectively) denote the left endpoint (right endpoint, respectively) of interval I_i . We order the vertices of G by u_1, u_2, \dots, u_n in increasing order of their left endpoints. Then we have following two observations.

Observation 5 u_1, u_2, \dots, u_n is a ordering of an interval graph G by the increasing order of their left endpoints. If $u_i u_j \in E$ with $j < i$, then $u_j u_k \in E$ for every $j + 1 \leq k \leq i$.

Let $V_i = \{u_j \mid j \leq i\}$ and $G[V_i]$ be an induced subgraph of G . It is obvious that $G[V_n] = G$. Let $F(u_i) = u_j$, where $j = \min\{k \mid u_k u_i \in E \text{ and } k < i\}$. In particular, $F(u_1) = u_1$. Let $w(u_i) = u_j$, where $j = \max\{k \mid u_k u_i \notin E \text{ and } k < i\}$. In particular, If $w(u_i)$ does not exist, we assume that $w(u_i) = u_0$ ($u_0 \notin V$). PD_i denotes a minimum paired-dominating set of $G[V_i]$. In this paper, we only consider connected interval graph.

Observation 6 If G is a connected interval graph, then $G[V_i]$ is also connected.

Lemma 7 If $F(u_i) \neq u_i$ and $F(F(u_i)) = F(u_i)$, then $PD_i = \{u_i, F(u_i)\}$.

Proof Since $F(F(u_i)) = F(u_i)$, hence $F(u_i) = u_1$. By Observation 5, for every $1 < j \leq i$, $u_1 u_j \in E$. So $\{u_i, u_1\} = \{u_i, F(u_i)\}$ is a minimum paired-dominating set of $G[V_i]$. Hence, $PD_i = \{u_i, F(u_i)\}$. \square

Lemma 8 $|PD_{i+1}| \geq |PD_i|$ for $2 \leq i \leq n - 1$.

Proof If $u_{i+1} \notin PD_{i+1}$, then PD_{i+1} is also a paired-dominating set of $G[V_i]$. So $|PD_{i+1}| \geq |PD_i|$. If $u_{i+1} \in PD_{i+1}$ and u_k, u_{i+1} are paired in PD_{i+1} . We consider two cases.

Case 1: $k = i$.

That is u_i, u_{i+1} are paired in PD_{i+1} . If $N_{G[V_i]}(u_i) \subseteq PD_{i+1}$, then $PD_{i+1} - \{u_i, u_{i+1}\}$ is a paired-dominating set of $G[V_i]$. So $|PD_{i+1}| > |PD_{i+1}| - 2 \geq |PD_i|$. If there is a vertex $w \in N_{G[V_i]}(u_i)$ with $w \notin PD_{i+1}$, then $PD_{i+1} - \{u_{i+1}\} \cup \{w\}$ is also a paired-dominating set of $G[V_i]$. We also get $|PD_{i+1}| \geq |PD_i|$.

Case 2: $k < i$.

If $u_i \notin PD_{i+1}$, then $PD_{i+1} - \{u_{i+1}\} \cup \{u_i\}$ is a paired-dominating set of $G[V_i]$, so $|PD_{i+1}| \geq |PD_i|$. If $u_i \in PD_{i+1}$ and u_i is paired with u_l , then $u_l u_k \in E$ and $PD_{i+1} - \{u_i, u_{i+1}\}$ is a paired-dominating set of $G[V_i]$, so $|PD_{i+1}| > |PD_{i+1}| - 2 \geq |PD_i|$. \square

Lemma 9 Let $F(u_i) = u_k$ and $F(u_k) = u_j$ with $j < k < i$.

- (1) $PD_i = PD_l \cup \{u_j, u_k\}$ if $w(u_j) = u_l$ with $l \geq 2$.
- (2) $PD_i = \{u_1, u_2, u_j, u_k\}$ if $w(u_j) = u_1$.
- (3) $PD_i = \{u_j, u_k\}$ if $w(u_j) = u_0$.

Proof (1) It is obvious that $PD_l \cup \{u_j, u_k\}$ is a paired-dominating set of $G[V_i]$. It is sufficient to prove $|PD_i| \geq |PD_l| + 2$. Since $F(u_i) = u_k$, there must exist a vertex $u_{i_1} \in PD_i$ with $k \leq i_1 \leq i$, which dominates u_i . We may assume that u_{i_1} is the last vertex in PD_i which dominates u_i and u_{i_1} is paired with u_{k_1} . It is obvious that $k_1 \geq j$. Let $l' = \min\{a, b\}$, where $w(u_{i_1}) = u_a$ and $w(u_{k_1}) = u_b$. Since $u_l u_{k_1} \notin E$ and $u_l u_{i_1} \notin E$ (otherwise $u_l u_j \in E$, a contradiction), so $l' \geq l \geq 2$. Let u_c is the last vertex in $PD_i - \{u_{i_1}, u_{k_1}\}$. If $c \geq l'$, then $PD_i - \{u_{i_1}, u_{k_1}\}$ is a paired-dominating set of $G[V_c]$. So $|PD_i| - 2 \geq |PD_c|$. On the other hand, since $c \geq l' \geq l$, by Lemma 8, $|PD_c| \geq |PD_l|$. Then $|PD_i| \geq |PD_l| + 2$. If $c < l'$, then $PD_i - \{u_{i_1}, u_{k_1}\}$ is a paired-domination set of $G[V_{l'}]$. Since $l' \geq l$, $|PD_i| - 2 \geq |PD_{l'}| \geq |PD_l|$. So $|PD_i| \geq |PD_l| + 2$. Thus $PD_i = PD_l \cup \{u_j, u_k\}$.

(2) Note that $u_1 u_2 \in E$ and $j \geq 3$ in this situation. So it is easy to know that $PD_i = \{u_1, u_2, u_j, u_k\}$ if $w(u_j) = u_1$.

(3) It is obvious. \square

Now we give an intuitive algorithm for determining a minimum paired-dominating set in interval graphs.

Algorithm MPDI. Find a minimum paired-dominating set of an interval graph.

Input. An interval graph $G = (V, E)$ with a vertex ordering u_1, u_2, \dots, u_n ordered by the increasing order of left endpoints, in which each vertex u_i has a label $D(u_i) = 0$. Let $F(u_i) = u_j$ ($F(u_1) = u_1$) such that $j = \min\{k \mid u_k u_i \in E \text{ and } k < i\}$.

Output. A minimum paired-dominating set PD of G .

Method.

$PD = \emptyset$;

For $i = n$ to 1 do

 If $(D(u_i) = 0)$ then

 If $(F(u_i) \neq u_i \text{ and } F(F(u_i)) \neq F(u_i))$ then

$PD = PD \cup \{F(u_i), F(F(u_i))\}$;

$D(u) = 1$ for every vertex $u \in N[F(u_i)]$;

$D(w) = 1$ for every vertex $w \in N[F(F(u_i))]$;

 else if $(F(u_i) \neq u_i)$ then

$PD = PD \cup \{u_i, F(u_i)\}$;

$D(u) = 1$ for every vertex $u \in N[F(u_i)]$;

 else

$PD = PD \cup \{u_i, u_2\}$;

```

         $D(u_i) = 1;$ 
    endif
endif
endif
endfor

```

Theorem 10 *Given a vertex ordering ordered by the increasing order of left endpoints, the algorithm MPDI can produce a minimum paired-dominating set of G in $O(m + n)$, where $m = |E(G)|$ and $n = |V(G)|$.*

Proof By Lemmas 7 and 9, we know that the algorithm MPDI can produce a minimum paired-dominating set of an interval graph G . Since each vertex and edge are scanned in a constant number, hence the algorithm MPDI can finish in $O(m + n)$, where $m = |E(G)|$ and $n = |V(G)|$. \square

4 NP-completeness of paired-domination problem

A graph is *chordal* if every cycle of length greater than three has a chord, i.e. an edge jointing two nonconsecutive vertices in the cycle. A graph is *split* if its vertex set can be partitioned into a stable set and a clique. Obviously, block graphs, interval graphs and split graphs are three subclasses of chordal graphs. This section establishes *NP*-complete results for the paired-domination problem in bipartite graphs and chordal graphs. The transformation is from the vertex cover problem, which is known to be *NP*-complete. The *vertex cover problem* is for a given nontrivial graph and a positive integer k to answer if there is a vertex set of size at most k such that each edge of the graph has at least one end vertex in this set.

Theorem 11 *Paired-domination problem is NP-complete for bipartite graphs.*

Proof For a bipartite graph $G = (V, E)$, a positive even integer k , and an arbitrary subset $S \subseteq V$ with $|S| \leq k$, it is easy to verify in polynomial time whether S is a paired-dominating set of G . Hence, paired-domination problem is in *NP*.

We construct a reduction from the vertex cover problem. Given a nontrivial graph $G = (V, E)$, where $V = \{v_1, v_2, \dots, v_n\}$ and $E = \{e_1, e_2, \dots, e_m\}$, Let $V_i = \{v_1^i, v_2^i, \dots, v_n^i\}$ and $E_i = \{e_1^i, e_2^i, \dots, e_m^i\}$ ($i = 1, 2$). Construct the graph $G' = (V', E')$ with vertex set $V' = V_1 \cup V_2 \cup E_1 \cup E_2$, and edge set $E' = \{uv \mid u \in V_1 \text{ and } v \in V_2\} \cup \{v_j^i e_k^i \mid i = 1, 2 \text{ and } v_j \text{ is incident to } e_k \text{ in } G\}$. Note that G' is a bipartite graph.

Next, we will show that G has a vertex cover of size at most k if and only if G' has a paired-dominating set of size at most $2k$. Let $VC = \{v_{i_1}, v_{i_2}, \dots, v_{i_k}\}$ be a vertex cover of G . Then it is obvious that $\{v_{i_1}^1, v_{i_2}^1, \dots, v_{i_k}^1\} \cup \{v_{i_1}^2, v_{i_2}^2, \dots, v_{i_k}^2\}$ is a paired-dominating set of G' and its size is $2k$. For the converse, let PD be a paired-dominating set of G' with $|PD| \leq 2k$. Obviously,

we can assume that $k \leq n - 1$ since $V = \{v_1, v_2, \dots, v_n\}$ is a vertex cover of G . If $PD \cap E_1 \neq \emptyset$, without loss of generality, we assume that $e_1^1 \in PD$ and its paired vertex in PD is v_1^1 . Since $k \leq n - 1$, there exists a vertex $v_j^2 \notin PD$. Hence, $PD \cup \{v_j^2\} - \{e_1^1\}$ is also a paired-dominating set of G' . Then we may assume that $PD \cap (E_1 \cup E_2) = \emptyset$. Suppose that $VC_1 = PD \cap V_1$, and note that $|VC_1| \leq k$. Let $VC = \{v_i | v_i^1 \in VC_1\}$ and VC is a vertex cover of G such that $|VC| \leq k$.

Finally, one can construct G' from G in polynomial time. This implies that paired-domination problem is *NP*-complete for bipartite graphs. \square

Theorem 12 *Paired-domination problem is NP-complete for chordal graphs.*

Proof We still construct a reduction from the vertex cover problem. Given a nontrivial graph $G = (V, E)$, where $V = \{v_1, v_2, \dots, v_n\}$ and $E = \{e_1, e_2, \dots, e_m\}$, Let $V_i = \{v_1^i, v_2^i, \dots, v_n^i\}$ ($i = 1, 2$) and $E_i = \{e_1^i, e_2^i, \dots, e_m^i\}$ ($i = 1, 2$). Construct the graph $G' = (V', E')$ with vertex set $V' = V_1 \cup V_2 \cup E_1 \cup E_2$, and edge set $E' = \{uv | u \in V_1 \cup V_2 \text{ and } u \neq v\} \cup \{v_j^i e_k^i | i = 1, 2 \text{ and } v_j \text{ is incident to } e_k \text{ in } G\}$. Note that G' is a chordal graph.

It is straightforward to show that G has a vertex cover of size at most k if and only if G' has a paired-dominating set of size at most $2k$. The proof is almost similar with that of Theorem 11. In here, we can also assume that $k \leq n - 1$ and $PD \cap (E_1 \cup E_2) = \emptyset$. So, either $VC_1 = PD \cap V_1$ or $VC_2 = PD \cap V_2$ has size at most k . The detail is left to readers. \square

Note that G' in Theorem 12 is also a split graph. Hence we get a stronger result as follows.

Corollary 13 *Paired-domination problem is NP-complete for split graphs.*

References

- [1] T. W. Haynes, S. T. Hedetniemi and P. J. Slater (eds), *Fundamentals of Domination in Graphs*, New York, Marcel Dekker 1998.
- [2] T. W. Haynes, S. T. Hedetniemi and P. J. Slater (eds), *Domination in Graphs: Advanced Topics*, New York, Marcel Dekker 1998.
- [3] T. W. Haynes and P. J. Slater, Paired-domination in graphs, *Networks* 32(1998), 199-206.
- [4] M. A. Henning, Graphs with large paired-domination number, *J. Comb. Optim.* 13(2007), 61-78.
- [5] T. C. E. Chen, L. Y. Kang and C. T. Ng, Paired domination on interval and circular-arc graphs, *Discrete Appl. Math.* 155(2007), 2077-2086.
- [6] H. Qiao, L. Y. Kang, M. Caedei and D. Z. Du, Paired-domination of trees, *J. Global Optim.* 25(2003), 43-54.
- [7] K. S. Booth and G. S. Lueker, Testing for consecutive ones property, interval graphs and graph planarity using *PQ*-tree algorithms, *J. Comput. Syst. Sci.* 13(1976), 335-379.

- [8] P. D. Sylvain Gravier and M. A. Henning, Paired-domination in generalized claw-free graphs, *J. Comb. Optim.* 14(2007), 1-7.
- [9] D. B. West, *Introduction to Graph Theory*, 2nd ed., Prentice Hall, Inc., NJ, 2001.