

Simulation of high-spin Heisenberg chains in coupled cavities

Jaeyoon Cho,¹ Dimitris G. Angelakis,^{2,3} and Sougato Bose¹

¹*Department of Physics and Astronomy, University College London, Gower St., London WC1E 6BT, UK*

²*Centre for Quantum Technologies, National University of Singapore, 2 Science Drive 3, Singapore 117542*

³*Science Department, Technical University of Crete, Chania, Crete, Greece, 73100*

(Dated: February 6, 2020)

We propose a scheme to realize the antiferromagnetic Heisenberg model of any spin in an array of coupled cavities. Our scheme is based on a fixed number of atoms confined in each cavity and collectively applied constant laser fields, and is in a regime where both atomic and cavity excitations are suppressed. It is shown that as well as optically controlling the effective spin-chain Hamiltonian, it is also possible to engineer the magnitude of the spin. Our scheme would open up an unprecedented way to simulate otherwise intractable high-spin problems in many-body physics.

The quantum spin chain has played a crucial role as a basic model accounting for the magnetic and thermodynamic natures of many-body systems. Despite extensive investigations, however, many aspects of the model are still largely unexplored both analytically and numerically, especially for the cases of higher spins. The main difficulty in the numerical treatment originates from the fact that the Hilbert-space dimension blows up exponentially as the number of spins increases. As Feynman first noted [1], this difficulty would be overcome in terms of quantum simulation based on precisely controlled quantum systems. Realization of quantum simulation, expected in a near future, will mark a milestone towards the realization of sophisticated quantum computation.

In the context of quantum information processing, a qubit is identical to an $s = \frac{1}{2}$ spin, and in a few implementations, such as the arrays of Josephson junctions [2] or quantum dots [3], the spin-chain Hamiltonian naturally emerges from the spin-like coupling between qubits, albeit with limited control of the coupling constants. On the other hand, in optical lattices, perturbative evolution with respect to the Mott-insulator state can be described by an effective spin-chain Hamiltonian [4, 5]. This approach has its own merit in that the spin-coupling constants can be optically controlled to a great extent. An alternative approach, recently under active investigation, is to use the array of coupled cavities, which are ideally suited to addressing individual spins [6, 7, 8, 9, 10, 11, 12, 13, 14]. In this approach, a spin is represented by either polaritons or hyperfine ground levels. The former, proposed in Refs. [8] and [14], allows a stronger spin-spin coupling than the latter, but lacks the optical control of the coupling. On the other hand, the latter, proposed in Ref. [9], retains the optical controllability, but relies on rapid switching of optical pulses and the consequent Trotter expansion, which unavoidably involves additional errors and makes error-free implementation more difficult. More importantly, the question of simulating chains of higher spins, which may have a completely different phase diagram, remains open. In some sense, these are more important to simulate because unlike spin- $\frac{1}{2}$ chains, they do not have exact




FIG. 1: Involved atomic levels and transitions. Both transitions $|a\rangle \leftrightarrow |e\rangle$ and $|b\rangle \leftrightarrow |e\rangle$ are coupled to the same cavity mode with coupling rates g_1 and g_2 and with detunings Δ_1 and Δ_2 , respectively. Two laser fields with Rabi frequency Ω_1 and Ω_2 are also applied with detunings Δ_1 and Δ_2 , respectively, and the transition between ground levels $|a\rangle$ and $|b\rangle$ is driven with Rabi frequency $\omega/2$ by Raman lasers. γ denotes the atomic spontaneous decay rate.

analytical solutions for a wide range of parameters including the XXX case, except for special kinds of models [15]. Additionally, going to higher spins should make perturbative spin-wave theory more accurate, whose predictions can be tested.

In this paper, we propose a scheme to realize the anisotropic (XXZ) or isotropic (XXX) antiferromagnetic Heisenberg spin-chain model of any spin in an array of coupled cavities. Our scheme is experimentally feasible in that simply applying a small number of constant laser fields suffices for our purpose. If the number of lasers is increased, the individual constants of the spin-chain Hamiltonian are controlled more flexibly. Most of all, a strong advantage of our scheme is that the magnitude of the spin itself can be engineered arbitrarily. This advantage contrasts with all the earlier schemes mentioned above including those for optical lattices, in which the spin is fixed in nature mostly as $s = \frac{1}{2}$ ($s = 1$ in Ref. [5]). $s > \frac{1}{2}$ spin chains exhibit fascinating physics that $s = \frac{1}{2}$ spin chains can not have. A well-known example is Haldane's conjecture that antiferromagnetic Heisenberg integral-spin chains have a unique disordered ground state with a finite excitation gap, whereas half-

integral-spin chains are gapless [16, 17]. Our scheme could be used to prepare a ground state, for example, through an adiabatic evolution, and measure its excitation gap and spin correlation functions [5]. Spin chains also play an important role as a quantum channel for short-distance quantum communication [18, 19]. The property of an $s = 1$ antiferromagnetic spin chain as a quantum channel strongly depends on its phase [20]. In some phases, it provides an efficient channel, outperforming that of a ferromagnetic chain. It has been also shown that a ground state with an excitation gap, as is the case for the spin-1 chain, can serve as a more efficient quantum channel [21]. Although the idea of communicating using spin chains is ultimately meant for solid-state applications, our system can serve as a preliminary test for comparing and contrasting the performance of various spin- s chains.

We use two ground levels of a three-level atom to represent an $s = \frac{1}{2}$ spin (in a rotated basis, as will be seen later). We start by recalling that in terms of two states $|\downarrow\rangle$ and $|\uparrow\rangle$ of one atom, the $s = \frac{1}{2}$ spin is described in terms of operators $s^Z = \frac{1}{2}(|\uparrow\rangle\langle\uparrow| - |\downarrow\rangle\langle\downarrow|)$, $s^+ = |\uparrow\rangle\langle\downarrow|$, and $s^- = |\downarrow\rangle\langle\uparrow|$. Our starting point is an observation that if there are M identical atoms, one can straightforwardly define total spin operator $S^Z = \sum_{j=1}^M s_j^Z$ with $S^\pm = \sum_{j=1}^M s_j^\pm$ (j is the index for the atoms), by which the atoms represent $S = \frac{M}{2}$, $S = \frac{M}{2} - 1$, and so on. Keeping this in mind, let us consider a coupled array of N identical cavities, each of which contains M identical single atoms. For simplicity, we assume the periodic boundary condition. Let us first consider a simple case, as depicted in Fig. 1. Let us denote by $|\psi\rangle_{jk}$ the state $|\psi\rangle$ of the k th atom in the j th cavity. In the rotating frame, the Hamiltonian reads

$$\begin{aligned}
H = & \sum_{j=1}^N [e^{i\Delta_1 t} \Omega_1 \Lambda_j^{eb} + e^{i\Delta_2 t} \Omega_2 \Lambda_j^{ea} + h.c.] \\
& + \sum_{j=1}^N [(e^{i\Delta_1 t} g_1 \Lambda_j^{ea} + e^{i\Delta_2 t} g_2 \Lambda_j^{eb}) a_j + h.c.] \quad (1) \\
& + \sum_{j=1}^N \left[\frac{\omega}{2} (\Lambda_j^{ab} + \Lambda_j^{ba}) + J (a_j^\dagger a_{j+1} + a_j a_{j+1}^\dagger) \right],
\end{aligned}$$

where $\Lambda_j^{xy} = \sum_{k=1}^M (|x\rangle\langle y|)_{jk}$ ($x, y = a, b, e$), a_j is the annihilation operator for the j th cavity mode, Δ_j is the corresponding detuning, Ω_j and $\frac{\omega}{2}$ are the corresponding Rabi frequencies of the classical fields, g_j is the corresponding atom-cavity coupling rate, and J is the inter-cavity hopping rate of photons. Both the transitions are coupled to the same cavity mode. The transition between $|a\rangle$ and $|b\rangle$ is induced by two-photon Raman transition using far-detuned lasers. For now, we ignore the spontaneous decay rate γ of the atom.

Before we proceed, it is instructive to write down our

parameter regime:

$$\frac{g_1^2}{\Delta_1} = \frac{g_2^2}{\Delta_2}, \quad (2)$$

$$\Delta_j, \Delta_1 - \Delta_2 \gg \sqrt{\frac{M}{2}} g_j \gg J \gtrsim |\Omega_j|, \quad (3)$$

$$M \frac{g_1^2}{\Delta_1} \sim \left| M \frac{g_1^2}{\Delta_1} \pm \omega \right| \sim |\omega| \gg 2J. \quad (4)$$

The condition (2) can be fulfilled with conventionally used alkali-metal atoms, such as rubidium and caesium. For example, one may choose ground hyperfine levels $|F = 1, m_F = -1\rangle$ and $|F = 2, m_F = -1\rangle$ of a ^{87}Rb atom to represent $|a\rangle$ and $|b\rangle$, respectively, and use σ^+ -polarized light, for which $g_1 > g_2$. The detuning Δ_j is then comparable to the hyperfine splitting between the two levels. Although there are multiple excited levels, their contributions can be summed up and denoted by single parameters in what follows. The other conditions (3) and (4) can be satisfied simultaneously when $\sqrt{\frac{M}{2}} g_j / \Delta_j \gg J / \sqrt{\frac{M}{2}} g_j$. For example, our scheme works well in case $\Delta_j / 1000 \sim \sqrt{\frac{M}{2}} g_j / 100 \sim J$, which is allowed by strong atom-cavity coupling.

Our regime is chosen so that the excitation of the atom or the cavity photon is suppressed (condition (3)), while the communication between atoms is mediated by virtual cavity photons. The first step is to perform the adiabatic elimination [22]. For this, we diagonalize the cavity-hopping terms by employing a Fourier-transformed basis as $a_j = \frac{1}{\sqrt{N}} \sum_{k=1}^N F_{jk} b_k$, where $F_{jk} = \exp(-i \frac{2\pi}{N} jk)$ and $\sum_{j=1}^N F_{jk} F_{jl}^* = N \delta_{kl}$. The diagonalized form reads $J \sum_j (a_j^\dagger a_{j+1} + a_j a_{j+1}^\dagger) = \sum_j \nu_j b_j^\dagger b_j$, where $\nu_j = 2J \cos(\frac{2\pi}{N} j)$. Substituting this into the Hamiltonian and performing the adiabatic elimination, the effective Hamiltonian is given by

$$\begin{aligned}
H = & - \sum_{j=1}^N \frac{g_1^2}{\Delta_1} (\Lambda_j^{aa} + \Lambda_j^{bb}) a_j^\dagger a_j \\
& + \sum_{j=1}^N \left[\frac{\omega}{2} (\Lambda_j^{ab} + \Lambda_j^{ba}) + \nu_j b_j^\dagger b_j \right] \quad (5) \\
& - \sum_{j=1}^N [(\mu_1 \Lambda_j^{ba} + \mu_2 \Lambda_j^{ab}) a_j + h.c.],
\end{aligned}$$

where $\mu_j = \frac{g_j \Omega_j^*}{\Delta_j}$.

Now let us introduce spin operators. We use a rotated basis

$$\left\{ |\uparrow\rangle = \frac{1}{\sqrt{2}}(|a\rangle + |b\rangle), |\downarrow\rangle = \frac{1}{\sqrt{2}}(|a\rangle - |b\rangle) \right\} \quad (6)$$

to represent an $s = \frac{1}{2}$ spin. Note that these are the eigenstates of $\frac{\omega}{2}(|a\rangle\langle b| + |b\rangle\langle a|)$. The underlying idea is

to apply the Raman lasers with Rabi frequency $\frac{\omega}{2}$ constantly, introducing a fixed amount of energy splitting $|\omega|$ between the two spin states. The total spin is then defined in terms of the operators

$$S_j^Z = \sum_{k=1}^M s_{jk}^Z \text{ and } S_j^\pm = \sum_{k=1}^M s_{jk}^\pm, \quad (7)$$

where $s_{jk}^Z = \frac{1}{2}(|\uparrow\rangle\langle\uparrow| - |\downarrow\rangle\langle\downarrow|)_{jk}$, $s_{jk}^+ = (|\uparrow\rangle\langle\downarrow|)_{jk}$, and $s_{jk}^- = (|\downarrow\rangle\langle\uparrow|)_{jk}$. The total spin is given by $S_j^2 = (S_j^Z)^2 + \frac{1}{2}(S_j^+ S_j^- + S_j^- S_j^+)$. If M is even (odd), the atoms represent integral (half-integral) spins up to $\frac{M}{2}$. The atomic operators are now written as $\Lambda_j^{\downarrow\downarrow} = \sum_k s_{jk}^- s_{jk}^+ = \frac{M}{2} - S_j^Z$, $\Lambda_j^{\uparrow\uparrow} = \sum_k s_{jk}^+ s_{jk}^- = \frac{M}{2} + S_j^Z$, $\Lambda_j^{\uparrow\downarrow} = S_j^+$, and $\Lambda_j^{\downarrow\uparrow} = S_j^-$. Substituting these operators, the Hamiltonian in the rotating frame reads

$$H = - \sum_{j,k=1}^N \left[\left\{ e^{i\lambda t} \mu_{12}^+ S_j^Z + e^{i(\lambda+\omega)t} \frac{\mu_{12}^-}{2} S_j^+ - e^{i(\lambda-\omega)t} \frac{\mu_{12}^-}{2} S_j^- \right\} \frac{e^{-i\nu_k t}}{\sqrt{N}} F_{jk} b_k + h.c. \right], \quad (8)$$

where $\lambda = M \frac{g_1^2}{\Delta_1}$ and $\mu_{12}^\pm = \mu_1 \pm \mu_2$. Note that in view of conditions (3) and (4), the effective Rabi frequency $\left| \sqrt{\frac{M}{2}} \mu_{12}^\pm \right|$ is much smaller than λ , $|\lambda \pm \omega|$, and $|\omega|$. This allows us to make use of the adiabatic elimination once more. Taking the subspace without real photons, the effective Hamiltonian is given by

$$H = \sum_{j,k,l=1}^N \left[\left\{ \frac{|\mu_{12}^+|^2}{\lambda - \nu_k} S_j^Z S_l^Z + \frac{|\mu_{12}^-|^2}{4(\lambda + \omega - \nu_k)} S_j^+ S_l^- + \frac{|\mu_{12}^-|^2}{4(\lambda - \omega - \nu_k)} S_j^- S_l^+ \right\} \frac{F_{jk} F_{lk}^*}{N} \right]. \quad (9)$$

In view of the condition (4) and $\nu_k = J(F_{1k} + F_{1k}^*)$, we expand the first term as

$$\begin{aligned} \frac{F_{jk} F_{lk}^*}{\lambda - \nu_k} &\simeq \frac{F_{jk} F_{lk}^*}{\lambda} \left(1 + \frac{J}{\lambda} F_{1k} + \frac{J}{\lambda} F_{1k}^* \right) \\ &= \frac{1}{\lambda} (F_{jk} F_{lk}^* + \frac{J}{\lambda} F_{(j+1)k} F_{lk}^* + \frac{J}{\lambda} F_{jk} F_{(l+1)k}^*), \end{aligned} \quad (10)$$

and similarly for the others. This approximation is valid when $J \gg \left| \sqrt{\frac{M}{2}} \mu_{12}^\pm \right|$, since otherwise the next higher order correction to the effective Hamiltonian would be comparable to the terms with J . This condition is met by the condition (3). Since $\sum_{k=1}^N F_{jk} F_{lk}^* = N \delta_{jl}$, the first term in Eq. (10) leads to the single-ion anisotropy $(S_j^Z)^2$, while the other two terms lead to spin-spin coupling $S_j^Z S_{j+1}^Z$ between adjacent sites. Putting every summation in the Hamiltonian together, we end up with our




FIG. 2: Additional lasers to get full control of the spin-chain Hamiltonian. These lasers are applied in addition to the set up of Fig. 1.

final spin-chain Hamiltonian

$$H = \sum_{j=1}^N [A(S_j)^2 + B(S_j^Z)^2 + CS_j^Z] + \sum_{j=1}^N [D(S_j^X S_{j+1}^X + S_j^Y S_{j+1}^Y) + ES_j^Z S_{j+1}^Z], \quad (11)$$

where $A = \frac{\lambda}{\lambda^2 - \omega^2} \frac{|\mu_{12}^-|^2}{2}$, $B = \frac{|\mu_{12}^+|^2}{\lambda} - \frac{\lambda}{\lambda^2 - \omega^2} \frac{|\mu_{12}^-|^2}{2}$, $C = -\frac{\omega}{\lambda^2 - \omega^2} \frac{|\mu_{12}^-|^2}{2}$, $D = \frac{J}{2} (|\frac{\mu_{12}^-}{\lambda + \omega}|^2 + |\frac{\mu_{12}^-}{\lambda - \omega}|^2)$, $E = 2J |\frac{\mu_{12}^+}{\lambda}|^2$. This Hamiltonian already covers a wide range of parametric regimes for the antiferromagnetic ($D, E > 0$) Heisenberg spin-chain model, although individual control of the parameters is limited owing to their mutual dependency. Interestingly, the Hamiltonian also contains the single-ion anisotropy $(S_j^Z)^2$, which is of essential importance in high-spin cases [23], whereas it is merely a meaningless constant in the spin- $\frac{1}{2}$ case.

Full control of the individual parameters is allowed by bringing in more lasers, shown in FIG. 2, in addition to the set up of FIG. 1. The classical fields with Rabi frequency Ω_3 and Ω_4 , which are applied with an additional detuning $\delta \sim \lambda$, make a similar contribution to the effective Hamiltonian as those with Ω_1 and Ω_2 . This can be reflected in Hamiltonian (8) by adding the same terms with λ and μ_{12}^\pm replaced by $\lambda - \delta$ and μ_{34}^\pm , respectively, where $\mu_{34}^\pm = \mu_3 \pm \mu_4$, $\mu_3 = \frac{g_1 \Omega_3^*}{2} (\frac{1}{\Delta_1} + \frac{1}{\Delta_1 + \delta})$, and $\mu_4 = \frac{g_2 \Omega_4^*}{2} (\frac{1}{\Delta_2} + \frac{1}{\Delta_2 + \delta})$. The Stark shift $-\mu_z |b\rangle\langle b|$ induced by another far-detuned laser field (using a different level and polarization) results in adding $\sum_{j=1}^N \frac{\mu_z}{2} (e^{i\omega t} S_j^+ + e^{-i\omega t} S_j^-)$ in Hamiltonian (8). Remind that in our previous derivation, we have adjusted $\{0, \lambda, \lambda \pm \omega\}$ so that they are distinct in frequency with the similar frequency spacing (condition (4)), thereby causing each summation in the Hamiltonian (8) to contribute independently to the constants in the final Hamiltonian (11). We adjust ω , λ , $\lambda \pm \omega$, $\lambda - \delta$, and $\lambda - \delta \pm \omega$ in the same spirit. For the ease of presentation, let us take a particular situation where $\omega > 0$, $\lambda = 3\omega$, and $\lambda - \delta = -6\omega$, although this is not a necessary condi-

tion. We then obtain the same Hamiltonian (11) with parameters given by $A = \frac{1}{\lambda}(\frac{9}{16}|\mu_{12}^-|^2 - \frac{9}{35}|\mu_{34}^-|^2)$, $B = \frac{1}{\lambda}(|\mu_{12}^+|^2 - \frac{9}{16}|\mu_{12}^-|^2 - \frac{1}{2}|\mu_{34}^+|^2 + \frac{9}{35}|\mu_{34}^-|^2)$, $C = \frac{1}{\lambda}(\frac{3}{2}|\mu_z|^2 - \frac{3}{16}|\mu_{12}^-|^2 - \frac{3}{70}|\mu_{34}^-|^2)$, $D = \frac{J}{\lambda^2}(\frac{45}{32}|\mu_{12}^-|^2 + \frac{333}{1225}|\mu_{34}^-|^2)$, and $E = \frac{J}{\lambda^2}(2|\mu_{12}^+|^2 + \frac{1}{2}|\mu_{34}^+|^2)$. Note that C is determined independently thanks to the term $|\mu_z|^2/\lambda$, while other terms are also determined freely. Hence, this parameter set covers any anisotropic or isotropic Heisenberg spin chains, with the single-ion anisotropy turned on or off.

Although atomic excitation is heavily suppressed, the main source of decoherence in our system is the spontaneous decay of atoms. In relation to the effective spin chain, the atomic spontaneous decay results in depolarization of the spins. This effect can be accounted for by considering a conditional Hamiltonian $H_C = H - \frac{i}{2} \sum_{j=1}^N (\gamma'_A \Lambda_j^{aa} + \gamma'_B \Lambda_j^{bb})$, where the effective decay rates are approximately given by $\gamma'_A = \gamma \left(\frac{|\Omega_1|^2}{\Delta_1^2} + \frac{|\Omega_3|^2}{(\Delta_1 + \delta)^2} \right)$ and $\gamma'_B = \gamma \left(\frac{|\Omega_2|^2}{\Delta_2^2} + \frac{|\Omega_4|^2}{(\Delta_2 + \delta)^2} \right)$, assuming other lasers are sufficiently detuned and thus make a negligible contribution to the decay. In particular, if Ω_j s are chosen in such a way that the two contributions are balanced, i.e., $\gamma'_A = \gamma'_B = \gamma'$, the depolarization is nearly independent of the spin state. In this case, the conditional Hamiltonian is approximately given by $H_C = H - iNM\frac{\gamma'}{2}$. Consequently, the state of the system at time t may be written as $\rho(t) = e^{-NM\gamma't} \rho_Q(t) + (1 - e^{-NM\gamma't}) \rho_M$, where $\rho_Q(t)$ is the desired quantum state evolved by the spin-chain Hamiltonian and ρ_M is the fully mixed state. This property is useful for testing condensed-matter theories (for example, by adiabatically preparing a ground state [5]), since even under depolarization, the quantum nature retained in the coherent portion $\rho_Q(t)$ could be observed over a time scale $\sim 1/NM\gamma'$. One requirement is that the spin-spin coupling rate multiplied by $(M/2)^2$ should be much larger than the global decoherence rate. Reminding that the coupling rate is given by $\sim 2J|\mu_j|^2/\lambda^2 \sim (J/M)(\Omega_j/\sqrt{M/2g_j})^2$, we require $\gamma \ll \frac{J}{2N} \left(\frac{\Delta_j}{\sqrt{M/2g_j}} \right)^2$. Since $\Delta_j \gg \sqrt{M/2g_j}$ from condition (3), this requirement can be met for a moderate N if $J \gtrsim \gamma$ is satisfied along with our previous assumption of strong atom-cavity coupling. Note, however, that testing Haldane's conjecture for higher-spin chains is more demanding, since the lowest excitation gap is expected, from its asymptotic behavior, to decrease rapidly with increasing M , while the spin correlation length increases rapidly [16]. There are several micro-cavity models under active development which are expected to fall into our regime of strong atom-cavity coupling, although achieving them is still experimentally challenging [24, 25, 26, 27]. Another experimental challenge is to have a fixed (and known) number of atoms in a cavity [28]. Although this is difficult to realize experimentally, it is unavoidable so as to simulate higher spins.

The above mentioned micro-cavity models are expected to be suitable for this purpose, in virtue of the progress in the micro-fabrication techniques.

This work was supported by the Korea Research Foundation Grant (KRF-2007-357-C00016) funded by the Korean Government (MOEHRD). SB thanks the Engineering and Physical Sciences Research Council (EPSRC) UK for an Advanced Research Fellowship and for support through the Quantum Information Processing IRC (GR/S82176/01) and the Royal Society and the Wolfson Foundation.

-
- [1] R. P. Feynman, *Int. J. Theor. Phys.* **21**, 467 (1982).
 - [2] C. Bruder, R. Fazio, and G. Schön, *Phys. Rev. B* **47**, 342 (1993).
 - [3] D. Loss and D. P. DiVincenzo, *Phys. Rev. A* **57**, 120 (1998).
 - [4] L.-M. Duan, E. Delmer, and M. D. Lukin, *Phys. Rev. Lett.* **91**, 090402 (2003).
 - [5] J. J. Garcia-Ripoll, M. A. Martin-Delgado, and J. I. Cirac, *Phys. Rev. Lett.* **93**, 250405 (2004).
 - [6] M. J. Hartmann, F. G. S. L. Brandão, and M. B. Plenio, *Nature Phys.* **2**, 849 (2006).
 - [7] A. D. Greentree, C. Tahan, J. H. Cole, and L. C. L. Hollenberg, *Nature Phys.* **2**, 856 (2006).
 - [8] D. G. Angelakis, M. F. Santos, and S. Bose, *Phys. Rev. A* **76**, 031805(R) (2007).
 - [9] M. J. Hartmann, F. G. S. L. Brandão, and M. B. Plenio, *Phys. Rev. Lett.* **99**, 160501 (2007).
 - [10] D. Rossini and R. Fazio, *Phys. Rev. Lett.* **99**, 186401 (2007).
 - [11] M. Paternostro, G. S. Agarwal, and M. S. Kim, arXiv:0707.0846.
 - [12] J. Cho, D. G. Angelakis, and S. Bose, arXiv:0712.2413.
 - [13] Y. Li, M. X. Huo, Z. Song, and C. P. Sun, arXiv:0802.0079.
 - [14] A. Kay and D. G. Angelakis, arXiv:0802.0488.
 - [15] I. Affleck, *J. Phys.: Condens. Matter.* **1**, 3047 (1989).
 - [16] F. D. M. Haldane, *Phys. Lett.* **93A**, 464 (1983).
 - [17] F. D. M. Haldane, *Phys. Rev. Lett.* **50**, 1153 (1983).
 - [18] S. Bose, *Phys. Rev. Lett.* **91**, 207901 (2003).
 - [19] A. Bayat and S. Bose, arXiv:0706.4176.
 - [20] O. Romero-Isart, K. Eckert, and A. Sanpera, *Phys. Rev. A* **75**, 050303(R) (2007).
 - [21] M. J. Hartmann, M. E. Reuter, and M. B. Plenio, *New J. Phys.* **8**, 94 (2006).
 - [22] D. F. V. James and J. Jerke, *Can. J. Phys.* **85**, 625 (2007).
 - [23] R. Botet, R. Jullien, and M. Kolb, *Phys. Rev. B* **28**, 3914 (1983).
 - [24] S. M. Spillane *et al.*, *Phys. Rev. A* **71**, 013810 (2005).
 - [25] J. Majer *et al.*, *Nature* **449**, 443 (2007).
 - [26] M. A. Sillanpää, J. I. Park, and R. W. Simmonds, *Nature* **449**, 438 (2007).
 - [27] T. Aoki *et al.*, *Nature* **443**, 671 (2006).
 - [28] One possibility is using NV centers in diamond where single atom implantation is close to be possible through the isotropic control of nitrogen [J. Meijer *et al.*, *Appl. Phys. A* **83**, 321 (2006)].