

Moufang symmetry I. Generalized Lie and Maurer-Cartan equations

Eugen Paal

Abstract

The continuous Moufang loops are characterized as the algebraic systems where the associativity law is perturbed minimally. The minimal perturbation of associativity is characterized by the first-order partial differential equations, which in a natural way generalize the Lie and Maurer-Cartan equations from the theory of Lie groups.

1 Introduction and outline of the paper

In 1972, Sabinin stated [15, 16] equivalence of algebraic loops and homogeneous spaces. Following his construction, one may characterize loops as *groups with hidden associativity*. Due to perturbation of associativity, the particular loops are too difficult to describe. In such a situation, the Moufang loops are distinguished by their *minimal* perturbation of associativity. Thus, one may expect that a Moufang loop represents a minimally perturbed symmetry that is concisely referred as the *Moufang symmetry*.

In this paper, the minimal perturbation of associativity in the continuous Mofang loops is characterized by the first-order partial differential equations, which in a natural way generalize the Lie and Maurer-Cartan equations from the theory of Lie groups.

2 Moufang loops

A *Moufang loop* [9] (see also [4, 3, 12]) is a set G with a binary operation (multiplication) $\cdot : G \times G \rightarrow G$, denoted also by juxtaposition, so that the following three axioms are satisfied.

- 1) In the equation $gh = k$, the knowledge of any two of $g, h, k \in G$ specifies the third one *uniquely*.
- 2) There is a distinguished element $e \in G$ with the property $eg = ge = g$ for all $g \in G$.
- 3) The *Moufang identities* hold in G ,

$$g(h \cdot gk) = (gh \cdot g)k, \quad (kg \cdot h)g = k(g \cdot hg), \quad (gh)(kg) = g(hk \cdot g) \quad (2.1)$$

It must be noted that the Moufang identities (2.1) are equivalent (see e.g [4, 3]), which means that each of the above identities may be considered as a defining identity of the Moufang loop.

Recall that a set with a binary operation is called a *groupoid*. A groupoid G with axiom 1) is called a *quasigroup*. If axioms 1) and 2) are satisfied, the groupoid (quasigroup) G is called a *loop*. The element e in axiom 2) is called the *unit* (element) of the (Moufang) loop G .

In a (Moufang) loop, the multiplication need not be neither associative nor commutative. The *associativity* and *commutativity* laws read, respectively,

$$g \cdot hk = gh \cdot k, \quad gh = hg, \quad \forall g, h, k \in G$$

The most familiar kind of loops are the ones with the *associative* law, and these are called *groups*. A loop G is called *commutative* if the commutativity law holds in G , and (only) the commutative associative loops are said to be *Abelian*.

A characteristic property of the Moufang loops is their *diassociativity*: in a Moufang loop every two elements generate an associative subloop (group) [9]. In particular, from this it follows that

$$g \cdot gh = g^2h, \quad hg \cdot g = hg^2, \quad gh \cdot g = g \cdot hg, \quad \forall g, h \in G$$

Here, the first two identities are called the left and right *alternativity*, respectively, and the third one is said to be *flexibility*. Note that these identities follow from the Moufang identities. Due to flexibility, the Moufang identity can be rewritten in a nice symmetric form

$$gh \cdot kg = g \cdot hk \cdot g, \quad \forall g, h, k \in G$$

The unique solution of the equation $xg = e$ ($gx = e$) is called the left (right) *inverse* element of $g \in G$ and is denoted as g_R^{-1} (g_L^{-1}). It also follows from the diassociativity of the Moufang loop that

$$g_R^{-1} = g_L^{-1} \doteq g^{-1}, \quad g^{-1} \cdot gh = hg \cdot g^{-1} = h, \quad (g^{-1})^{-1} = g, \quad (gh)^{-1} = h^{-1}g^{-1}$$

for all $g, h \in G$.

The *smallest* non-associative Moufang loop was described by Chein and Pflugfelder in [6], its order is 12. One can find an elaborated list of the low-order non-associative Moufang loops in [5].

3 Analytic Moufang loops and tangent Malcev algebras

A Moufang loop G is said to be *analytic* [8] if G is a finite dimensional real, analytic manifold so that both the Moufang loop operation $G \times G \rightarrow G: (g, h) \mapsto gh$ and the inversion map $G \rightarrow G: g \mapsto g^{-1}$ are analytic ones. The dimension of G will be denoted as $\dim G \doteq r$.

In this paper, the global aspects are not considered and all considerations are local, i.e. in a fixed chart U_e of the unit $e \in U_e \subset G$ where all subsequent constructions are expected to be well defined. The triple (G, e, U_e) is called a local analytic Moufang loop. By keeping this in mind we often call G the *local* analytic Moufang loop as well.

The local coordinates of $g \in G$ are denoted (in selected fixed chart $U_e \subset G$ of the unit $e \in G$) by g^1, \dots, g^r so that the local coordinates of the unit e are supposed to be zero: $e^i = 0$, $i = 1, \dots, r$. One has the evident initial conditions

$$(ge)^i = (eg)^i = g^i, \quad i = 1, \dots, r$$

As in the case of the Lie groups [13], we can use the Taylor expansions

$$\begin{aligned} (gh)^i &= h^i + u_j^i(h)g^j + \frac{1}{2!}u_{jk}^i(h)g^j h^k \dots \\ &= g^i + v_j^i(g)h^j + \frac{1}{2!}v_{jk}^i(g)g^j h^k \dots \\ &= g^i + h^i + a_{jk}^i g^j h^k + b_{jkl}^i g^j g^k h^l + d_{jkl}^i g^j h^k h^l + \dots \end{aligned}$$

to introduce the *auxiliary functions* $u_j^i, v_j^i, u_{jk}^i, v_{jk}^i$ and the *structure constants*

$$c_{jk}^i \doteq a_{jk}^i - a_{kj}^i = -c_{kj}^i$$

Due to such a parametrization, one has the initial conditions

$$u_j^i(e) = \delta_j^i = v_j^i(e), \quad u_{jk}^i(e) = 0 = v_{jk}^i(e),$$

and symmetry with respect to the lower indices

$$u_{jk}^i = u_{kj}^i, \quad v_{jk}^i = v_{kj}^i$$

Also, it follows from axiom 1) of the Moufang loop that

$$\det[u_j^i] \neq 0 \neq \det[v_j^i]$$

The *tangent algebra* of G can be defined similarly to the tangent algebra of the Lie group [13]. Geometrically, this algebra is the tangent space $T_e(G)$ of G at e . The product of $x, y \in T_e(G)$ will be denoted by $[x, y] \in T_e(G)$. In coordinate form,

$$[x, y]^i \doteq c_{jk}^i x^j y^k = -[y, x]^i, \quad i = 1, \dots, r$$

The tangent algebra of G will be denoted by $\Gamma_G \doteq (T_e(G), [\cdot, \cdot])$. The latter need not be a Lie algebra. In other words, there may exist a triple $x, y, z \in T_e(G)$, such that the Jacobi identity fails,

$$J(x, y, z) \doteq [x, [y, z]] + [y, [z, x]] + [z, [x, y]] \neq 0$$

Instead, for all $x, y, z \in T_e(G)$, one has [8] a more general identity

$$[[x, y], [z, x]] + [[[x, y], z], x] + [[[y, z], x], x] + [[[z, x], x], y] = 0$$

called the *Malcev identity*. The tangent algebra Γ_G is hence said to be the *Malcev algebra*. The Malcev identity concisely reads [14]

$$[J(x, y, x), x] = J(x, y, [x, z])$$

from which it can be easily seen that every Lie algebra is a Malcev algebra as well. It has been shown in [7] that every finite-dimensional real Malcev algebra is the tangent algebra of some local analytic Moufang loop.

The *smallest* non-associative real analytic Moufang loop and its Malcev algebra were described by Akiwis in [1], their dimension is 4.

4 Associators

In a (Moufang) loop, due to non-associativity, the elements $g \cdot hk$ and $gh \cdot k$ need not coincide. The non-associativity of G can be measured by the formal non-invariant functions

$$A^i(g, h, k) \doteq (g \cdot hk)^i - (gh \cdot k)^i, \quad i = 1, \dots, r$$

which are called *associators* of G . One has the evident initial conditions

$$A^i(e, h, k) = A^i(g, e, k) = A^i(g, h, e) = 0, \quad i = 1, \dots, r$$

The associators A^i are considered as generating functions in the following sense. At first, define the *first-order* associators l_j^i, r_j^i, m_j^i by

$$\begin{aligned} A^i(g, h, k) &\doteq l_j^i(h, k)g^j + O(g^2) \\ &\doteq r_j^i(g, h)k^j + O(k^2) \\ &\doteq m_j^i(g, k)h^j + O(h^2) \end{aligned}$$

These can be easily calculated and the result reads

$$\begin{aligned} l_j^i(g, h) &= -u_j^s(g) \frac{\partial (gh)^i}{\partial g^s} + u_j^i(gh) \\ r_j^i(g, h) &= -v_j^i(gh) + v_j^s(h) \frac{\partial (gh)^i}{\partial h^s} \\ m_j^i(g, h) &= -v_j^s(g) \frac{\partial (gh)^i}{\partial g^s} + u_j^s(h) \frac{\partial (gh)^i}{\partial h^s} \end{aligned}$$

Next one can check the initial conditions

$$l_j^i(e, g) = r_j^i(e, g) = m_j^i(e, g) = 0 = l_j^i(g, e) = r_j^i(g, e) = m_j^i(g, e)$$

and define the *second-order* associators $l_{jk}^i, \tilde{l}_{jk}^i, m_{jk}^i, \tilde{m}_{jk}^i, r_{jk}^i, \tilde{r}_{jk}^i$ by

$$\begin{aligned} l_j^i(g, h) &\doteq l_{jk}^i(h)g^k + O(g^2) \\ &\doteq \tilde{l}_{jk}^i(g)h^k + O(h^2) \\ r_j^i(g, h) &\doteq r_{jk}^i(h)g^k + O(g^2) \end{aligned}$$

$$\begin{aligned}
&\doteq \tilde{r}_{jk}^i(g)h^k + O(h^2) \\
m_j^i(g, h) &\doteq \tilde{m}_{jk}^i(h)g^k + O(g^2) \\
&\doteq m_{jk}^i(g)h^k + O(h^2)
\end{aligned}$$

By calculating, the result reads

$$\begin{aligned}
l_{jk}^i(g) &= \tilde{m}_{kj}^i(g) = -u_{kj}^i(g) - a_{jk}^s u_s^i(g) + u_k^s(g) \frac{\partial u_j^i(g)}{\partial g^s} \\
r_{jk}^i(g) &= \tilde{l}_{kj}^i(g) = v_j^s(g) \frac{\partial u_k^i(g)}{\partial g^s} - u_k^s(g) \frac{\partial v_j^i(g)}{\partial g^s} \\
m_{jk}^i(g) &= \tilde{r}_{kj}^i(g) = v_{jk}^i(g) + a_{jk}^s v_s^i(g) - v_j^s(g) \frac{\partial v_k^i(g)}{\partial g^s}
\end{aligned}$$

Finally one has the initial conditions

$$l_{jk}^i(e) = r_{jk}^i(e) = m_{jk}^i(e) = 0 = \tilde{l}_{jk}^i(e) = \tilde{r}_{jk}^i(e) = \tilde{m}_{jk}^i(e)$$

and define the *third-order* associators $l_{jkl}^i, \tilde{l}_{jkl}^i, m_{jkl}^i, \tilde{m}_{jkl}^i, r_{jkl}^i, \tilde{r}_{jkl}^i$ by

$$\begin{aligned}
l_{jk}^i(g) &\doteq l_{jkl}^i g^l + O(g^2), & \tilde{l}_{jk}^i(g) &\doteq \tilde{l}_{jkl}^i g^l + O(g^2) \\
r_{jk}^i(g) &\doteq r_{jkl}^i g^l + O(g^2), & \tilde{r}_{jk}^i(g) &\doteq \tilde{r}_{jkl}^i g^l + O(g^2) \\
m_{jk}^i(g) &\doteq m_{jkl}^i g^l + O(g^2), & \tilde{m}_{jk}^i(g) &\doteq \tilde{m}_{jkl}^i g^l + O(g^2)
\end{aligned}$$

By calculating, one gets

$$l_{jkl}^i = m_{kjl}^i = r_{ljk}^i = \tilde{l}_{jlk}^i = \tilde{r}_{lkj}^i = \tilde{m}_{kjl}^i = a_{js}^i a_{kl}^s - a_{jk}^s a_{sl}^i + 2(d_{jkl}^i - b_{jkl}^i)$$

Thus, all third-order associators can be obtained from l_{jkl}^i via permutations of the lower indices. One can check the *Akivis identity* [2]

$$\begin{aligned}
l_{jkl}^i + r_{jkl}^i + m_{jkl}^i - \tilde{l}_{jkl}^i - \tilde{r}_{jkl}^i - \tilde{m}_{jkl}^i &= l_{jkl}^i + l_{klj}^i + l_{ljk}^i - l_{jlk}^i - l_{lkj}^i - l_{kjl}^i \\
&= c_{js}^i c_{kl}^s + c_{ks}^i c_{lj}^s + c_{ls}^i c_{jk}^s
\end{aligned}$$

For $x, y, z \in T_e(G)$ define [2] their trilinear product $(x, y, z) \in T_e(G)$ by

$$(x, y, z)^i \doteq l_{jkl}^i x^j y^k z^l, \quad i = 1 \dots, r$$

Then the Akivis identity reads [2]

$$J(x, y, z) = (x, y, z) + (y, z, x) + (z, x, y) - (x, z, y) - (z, y, x) - (y, x, z)$$

5 Generalized Lie equations

Differentiate the Moufang identities (2.1) with respect to g^j at $g = e$. Then, suitable re-denoting variables we obtain for the first-order associators the constraints

$$l_j^i(g, h) = -m_j^i(g, h), \quad r_j^i(g, h) = -m_j^i(g, h), \quad l_j^i(g, h) = r_j^i(g, h)$$

Thus, the Moufang identities give rise to the first-order constraints

$$l_j^i(g, h) = r_j^i(g, h) = -m_j^i(g, h) \tag{5.1}$$

The latter in a natural way generalize the Lie equations. If G is associative, then one has the Lie equations (see e.g [13]):

$$l_j^i(g, h) = r_j^i(g, h) = -m_j^i(g, h) = 0$$

By comparing the latter with (5.1) one can say that in the variety of continuous non-associative loops the local analytic Moufang loops have the property that their associativity is perturbed *minimally*. Constraints (5.1) are hence called the *first-order minimality conditions*.

Define the new auxiliary functions w_j^i by

$$u_j^i(g) + v_j^i(g) + w_j^i(g) = 0 \quad (5.2)$$

Then constraints (5.1) can be rewritten as the differential equations of the 1st order, called the *generalized Lie equations*

$$\begin{aligned} w_j^s(g) \frac{\partial(gh)^i}{\partial g^s} + u_j^s(h) \frac{\partial(gh)^i}{\partial h^s} + u_j^i(gh) &= 0 \\ v_j^s(g) \frac{\partial(gh)^i}{\partial g^s} + w_j^s(h) \frac{\partial(gh)^i}{\partial h^s} + v_j^i(gh) &= 0 \\ u_j^s(g) \frac{\partial(gh)^i}{\partial g^s} + v_j^s(h) \frac{\partial(gh)^i}{\partial h^s} + w_j^i(gh) &= 0 \end{aligned}$$

In view of (5.2) these equations are linearly dependent.

6 Generalized Maurer-Cartan equations

Differentiate constraints (5.1) with respect to g^k and h^k at $g = e = h$, respectively. Then, re-denoting variables one obtains the second-order constraints

$$l_{jk}^i(g) = r_{jk}^i(g) = m_{jk}^i(g) = -m_{kj}^i(g) \quad (6.1)$$

Again, if G is associative, i.e a local Lie group, we have

$$l_{jk}^i(g) = r_{jk}^i(g) = m_{jk}^i(g) = -m_{kj}^i(g) = 0$$

Constraints (6.1) are called the *second-order minimality conditions* for G . Lets consider the latter more closely.

It follows from the skew-symmetry $l_{jk}^i = -l_{kj}^i$ and $m_{jk}^i = -m_{kj}^i$, respectively, that

$$2u_{jk}^i = u_k^s \frac{\partial u_j^i}{\partial g^s} + u_j^s \frac{\partial u_k^i}{\partial g^s} - (a_{jk}^i + a_{kj}^i) u_j^s, \quad 2v_{jk}^i = v_k^s \frac{\partial v_j^i}{\partial g^s} + v_j^s \frac{\partial v_k^i}{\partial g^s} - (a_{jk}^i + a_{kj}^i) v_j^s$$

By using u_{jk}^i and v_{jk}^i one can see that

$$u_k^s \frac{\partial u_j^i}{\partial g^s} - u_j^s \frac{\partial u_k^i}{\partial g^s} = c_{jk}^s u_s^i + 2l_{jk}^i, \quad v_k^s \frac{\partial v_j^i}{\partial g^s} - v_j^s \frac{\partial v_k^i}{\partial g^s} = c_{kj}^s v_s^i + 2m_{jk}^i$$

Now use $l_{jk}^i = r_{jk}^i$, $m_{jk}^i = -r_{kj}^i$ and r_{jk}^i to obtain the differential equations for the first order auxiliary functions u_j^i and v_j^i ,

$$u_k^s \frac{\partial u_j^i}{\partial g^s} - u_j^s \frac{\partial u_k^i}{\partial g^s} = c_{jk}^s u_s^i + 2 \left(v_j^s \frac{\partial u_k^i}{\partial g^s} - u_k^s \frac{\partial v_j^i}{\partial g^s} \right) \quad (6.2a)$$

$$v_k^s \frac{\partial v_j^i}{\partial g^s} - v_j^s \frac{\partial v_k^i}{\partial g^s} = c_{kj}^s v_s^i + 2 \left(u_j^s \frac{\partial v_k^i}{\partial g^s} - v_k^s \frac{\partial u_j^i}{\partial g^s} \right) \quad (6.2b)$$

called the *generalized Maurer-Cartan equations* of G , thees generalize the Maurer-Cartan equations (see e.g [13]) in a minimal way.

Finally, note that the constraints $r_{jk}^i = -r_{kj}^i$ read

$$v_j^s \frac{\partial u_k^i}{\partial g^s} - u_k^s \frac{\partial v_j^i}{\partial g^s} = u_j^s \frac{\partial v_k^i}{\partial g^s} - v_k^s \frac{\partial u_j^i}{\partial g^s} \quad (6.3)$$

The generalized Maurer-Cartan differential equations can be rewritten more concisely. For $x \in T_e(G)$ introduce the *infinitesimal translations* as the local vector fields in the vicinity of the unit $e \in G$,

$$L_x \doteq L_x(g) \doteq x^j u_j^i(g) \frac{\partial}{\partial g^i}, \quad R_x \doteq R_x(g) \doteq x^j v_j^i(g) \frac{\partial}{\partial g^i} \quad \in T_g(G)$$

Define the Lie bracketing $[A, B]$ of the (local) vector fields A and B (on G) in the usual way $[A, B] \doteq AB - BA$. Then, the generalized Maurer-Cartan equations (6.2a,b) and differential equations (6.3) can be rewritten, respectively, as the commutation relations (CR) called a *birepresentation* [11] of the tangent Malcev algebra Γ_G as follows:

$$[L_x, L_y] = L_{[x,y]} - 2[L_x, R_y], \quad [R_x, R_y] = R_{[y,x]} - 2[R_x, L_y], \quad [L_x, R_y] = [R_x, L_y]$$

Here, one can easily see relation to the structure equations of *alternative* algebras [17]. In the case of associativity of G , the (generalized) Maurer-Cartan equations read

$$2[L_x, R_y] = L_{[x,y]} - [L_x, L_y] = R_{[y,x]} - [R_x, R_y] = 2[R_x, L_y] = 0$$

7 Akivis identity

Differentiate (6.1) with respect to g^l ($l = 1, \dots, n$) at $g = e$. Then one gets the 3rd order constraints, called the *third-order minimality conditions*,

$$l_{jkl}^i = l_{klj}^i = -l_{lkj}^i = r_{jkl}^i = m_{jkl}^i = -\tilde{l}_{jkl}^i = -\tilde{r}_{jkl}^i = -\tilde{m}_{jkl}^i$$

Thus the third-order associators of a local analytic Moufang loop have the property of their total anti-symmetry with respect of the lower indexes. From this it follows that for a continuous Moufang loop G the Akivis identity reads

$$J(x, y, z) = 6(x, y, z), \quad \forall x, y, z \in T_e(G)$$

Acknowledgements

The research was in part supported by the Estonian Research Council (Grant ETF9038) and Astralgo Tallinn. Author also partially used, modified and extended material from [10].

References

- [1] Akivis M A 1977 In *Differential Geometry* (Kalinin State University: Kalinin, in Russian) 3–9
- [2] Akivis M A 1976 *Sibirski Matem J* **17** 5–11 (in Russian)
- [3] Belousov V D 1967 *Foundations of the Theory of Quasigroups and Loops* (Nauka: Moscow, in Russian)
- [4] Bruck R H 1971 *A Survey of Binary Systems* (Springer: Berlin-Heidelberg-New York)
- [5] Chein O 1978 *Moufang Loops of Small Order* Memoirs AMS.
- [6] Chein O and Orlik Pflugfelder H 1971 *Archiv Math* **XXII** 573–76
- [7] Kuz'min E N 1971 *Algebra i Logika* **10** 3–22 (in Russian)
- [8] Mal'tsev A 1955 *Matem Sb* **36** 569–76 (in Russian)
- [9] Moufang R 1935 *Math Ann* **B110** 416–30
- [10] Paal E 1987 *An Introduction to the Moufang Symmetry* (Estonian Acad Sci: Tallinn-Tartu, in Russian)
- [11] Paal E 1993 *Proc Estonian Acad Sci Phys Math* **42** 157–65 (in Russian)
- [12] Pflugfelder H 1990 *Quasigroups and Loops: Introduction* (Heldermann Verlag: Berlin)
- [13] Pontryagin L S 1966 *Topological Groups* (Gordon and Breach: New York)
- [14] Sagle A A 1961 *Trans Amer Math Soc* **101** 426–58
- [15] Sabinin L V 1972 *Dokl Akad Nauk SSSR* **205** 533–36 (in Russian)
- [16] Sabinin L V 1972 *Matem Zametki* **12** 605–16 (in Russian)
- [17] Schafer R D 1966 *An Introduction to Nonassociative Algebras* (Academic Press: New York)