

# Dynamical Structure Factor and Spin-Density Separation for a Weakly-Interacting Two-Component Bose Gas

M.-C. Chung and A. B. Bhattacharjee

Max-Planck-Institut für Physik komplexer Systeme, 01187 Dresden, Germany

(Dated: May 28, 2019)

We show that spin-density separation in a Bose gas is not restricted to 1D but also occurs in higher dimension. The ratio ( $\alpha$ ) of the intra-species atom-atom interaction strength to the inter-species interaction strength, strongly influences the dynamics of spin-density separation and the elementary excitations. The density wave is phonon-like for all values of  $\alpha$ . For  $\alpha < 1$ , spin wave is also phonon-like. The spin waves have a quadratic dispersion in the  $\alpha = 1$  coupling regime, while in the phase separated regime ( $\alpha > 1$ ) the spin waves are found to be damped. The dynamical structure factor (DSF) reveals two distinct peaks corresponding to the density and spin waves for  $\alpha \leq 1$ . For  $\alpha > 1$  there is only one DSF peak corresponding to the density wave.

Experimental progress in the ability to coherently manipulate ultracold atomic gases motivates to study fundamental properties of these systems. Current technology allows one to create two-component Bose gases which has been demonstrated in experimental studies [1]. Relative spatial oscillations of the two-components can be viewed as spin waves. Many theoretical as well as experimental spin related effects have been predicted and observed in these two-component Bose gas, such as phase separation [2], exotic ground states [3] and suppression of spin transport in strong coupling regime [4]. The capability of a system to respond to an excitation probe transferring momentum and energy is described by the dynamic structure factor (DSF), which characterizes the collective excitations in these Bose gases. The DSF is now easily accessible experimentally using the technique of Bragg spectroscopy [5]. However, all these theoretical studies related to spin-density separation (motivated by analogous effect in Fermi systems in 1D) are restricted to 1D. There is a fundamental question to what extent spin-density separation survives in higher dimension. The purpose of the present Letter is to seek an answer to this question and to show that spin-density separation in a Bose gas within the hydrodynamic approximation and at zero temperature is more generic and occurs unlike in a Fermi system, in all dimensions. We will always work with  $\hbar = 1$ . We also compute the dynamical structure factor (DSF) which reveals distinct features of spin-charge separation in all dimensions.

We start with the Lagrangian density for two-component Bose gas at zero temperature:

$$\mathcal{L} = \frac{i}{2} \sum_{i=1,2} (\varphi_i^* \partial_t \varphi_i - \varphi_i \partial_t \varphi_i^*) - \frac{1}{2m} (\nabla_r \varphi_i)^2 - \mu_i n_i - \frac{1}{2} \sum_{i,j=1,2} g_{d,ij} n_i n_j, \quad (1)$$

where  $\varphi_i = \varphi_i(r, t)$ ,  $i = 1, 2$  is the field representing two different Bose particles,  $r$  is the space coordinate,  $t$  is the real time. Also  $\mu_i$ ,  $n_i = |\varphi_i|^2$  is the chemical potential and the particle density of the  $i^{\text{th}}$  component and

$g_{d,ij}$  is the effective atom-atom interaction between the  $i^{\text{th}}$  and  $j^{\text{th}}$  components. Here we consider bosonic atoms of the same isotope of mass  $m$  but having different internal spin states, therefore we have  $g_{d,11} = g_{d,22} \equiv g_d$  and  $g_{d,12} = g_{d,21} \equiv g'_d$ . For simplicity, we consider the same average atom number density for the two components, i.e.  $\bar{n}_1 = \bar{n}_2 = \bar{n}$ . For a 3D Bose gas,  $g_3 = 4\pi a_3/m$  [6, 7]. Here  $a_3$  is the 3D scattering length. For lower dimensional Bose gas in a 3D trap with longitudinal harmonic trapping frequency  $\omega_\perp$ ,  $g_2 = 4\pi/(m \ln \bar{n} a_2)$  [8, 9, 10] with the 2D scattering length given as  $a_2 \simeq 7.41 e^{-\sqrt{\pi/a_3^2 m \omega_\perp}}$  [11] and  $g_1 = 2\omega_0 a_3$  with  $a_3 \ll 1/\sqrt{m\omega_\perp}$  [12]. The behavior of the system depends crucially on the dimensionless parameter  $\gamma_d(\gamma'_d) = mg_d n^{1-2/d} (mg'_d n^{1-2/d})$ . For the gas to be weakly interacting, we must have  $\gamma_d(\gamma'_d) \ll 1$ . The chemical potential  $\mu_i$  of the  $i^{\text{th}}$  component is determined by the condition  $\sum_{ij} g_{d,ij} n_j = \mu_i n_i$ .

To understand the low-energy excitations in two-component Bose gas, one can derive a low-energy effective hydrodynamical Lagrangian that contains only modes related to the low-energy excitations [8, 13]. We write the Boson field  $\varphi_i$  in the terms of the number density  $n_i$  and the phase  $\theta_i$  as  $\varphi_i = n_i e^{i\theta_i}$ . In the weak coupling regime the phase changes slowly in space while the density fluctuates fast [7], therefore one can integrate out the high energy fast density fluctuation [13] to obtain the effective hydrodynamic action. We introduce the density fluctuation  $\delta n_i$  as  $n_i = \bar{n} + \delta n_i$ . In terms of the new basis,  $\delta n_{\rho(\sigma)} = (\delta n_1 \pm \delta n_2)/\sqrt{2}$  and  $\theta_{\rho(\sigma)} = (\theta_1 \pm \theta_2)/\sqrt{2}$ , the action obtained from the Lagrangian density can be rewritten as

$$S = \int d^d \mathbf{x} dt - \sum_{\lambda=\rho,\sigma} [(\bar{n}_\lambda + \delta n_\lambda) \partial_t \theta_\lambda + \frac{\bar{n} (\nabla_r \theta_\lambda)^2}{2m} + \frac{(\nabla_r \delta n_\lambda)^2}{8m\bar{n}} + \frac{g_{d,\lambda}}{2} (\delta n_\lambda)^2], \quad (2)$$

where  $\bar{n}_\rho(\bar{n}_\sigma) = \sqrt{2}\bar{n}(0)$  and  $g_{d,\rho(\sigma)} = g(1 \pm \alpha)$  with  $\alpha = g'_d/g_d$ .

For  $\alpha < 1$ , after performing two Gaussian integrals,




FIG. 1: Dynamic structure factor for all dimensions in the hydrodynamic regime and at zero temperature for  $\alpha < 1$ , shows two distinct peaks corresponding to the density and the spin waves, centered at  $v_{\rho}q$  and  $v_{\sigma}q$ , respectively. The one dimensional structure factor is found to be a delta function, while the two- and three dimensional DSF is broadened because of Beliaev damping. Clearly, the DSF for the density wave is broader compared to that of spin waves. These are interesting signatures of spin-density separation to look for using Bragg spectroscopy.

the effective action has the form[13]

$$S_{\text{eff}} = \int d^d \mathbf{x} dt \sum_{\lambda=\rho,\sigma} \left[ \frac{1}{2g_{d,\lambda}} |\partial_t \phi_\lambda|^2 - \frac{\bar{n}}{2m} |\nabla_r \phi_\lambda|^2 \right], \quad (3)$$

where  $\phi_{\rho(\sigma)} = e^{i\theta_{\rho(\sigma)}}$ . Here we assumed that the fields  $\theta_{\rho(\sigma)}$  vary slowly in space and we have dropped the  $\frac{\nabla_r^2}{8m\bar{n}}$  term. The effective Lagrangian density can be related to the physically measurable parameters as follows

$$\mathcal{L}_{\text{eff}} = \sum_{\lambda=\rho,\sigma} \frac{\chi_\lambda}{2} (|\partial_t \phi_\lambda|^2 - v_\lambda^2 |\nabla_r \phi_\lambda|^2), \quad (4)$$

where  $\chi_{\rho(\sigma)} = 1/g_{d,\rho(\sigma)}$  is the density (spin) compressibility and  $v_{\rho(\sigma)} = \sqrt{\bar{n}g_{d,\rho(\sigma)}/m}$  is the sound velocity of the density (spin) mode. The effective Lagrangian (4) describes the low-energy excitations of two sound waves with linear dispersions  $\omega_{\rho(\sigma)} = v_{\rho(\sigma)}k$ . The bosons split into two gapless modes, namely density mode and spin mode, propagating with different velocities. The density wave propagates faster than the spin wave, which can be seen by the relation  $v_\sigma/v_\rho \approx \sqrt{(1-\alpha)/(1+\alpha)}$ . In this regime the system lies at the quantum critical point because of the gapless dispersion relations.

The meaning of the low-energy effective Lagrangian (4) is that the bosonic system separates into two independent degree of freedom, i.e. spin and density. In contrast to the spin-charge separation for fermionic systems in 1D, spin-density separation in bosonic systems occurs not only in 1D but also in higher dimensions. The Bogoliubov energy dispersion relation of one-component interacting

Bose gas is  $\epsilon(k) = \sqrt{((k^2/2m)^2 + g_d \bar{n} k^2/m)}$  [14]. For the two component Bose gas, replacing the interaction  $g_d$  with  $g_{d,\rho(\sigma)}$ , we obtain two branches of the excitations

$$\epsilon_{\rho(\sigma)}(k) = \sqrt{(k^2/2m)^2 + \frac{g_{d,\rho(\sigma)}(1 \pm \alpha)\bar{n}}{m} k^2}, \quad (5)$$

which is in agreement with the result obtained by the semiclassical method [15]. From the dispersion relations (5) we can define the chemical potential for the density and spin waves as  $\mu_{\rho(\sigma)} = g_{d,\rho(\sigma)}\bar{n}$ .

For  $\alpha = 1$  ( $g_d = g'_d$ ), only one Gaussian integral can be performed in action (2) giving the gapless density wave with linear dispersion. However, one obtains a quadratic dispersion for the spin-wave excitations, in agreement with  $SU(2)$  symmetry [4, 16]. This effect can also be seen from the Bogoliubov excitations  $\epsilon_\rho = \sqrt{(k^2/2m)^2 + 2g\bar{n}k^2/m}$  and  $\epsilon_\sigma = k^2/2m$  by replacing  $g_{d,\rho} = 2g_d$  and  $g_{d,\sigma} = 0$  in (5). In this case, due to the  $SU(2)$  symmetry, the eigenstates are classified according to their total spin  $S$  ranging from 0 to  $N/2$ , and according to recent result by Li et.al [17], the ground state is fully polarized ( $S = N/2$ ). The ground state is describe by Lieb-Liniger(LL) model of one-component Bose gas [18], for which the elementary excitations in the weak-coupling regime are density waves [19], and the system is ferromagnetic. Spin excitations above the ground state corresponding to relative oscillations of the two gas components, are gapped, as in the case of the ferromagnetic Heisenberg model for spin systems. This gap gives the spin-wave excitation an effective mass.

In the case of  $\alpha > 1$  ( $g_d < g'_d$ ), we found  $g_{d,\sigma} < 0$ . This implies,  $v_\sigma (= \sqrt{g_{d,\sigma}\bar{n}/m})$  in the long wave length limit is imaginary. The spin waves become unstable and damp out in the thermodynamic limit. Therefore we obtain a phase separation of the two-component Bose gas [15].

The dynamical structure factor (DSF) of many-body system is defined as follows

$$S_{\rho(\sigma)}(q, \omega) = \int d^d \mathbf{x} dt e^{i(\omega t - \mathbf{q} \cdot \mathbf{x})} \langle \delta n_{\rho(\sigma)}(\mathbf{x}, t) \delta n_{\rho(\sigma)}(\mathbf{0}, 0) \rangle, \quad (6)$$

where  $\langle \dots \rangle$  can be calculated using path integral with the effective action. Experimentally, one can measure the dynamical structure factor using Bragg spectroscopy[5].

For  $\alpha < 1$ , from the action 2, one can get the equation of motion for  $\delta n_{\rho(\sigma)}$  as  $\delta n_{\rho(\sigma)}(\mathbf{x}, t) = -1/g_{d,\rho(\sigma)} \partial_t \theta_{\rho(\sigma)}(\mathbf{x}, t)$ . From the quadratic Lagrangian density, the DSF (6) can be obtained as

$$S_{\rho(\sigma)}(q, \omega) = \text{Im} \frac{\chi_{\rho(\sigma)} v_{\rho(\sigma)}^2 q^2}{\omega^2 - \omega_{\rho(\sigma)}^2(q)}, \quad (7)$$

where  $\omega_{\rho(\sigma)}(q) = v_{\rho(\sigma)}q + i\Gamma_{\rho(\sigma)}(q)$  with the quasiparticle decay rate  $\Gamma_{\rho(\sigma)}(q)$ .

In order to obtain the DSF, one has to find the compressibility  $\chi_{\rho(\sigma)}$ , velocity  $v_{\rho(\sigma)}$  and decay rate  $\Gamma_{\rho(\sigma)}(q)$

in terms of the dimensionless parameters  $\gamma_{d,\rho(\sigma)}$ . Using the macroscopic argument, the compressibility  $\chi_{\rho(\sigma)}$  is related to the energy  $E(g_{\rho(\sigma)}, n)$  as  $\chi_{\rho(\sigma)}^{-1} = \frac{1}{V} \frac{\partial^2 E}{\partial n^2}$  with the constant system size:  $V = L^d$  and density:  $n = N/V$ . Similarly, the sound velocity can also be obtained using the macroscopic energy spectrum as  $v_{\rho(\sigma)} = (\frac{V}{mn} \frac{\partial^2 E}{\partial V^2})^{1/2}$  with constant particle number  $N$  [21]. The way to obtain the ground state energy spectrum is diverse and depends on the dimension. As indicated by Beliaev [22], the decay rate  $\Gamma_{\rho(\sigma)}(q)$  is caused by the process of a long wavelength phonon decaying into two phonons and it can be calculated for small momenta using the formula [8]

$$\Gamma_{\rho(\sigma)}(q) = \frac{9v_{\rho(\sigma)}}{128\pi^2 \bar{n}m} \int d^d k |\mathbf{q}||\mathbf{k}||\mathbf{q}-\mathbf{k}| \delta(\epsilon_{\rho(\sigma)}(\mathbf{q}) - \epsilon_{\rho(\sigma)}(\mathbf{k}) - \epsilon_{\rho(\sigma)}(\mathbf{q}-\mathbf{k})). \quad (8)$$

Therefore the Beliaev decay rate is also dimensional dependent.

In 3D, the dimensionless parameter  $\gamma_3 = 2\pi a_3 \bar{n}^{1/3}/m$ . In this case, the requirement for a dilute gas  $\bar{n}a^3 \ll 1$  corresponds to the weak-coupling condition  $\gamma_3 \ll 1$ . The ground state energy was given for the first time by Lee, Huang and Yang [6] as  $E = N\bar{n}^{2/3}/(2m)\gamma_3(1 + 16\gamma_3^{3/2}/5\pi^2)$ . The ground state compressibility and velocity are given by  $\chi_{\rho(\sigma)}^{-1} = g_{3,\rho(\sigma)}(1 + \frac{2}{\pi^2}\gamma_{3,\rho(\sigma)}^{\frac{3}{2}})$  and  $v_{\rho(\sigma)} = \sqrt{\frac{g_{3,\rho(\sigma)}\bar{n}}{m}}(1 + \frac{2}{\pi^2}\gamma_{3,\rho(\sigma)}^{\frac{3}{2}})^{\frac{1}{2}}$ , respectively. The decay rate for 3D system is obtained from eq(8):  $\Gamma_{\rho(\sigma)}(q) = \Gamma(q) = \frac{3\sigma^5}{640\pi m \bar{n}}$  [22]. We can see that the decay rates for density and spin waves are equal and proportional to  $q^5$ . The DSF for  $\omega > 0$  can be approximated as

$$S_{\rho(\sigma)}(q, \omega) \approx \frac{\chi_{\rho(\sigma)} v_{\rho(\sigma)} q \Gamma_{\rho(\sigma)}(q)}{2 \left[ (\omega - v_{\rho(\sigma)} q)^2 + \Gamma_{\rho(\sigma)}^2(q) \right]}, \quad (9)$$

In the Bragg scattering experiment, one should obtain two peaks centered at  $v_{\rho(\sigma)}q$  for the cross section with the width  $\Gamma(q)$ .

For 2D Bose gas, renormalization-group analysis [10, 23] shows that the interaction of the 2D dilute gas is marginally irrelevant only in a dilute limit specified by  $\ln \ln \gamma_2 \gg 1$ . The corresponding ground state energy for a weak-interacting gas is given by  $E = N\bar{n}/(2m)\gamma_2(1 - C\gamma_2)$  where constant  $C \ll 1$  is not universal but model-dependent due to the marginal interaction[23]. The compressibility and velocity for spin and density-wave excitations are  $\chi_{\rho(\sigma)}^{-1} = g_{2,\rho(\sigma)}(1 - (C - \frac{3}{8\pi})\gamma_{2,\rho(\sigma)})$  and  $v_{\rho(\sigma)} = \sqrt{\frac{\bar{n}\gamma_{2,\rho(\sigma)}}{m}}(1 - (C - \frac{3}{8\pi})\gamma_{2,\rho(\sigma)})^{1/2}$ . The Beliaev decay rate can be obtained by the integral (8):  $\Gamma_{\rho(\sigma)}(q) = \frac{\sqrt{3}v_{\rho(\sigma)}q^3}{64\pi}$ . Therefore the DSF (9) has a broader width for density waves than spin waves.

In the case of one dimension, contrary to 2D and 3D systems, the weak coupling means that the system is




FIG. 2: Dynamic structure factor for  $SU(2)$  symmetric Hamiltonian, which can be measured by Bragg spectroscopy. Note that  $\omega$  is in the units of  $g_d \bar{n}$  and  $q$  is in the units of  $\sqrt{m g_d \bar{n}}$ . The DSF of the density waves varies linearly with  $q$ , while the DSF of the spin waves shows the quadratic dependence on  $q$ .

in the high density regime because  $\gamma_1 = m g_1 / \bar{n}$ . In this regime, Lieb and Liniger [18] first gave the ground state energy as:  $E = \frac{N n^2}{2m} \gamma_1 (1 - \frac{4}{3\pi} \sqrt{\gamma_1})$ . A few algebra leads to the compressibility and sound velocities as  $\chi_{\rho(\sigma)}^{-1} = g_{1,\rho(\sigma)}(1 - \frac{1}{2\pi} \sqrt{\gamma_{1,\rho(\sigma)}})$  and  $v_{\rho(\sigma)} = \sqrt{\frac{g_{1,\rho(\sigma)}\bar{n}}{m}}(1 - \frac{1}{2\pi} \sqrt{\gamma_{1,\rho(\sigma)}})^{1/2}$ . For 1D, one obtains no decay rate. The reason is that the scenario for one phonon decaying into two phonons cannot exist due to the fact that energy conservation law in eq(8) cannot be fulfilled in 1D. Therefore two sharp peaks should be observed in the Bragg scattering experiments. Figure 1 illustrates and summarizes the results obtained above for the DSF in all the three dimensions.

In the case of  $\alpha = 1$ , the situation changes. For the density waves the dynamic structure factor remains the same as that in two-sound regime, while the DSF for spin-wave excitation alters due to the dramatic changing of the dispersion from linear to quadratic. In order to calculate the DSF one can use the effective Hamiltonian in the weak-coupling regime:

$$H = \sum_p \epsilon_p a_p^\dagger a_p + \sum_p \epsilon_p b_p^\dagger b_p + g_d \sqrt{\frac{\bar{n}}{V}} \sum_{k,q \neq 0} \sqrt{\frac{\epsilon_q}{\epsilon_q}} (a_q^\dagger + a_q) b_{k-q}^\dagger b_k \quad (10)$$

with the spectrum of free spin waves  $\epsilon_p = p^2/2m$ , the Bogoliubov spectrum  $\epsilon_p = \sqrt{\epsilon_p^2 + 2\mu_d n \epsilon_p}$  [4] and the chemical potential:  $\mu_d = 2g_d \bar{n}$ . Using  $\delta n_\sigma = \sqrt{\bar{n}}(b^\dagger + b)$ , the DSF can be related to the imaginary part of the Green

function as  $S(q, \omega) = \bar{n} \text{Im}G(q, \omega)$  where  $G(q, \omega)$  is the single particle Green function of the spin operators  $b_q$  and  $b_q^\dagger$ . Therefore the DSF for the spin waves reads

$$S(q, \omega) = \frac{\bar{n} \Gamma_{d, \sigma}(q)}{(\omega - \frac{q^2}{2m_d^*})^2 + \Gamma_{d, \sigma}^2(q)}, \quad (11)$$

where the effective mass  $m^*$  is determined by the equation:  $m/m_d^* = (1 + 2/m\partial^2 \text{Re}\Sigma(p)/\partial p^2)(p=0)$  with the self energy defined as  $\Sigma = G^{-1}(g_d) - G^{-1}(g_d = 0)$ , and the decay rate:  $\Gamma_{d, \sigma}(q) = \text{Im}\Sigma(q)$ . To the second order diagram for the self energy  $\Sigma$ , one obtains the inverse effective mass related to the dimensionless parameter  $\gamma_d = \mu_d \bar{n}^{-2/d}$  as  $m/m_d^* = 1 - \alpha_d \gamma_d^{d/2}$  with  $\alpha_d = 2/3\pi, 1/2\pi, 1/8\pi$  for one, two and three dimensions, respectively. The decay process depends on the spin-phonon interaction which requires the energy conservation:  $e_{q-k} + \epsilon_k = e_q$  with the spin momentum  $q$  and phonon momentum  $k$ . For  $q < \sqrt{m\mu_d}$ , this condition cannot be fulfilled, therefore  $\Gamma_{d\sigma} = 0$ , i.e.  $S(q, \omega) = \bar{n}\delta(\omega - \omega_q)$ . For  $q \gtrsim \sqrt{m\mu_d}$ , an approximation can be obtained as follows:  $\Gamma_{d\sigma}(\sqrt{m\mu_d}(1 + \delta)) = \beta_d \mu_d \delta^{3/2(d-1)}$  for  $\delta \ll 1$  with  $\beta_d = 0, 1/8\pi, 2/3\pi$  for  $d = 1, 2, 3$ , respectively. The eqn. (11) shows the fact that the excitations for spin waves for the Hamiltonian with  $SU(2)$  symmetry are not sound-like, but particle-like, with the DSF centered at the position proportional to  $q^2$  instead of  $q$ . Similar to the two-sound mode regime ( $\alpha < 1$ ), the DSF for one dimension is a delta function due to the fact that the energy conservation relation for a particle emitting a phonon cannot be fulfilled in one dimension at zero temperature. Fig. 2 shows the linear dispersion of the DSF for the density waves and the quadratic dispersion of the DSF for the spin waves. The delta function behavior are shown for low momenta.

For phase-separation regime ( $\alpha > 1$ ), the spin waves are thermodynamically unstable, therefore only density waves exist. The dynamic structure factor of spin waves is smeared out and there exists only one peak in the DSF, which is different from the other regimes. This property can be a prominent signature for checking whether the system is phase separated or not.

In this Letter we have shown that, unlike fermionic systems, spin-density separation in two-component bosonic system is a more generic feature and occurs in all dimensions. The density wave is found to be phonon like for all dimensions and coupling regimes. However, the spin waves show a linear dispersion (phonon like) in the  $\alpha < 1$  regime, quadratic dispersion in the  $\alpha = 1$  regime and is damped in the phase separated regime ( $\alpha > 1$ ). In the  $\alpha < 1$  regime, the DSF for all dimensions show two distinct peaks corresponding to the density and the spin waves, centered at  $v_\rho q$  and  $v_\sigma q$ , respectively. In the same regime, the one-dimensional structure factors are found to be delta functions, while the two- and three-dimensional DSF is broadened because of Beliaev damp-

ing. The DSF for the density wave is found to be broad compared to that of spin waves in 2D, while for 3D both have the same width. For  $\alpha = 1$ , the DSF for all dimensions also show two distinct peaks centered at  $v_\rho q$  and  $q^2/2m_d^*$ . For  $\alpha > 1$  there is only one peak corresponding to the density wave. These are interesting signatures of spin-density separation to look for using Bragg spectroscopy where, the response of the condensate to a two-photon Bragg pulse is measured. The momentum transferred would reflect the structure of the underlying excitation spectrum of spin and density wave.

M. C. Chung thanks B. Dora for valuable discussions.

- 
- [1] D. S. Hall et al., Phys. Rev. Lett. **81**, 1539 (1998); **81**, 1543 (1998).
  - [2] W. B. Colson and A. Fetter, J. Low Temp. Phys. **33**, 231 (1978); T-L. Ho and V. B. Shenoy, Phys. Rev. Lett. **77**, 3276 (1996); B. D. Esry et al., Phys. Rev. Lett. **78**, 3594 (1997); C.K. Law et al., Phys. Rev. Lett. **79**, 3105 (1997); H. Pu and N. P. Bigelow, Phys. Rev. Lett., **80**, 1130 (1998).
  - [3] T-L. Ho and S. K. Yip, Phys. Rev. Lett. **84**, 4031 (2000); A. B. Kuklov and B. V. Svistunov, **89**, 170403 (2002).
  - [4] J. N. Fuchs et al. Phys. Rev. Lett. **95**, 150402 (2005).
  - [5] J. Steinhauer et al., Phys. Rev. Lett. **90**, 060404 (2003).
  - [6] T. D. Lee, K. Huang, and C. N. Yang, Phys. Rev. Lett. **106**, 1135 (1957).
  - [7] Anthony. J. Leggett, Rev. Mod. Phys. **37**, 307 (2001).
  - [8] V. N. Popov, *Functional Integrals in Quantum Field Theory and Statistical Physics* (D. Riedel 1983).
  - [9] M. Schick, Phys. Rev. A, **3**, 1067 (1971). D. S. Fisher. For a review, see A. Posazhennikova, Rev. Mod. Phys. **78**, 1111(2006).
  - [10] D. S. Fisher and P. C. Hohenberg, Phys. Rev. B, **37**, 4936 (1988).
  - [11] M. D. Lee et al, Phys. Rev. A, **65**, 043617 (2002).
  - [12] M. Olshanii, Phys. Rev. Lett. **81**, 938 (1998).
  - [13] Xiao-Gang Wen *Quantum Field Theory of Many-Body Systems* (Oxford University Press 2004).
  - [14] N. N. Bogoliubov, J. Phys. (USSR), **11**, 23 (1947).
  - [15] J. Phys.: Condensed Matter **14** L327 (2002).
  - [16] B. I. Halperin and P. C. Hohenberg, Phys. Rev. **188**, 898 (1969); B. I. Halperin, Phys. Rev. B **11**, 178 (1975).
  - [17] Y. Q. Li, Europhys. Lett. **61**, 368 (2003).
  - [18] E. H. Lieb and W. Liniger, Phys. Rev. **130**, 1605 (1963).
  - [19] E. H. Lieb, Phys. Rev. **130**, 1616 (1963).
  - [20] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, New York, 1971).
  - [21] See, for example, F. London, *Superfluids* (John Wiley & Sons, Inc., New York), Vol. II p.83 and J. W. Negele and H. Orland, *Quantum Many-Particle Systems* (Addison Wesley), Chap. 6 p.303.
  - [22] S. T. Beliaev Sov. Phys. JETP **7**, 104(1958); **7**, 289 (1958).
  - [23] E. B. Kolomeisky and J. P. Straley, Phys. Rev. B, **46**, 11749 (1992).