Propagation Rules of Subsystem Codes

Salah A. Aly

Department of Computer Science Texas A&M University College Station, TX 77843, USA

Email: salah@cs.tamu.edu

Abstract—We demonstrate propagation rules of subsystem code constructions by extending, shortening and combining given subsystem codes. Given an $[[n,k,r,d]]_q$ subsystem code, we drive new subsystem codes with parameters $[[n+1,k,r,\geq d]]_q$, $[[n-1,k+1,r,\geq d-1]]_q$, $[[n,k-1,r+1,d]]_q$. We present the short subsystem codes. The interested readers shall consult our companion papers for upper and lower bounds on subsystem codes parameters, and introduction, trading dimensions, families, and references on subsystem codes [1], [2], [3] and references therein.

Subsystem Codes. Let \mathcal{H} be the Hilbert space $C^{q^n} = \mathbf{C}^q \otimes \mathbf{C}^q \otimes ... \otimes \mathbf{C}^q$. Let Q be a quantum code such that $\mathcal{H} = Q \oplus Q^{\perp}$, where Q^{\perp} is the orthogonal complement of Q. Recall definition of the error model acting in qubits [4], [3]. We can define the subsystem code Q as follows

Definition 1: An $[[n,k,r,d]]_q$ subsystem code is a decomposition of the subspace Q into a tensor product of two vector spaces A and B such that $Q = A \otimes B$, where $\dim A = k$ and $\dim B = r$. The code Q is able to detect all errors of weight less than d on subsystem A.

Subsystem codes can be constructed from the classical codes over \mathbf{F}_q and \mathbf{F}_{q^2} . The Euclidean construction of subsystem code is given as follows [1], [3].

Lemma 2 (Euclidean Construction): If C is a k'-dimensional \mathbf{F}_q -linear code of length n that has a k''-dimensional subcode $D = C \cap C^{\perp}$ and k' + k'' < n, then there exists an

$$[[n, n - (k' + k''), k' - k'', \operatorname{wt}(D^{\perp} \setminus C)]]_q$$

subsystem code.

I. Subsystem Codes vers. Co-subsystem Codes

In this section we show how one can trade the dimensions of subsystem and co-subsystem to obtain new codes from a given subsystem or stabilizer code. The results are obtained by exploiting the symplectic geometry of the space. A remarkable consequence is that nearly any stabilizer code yields a series of subsystem codes.

Our first result shows that one can decrease the dimension of the subsystem and increase at the same time the dimension of the co-subsystem while keeping or increasing the minimum distance of the subsystem code.

Theorem 3: Let q be a power of a prime p. If there exists an $((n, K, R, d))_q$ subsystem code with K > p that is pure to d', then there exists an $((n, K/p, pR, \ge d))_q$ subsystem code that

is pure to $\min\{d, d'\}$. If a pure $((n, p, R, d))_q$ subsystem code exists, then there exists a $((n, 1, pR, d))_q$ subsystem code.

Replacing \mathbf{F}_p -bases by \mathbf{F}_q -bases in the proof of the previous theorem yields the following variation of the previous theorem for \mathbf{F}_q -linear subsystem codes.

Theorem 4: Let q be a power of a prime p. If there exists a pure \mathbf{F}_q -linear $[[n,k,r,d]]_q$ subsystem code with r>0, then there exists a pure \mathbf{F}_q -linear $[[n,k+1,r-1,d]]_q$ subsystem code.

Theorem 5 (Generic methos): If there exists an $(\mathbf{F}_q$ -linear) $[[n,k,d]]_q$ stabilizer code that is pure to d', then there exists for all r in the range $0 \le r < k$ an $(\mathbf{F}_q$ -linear) $[[n,k-r,r,\ge d]]_q$ subsystem code that is pure to $\min\{d,d'\}$. If a pure $(\mathbf{F}_q$ -linear) $[[n,k,r,d]]_q$ subsystem code exists, then a pure $(\mathbf{F}_q$ -linear) $[[n,k+r,d]]_q$ stabilizer code exists.

Using this theorem we can derive many families of subsystem codes derived from families of stabilizer codes as shown in Table 1

II. PROPAGATION RULES

Let $C_1 \leq \mathbf{F}_q^n$ and $C_2\mathbf{F}_q^n$ be two classical codes defined over F_q . The direct sum of C_1 and C_2 is a code $C \leq \mathbf{F}_q^{2n}$ defined as follows

$$C = C_1 \oplus C_2 = \{ uv \mid u \in C_1, v \in C_2 \}. \tag{1}$$

In a matrix form the code C can be described as

$$C = \left(\begin{array}{cc} C_1 & 0 \\ 0 & C_2 \end{array} \right)$$

An $[n,k_1,d_1]_q$ classical code C_1 is a subcode in an $[c,k_2,d_2]_q$ if every codeword v in C_1 is also a codeword in C_2 , hence $k_1 \leq k_2$. We say that an $[[n,k_1,r_1,d_1]]_q$ subsystem code Q_1 is a subcode in an $[[n,k_2,r_2,d_2]]_q$ subsystem code Q_2 if every codeword $|v\rangle$ in Q_1 is also a codeword in Q_2 and $k_1+r_1\leq k_2+r_1$.

Notation. Let q be a power of a prime integer p. We denote by \mathbf{F}_q the finite field with q elements. We use the notation $(x|y)=(x_1,\ldots,x_n|y_1,\ldots,y_n)$ to denote the concatenation of two vectors x and y in \mathbf{F}_q^n . The symplectic weight of $(x|y) \in \mathbf{F}_q^{2n}$ is defined as

$$\operatorname{swt}(x|y) = \{(x_i, y_i) \neq (0, 0) \mid 1 \le i \le n\}.$$

Family	Stabilizer $[[n, k, d]]_q$	Subsystem $[[n, k-r, r, d]]_q, k > r \ge 0$
Short MDS	$[[n, n-2d+2, d]]_q$	$[[n, n-2d+2-r, r, d]]_q$
Hermitian Hamming	$[[n, n-2m, 3]]_q$	$m \ge 2, [[n, n-2m-r, r, 3]]_q$
Euclidean Hamming	$[[n, n-2m, 3]]_q$	$[[n, n-2m-r, r, 3]]_q$
Melas	$[[n, n-2m, \ge 3]]_q$	$[[n, n-2m-r, r, \ge 3]]_q$
Euclidean BCH	$[[n, n-2m\lceil(\delta-1)(1-1/q)\rceil, \geq \delta]]_q$	$[[n, n-2m\lceil(\delta-1)(1-1/q)\rceil-r, r, \geq \delta]]_q$
Hermitian BCH	$[[n, n-2m\lceil(\delta-1)(1-1/q^2)\rceil, \geq \delta]]_q$	$[[n, n-2m\lceil(\delta-1)(1-1/q^2)\rceil-r, r, \geq \delta]]_q$
Punctured MDS	$[[q^2 - q\alpha, q^2 - q\alpha - 2\nu - 2, \nu + 2]]_q$	$[[q^2 - q\alpha, q^2 - q\alpha - 2\nu - 2 - r, r, \nu + 2]]_q$
Euclidean MDS	$[[n, n-2d+2]]_q$	$[[n, n-2d+2-r, r]]_q$
Hermitian MDS	$[[q^2 - s, q^2 - s - 2d + 2, d]]_q$	$[[q^2 - s, q^2 - s - 2d + 2 - r, r, d]]_q$
Twisted	$[[q^r, q^r - r - 2, 3]]_q$	$[[q^r, q^r - r - 2 - r, r, 3]]_q$
Extended twisted	$[[q^2+1,q^2-3,3]]_q$	$[[q^2+1,q^2-3-r,r,3]]_q$
Perfect	$[[n, n-s-2, 3]]_q$	$[[n, n-s-2-r, r, 3]]_q$
	$[[n, n-s-2, 3]]_q$	$[[n, n-s-2-r, r, 3]]_q$

Fig. 1. Families of subsystem codes from stabilizer codes

We define $swt(X) = min\{swt(x) | x \in X, x \neq 0\}$ for any nonempty subset $X \neq \{0\}$ of \mathbf{F}_a^{2n} .

The trace-symplectic product of two vectors u = (a|b) and v = (a'|b') in \mathbf{F}_q^{2n} is defined as

$$\langle u|v\rangle_s = \operatorname{tr}_{q/p}(a'\cdot b - a\cdot b'),$$

where $x \cdot y$ denotes the dot product and $\operatorname{tr}_{q/p}$ denotes the trace from \mathbf{F}_q to the subfield \mathbf{F}_p . The trace-symplectic dual of a code $C \subseteq \mathbf{F}_q^{2n}$ is defined as

$$C^{\perp_s} = \{ v \in \mathbf{F}_q^{2n} \mid \langle v | w \rangle_s = 0 \text{ for all } w \in C \}.$$

We define the Euclidean inner product $\langle x|y\rangle=\sum_{i=1}^n x_iy_i$ and the Euclidean dual of $C \subseteq \mathbf{F}_q^n$ as

$$C^{\perp} = \{x \in \mathbf{F}_q^n \mid \langle x | y \rangle = 0 \text{ for all } y \in C\}.$$

We also define the Hermitian inner product for vectors x, yin $\mathbf{F}_{q^2}^n$ as $\langle x|y\rangle_h = \sum_{i=1}^n x_i^q y_i$ and the Hermitian dual of $C \subseteq \mathbf{F}_{q^2}^n$ as

$$C^{\perp_h} = \{ x \in \mathbf{F}_{q^2}^n \mid \langle x | y \rangle_h = 0 \text{ for all } y \in C \}.$$

Theorem 6: Let C be a classical additive subcode of \mathbf{F}_a^{2n} such that $C \neq \{0\}$ and let D denote its subcode $D = C \cap C^{\perp_s}$. If x = |C| and y = |D|, then there exists a subsystem code $Q = A \otimes B$ such that

- i) dim $A = q^n/(xy)^{1/2}$,
- ii) dim $B = (x/y)^{1/2}$.

The minimum distance of subsystem A is given by

(a)
$$d = \text{swt}((C + C^{\perp_s}) - C) = \text{swt}(D^{\perp_s} - C)$$
 if $D^{\perp_s} \neq C$;
(b) $d = \text{swt}(D^{\perp_s})$ if $D^{\perp_s} = C$.

Thus, the subsystem A can detect all errors in E of weight less than d, and can correct all errors in E of weight $\leq \lfloor (d-1)/2 \rfloor$.

A. Extending Subsystem Codes

We derive new subsystem codes from known ones by extending and shortening the length of the code.

Theorem 7: If there exists an $((n, K, R, d))_a$ Clifford subsystem code with K > 1, then there exists an $((n+1, K, R, \ge$ $d)_q$ subsystem code that is pure to 1.

Proof: We first note that for any additive subcode $X \leq$ \mathbf{F}_q^{2n} , we can define an additive code $X' \leq \mathbf{F}_q^{2n+2}$ by

$$X' = \{(a\alpha|b0) \mid (a|b) \in X, \alpha \in \mathbf{F}_q\}.$$

We have |X'| = q|X|. Furthermore, if $(c|e) \in X^{\perp_s}$, then $(c\alpha|e0)$ is contained in $(X')^{\perp_s}$ for all α in \mathbf{F}_q , whence $(X^{\perp_s})' \subseteq (X')^{\perp_s}$. By comparing cardinalities we find that equality must hold; in other words, we have

$$(X^{\perp_s})' = (X')^{\perp_s}.$$

By Theorem 6, there are two additive codes C and Dassociated with an $((n, K, R, d))_q$ Clifford subsystem code such that

$$|C| = q^n R/K$$

and

$$|D| = |C \cap C^{\perp_s}| = q^n/(KR).$$

We can derive from the code C two new additive codes of length 2n+2 over \mathbf{F}_q , namely C' and $D'=C'\cap (C')^{\perp_s}$. The codes C' and D' determine a $((n+1,K',R',d'))_q$ Clifford subsystem code. Since

$$D' = C' \cap (C')^{\perp_s} = C' \cap (C^{\perp_s})'$$

= $(C \cap C^{\perp_s})'$,

we have |D'| = q|D|. Furthermore, we have |C'| = q|C|. It follows from Theorem 6 that

- $\begin{array}{ll} \text{(i)} \ \ K'=q^{n+1}/\sqrt{|C'||D'|}=q^n/\sqrt{|C||D|}=K,\\ \text{(ii)} \ \ R'=(|C'|/|D'|)^{1/2}=(|C|/|D|)^{1/2}=R, \end{array}$
- (iii) $d' = \operatorname{swt}((D')^{\perp_s} \setminus C') \ge \operatorname{swt}((D^{\perp_s} \setminus C)') = d.$

Since C' contains a vector $(\mathbf{0}\alpha|\mathbf{0}0)$ of weight 1, the resulting subsystem code is pure to 1.

Corollary 8: If there exists an $[[n, k, r, d]]_q$ subsystem code with k > 0 and $0 \le r < k$, then there exists an $[n+1, k, r, \ge n]$ $d]_q$ subsystem code that is pure to 1.

B. Shortening Subsystem Codes

We can also shorten the length of a subsystem code and still trade the dimensions of the new subsystem code and its co-subsystem code as shown in the following Lemma.

Theorem 9: If an $((n, K, R, d))_q$ pure subsystem code Q exists, then there is a pure subsystem code Q_p with parameters $((n-1, qK, R, \geq d-1))_q$.

Proof: We know that existence of the pure subsystem code Q with parameters $((n, K, R, d))_q$ implies existence of a pure stabilizer code with parameters $((n, KR, \geq d))_q$ for $n \geq$ 2 and $d \ge 2$ from [2, Theorem 2.]. By [5, Theorem 70], there exist a pure stabilizer code with parameters $((n-1, qKR, \geq$ $(d-1)_q$. This stabilizer code can be seen as $((n-1, qKR, 0, \geq 1)_q)_q$. $(d-1)_q$ subsystem code. By using [2, Theorem 2.], there exists a pure \mathbf{F}_q -linear subsystem code with parameters ((n- $(1, qK, R, \geq d-1)_q$ that proves the claim. Analog of the previous Theorem is the following Lemma.

Lemma 10: If an \mathbf{F}_q -linear $[[n, k, r, d]]_q$ pure subsystem code Q exists, then there is a pure subsystem code Q_p with parameters $[[n-1, k+1, r, \geq d-1]]_q$.

Proof: We know that existence of the pure subsystem code Q implies existence of a pure stabilizer code with parameters $[[n, k + r, \geq d]]_q$ for $n \geq 2$ and $d \geq 2$ by using [2, Theorem 2. and Theorem 5.]. By [5, Theorem 70], there exist a pure stabilizer code with parameters [n-1, k+1] $r+1, \geq d-1$]_q. This stabilizer code can be seen as an $[[n-1,k+r+1,0,\geq d-1]]_q$ subsystem code. By using [2, Theorem 3.], there exists a pure \mathbf{F}_q -linear subsystem code with parameters $[[n-1,k+1,r,\geq d-1]]_q$ that proves the

We can also prove the previous Theorem by defining a new code C_p from the code C as follows.

Theorem 11: If there exists a pure subsystem code Q = $A \otimes B$ with parameters $((n, K, R, d))_q$ with $n \geq 2$ and $d \geq 2$, then there is a subsystem code Q_p with parameters ((n - $1, K, qR, \geq d-1)_q$

Proof: By Theorem 6, if an $((n, K, R, d))_q$ subsystem code Q exists for K > 1 and $1 \le R < K$, then there exists an additive code $C\in \mathbf{F}_q^{2n}$ and its subcode $D\leq \mathbf{F}_q^{2n}$ such that $|C|=q^nR/K$ and $|D|=|C\cap C^{\perp_s}|=q^n/KR$. Furthermore, $d = \min \operatorname{swt}(D^{\perp_s} \setminus C)$. Let $w = (w_1, w_2, \dots, w_n)$ and u = (u_1, u_2, \dots, u_n) be two vectors in \mathbf{F}_q^n . W.l.g., we can assume that the code D^{\perp_s} is defined as

$$D^{\perp_s} = \{(u|w) \in \mathbf{F}_q^{2n} \mid w, u \in \mathbf{F}_q^n\}.$$

Let $w_{-1}=(w_1,w_2,\ldots,w_{n-1})$ and $u_{-1}=(u_1,u_2,\ldots,u_{n-1})$ be two vectors in \mathbf{F}_q^{n-1} . Also, let $D_p^{\perp_s}$ be the code obtained by puncturing the first coordinate of D^{\perp_s} , hence

$$D_p^{\perp_s} = \{(u_{-1}|w_{-1}) \in \mathbf{F}_q^{2n-2} \mid w_{-1}, u_{-1} \in \mathbf{F}_q^{n-1}\}.$$

since the minimum distance of D^{\perp_s} is at least 2, it follows that $|D_p^{\perp_s}| = |D^{\perp_s}| = K^2|C| = K^2q^nR/K = q^nRK$ and the minimum distance of $D_p^{\perp s}$ is at least d-1. Now, let us

construct the dual code of $D_p^{\perp_s}$ as follows.

$$\begin{array}{rcl} (D_p^{\perp_s})^{\perp_s} & = & \{(u_{-1}|w_{-1}) \in \mathbf{F}_q^{2n-2} \mid \\ & & (0u_{-1}|0w_{-1}) \in D, w_{-1}, u_{-1} \in \mathbf{F}_q^{n-1} \}. \end{array}$$

Furthermore, if $(u_{-1}|w_{-1}) \in D_p$, then $(0u_{-1}|0w_{-1}) \in D$. Therefore, D_p is a self-orthogonal code and it has size given

$$|D_p| = q^{2n-2}/|D_p^{\perp_s}| = q^{n-2}/RK.$$

We can also puncture the code C to the code C_p at the first coordinate, hence

$$C_p = \{(u_{-1}|w_{-1}) \in \mathbf{F}_q^{2n-2} \mid w_{-1}, u_{-1} \in \mathbf{F}_q^{n-1}, (aw_{-1}|bu_{-1}) \in C, a, b \in F_q\}.$$

Clearly, $D \subseteq C$ and if a = b = 0, then the vector $(0u_{-1}|0w_{-1}) \in D$, therefore, $(u_{-1}, w_{-1}) \in D_p$. This gives us that $D_p \subseteq C_p$. Furthermore, hence $|C| = |C_p|$. The dual code $C_p^{\perp_s}$ can be defined as

$$\begin{array}{rcl} C_p^{\perp_s} & = & \{(u_{-1}|w_{-1}) \in \mathbf{F}_q^{2n-2} \mid w_{-1}, u_{-1} \in \mathbf{F}_q^{n-1}, \\ & (ew_{-1}|fu_{-1}) \in C^{\perp_s}, e, f \in F_q\}. \end{array}$$

Also, if e = f = 0, then $D_p \subseteq C_p^{\perp_s}$, furthermore,

$$D_p^{\perp_s} = C_p \cup C_p^{\perp_s} = \{(u_{-1}|w_{-1}) \in \mathbf{F}_q^{2n-2} \mid (0u_{-1}|0w_{-1}) \in D\}$$
 (2)

Therefore there exists a subsystem code $Q_p = A_p \otimes B_p$. Also, the code $D_p^{\perp_s}$ is pure and has minimum distance at least d-1. We can proceed and compute the dimension of subsystem A_p and co-subsystem B_p from Theorem 6 as

$$\begin{array}{ll} \text{(i)} \ \ K_p &= q^{n-1}/\sqrt{|C_p||D_p|} \\ q^{n-1}/\sqrt{(q^nR/K)(q^{n-2}/RK)} = K, \\ \text{(ii)} \ \ R_p = (|C_p|/|D_p'|)^{1/2} = ((q^nR/K)/(q^{n-2}/RK))^{1/2} = \end{array}$$

(ii)
$$R_p = (|C_p|/|D_p'|)^{1/2} = ((q^n R/K)/(q^{n-2}/RK))^{1/2} = qR,$$

(iii)
$$d_p = \operatorname{swt}((D_p)^{\perp_s} \setminus C_p) = \operatorname{swt}((D^{\perp_s} \setminus C_p)) \ge d - 1.$$

Therefore, there exists a subsystem cod with parameters $((n-1, K, qR, \geq d-1))_q$.

The minimum distance condition follows since the code Q has $d = \min \operatorname{swt}(D^{\perp_s} \setminus C)$ and the code Q_p has minimum distance as Q reduced by one. So, the minimum weight of $D_p^{\perp_s}\backslash C_p$ is at least the minimum weight of $(D^{\perp_s}\backslash C)-1$

$$d_p = \min \operatorname{swt}(D_p^{\perp_s} \backslash C_p)$$

 $\geq \min \operatorname{swt}(D^{\perp_s} \backslash C) - 1 = d - 1$

If the code Q is pure, then $\min \operatorname{swt}(D^{\perp_s}) = d$, therefore, the new code Q_p is pure since $d_p = \min \operatorname{swt}(D_p^{\perp_s}) \geq d$.

We conclude that if there is a subsystem code with parameters $((n-1, K, qR, \ge d-1))_q$, using [2, Theorem 2.], there exists a code with parameters $((n-1, qK, R, \ge d-1))_q$.

C. Reducing Dimension

We also can reduce dimension of the subsystem code for fixed length n and minimum distance d, and still obtain a new subsystem code with improved minimum distance as shown in the following results.

Theorem 12: If a (pure) \mathbf{F}_q -linear $[[n, k, r, d]]_q$ subsystem code Q exists for $d \ge 2$, then there exists an \mathbf{F}_q -linear [n, k-1] $[1, r, d_e]_q$ subsystem code Q_e (pure to d) such that $d_e \geq d$.

Proof: Existence of the $[[n, k, r, d]]_q$ subsystem code Q, implies existence of two additive codes $C \leq \mathbf{F}_q^{2n}$ and $D \leq \mathbf{F}_q^{2n}$ such that $|C| = q^{n-k+r}$ and $|D| = |C \cap C^{\perp_s}| = q^{n-k-r}$. Furthermore, $d = \min \operatorname{swt}(D^{\perp_s} \setminus C)$ and $D \subseteq D^{\perp_s}$.

The idea of the proof comes by extending the code D by some vectors from $D^{\perp_s} \setminus (C \cup C^{\perp_s})$. Let us choose a code D_e of size $|q^{n+1-r-k}| = q|D|$. We also ensure that the code D_e is self-orthogonal. Clearly extending the code D to D_e will extend both the codes C and C^{\perp_s} to C_e and $C^{\perp_s}_e$, respectively. Hence $C_e=q|C|=q^{n+1+r-k}$ and $D_e=C_e\cap C^{\perp_s}_e$.

There exists a subsystem code Q_e stabilized by the code C_e . The result follows by computing parameters of the subsystem code $Q_e = A_e \otimes B_e$.

(i)
$$K_e = q^n / \sqrt{|C_e||D_e|} = q^n / ((q^{n+1+r-k})(q^{n+1-k-r}))^{1/2} = q^{k-1},$$

(ii)
$$R_e = (|C_e|/|D_e|)^{1/2} = ((q^{n+1}R/K)/(q^{n+1}/RK))^{1/2} = q^r,$$

(iii) $d_e = \operatorname{swt}((D_e)^{\perp_s} \setminus C_e) \ge \operatorname{swt}((D^{\perp_s} \setminus C_e)) = d.$ If the

inequality holds, then the code is pure to d.

Arguably, It follows that the set $(D_e^{\perp_s} \setminus C_e)$ is a subset of the set $D^{\perp_s}\setminus C$ because $C\leq C_e$, hence the minimum weight d_e is at least d.

Lemma 13: Suppose an $[[n, k, r, d]]_q$ linear pure subsystem code Q exists generated by the two codes $C, D \leq \mathbf{F}_q^{2n}$. Then there exist linear $[[n-m,k',r',d']]_q$ and $[[n-m,k'+r'-m]]_q$ r'', r'', d']_q subsystem codes with $k' \geq k - m$, $r' \geq r$, $0 \leq r''$ r'' < k' + r', and $d' \ge d$ for any integer m such that there exists a codeword of weight m in $(D^{\perp_s} \setminus C)$.

Proof: [Sketch] This lemma 13 can be proved easily by mapping the subsystem code Q into a stabilizer code. By using [4, Theorem 7.], and the new resulting stabilizer code can be mapped again to a subsystem code with the required parameters.

D. Combining Subsystem Codes

We can also construct new subsystem codes from given two subsystem codes. The following theorem shows that two subsystem codes can be merged together into one subsystem code with possibly improved distance or dimension.

Theorem 14: Let Q_1 and Q_2 be two pure subsystem codes with parameters $[[n_1, k_1, r_1, d_1]]_2$ and $[[n_2, k_2, r_2, d_2]]_2$ for $k_2 + r_2 \leq n_1$, respectively. Then there exists a subsystem code with parameters $[[n_1 + n_2 - k_2 - r_2, k_1 + r_1 - r, r, d]]_2$, where $d \ge min\{d_1, d_1 + d_2 - k_2 - r_2\}$ and $0 \le r < k_1 + r_1$.

Proof: Existence of an $[[n_i, k_i, r_i, d_i]]_2$ pure subsystem code Q_i for $i \in \{1, 2\}$, implies existence of a pure stabilizer code S_i with parameters $[[n_i, k_i + r_i, d_i]]_2$ with $k_2 + r_2 \leq n_1$, see [2]. Therefore, by [4, Theorem 8.], there exists a stabilizer code with parameters $[[n_1 + n_2 - k_2 - r_2, k_1 + r_1, d]]_2, d \ge 1$ $\min\{d_1, d_1+d_2-k_2-r_2\}$. But this code gives us a subsystem code with parameters $[[n_1 + n_2 - k_2 - r_2, k_1 + r_1 - r, r, \ge d]]_2$ with $k_2 + r_2 \le n_1$ and $0 \le r < k_1 + r_1$ that proves the claim.

Theorem 15: Let Q_1 and Q_2 be two pure subsystem codes with parameters $[[n, k_1, r_1, d_1]]_q$ and $[[n, k_2, r_2, d_2]]_q$, respectively. If $Q_2 \subseteq Q_1$, then there exists an $[2n, k_1 + k_2 + r_1 +$ $[r_2 - r, r, d]_q$ pure subsystem code with minimum distance $d \ge \min\{d_1, 2d_2\}$ and $0 \le r < k_1 + k_2 + r_1 + r_2$.

Proof: Existence of a pure subsystem code with parameters $[[n, k_i, r_i, d_i]]_q$ implies existence of a pure stabilizer code with parameters $[[n, k_i+r_i, d_i]]_q$ using [2, Theorem 4.]. But by using [5, Lemma 74.], there exists a pure stabilizer code with parameters $[2n, k_1+k_2+r_1+r_2, d]_q$ with $d \ge \min\{2d_2, d_1\}$. By [2, Theorem 2., Corollary 6.], there must exist a pure subsystem code with parameters $[2n, k_1 + k_2 + r_1 + r_2 - r, r, d]_q$ where $d \ge \min\{2d_2, d_1\}$ and $0 \le r < k_1 + k_2 + r_1 + r_2$, which proves the claim.

We can recall the trace alternative product between two codewords of a classical code and the proof of Theorem 15 can be stated as follows.

Lemma 16: Let Q_1 and Q_2 be two pure subsystem codes with parameters $[[n, k_1, r_1, d_1]]_q$ and $[[n, k_2, r_2, d_2]]_q$, respectively. If $Q_2 \subseteq Q_1$, then there exists an $[2n, k_1 + k_2, r_1 +$ $[r_2,d]_q$ pure subsystem code with minimum distance $d \geq 1$ $\min\{d_1, 2d_2\}.$

Proof: Existence of the code Q_i with parameters $[[n, K_i, R_i, d_i]]_q$ implies existence of two additive codes C_i and D_i for $i \in \{1,2\}$ such that $|C_i| = q^n R_i/K_i$ and $|D_i| = |C \cup C^{\perp_s}| = q^n / R_i K_i.$

We know that there exist additive linear codes $D_i \subseteq D_i^{\perp_a}$, $D_i \subseteq C_i$, and $D_i \subseteq C_i^{\perp_a}$. Furthermore, $D_i = C_i \cap C_i^{\perp_a}$ and $d_i = wt(D_i^{\perp_a} \setminus C_i)$. Also, $C_i = q^{n+r_i-k_i}$ and $|D| = q^{n-r_i-k_i}$.

Using the direct sum definition between to linear codes, let us construct a code D based on D_1 and D_2 as

$$D = \{(u, u + v) \mid u \in D_1, v \in D_2\} \le \mathbf{F}_{a^2}^{2n}.$$

The code D has size of $|D| = q^{2n - (r_1 + r_2 + k_1 + k_2) = |D_1||D_2|}$. Also, we can define the code C based on the codes C_1 and C_2 as

$$C = \{(a, a + b) \mid a \in C_1, b \in C_2\} \le \mathbf{F}_{q^2}^{2n}.$$

The code C is of size $|C| = |C_1||C_2| = q^{2n+r_1+r_2-k_1-k_2}$. But the trace-alternating dual of the code D is

$$D^{\perp_a} = \{ (u' + v'|, v') \mid u' \in D_1^{\perp_a}, v' \in D_2^{\perp_a} \}.$$

We notice that (u'+v',v') is orthogonal to (u,u+v) because, from properties of the product,

$$\langle (u, u + v) \mid (u' + v', v') \rangle_a = \langle u \mid u' + v' \rangle_a + \langle u + v \mid v' \rangle_a$$

= 0

holds for $u \in D_1, v \in D_2, u' \in D_1^{\perp_a}$, and $v' \in D_2^{\perp_a}$.

Therefore, $D\subseteq D^{\perp_a}$ is a self-orthogonal code with respect to the trace alternating product. Furthermore, $C^{\perp_a}=\{(a'+b',b')\mid a'\in C_1^{\perp_a},b'\in C_2^{\perp_a}\}$. Hence, $C\cap C^{\perp_a}=\{(a,a+b)\cap (aa+b',b')\}=D$. Therefore, there exists an \mathbf{F}_q -linear subsystem code $Q=A\otimes B$ with the following parameters.

$$K = |A| = q^{2n}/(|C||D|)^{1/2}$$

$$= \frac{q^{2n}}{\sqrt{(q^{2n}R_1R_2/K_1K_2)(q^{2n}/K_1K_2R_1R_2)}}$$

$$= \frac{q^{2n}}{\sqrt{q^{2n+r_1+r_2-k_1-k_2}q^{2n-r_1-r_2-k_1-k_2}}}$$

$$= q^{k_1k_2} = K_1K_2.$$

- ii) $R = (\frac{|C|}{|D|})^{1/2} = R_1 R_2$.
- iii) the minimum distance is a direct consequence.

Theorem 17: If there exist two pure subsystem quantum codes Q_1 and Q_2 with parameters $[[n_1,k_1,r_1,d_1]]_q$ and $[[n_2,k_2,r_2,d_2]]_q$, respectively. Then there exists a pure subsystem code Q' with parameters $[[n_1+n_2,k_1+k_2+r_1+r_2-r,r,\geq \min(d_1,d_2)]]_q$.

Proof: This Lemma can be proved easily from [2, Theorem 5.] and [5, Lemma 73.]. The idea is to map a pure subsystem code to a pure stabilizer code, and once again map the pure stabilizer code to a pure subsystem code.

Theorem 18: If there exist two pure subsystem quantum codes Q_1 and Q_2 with parameters $[[n_1,k_1,r_1,d_1]]_q$ and $[[n_2,k_2,r_2,d_2]]_q$, respectively. Then there exists a pure subsystem code Q' with parameters $[[n_1+n_2,k_1+k_2,r_1+r_2,\geq \min(d_1,d_2)]]_q$.

Proof: Existence of the code Q_i with parameters $[[n,K_i,R_i,d_i]]_q$ implies existence of two additive codes C_i and D_i for $i\in\{1,2\}$ such that $|C_i|=q^nR_i/K_i$ and $|D_i|=|C\cup C^{\perp_s}|=q^n/R_iK_i$.

Let us choose the codes C and D as follows.

$$C = C_1 \oplus C_2 = \{uv \mid v \in C_1, v \in C_2\},\$$

and

$$D = D_1 \oplus D_2 = \{ab \mid a \in D_1, b \in C_2\},\$$

respectively. From this construction, and since D_1 and D_2 are self-orthogonal codes, it follows that D is also a self-orthogonal code. Furthermore, $D_1 \subseteq C_1$ and $D_2 \subseteq C_2$, then

$$D_1 \oplus D_2 \subseteq C_1 \oplus C_2$$
,

hence $D \subseteq C$. The code C is of size

$$|C| = |C_1||C_2| = q^{(n_1+n_2)-(k_1+k_2)+(r_1+r_2)}$$

= $q^{n_1}q^{n_2}R_1R_2/K_1K_2$

and D is of size

$$|D| = |D_1||D_2| = q^{(n_1+n_2)-(k_1+k_2)-(r_1+r_2)}$$

= $q^{n_1}q^{n_2}/R_1R_2K_1K_2$.

On the other hand.

ii)

$$C^{\perp_s} = (C_1 \oplus C_2)^{\perp_s} = C_2^{\perp_s} \oplus C_1^{\perp_s} \supseteq D_2 \oplus D_1.$$

Furthermore, $C \cap C^{\perp_s} = (C_1 \oplus C_2) \cap (C_2^{\perp_s} \cap C_1^{\perp_s}) = D$.

Therefore, there exists a subsystem code $Q = A \otimes B$ with the following parameters.

$$K = |A| = q^{n_1+n_2}/(|C||D|)^{1/2}$$

$$= \frac{q^{n_1+n_2}}{\sqrt{(q^{n_1+n_2}R_1R_2/K_1K_2)(q^{n_1+n_2}/K_1K_2R_1R_2)}}$$

$$= \frac{q^{n_1+n_2}}{\sqrt{q^{n_1+n_2+r_1+r_2-k_1-k_2}q^{n_1+n_2-r_1-r_2-k_1-k_2}}}$$

$$= q^{k_1k_2} = K_1K_2 = |A_1||A_2|.$$

 $R = \left(\frac{|C|}{|D|}\right)^{1/2} = \sqrt{\frac{q^{n_1}q^{n_2}R_1R_2/K_1K_2}{q^{n_1}q^{n_2}/R_1R_2K_1K_2}}$ $= R_1R_2 = |B_1||B_2|.$

iii) the minimum weight of $D^{\perp_s} \setminus C$ is at least the minimum weight of $D_1^{\perp_s} \setminus C_1$ or $D_2^{\perp_s} \setminus C_2$.

$$d = \min\{\operatorname{swt}(D_1^{\perp_s} \setminus C_1), (D_2^{\perp_s} \setminus C_2)\}$$

$$\geq \min\{d_1, d_2\}.$$

Theorem 19: Given two pure subsystem codes Q_1 and Q_2 with parameters $[[n_1,k_1,r_1,d_1]]_q$ and $[[n_2,k_2,r_2,d_2]]_q$, respectively, with $k_2 \leq n_1$. An $[[n_1+n_2-k_2,k_1+r_1+r_2-r,r,d]]_q$ subsystem code exists such that $d \geq \min\{d_1,d_1+d_2-k_2\}$ and $0 \leq r < k_1+r_1+r_2$.

Proof: The proof is a direct consequence as shown in the previous theorems.

Theorem 20: If an $((n, K, R, d))_{q^m}$ pure subsystem code exists, then there exists a pure subsystem code with parameters $((nm, K, R, \geq d))_q$. Consequently, if a pure subsystem code with parameters $((nm, K, R, \geq d))_q$ exists, then there exist a subsystem code with parameters $((n, K, R, \geq |d/m|))_{q^m}$..

Proof: Existence of a pure subsystem code with parameters $((n,K,R,d))_{q^m}$ implies existence of a pure stabilizer code with parameters $((n,KR,d))_{q^m}$ using [2, Theorem 5.]. By [5, Lemma 76.], there exists a stabilizer code with parameters $((nm,KR,\geq d))_q$. From [2, Theorem 2,5.], there exists a pure subsystem code with parameters $((nm,K,R,\geq d))_q$ that proves the first claim. By [5, Lemma 76.] and [2, Theorem 2,5.], and repeating the same proof, the second claim is a consequence.

III. Special and Short Subsystem Codes
$$[[8,1,2,3]]_2$$
 and $[[6,1,1,3]]_3$

In this section we present the shortest subsystem codes over \mathbf{F}_2 and \mathbf{F}_3 fields. Theorem 5 implies that a stabilizer code with parameters $[[n,k,d]]_q$ gives subsystem codes with parameters $[[n,k-r,r,d]]_q$, see the tables in [1].

$n \setminus k$	k-1	k	k+1
n-1	$[r+2, d-1]_q$	$[\leq r+2,d]_q, [r+1,d-1]_q$	$[r,d-1]_q$
n	$[r+1,d]_q, [r+1,\geq d]_q$	$[r,d]_q \to [\leq r, \geq d]_q$	$[r-1,d]_q$
		$\rightarrow [\geq r, \leq d]_q$	
n+1	$[\geq r, \geq d]_q$	$[\geq r, d]_q$, $[r, \geq d]_q$	_

Fig. 2. Existence of subsystem propagation rules

Consider a stabilizer code with parameters $[[8,3,3]]_2$. This code can be used to derive $[[8,2,1,3]]_2$ and $[[8,1,2,3]]_2$ subsystem codes. We give an explicit construction of these codes. Further, we claim that $[[8,1,2,3]]_2$ and $[[8,2,1,3]]_2$ are the shortest nontrivial binary subsystem codes. We show the stabilizer and normalizer matrices for these codes. Also, we prove their minimum distances using the weight enumeration of these codes. We present two codes with less length, however we can not tolerate more than 2 gauge qubits. The following example shows [[8,1,2,3]] short subsystem code over \mathbf{F}_2 .

Example 21:

$$D_{S} = \begin{bmatrix} X & I & Y & I & Z & Y & X & Z \\ Y & I & Y & X & I & Z & Z & X \\ I & X & Y & Y & Z & X & Z & I \\ I & Y & I & Z & Y & X & X & Z \\ I & I & X & Z & X & Y & Z & Y \end{bmatrix}$$
(4)

$$D_{S}^{\perp} = \begin{bmatrix} X & I & I & I & I & I & Z & Y \\ Y & I & I & I & I & Y & X & X \\ I & X & I & I & I & Y & Y & X \\ I & X & I & I & I & Y & Y & X \\ I & Y & I & I & I & I & X & Z \\ I & I & X & I & I & Y & Z & I \\ I & I & X & I & Y & I & Z & X \\ I & I & I & X & I & Y & Y & Y \\ I & I & I & I & X & I & Y & Z & Z \\ I & I & I & I & Y & Y & Z & Z \\ I & I & I & I & I & Z & X & Y \end{bmatrix}$$

$$(5)$$

$$C_{S} = \begin{bmatrix} X & I & Y & I & Z & Y & X & Z \\ Y & I & Y & X & I & Z & Z & X \\ I & X & Y & Y & Z & X & Z & I \\ I & Y & I & Z & Y & X & X & Z \\ I & I & X & Z & X & Y & Z & Y \\ \hline Y & I & I & I & I & Y & X & X \\ I & X & I & I & I & Y & Y & X \end{bmatrix}$$
(6)

$$C_{S}^{\perp} = \begin{bmatrix} X & I & Y & I & Z & Y & X & Z \\ Y & I & Y & X & I & Z & Z & X \\ I & X & Y & Y & Z & X & Z & I \\ I & Y & I & Z & Y & X & X & Z \\ I & I & X & Z & X & Y & Z & Y \\ \hline X & I & I & I & I & I & Z & Y \\ I & I & I & Y & I & Y & Y & Y \end{bmatrix}$$
(7)

We notice that the matrix D_S generates the code $D=C\cap C^{\perp_s}$. Furthermore, dimensions of the subsystems A and B are

given by $k = \dim D^{\perp_s} - \dim C = (11-7)/2 = 2$ and $r = \dim C - \dim D = (7-5)/2 = 1$. Hence we have $[[8, 2, 1, 3]]_2$ and $[[8, 1, 2, 3]]_2$ subsystem codes.

We show that the subsystem codes $[[8,1,2,3]]_2$ is not better than the stabilizer code $[[8,3,3]]_2$ in terms of syndrome measurement. The reason is that the former needs 8-1-2=5 syndrome measurements, while the later needs also 8-3=5 measurements. This is an obvious example where subsystem codes have no superiority in terms of syndrome measurements.

We post an open question regarding the threshold value and fault tolerant gate operations for this code. We do not know at this time if the code $[[8,1,2,3]]_2$ has better threshold value and less fault-tolerant operations. Also, does the subsystem code with parameters $[[8,1,3,3]]_2$ exist?

No nontrivial $[[7,1,1,3]]_2$ **exists.** There exists a trivial $[[7,1,1,3]]_2$ code obtained by simply extending the $[[7,1,3]]_2$ code as the $[[5,1,3]]_2$ code. We show the smallest subsystem code with length 7 must have at most minimum weight equals to 2. Since $[[7,2,2]]_2$ exists, then we can construct the stabilizer and normalizer matrices as follows.

$$D_{S} = \begin{bmatrix} X & X & X & X & I & I & I \\ Y & Y & Y & Y & I & I & I \\ I & I & I & I & X & I & I \\ I & I & I & I & I & X & I \\ I & I & I & I & I & I & X \end{bmatrix}$$
(8)

$$D_{S}^{\perp} = \begin{bmatrix} X & I & I & X & I & I & I \\ Y & I & I & Y & I & I & I \\ I & X & I & X & I & I & I \\ I & Y & I & Y & I & I & I \\ I & I & X & X & I & I & I \\ I & I & X & X & I & I & I \\ I & I & I & I & X & I & I \\ I & I & I & I & I & X & I \\ I & I & I & I & I & I & X \end{bmatrix}$$
(9)

Clearly, from our construction and using Theorem 5, there must exist a subsystem code with parameters k and r given as follows. $\dim D^{\perp_s} = 9/2$ and $\dim C = 7/2$. Also, $\dim D = 5/2$ and $\min(D^{\perp_s} \setminus C) = 2$. Therefore, , k = (9-7)/2 = 1 and r = (7-5)/2 = 1. Consequently, the parameters of the subsystem code are $[[7,1,1,2]]_2$.

This example shows [[6, 1, 1, 3]] short subsystem code over \mathbf{F}_3 .

Example 22: We give a nontrivial short subsystem code over \mathbf{F}_3 . This is derived from the $[[6, 2, 3]]_3$ graph quantum

code. Also, we show in [1] an example for an $[[6,1,1,3]]_7$ subsystem code over \mathbf{F}_7 . Consider the field \mathbf{F}_3 and let $C \subseteq \mathbf{F}_3^{12}$ be a linear code defined by the following generator matrix.

$$C = \begin{bmatrix} 1 & 0 & 0 & 0 & 2 & 0 & 0 & 2 & 0 & 2 & 0 & 2 \\ 0 & 1 & 0 & 0 & 0 & 2 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 2 & 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 2 & 2 & 0 & 2 & 0 & 2 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 & 0 & 0 & 2 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} \underline{S} \\ \overline{X_1} \\ Z_1 \end{bmatrix}.$$

Let the symplectic inner product $\langle (a|b)|(c|d)\rangle_s = a\cdot d - b\cdot c$. Then the symplectic dual of C is generated by

$$C^{\perp_s} = \begin{bmatrix} S \\ \overline{X_2} \\ \overline{Z_2} \end{bmatrix},$$

where $X_2 = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 2 & 0 & 0 & 0 \end{bmatrix}$ and

We can also have a trivial $[[6,1,1,3]]_2$ code. This trivial extension seems to argue against the usefulness of subsystem codes and if they will really lead to improvement in performance. An obvious open question is if there exist nontrivial $[[6,1,1,3]]_2$ or $[[7,1,1,3]]_2$ subsystem codes.

REFERENCES

- [1] S. A. Aly and A. Klappenecker. Structures and constructions of subsysem codes over finite fields. *Phys. Rev. A.*, 2008. on submission.
- [2] S. A. Aly and A. Klappenecker. Subsystem code constructions. In Proc. 2008 IEEE International Symposium on Information Theory, Toronto, Canada, Submitted, 2008.
- [3] S. A. Aly, A. Klappenecker, and P. K. Sarvepalli. Subsystem codes. In 44th Annual Allerton Conference on Communication, Control, and Computing, Monticello, Illinois, September, 2006, 2006.
- [4] A.R. Calderbank, E.M. Rains, P.W. Shor, and N.J.A. Sloane. Quantum error correction via codes over GF(4). *IEEE Trans. Inform. Theory*, 44:1369–1387, 1998.
- [5] A. Ketkar, A. Klappenecker, S. Kumar, and P. K. Sarvepalli. Nonbinary stabilizer codes over finite fields. *IEEE Trans. Inform. Theory*, 52(11):4892–4914, 2006.