Reformulating SU(N) Yang-Mills theory based on change of variables

Kei-Ichi Kondo,^{\dagger ,1} Toru Shinohara,^{\ddagger ,2} and Takeharu Murakami^{\ddagger ,3}

[†]Department of Physics, Graduate School of Science, Chiba University, Chiba 263-8522, Japan

[†]Graduate School of Science and Technology, Chiba University, Chiba 263-8522, Japan

Abstract

We propose a new version of SU(N) Yang-Mills theory reformulated in terms of new field variables which are obtained by a nonlinear change of variables from the original Yang-Mills gauge field. The reformulated Yang-Mills theory enables us to study the low-energy dynamics by explicitly extracting the topological degrees of freedom such as magnetic monopoles and vortices to clarify the mechanism for quark confinement. The dual superconductivity in Yang-Mills theory is understood in a gauge-invariant manner, as demonstrated recently by a non-Abelian Stokes theorem for the Wilson loop operator, although the basic idea of this reformulation is based on the Cho-Faddeev-Niemi decomposition of the gauge potential.

Key words: Yang-Mills theory, quark confinement, Abelian dominance, magnetic monopole, gluon mass

PACS: 12.38.Aw, 12.38.Lg

³ E-mail: tom@fullmoon.sakura.ne.jp

¹ E-mail: kondok@faculty.chiba-u.jp

² E-mail: sinohara@graduate.chiba-u.jp

Contents

1	Introduction	1
2	Construction of the color vector field2.1 $SU(2)$ case2.2 $SU(3)$ case2.2.1Independent degrees of freedom carried by the color field \boldsymbol{n} 2.2.2Maximal and minimal cases defined by degeneracies2.2.3Minimal case2.2.4Maximal case2.3 $SU(N)$ case2.3.1General consideration2.3.2 $SU(4)$	3 4 6 8 9 10 11 12 12 12 14
3	Reduction condition and change of variables	18
0	3.1Maximal case for $SU(N)$ 3.1.1Decomposition3.1.2Reduction in the maximal case3.2Minimal case for $SU(N)$ 3.2.1Decomposition3.2.2Reduction in the minimal case	20 20 23 26 26 29
4	SU(3) case	31
	 4.1 Maximal case for SU(3)	31 31 33 34 34
	4.2.1 Decomposition for $SU(3)$: minimal case $\ldots \ldots \ldots$	35 36
5	Quantization of Yang-Mills theory5.1Jacobian for the nonlinear change of variables5.2Functional integration	37 37 39
6	Conclusion and discussion	41
\mathbf{A}	SU(N) group	42
в	Derivation of identities B.1 Maximal case B.2 Minimal case	44 44 45
\mathbf{C}	More $SU(3)$ cases	46

1 Introduction

The original Yang-Mills theory [1] is formulated in terms of the Yang-Mills gauge field. This formulation is suitable for studying the high-energy dynamics of Yang-Mills theory. For instance, it is well known that the perturbation theory in the coupling constant developed in terms of the Yang-Mills field is very powerful in the ultraviolet region due to its asymptotic freedom. In the low-energy region, however, we encounter the strong coupling problem, and its description in terms of the Yang-Mills gauge field is no longer valid. In studying the low-energy dynamics of Yang-Mills theory and quantum chromodynamics (QCD), it is important to specify the most relevant degrees of freedom for the phenomenon in question. Quark confinement as a typical phenomenon in the low-energy dynamics caused by strong interactions is believed to be explained by topological defects including magnetic monopoles, vortices and merons. This motivates us to devise another formulation of Yang-Mills theory in terms of new variables reflecting the topological degrees of freedom.

For quark confinement, magnetic monopoles and center vortices are believed to be the dominant topological degrees of freedom. However, Abelian magnetic monopoles [2] or center vortices in SU(N) Yang-Mills theory have been obtained as gauge-fixing defects by partial gauge fixings called the maximal Abelian gauge [2,3] $SU(N) \rightarrow U(1)^{N-1}$ or the maximal center gauge [4] $SU(N) \rightarrow Z(N)$, where $U(1)^{N-1}$ is the maximal torus (Cartan) subgroup and Z(N) is the center subgroup of SU(N). Therefore, the current method of constructing Abelian magnetic monopoles and center vortices cannot avoid charge of the gauge artifact.

In view of this, we take up the Cho-Faddeev-Niemi-Shabanov (CFNS) decomposition of the Yang-Mills field, which was first proposed in 1981 by Cho [5] and has been recently readdressed by Faddeev and Niemi [6] and Shabanov [7]. This machinery enables one to explain and understand some of the low-energy phenomena by separating the contributions due to topological defects in a *gauge-invariant manner*. This feature should be compared with other methods.

By developing the approach founded in CFNS [5–7], we have succeeded in giving a gauge-invariant description of the dual superconductivity in SU(2) Yang-Mills theory in a continuum [8–10] and on a lattice [11–13]. In particular, the Wilson loop operator [14] is expressed exactly in terms of the gauge-invariant magnetic current [15], and the magnetic monopole can be defined in a gauge-invariant manner, which follows from a non-Abelian Stokes theorem for the Wilson loop operator, see [15–17] and references therein. These results enable us to determine the role of a magnetic monopole in confinement.

For the SU(2) gauge group, the CFNS decomposition of the Yang-Mills field and the reformulation of Yang-Mills theory in terms of the resulting new variables are essentially unique. Therefore, we have nothing new to be added for SU(2) gauge group. It is important to note that a unit vector field $\boldsymbol{n}(x)$ called the color field hereafter plays the key role.

For the SU(N) gauge group, the conventional approaches [18–21] introduce r unit vector fields $\mathbf{n}_j(x)$ from the beginning, where r = N - 1 is the rank of the SU(N)group. These r color fields are used to define r gauge-invariant magnetic monopoles. This feature should be compared with the Abelian projection in which r Abelian magnetic monopoles corresponding to every U(1) of the maximal torus subgroup $U(1)^r$ are generated due to the partial gauge fixing $SU(N) \to U(1)^r$. In this paper we point out that such an approach is sufficient, but not necessary for reformulating Yang-Mills theory and that there are other options. In fact, we explicitly construct an option called the *minimal case*. In connection to the minimal case, we call the conventional option [option corresponding to the conventional case] the *maximal case*. Even in the maximal case, we show that there are new options that yield the same conclusion.

The conventional approach suggests that r magnetic monopoles can be the dominant topological degrees of freedom for confinement. However, this is not the inevitable conclusion. As conjectured in [17], it is shown [15] using a non-Abelian Stokes theorem that the SU(N) Wilson loop operator can be rewritten in terms of the gauge-invariant magnetic current corresponding to a single non-Abelian mag*netic monopole* [22] irrespective of N. Therefore, the Wilson loop operator does not require r gauge-dependent Abelian and gauge-invariant Abelian-like magnetic monopoles. Rather, the Wilson loop operator can probe only a single non-Abelian magnetic monopole. From this viewpoint, the minimal option is the most economical way of reformulating Yang-Mills theory, where a single unit color field h is introduced and the gauge-invariant magnetic monopole constructed in this option agrees exactly with the magnetic monopole derived from the Wilson loop operator through a non-Abelian Stokes theorem. The lattice versions of new reformulations for the SU(N)gauge group given in this paper are constructed in |23, 24|, which enable us to perform a numerical simulation on the lattice to study the nonperturbative phenomena in question.

This paper is organized as follows. In section 2, we discuss how to construct the color field for SU(N) gauge group. In this construction we introduce the maximal and minimal options for SU(N), $N \ge 3$. The distinction originates from the degeneracy associated with the diagonalization of the scalar field introduced for determining the color field, which undergoes an adjoint transformation under the gauge transformation. We examine how many degrees of freedom the color field carries from the viewpoint of the change of variables. Moreover, we point out that there are intermediate options other than the maximal and minimal options for SU(N), $N \ge 4$.

In section 3, we show how such a color field defined in section 2 is realized as a functional of the original Yang-Mills field for the maximal and minimal options. We obtain the defining equations, which enable us to specify the new variables as functionals of the original Yang-Mills field. This is important for identifying the color field as one of the new variables obtained by the change of variables from the original Yang-Mills field. We give a prescription called the reduction condition for obtaining the color field as a functional of the original Yang-Mills field.

In section 4, we discuss the physically interesting SU(3) case to explicitly demonstrate the general procedure given in the previous section.

In section 5, we consider a quantum version of the reformulated Yang-Mills theory based on a functional integral method. The key issue is to obtain the Jacobian associated with the change of variables from the original Yang-Mills field to the new variables.

The final section is devoted to a conclusion and discussion. Some of the technical content is summarized in the Appendices.

2 Construction of the color vector field

In the conventional approach, the magnetic monopole in Yang-Mills theory is extracted using the *Abelian projection* [2]. The Abelian projection selects a special direction in color space. Consequently, partial gauge fixing, e.g., the maximal Abelian (MA) gauge [2,3] used so far for realizing the Abelian projection, breaks the color symmetry explicitly. In our reformulation of Yang-Mills theory, the color field $\mathbf{n}(x)$ plays the role of recovering the color symmetry that is lost in the conventional treatment of the Abelian projection, in which the maximal torus subgroup is selected from the original gauge group G.

First of all, we consider how to define the *color field* $\mathbf{n}(x)$ in SU(N) Yang-Mills gauge theory. The color field $\mathbf{n}(x)$ is used to specify only the color direction in the color space at each space-time point $x \in \mathbb{R}^D$ in *D*-dimensional space-time, and its magnitude is irrelevant for our purposes. Therefore, the color field $\mathbf{n}(x)$ is introduced as a unit field; its length is equal to one, i.e., $\mathbf{n}(x) \cdot \mathbf{n}(x) = 1$.

To express the color vector, we use two different notations in this paper: <u>the vector form</u>:

$$\mathbf{n}(x) = (n_A(x))_{A=1}^d = (n_1(x), n_2(x), \cdots, n_d(x)), \quad d \equiv \dim SU(N) = N^2 - 1, \quad (2.1)$$

where d is the dimension of the gauge group G = SU(N). Lie algebra form:

$$\mathbf{n}(x) = n_A(x)T_A \ (A = 1, 2, \cdots, d),$$
 (2.2)

where T_A are the generators of the Lie algebra $\mathscr{G} = su(N)$ of the Lie group G = SU(N). The two notations are equivalent, since the color field is constructed in such a way that it transforms according to the adjoint representation under the action of the gauge group. Hereafter, we use these notations for other fields that transform according to the adjoint representation under the action of the gauge group, e.g.,

$$\dot{\phi}(x) = (\phi_A(x))_{A=1}^d = (\phi_1(x), \phi_2(x), \cdots, \phi_d(x)),$$
(2.3)

$$\phi(x) = \phi_A(x)T_A \ (A = 1, 2, \cdots, d).$$
 (2.4)

It should be understood that T_A denotes the generator in the fundamental representation, unless otherwise stated.

The product of two Lie algebra valued functions, $\mathscr{X} = \mathscr{X}^A T^A$ and $\mathscr{Y} = \mathscr{Y}^A T^A$, is rewritten as the sum of three types of products, which is written in the vector form as

$$\mathscr{X}\mathscr{Y} = \mathscr{X}^{A}\mathscr{Y}^{B}T^{A}T^{B} = \frac{1}{2N}(\mathbf{X}\cdot\mathbf{Y})\mathbf{1} + \frac{1}{2}i\mathbf{X}\times\mathbf{Y} + \frac{1}{2}\mathbf{X}*\mathbf{Y}, \qquad (2.5)$$

where the three types of products are defined by

$$\mathbf{X} \cdot \mathbf{Y} := X^A Y^A, \tag{2.6a}$$

$$\mathbf{X} \times \mathbf{Y} := f^{ABC} X^B Y^C, \tag{2.6b}$$

$$\mathbf{X} * \mathbf{Y} := d^{ABC} X^B Y^C. \tag{2.6c}$$

Here the structure constants f^{ABC} of the Lie algebra su(N) are defined by

$$f^{ABC} = -2i \operatorname{Tr}([T^A, T^B]T^C), \quad (A, B, C \in \{1, 2, \dots, N^2 - 1\})$$
(2.7)

from the commutators among the generators:

$$[T^A, T^B] = i f^{ABC} T^C. (2.8)$$

On the other hand, the anticommutator of the generators:

$$\{T_A, T_B\} = \frac{1}{N} \delta^{AB} \mathbf{1} + d^{ABC} T_C, \qquad (2.9)$$

defines completely symmetric symbols:

$$d^{ABC} = 2\text{Tr}(\{T^A, T^B\}T^C).$$
 (2.10)

In this paper we adopt the following normalization for the generators of su(N):

$$\operatorname{Tr}(T^{A}T^{B}) = \frac{1}{2}\delta^{AB}. \quad (A, B \in \{1, 2, \dots, N^{2} - 1\})$$
 (2.11)

In other words, the Lie algebra is closed under the products \cdot, \times and *.

2.1 SU(2) case

First of all, we consider the SU(2) case as an introductory problem. For SU(2), a Hermitian matrix

$$\phi(x) = \phi_A(x)T_A, \quad T_A = \frac{1}{2}\sigma_A, \ (A = 1, 2, 3)$$
 (2.12)

with 2 by 2 Pauli matrices σ_A can be cast into the diagonal form using a suitable unitary matrix $U \in SU(2)(\subset U(2))$ as

$$U(x)\phi(x)U^{\dagger}(x) = \operatorname{diag}(\Lambda_1(x), \Lambda_2(x)) := \mathbf{\Lambda}(x).$$
(2.13)

The eigenvalues Λ_1 and Λ_2 must be real due to the Hermiticity of ϕ , i.e., $\phi^{\dagger} = \phi$. The traceless condition $\operatorname{tr}(\phi) = \operatorname{tr}(\Lambda) = \Lambda_1 + \Lambda_2 = 0$ yields $\Lambda_1 = -\Lambda_2$. Hence, $\phi(x)$ is written as

$$\boldsymbol{\phi}(x) = U^{\dagger}(x) \operatorname{diag}(\Lambda_1(x), -\Lambda_1(x))U(x) = \Lambda_1(x)U^{\dagger}(x)\sigma_3U(x).$$
(2.14)

For the explicit representation of the matrix

$$\phi(x) = \phi_A(x)T_A = \frac{1}{2} \begin{pmatrix} \phi_3(x) & \phi_1(x) - i\phi_2(x) \\ \phi_1(x) + i\phi_2(x) & -\phi_3(x) \end{pmatrix},$$
(2.15)

its eigenvalues are easily obtained as

$$\Lambda_1(x), \Lambda_2(x) = \pm \frac{1}{2} \sqrt{\phi_1(x)^2 + \phi_2(x)^2 + \phi_3(x)^2} = \pm \frac{1}{2} \sqrt{\phi(x) \cdot \phi(x)} := \pm \frac{1}{2} |\phi(x)|,$$
(2.16)

since $\det(\phi - \lambda \mathbf{1}) = \lambda^2 - \frac{1}{4}(\phi_1^2 + \phi_2^2 + \phi_3^2).$

Imposing the condition of unit length $|\phi(x)| = 1$, we obtain $\Lambda_1(x) = \pm 1/2$, and therefore a unit color field $\mathbf{n}(x)$ is obtained in the form

$$\boldsymbol{n}(x) = \pm U^{\dagger}(x)T_3U(x). \tag{2.17}$$

The unit color field $\boldsymbol{n}(x)$ is also expressed as

$$n(x) = \phi(x)/|\phi(x)|, \quad |\phi(x)| := \sqrt{2\mathrm{tr}(\phi(x)^2)} = \sqrt{\phi(x) \cdot \phi(x)}.$$
 (2.18)

Note that $\phi(x)$ is invariant under the U(1) local transformation in the sense that for $U_{\theta}(x) = e^{iT_{3}\theta}U(x)$,

$$U_{\theta}(x)\phi(x)U_{\theta}^{\dagger}(x) = \Lambda_1(x)e^{iT_3\theta}\sigma_3 e^{-iT_3\theta} = \Lambda_1(x)\sigma_3.$$
(2.19)

Here we consider the Weyl group as a discrete subgroup of SU(2), i.e., the permutation group \mathscr{P}_2 with 2 elements corresponding to the permutations of the 2 bases in the fundamental representation. An element of the Weyl group is expressed as a constant (or uniform) off-diagonal matrix, e.g.,

$$W = \exp\left[i\pi\left(\frac{\sigma_1}{2}\cos\varphi + \frac{\sigma_2}{2}\sin\varphi\right)\right]$$
$$= i\left(\sigma_1\cos\varphi + \sigma_2\sin\varphi\right)$$
$$= i\left(\begin{array}{cc}0 & e^{-i\varphi}\\e^{+i\varphi} & 0\end{array}\right) \in \mathscr{P}_2 \subset SU(2).$$
(2.20)

Therefore, $\phi(x)$ changes its sign under the Weyl transformation: For $U_W(x) = WU(x)$, we have

$$U_W(x)\phi(x)U_W^{\dagger}(x) = \Lambda_1(x)W\sigma_3W^{\dagger} = -\Lambda_1(x)\sigma_3.$$
(2.21)

Changing the sign of $\Lambda(x)$ globally corresponds to changing the order of the diagonal components $\Lambda_k(x)$ globally. We can fix the discrete global symmetry for the Weyl group by imposing an additional condition such as $\Lambda(x) \ge 0$ or the ordering condition $\Lambda_1(x) \ge \Lambda_2(x)$.

Now we consider the situation in which the two eigenvalues become degenerate at a certain space-time point $x_0 \in \mathbb{R}^D$, i.e., $\Lambda_1(x_0) = \Lambda_2(x_0)$. This is realized only when $\Lambda_1(x_0) = \Lambda_2(x_0) = 0$. Therefore, the three conditions $\phi_1(x_0) = \phi_2(x_0) =$ $\phi_3(x_0) = 0$ must be satisfied simultaneously at the degenerate point x_0 . Hence, a set of degenerate points forms a 0-dimensional (pointlike) manifold in \mathbb{R}^3 or a onedimensional (linelike) manifold in \mathbb{R}^4 . Note that the conditions, $\phi_1(x_0) = \phi_2(x_0) =$ $\phi_3(x_0) = 0$, are SU(2) rotation-invariant. The rotation-invariant condition $\phi(x) = 0$ for $\phi(x)$ can be regarded as the singularity condition for the unit vector field $\mathbf{n}(x) =$ $\phi(x)/|\phi(x)|$. Therefore, the degenerate point of $\phi(x)$ appears as the singular point of $\mathbf{n}(x)$. In fact, it is known that a magnetic monopole in Yang-Mills theory can appear as the hedgehog configuration at such a singular point, see, for example, [15].

For G = SU(2), the color field is given by the three-dimensional unit vector

$$\mathbf{n}(x) = (n_1(x), n_2(x), n_3(x)).$$
(2.22)

The unit vector $\mathbf{n}(x) = (n_1(x), n_2(x), n_3(x))$ specifies a two-sphere S^2 at x, which is isomorphic to SU(2)/U(1), since $n_1(x)^2 + n_2(x)^2 + n_3(x)^2 = 1$. The maximal torus group of SU(2) is U(1) and the U(1) local rotation corresponds to the local rotation around the direction of the color field $\mathbf{n}(x)$, which does not change (the direction of) the color vector $\mathbf{n}(x)$ at x. Here, by the local rotation, we mean that the angle of rotation is arbitrary at each spacetime point. This enables us to decompose the group $SU(2)_{local}$ as

$$SU(2)_{\text{local}} = [SU(2)/U(1)]_{\text{local}} \times U(1)_{\text{local}}.$$
 (2.23)

With shown in the above, the color field in the SU(2) case can be constructed using the so-called adjoint orbit representation:

$$\mathbf{n}(x) = U^{\dagger}(x)T_3U(x), \quad U(x) \in SU(2).$$
 (2.24)

It is easy to check for example using the Euler-angle representation of the group element U(x) that the diagonal U(1) part of the group element U(x) does not affect the right-hand side of the adjoint orbit representation. Therefore, $\mathbf{n}(x)$ defined in this way indeed reflects the $[SU(2)/U(1)]_{\text{local}}$ symmetry. The composition of $U(x) \rightarrow$ $U(x)U_{\omega}(x)$ yields the adjoint local rotation

$$\boldsymbol{n}(x) \to \boldsymbol{n}_{\omega}(x) = U_{\omega}^{\dagger}(x)U^{\dagger}(x)T_{3}U(x)U_{\omega}(x) = U_{\omega}^{\dagger}(x)\boldsymbol{n}(x)U_{\omega}(x), \qquad (2.25)$$

whose infinitesimal form is given for $U_{\omega}(x) = \exp(ig\omega(x)) = \exp(igT_A\omega_A(x))$ by

$$\delta_{\omega}\boldsymbol{n}(x) := \boldsymbol{n}_{\omega}(x) - \boldsymbol{n}(x) = ig[\boldsymbol{n}(x), \boldsymbol{\omega}(x)].$$
(2.26)

This is indeed the desired transformation property for the color field under the local rotation.

The conventional Abelian projection in the SU(2) case corresponds to taking the special limit of the color field; the color field is chosen to be uniform in the third direction at whole space-time points:

$$\mathbf{n}(x) \equiv \mathbf{n}_{\infty} := (0, 0, 1) \Longleftrightarrow \mathbf{n}(x) \equiv \mathbf{n}_{\infty} := T_3.$$
(2.27)

Even in this limit, we have still the U(1) local symmetry for the U(1) local rotation around the uniform color field $\mathbf{n}_{\infty} := (0, 0, 1)$, which does not change the uniform color vector. This limit is identified with the partial gauge fixing corresponding to the Abelian projection

$$SU(2)_{\text{local}} \to U(1)_{\text{local}}.$$
 (2.28)

2.2 SU(3) case

We now consider the SU(3) case. We consider a real scalar field $\phi(x)$ taking its value in the Lie algebra su(3) of SU(3):

$$\phi(x) = \phi_A(x)T_A, \quad \phi_A(x) \in \mathbb{R}. \ (A = 1, 2, \cdots, 8)$$
 (2.29)

Thus, ϕ is a traceless and Hermitian matrix for real ϕ_A , i.e., $\operatorname{tr}(\phi) = 0$ and $\phi^{\dagger} = \phi$, since we have chosen the traceless generators to be Hermitian; $\operatorname{tr}(T_A) = 0$ and $(T_A)^{\dagger} = T_A$. The Hermitian matrix ϕ can be cast into the diagonal form Λ using a suitable unitary matrix $U \in SU(3)(\subset U(3))$:

$$U(x)\phi(x)U^{\dagger}(x) = \operatorname{diag}(\Lambda_1(x), \Lambda_2(x), \Lambda_3(x)) := \mathbf{\Lambda}(x), \qquad (2.30)$$

with real elements $\Lambda_1, \Lambda_2, \Lambda_3$. The traceless condition leads to

$$\operatorname{tr}(\boldsymbol{\phi}(x)) = \operatorname{tr}(\boldsymbol{\Lambda}(x)) = \Lambda_1(x) + \Lambda_2(x) + \Lambda_3(x) = 0.$$
(2.31)

The diagonal matrix Λ is expressed as a linear combination of two diagonal generators H_1 and H_2 belonging to the Cartan subalgebra as

$$U(x)\phi(x)U^{\dagger}(x) = a(x)H_1 + b(x)H_2.$$
(2.32)

From (2.30), we obtain the relation

$$2\operatorname{tr}(\boldsymbol{\phi}^{2}) = 2\phi_{A}\phi_{B}\operatorname{tr}(T_{A}T_{B}) = \phi_{A}\phi_{B}\delta^{AB} = \phi_{A}\phi_{A} = \vec{\phi}\cdot\vec{\phi} \quad (A, B = 1, 2, \cdots, 8)$$
$$= 2\operatorname{tr}(\boldsymbol{\Lambda}^{2}) = 2(\Lambda_{1}^{2} + \Lambda_{2}^{2} + \Lambda_{3}^{2}) = a^{2} + b^{2}.$$
(2.33)

At this stage, the color vector field $\mathbf{n}(x)$ defined by $\mathbf{n}(x) = \phi(x)/|\phi(x)|$ is an eight-dimensional unit vector, i.e.,

$$\mathbf{n}(x) \cdot \mathbf{n}(x) = n_A(x)n_A(x) = 2\mathrm{tr}(\mathbf{n}(x)^2) = 1 \ (A = 1, 2, \cdots, 8).$$
(2.34)

In other words, **n** belongs to the 7-sphere, $\mathbf{n} \in S^7$, or the target space of the map is S^7 ; $\mathbf{n} : \mathbb{R}^D \to S^7$. Hence, $\mathbf{n}(x)$ has 7 independent degrees of freedom at each x. However, the group G does not act transitively on the manifold of the target space S^7 . ¹ If the stationary subgroup \tilde{H} of G is nontrivial,² then the transitive target space M is identified with the left coset space, G/\tilde{H} , where $G/\tilde{H} \subset M$. Thus, we have

$$x \in \mathbb{R}^D \to \boldsymbol{n}(x) := \boldsymbol{\phi}(x)/|\boldsymbol{\phi}(x)| \in G/\tilde{H} \subset S^7.$$
 (2.35)

Then the unit color field $\boldsymbol{n}(x) \in su(3)$ is expressed as

$$\boldsymbol{n}(x) = (\cos \vartheta(x))\boldsymbol{n}_1(x) + (\sin \vartheta(x))\boldsymbol{n}_2(x),$$

$$\boldsymbol{n}_1(x) := U^{\dagger}(x)H_1U(x), \quad \boldsymbol{n}_2(x) := U^{\dagger}(x)H_2U(x),$$

(2.36)

where $a^2 + b^2 = 1$ is used to rewrite *a* and *b* in terms of an angle ϑ : $\cos \vartheta(x) = a(x)$, $\sin \vartheta(x) = b(x)$. Note that $n_1(x)$ and $n_2(x)$ are Hermitian and traceless unit fields:

$$\boldsymbol{n}_{j}^{\dagger}(x) = \boldsymbol{n}_{j}(x), \quad \operatorname{tr}(\boldsymbol{n}_{j}(x)) = 0, \quad 2\operatorname{tr}(\boldsymbol{n}_{j}(x)^{2}) = 1 \ (j \in \{1, 2\}).$$
 (2.37)

This is also the case for the color field n(x):

$$\mathbf{n}^{\dagger}(x) = \mathbf{n}(x), \quad \operatorname{tr}(\mathbf{n}(x)) = 0, \quad 2\operatorname{tr}(\mathbf{n}(x)^2) = 1.$$
 (2.38)

It should be remarked that U used in the expression for n_1 and n_2 is regarded as an element of SU(3) rather than U(3), since a diagonal generator of a unit matrix in U(3) commutes trivially with all the other generators of U(3). It is easy to see that the SU(2) case considered in the previous subsection is reproduced in a similar way to the SU(3) case we have just considered.

¹We say that the group G acts transitively on the manifold M if any two elements of M are connected by a group transformation.

²We define the stationary subgroup as a subgroup of G that consists of all the group elements h that leave the reference state (or highest weight state) $|\Lambda\rangle$ invariant up to a phase factor: $h|\Lambda\rangle = |\Lambda\rangle e^{i\phi(h)}$.

For concreteness, we can adopt the Gell-Mann matrices λ_A for SU(3) to denote the generators $T_A = \lambda_A/2$ $(A = 1, \dots, 8)$:

$$\lambda^{1} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda^{2} = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \lambda^{3} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$
$$\lambda^{4} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \lambda^{5} = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \quad \lambda^{5} = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix},$$
$$\lambda^{6} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \lambda^{7} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad \lambda^{8} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}. \quad (2.39)$$

For the Gell-Mann matrices, we have

$$f_{123} = 1,$$

$$f_{147} = f_{246} = f_{257} = f_{345} = -f_{156} = -f_{367} = \frac{1}{2},$$

$$f_{458} = f_{678} = \frac{\sqrt{3}}{2},$$

(2.40)

and

$$d_{118} = d_{228} = d_{338} = -d_{888} = \frac{1}{\sqrt{3}},$$

$$d_{146} = d_{157} = d_{256} = d_{344} = d_{355} = \frac{1}{2},$$

$$d_{247} = d_{366} = d_{377} = -\frac{1}{2},$$

$$d_{448} = d_{558} = d_{668} = d_{778} = -\frac{1}{2\sqrt{3}}.$$

(2.41)

For this choice: $H_1 = T_3 = \lambda_3/2$ and $H_2 = T_8 = \lambda_8/2$,

$$a(x) = 2\Lambda_1(x) + \Lambda_3(x) = -2\Lambda_2(x) - \Lambda_3(x), \quad b(x) = -\sqrt{3}\Lambda_3(x).$$
(2.42)

Here we use the notation n_3, n_8 instead of n_1, n_2 :

$$\boldsymbol{n}_3(x) = U^{\dagger}(x)T_3U(x) = \boldsymbol{n}_1(x), \quad \boldsymbol{n}_8(x) = U^{\dagger}(x)T_8U(x) = \boldsymbol{n}_2(x).$$
 (2.43)

2.2.1 Independent degrees of freedom carried by the color field n

From the viewpoint of reformulating the theory, it is important to recognize which are the independent degrees of freedom to describe the theory in question. For SU(3)with a rank of two, it appears to be convenient to introduce two unit vector fields $\mathbf{n}_3(x)$ and $\mathbf{n}_8(x)$. In fact, this choice has been adopted as a conventional approach by many authors. However, we point out that this choice is not necessarily unique from the viewpoint of reformulating the theory, as shown below.

First, we observe that under the three types of products, \mathbf{n}_k (k = 3, 8) satisfy

$$\mathbf{n}_3 \cdot \mathbf{n}_3 = 1, \quad \mathbf{n}_3 \cdot \mathbf{n}_8 = 0, \quad \mathbf{n}_8 \cdot \mathbf{n}_8 = 1, \tag{2.44a}$$

$$\mathbf{n}_3 \times \mathbf{n}_3 = 0, \quad \mathbf{n}_3 \times \mathbf{n}_8 = 0, \quad \mathbf{n}_8 \times \mathbf{n}_8 = 0, \tag{2.44b}$$

and

$$\mathbf{n}_3 * \mathbf{n}_3 = \frac{1}{\sqrt{3}} \mathbf{n}_8, \quad \mathbf{n}_3 * \mathbf{n}_8 = \frac{1}{\sqrt{3}} \mathbf{n}_3, \quad \mathbf{n}_8 * \mathbf{n}_8 = \frac{-1}{\sqrt{3}} \mathbf{n}_8,$$
 (2.44c)

where the equations (2.44c) are also written as $\mathbf{n}_i * \mathbf{n}_j = d^{ijk}\mathbf{n}_k$, $i, j, k \in \{3, 8\}$. ³ For SU(3), we find from (2.44a), (2.44b) and (2.44c) that \mathbf{n}_3 and \mathbf{n}_8 constitute a closed set of variables under all multiplications: \cdot, \times , and *. In particular, it is important to note that \mathbf{n}_8 can be constructed from \mathbf{n}_3 by the multiplication *, i.e., $\mathbf{n}_8 = \sqrt{3}\mathbf{n}_3 * \mathbf{n}_3$. (In other words, \mathbf{n}_8 is a composite of \mathbf{n}_3 .) In view of this, only the field \mathbf{n}_3 is an independent variable. This interpretation does not necessarily agree with the conventional approach. This case will be called the *maximal case*.

On the other hand, we find that the field \mathbf{n}_8 is closed under self-multiplication:

$$\mathbf{n}_8 \cdot \mathbf{n}_8 = 1, \tag{2.46a}$$

$$\mathbf{n}_8 \times \mathbf{n}_8 = 0, \tag{2.46b}$$

$$\mathbf{n}_8 * \mathbf{n}_8 = \frac{-1}{\sqrt{3}} \mathbf{n}_8. \tag{2.46c}$$

This constitutes a novel option, which will be called the *minimal case*.

This observation suggests that in the maximal or minimal case for SU(3), one of the fields, \mathbf{n}_3 or \mathbf{n}_8 , is sufficient as a representative of the color field \boldsymbol{n} to rewrite Yang-Mills theory based on the change of variables, and that the choice of \mathbf{n}_3 or \mathbf{n}_8 as a fundamental variable can be used an alternative for the equivalent reformulation of Yang-Mills theory. This statement will be verified in the next section.

It should be noted, however, that \mathbf{n}_3 carry 6 degrees of freedom, while \mathbf{n}_8 carry 4 degrees of freedom. Therefore, the other fields to be introduced for rewriting Yang-Mills theory must provide the remaining degrees of freedom in each case. In particular, for the SU(2) gauge group, \boldsymbol{n} carries always 2 degrees of freedom and there is no distinction between the maximal and minimal cases.

For $N \ge 4$, there exist intermediate cases other than the maximal and minimal cases. To see this, it is instructive to consider the SU(4) case explicitly, as outlined in the next section. This also clarifies the meaning of this type of classification and helps to correct a misunderstanding in the previous work [7] regarding the number of independent degrees of freedom.

2.2.2 Maximal and minimal cases defined by degeneracies

We now show that the unit color field n is classified into two categories, maximal and minimal, according to the degeneracies of the eigenvalues. By taking into account the fact that the three eigenvalues obey two equations,

$$\Lambda_1(x) + \Lambda_2(x) + \Lambda_3(x) = 0, \quad \Lambda_1(x)^2 + \Lambda_2(x)^2 + \Lambda_3(x)^2 = \frac{1}{2}, \quad (2.47)$$

³This is equivalent to the relationship between n_3 and n_8 of

$$\{\boldsymbol{n}_{i}, \boldsymbol{n}_{j}\} = \frac{1}{3}\delta^{ij}\mathbf{1} + d^{ijk}\boldsymbol{n}_{k}, \quad i, j, k \in \{3, 8\},$$
(2.45a)

following from the identity

$$\{T_i, T_j\} = \frac{1}{3}\delta^{ij}\mathbf{1} + d^{ijk}T_k, \quad i, j, k \in \{3, 8\},$$
(2.45b)

where we have used the identity for the anticommutator of the generators (2.9) and $d^{ija} = 2\text{Tr}(\{T_i, T_j\}T_a) = 0$ ($a \in \{1, 2, 4, 5, 6, 7\}$) from (2.10).

we find that only one degree of freedom is independent, which corresponds to the choice of ϑ . Therefore, the category to which n belongs is specified by the value of ϑ .

(I) Maximal case in which three eigenvalues $\Lambda_1, \Lambda_2, \Lambda_3$ of Λ are distinct. In the maximal case, the stationary subgroup of \boldsymbol{n} is

$$\tilde{H} = U(1) \times U(1), \qquad (2.48)$$

and \boldsymbol{n} covers the six-dimensional internal or target space $SU(3)/(U(1) \times U(1))$, i.e., the flag space F_2 :

$$\mathbf{n} \in G/\tilde{H} = SU(3)/(U(1) \times U(1)) = F_2.$$
 (2.49)

The maximal cases are realized for any angle $\vartheta \in [0, 2\pi)$ except for 6 values of ϑ in the minimal case, i.e., $\vartheta \notin \{\frac{1}{6}\pi, \frac{1}{2}\pi, \frac{5}{6}\pi, \frac{7}{6}\pi, \frac{3}{2}\pi, \frac{11}{6}\pi\}$.

(II) Minimal case in which two of the three eigenvalues are equal. In the minimal case, the stationary subgroup of n becomes

$$\tilde{H} = U(2), \tag{2.50}$$

and **n** covers only the four dimensional internal or target space SU(3)/U(2), i.e., the complex projective space $P^2(\mathbf{C})$:

$$\mathbf{n} \in G/\tilde{H} = SU(3)/U(2) = P^2(\mathbf{C}).$$
 (2.51)

The minimal cases are exhausted by 6 values of ϑ , i.e., $\vartheta = \frac{(2n-1)}{6}\pi$ $(n = 1, 2, \dots, 6)$ or $\vartheta \in \{\frac{1}{6}\pi, \frac{1}{2}\pi, \frac{5}{6}\pi, \frac{7}{6}\pi, \frac{3}{2}\pi, \frac{11}{6}\pi\}$

These definitions for maximal and minimal cases based on the degeneracies of eigenvalues are equivalent to those given in the previous subsection based on multiplication properties, as shown in the following.

2.2.3 Minimal case

According to (II) in the above definition, the minimal cases are given by $\vartheta = \frac{(2n-1)}{6}\pi$ $(n = 1, 2, \dots, 6)$.

• $\Lambda_2 = \Lambda_3$: $\Lambda_2 = \Lambda_3 = -\frac{1}{2\sqrt{3}} < \Lambda_1 = \frac{1}{\sqrt{3}} \Longrightarrow (a,b) = \left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right) \Longrightarrow \vartheta = \frac{1}{6}\pi$

•
$$\Lambda_3 = \Lambda_1$$
: $\Lambda_3 = \Lambda_1 = -\frac{1}{2\sqrt{3}} < \Lambda_2 = \frac{1}{\sqrt{3}} \Longrightarrow (a, b) = \left(-\frac{\sqrt{3}}{2}, \frac{1}{2}\right) \Longrightarrow \vartheta = \frac{5}{6}\pi$

•
$$\Lambda_1 = \Lambda_2$$
: $\Lambda_1 = \Lambda_2 = -\frac{1}{2\sqrt{3}} < \Lambda_3 = \frac{1}{\sqrt{3}} \Longrightarrow (a, b) = (0, -1) \Longrightarrow \vartheta = \frac{3}{2}\pi$

and their Weyl reflections (rotation by angle π) in the weight diagram:

• $\Lambda_2 = \Lambda_3$: $\Lambda_2 = \Lambda_3 = \frac{1}{2\sqrt{3}} > \Lambda_1 = -\frac{1}{\sqrt{3}} \Longrightarrow (a, b) = \left(-\frac{\sqrt{3}}{2}, -\frac{1}{2}\right) \Longrightarrow \vartheta = \frac{7}{6}\pi$

•
$$\Lambda_3 = \Lambda_1$$
: $\Lambda_3 = \Lambda_1 = \frac{1}{2\sqrt{3}} > \Lambda_2 = -\frac{1}{\sqrt{3}} \Longrightarrow (a,b) = \left(\frac{\sqrt{3}}{2}, -\frac{1}{2}\right) \Longrightarrow \vartheta = \frac{11}{6}\pi$

•
$$\Lambda_1 = \Lambda_2$$
: $\Lambda_1 = \Lambda_2 = \frac{1}{2\sqrt{3}} > \Lambda_3 = -\frac{1}{\sqrt{3}} \Longrightarrow (a, b) = (0, 1) \Longrightarrow \vartheta = \frac{\pi}{2}$.

Note that each of three two-dimensional vectors (a, b) is proportional to a weight vector for the fundamental representations of 3 and 3^{*} in the weight diagram, $(2/\sqrt{3})\Lambda_j$, see, for example [15]. The total set is invariant under the action of the Weyl group.

In particular, for $(a, b) = (0, \pm 1)$ or equivalently $\vartheta = \pi/2, 3\pi/2$, the color field $\mathbf{n}(x)$ can be written using only $\mathbf{n}_8(x)$, i.e., $\mathbf{n}(x) = \pm \mathbf{n}_8(x)$, and $\mathbf{n}_3(x)$ disappears from the color field. In other minimal cases, $\mathbf{n}_3(x)$ reappears in the color field $\mathbf{n}(x)$, and the $\vartheta = \pi/2, 3\pi/2$ cases may appear to be special and distinct from other cases. As the structure of the degenerate matrix Λ indicates, however, the color field $\mathbf{n}(x)$ has the same degrees of freedom in all the minimal cases. Therefore, the difference is apparent due to the special choice of the Gell-Mann matrices. In fact, we find that $\mathbf{n}_8(x)$ covers only the four dimensional internal space SU(3)/U(2), i.e., the complex projective space $P^2(\mathbf{C})$. Also,

$$\mathbf{n}_8 = U^{\dagger} T_8 U = U^{\dagger} H_2 U \in SU(3)/U(2) = P^2(\mathbf{C}),$$
 (2.52)

in agreement with the above definition (II). This originates from the fact that the $SU(2) \times U(1)_8 = U(2)$ rotations caused by four generators $\{T_1, T_2, T_3, T_8\}$ do not change (the direction of) $\boldsymbol{n}_8^{\infty} := T_8$, since the only generators commuting with T_8 are the four generators $\{T_1, T_2, T_3, T_8\}$ specified by the standard Gell-Mann matrices:

$$[T_8, T_A] = 0. \quad (A = 1, 2, 3, 8) \tag{2.53}$$

Thus, all choices of $\vartheta \in \{\frac{1}{6}\pi, \frac{1}{2}\pi, \frac{5}{6}\pi, \frac{7}{6}\pi, \frac{3}{2}\pi, \frac{11}{6}\pi\}$ should be treated on an equal footing, but the apparently simplest way to define the color field in the minimal case is to choose $\vartheta = \pi/2$, i.e., $\mathbf{n}(x) = \mathbf{n}_8(x)$.

2.2.4 Maximal case

In particular, for $(a, b) = (\pm 1, 0)$ or equivalently $\vartheta = 0, \pi$, the color field $\mathbf{n}(x)$ can be written using only $\mathbf{n}_3(x)$, i.e., $\mathbf{n}(x) = \pm \mathbf{n}_3(x)$, and $\mathbf{n}_8(x)$ disappears from the expression of the color field $\mathbf{n}(x)$. In the other maximal cases with $\vartheta \neq 0, \pi, \mathbf{n}(x)$ contains both $\mathbf{n}_3(x)$ and $\mathbf{n}_8(x)$. The appearances of the representation $\mathbf{n}(x)$ are considerably different from each other in the two cases. However, both reveal the same physical situation because $\mathbf{n}_8(x)$ is constructed from $\mathbf{n}_3(x)$ as $\mathbf{n}_8(x) = \sqrt{3}\mathbf{n}_3(x)*\mathbf{n}_3(x)$. Therefore, it is sufficient for us to consider $\mathbf{n}_3(x)$ to define the color field $\mathbf{n}(x)$. In fact, $\mathbf{n}_3(x)$ covers the six-dimensional internal or target space $SU(3)/(U(1) \times U(1))$, i.e., the flag space F_2 :

$$\mathbf{n}_3 = U^{\dagger} T_3 U = U^{\dagger} H_1 U \in SU(3) / (U(1) \times U(1)) = F_2, \qquad (2.54)$$

in agreement with the above definition (I). This is understood from the fact that the $U(1) \times U(1)$ rotations caused by the two diagonal generators $\{T_3, T_8\}$ do not change (the direction of) $\boldsymbol{n}_3^{\infty} := T_3$, since the only generators commuting with T_3 are the two generators $\{T_3, T_8\}$:

$$[T_3, T_3] = 0, \quad [T_3, T_8] = 0. \tag{2.55}$$

Thus the easiest way to define the color field in the maximal case is to choose $\vartheta = 0$, i.e., $\mathbf{n}(x) = \mathbf{n}_3(x)$. Of course, this does not prohibit the introduction of both $\mathbf{n}_3(x)$ and $\mathbf{n}_8(x)$ for convenience.⁴

⁴In the original approach for decomposing the SU(3) Yang-Mills gauge field [18], two fields n_3 and n_8 are introduced from the beginning as they were fundamental variables. Therefore, this option

2.3 SU(N) case

2.3.1 General consideration

We consider the su(N)-valued field

$$\phi(x) = \phi_A(x)T_A. \ (A = 1, 2, \cdots, N^2 - 1)$$
(2.56)

Thus, ϕ is a traceless and Hermitian matrix for real ϕ_A , i.e., $\operatorname{tr}(\phi) = 0$ and $\phi^{\dagger} = \phi$, since $\operatorname{tr}(T_A) = 0$ and $(T_A)^{\dagger} = T_A$. Therefore, it can be cast into the diagonal form Λ using a unitary matrix $U \in SU(N) \subset U(N)$:

$$U(x)\phi(x)U^{\dagger}(x) = \operatorname{diag}(\Lambda_1(x), \Lambda_2(x), \cdots, \Lambda_N(x)) := \mathbf{\Lambda}(x), \qquad (2.57)$$

where Λ_a $(a = 1, \dots, N)$ are real and satisfy

$$\operatorname{tr}(\boldsymbol{\phi}(x)) = \operatorname{tr}(\Lambda(x)) = \sum_{j=1}^{N} \Lambda_a(x) = 0.$$
(2.58)

Therefore, the diagonal matrix Λ is expressed as a linear combination of r = N - 1 diagonal generators H_j $(j = 1, 2, \dots, r)$ belonging to the Cartan subalgebra

$$U(x)\phi(x)U^{\dagger}(x) = \sum_{j=1}^{r} a_j(x)H_j,$$
(2.59)

where $a_i(x)$ is obtained as a linear combination of $\Lambda_i(x)$.

By taking into account the relation

$$\vec{\phi} \cdot \vec{\phi} = \phi_A \phi_A$$

$$= 2 \operatorname{tr}(\phi^2) = 2 \operatorname{tr}(U \phi U^{\dagger} U \phi U^{\dagger}) \sum_{j,k=1}^r 2a_j a_k \operatorname{tr}(H_j H_k) = \sum_{j=1}^r a_j^2$$

$$= 2 \operatorname{tr}(\Lambda^2) = 2 \sum_{j=1}^r \Lambda_j^2,$$
(2.60)

the unit color field $\boldsymbol{n}(x) \in su(N)$ satisfying

$$\mathbf{n}(x) \cdot \mathbf{n}(x) = n_A(x)n_A(x) = 2\mathrm{tr}(\mathbf{n}(x)^2) = 1$$
(2.61)

is expressed as

$$\boldsymbol{n}(x) = \sum_{j=1}^{r} a_j(x) \boldsymbol{n}_j(x), \quad \boldsymbol{n}_j(x) := U^{\dagger}(x) H_j U(x), \qquad (2.62)$$

where each coefficient a_j belongs to the (r-1)-sphere S^{r-1} :

$$\sum_{j=1}^{r} a_j(x)^2 = \mathbf{n}(x) \cdot \mathbf{n}(x) = n_A(x)n_A(x) = 2\operatorname{tr}(\mathbf{n}(x)^2) = 1.$$
(2.63)

is included in the maximal case.

Since each field $n_j(x)$ $(j = 1, \dots, r)$ is a traceless and Hermitian unit field:

$$\boldsymbol{n}_{j}^{\dagger}(x) = \boldsymbol{n}_{j}(x), \quad \operatorname{tr}(\boldsymbol{n}_{j}(x)) = 0, \quad 2\operatorname{tr}(\boldsymbol{n}_{j}(x)^{2}) = 1 \ (j = 1, \cdots, r),$$
 (2.64)

the color field n(x) satisfies the same property:

$$\mathbf{n}^{\dagger}(x) = \mathbf{n}(x), \quad \operatorname{tr}(\mathbf{n}(x)) = 0, \quad 2\operatorname{tr}(\mathbf{n}(x)^2) = 1.$$
 (2.65)

Note that the U used for the expression for n_j is regarded as an element of the SU(N) group rather than U(N), since a diagonal generator in U(N) commutes with all other generators of U(N).

At this stage, the color field $\boldsymbol{n}(x)$ is an (N^2-2) -dimensional vector, i.e., $n_A n_A = 1$ $(A = 1, 2, \dots, N^2 - 1)$, or $\boldsymbol{n}(x)$ has $(N^2 - 2)$ degrees of freedom at each x. Therefore, \boldsymbol{n}_j $(j = 1, \dots, r)$ must have $N^2 - 2 - (N - 2) = N^2 - N$ degrees of freedom in total, since a_j $(j = 1, \dots, r)$ are responsible for the remaining (N - 2) degree of freedom. See Appendix A for the fundamental properties of the SU(N) group.

The maximal and minimal cases for SU(N) are defined as follows.

(I) Maximal case in which all eigenvalues $\Lambda_1, \ldots, \Lambda_N$ of Λ are distinct. The stationary subgroup of \boldsymbol{n} is $\tilde{H} = U(1)^{N-1}$, and \boldsymbol{n} covers the $(N^2 - N)$ -dimensional internal or target space $SU(N)/U(1)^{N-1}$, i.e., the flag space F_2 :

$$\boldsymbol{n} \in SU(N)/U(1)^{N-1} = F_{N-1}.$$
 (2.66)

(II) Minimal case in which N-1 out of N eigenvalues of Λ are equal. The stationary subgroup of \boldsymbol{n} becomes $\tilde{H} = U(N-1)$, and \boldsymbol{n} covers only the 2(N-1)-dimensional internal or target space SU(N)/U(N-1), i.e., the complex projective space $P^{N-1}(\mathbb{C})$, which is a submanifold of F_{N-1} :

$$\mathbf{n} \in SU(N)/U(N-1) = CP^{N-1} = P^{N-1}(\mathbb{C}).$$
 (2.67)

For the minimal case of SU(N), we consider

$$\boldsymbol{n}_r = U^{\dagger} H_r U \in SU(N) / U(N-1) = CP^{N-1}, \quad \dim CP^{N-1} = 2(N-1).$$
 (2.68)

The $SU(N-1) \times U(1)_{N^2} = U(N-1)$ rotations caused by $(N-1)^2$ generators $\{T_1, T_2, \dots, T_{(N-1)^2-1}, T_{N^2-1}\}$ do not change the direction of \boldsymbol{n}_r , as only the $(N-1)^2$ generators commute with H_r . Consequently, \boldsymbol{n}_r runs over only the (2N-2)-dimensional internal space $SU(N)/U(N-1) = P^{N-1}(\mathbb{C})$. Therefore, \boldsymbol{n}_r covers the smallest target space. This is the minimal case.

To obtain the largest internal or target space of the color field \boldsymbol{n} , i.e., the (N^2-N) dimensional space $SU(N)/U(1)^{N-1}$, i.e., the flag space F_{N-1} , we must determine the diagonal matrix T in the form

$$\boldsymbol{n} = U^{\dagger}TU \in SU(N)/U(1)^{N-1} = F_{N-1}, \quad \dim F_{N-1} = N(N-1), \quad (2.69)$$

such that all the generators commuting with T are exhausted solely by r = N - 1 diagonal generators H_i (including itself):

$$[H_j, H_k] = 0, (2.70)$$

and the $U(1)^r = U(1)^{N-1}$ rotations caused by the r = N-1 diagonal generators H_j do not change the direction of \boldsymbol{n} (among the N^2-1 generators T_A for the general unitary transformation $U = \exp(i \sum_{A=1}^{N^2-1} \theta_A T_A)$). The diagonal matrix T can be constructed by a linear combination of all diagonal generators H_j such that all diagonal elements in T have distinct values. This is the maximal case.

In addition, there are intermediate cases in which some of the N eigenvalues are degenerate. To see the situation concretely, we examine the SU(4) case.

2.3.2 SU(4)

The SU(4) group has $4^2 - 1 = 15$ generators $T_A = \lambda_A/2$ $(A = 1, 2, \dots, 15)$. Among them, 3 generators T_3, T_8, T_{15} are diagonal in the Gell-Mann representation:

The weight vectors of the fundamental representation 4 of SU(4) are given by

$$\nu^{1} = \left(\frac{1}{2}, \frac{1}{2\sqrt{3}}, \frac{1}{2\sqrt{6}}\right), \quad \nu^{2} = \left(-\frac{1}{2}, \frac{1}{2\sqrt{3}}, \frac{1}{2\sqrt{6}}\right),$$
$$\nu^{3} = \left(0, -\frac{1}{\sqrt{3}}, \frac{1}{2\sqrt{6}}\right), \quad \nu^{4} = \left(0, 0, \frac{-3}{2\sqrt{6}}\right). \tag{2.72}$$

Since rank SU(4) = 3, the root space of SU(4) forms the three-dimensional space (T_3, T_8, T_{15}) in which there are 12 root vectors shown as follows. First, in the $T_3 - T_8$ plane, 6 root vectors are the same as those in SU(3):

$$(-\frac{1}{2}, \frac{\sqrt{3}}{2}, 0) =: \alpha^{2}, \quad (\frac{1}{2}, \frac{\sqrt{3}}{2}, 0) = \alpha^{1} + \alpha^{2},$$

$$(-1, 0, 0) = -\alpha^{1}, \qquad (1, 0, 0) =: \alpha^{1},$$

$$(-\frac{1}{2}, -\frac{\sqrt{3}}{2}, 0) = -\alpha^{1} - \alpha^{2}, \quad (\frac{1}{2}, -\frac{\sqrt{3}}{2}, 0) = -\alpha^{2}.$$

$$(2.73)$$

The other 6 root vectors have nonvanishing components in the T_{15} direction:

$$(\frac{1}{2}, \frac{1}{2\sqrt{3}}, \frac{2}{\sqrt{6}}) = \alpha^1 + \alpha^2 + \alpha^3, \qquad (-\frac{1}{2}, \frac{1}{2\sqrt{3}}, \frac{2}{\sqrt{6}}) = \alpha^2 + \alpha^3, (0, -\frac{1}{\sqrt{3}}, \frac{2}{\sqrt{6}}) =: \alpha^3, \qquad (2.74)$$

with their inversions with respect to the origin

$$(0, \frac{1}{\sqrt{3}}, -\frac{2}{\sqrt{6}}) = -\alpha^3,$$

$$(-\frac{1}{2}, -\frac{1}{2\sqrt{3}}, -\frac{2}{\sqrt{6}}) = -\alpha^2 - \alpha^3,$$

$$(-\frac{1}{2}, -\frac{1}{2\sqrt{3}}, -\frac{2}{\sqrt{6}}) = -\alpha^1 - \alpha^2 - \alpha^3.$$
(2.75)

The angle between two root vectors is $2\pi/3$ or $\pi/2$, as easily seen by calculating their inner product, since all the root vectors have the unit length. Which root vector is positive depends on the definition of the positive root, while the numbers of positive roots and simple roots do not depend on their definitions. For example, the positive roots according to our definition [17] are as follows:

$$(1,0,0), \quad (\frac{1}{2},\frac{\sqrt{3}}{2},0), \quad (-\frac{1}{2},\frac{\sqrt{3}}{2},0), \\ (\frac{1}{2},\frac{1}{2\sqrt{3}},\frac{2}{\sqrt{6}}), \quad (0,-\frac{1}{\sqrt{3}},\frac{2}{\sqrt{6}}), \quad (\frac{1}{2},\frac{1}{2\sqrt{3}},\frac{2}{\sqrt{6}}).$$
(2.76)

The simple roots are $\alpha^1, \alpha^2, \alpha^3$:

$$\alpha^1 := (1, 0, 0), \quad \alpha^2 := \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}, 0\right), \quad \alpha^3 := \left(0, -\frac{1}{\sqrt{3}}, \frac{2}{\sqrt{6}}\right),$$
(2.77)

and other roots are expressed as linear combinations of the simple roots as indicated above. In fact, α^1, α^2 and α^3 satisfy

$$\alpha^{1} \cdot \alpha^{1} = \alpha^{2} \cdot \alpha^{2} = \alpha^{3} \cdot \alpha^{3} = 1, \quad \alpha^{1} \cdot \alpha^{2} = \alpha^{2} \cdot \alpha^{3} = -\frac{1}{2}, \quad \alpha^{1} \cdot \alpha^{3} = 0.$$
 (2.78)

For SU(4), we find that the following six relations hold for the multiplication *:

$$\mathbf{n}_{1} * \mathbf{n}_{1} = \frac{1}{\sqrt{3}} \mathbf{n}_{2} + \frac{1}{\sqrt{6}} \mathbf{n}_{3},$$

$$\mathbf{n}_{1} * \mathbf{n}_{2} = \frac{1}{\sqrt{3}} \mathbf{n}_{1} = \mathbf{n}_{2} * \mathbf{n}_{1},$$

$$\mathbf{n}_{1} * \mathbf{n}_{3} = \frac{1}{\sqrt{6}} \mathbf{n}_{1} = \mathbf{n}_{3} * \mathbf{n}_{1},$$

$$\mathbf{n}_{2} * \mathbf{n}_{2} = \frac{-1}{\sqrt{3}} \mathbf{n}_{2} + \frac{1}{\sqrt{6}} \mathbf{n}_{3},$$

$$\mathbf{n}_{2} * \mathbf{n}_{3} = \frac{1}{\sqrt{6}} \mathbf{n}_{2} = \mathbf{n}_{3} * \mathbf{n}_{2},$$

$$\mathbf{n}_{3} * \mathbf{n}_{3} = \frac{-2}{\sqrt{6}} \mathbf{n}_{3}.$$
(2.79)

The last equation shows that the field \mathbf{n}_3 is closed under self-multiplication by *. This is the minimal case. The last three equations in (2.79) show that the field \mathbf{n}_2 and \mathbf{n}_3 are closed under multiplication. This corresponds to intermediate cases. The first three equations in (2.79) show that \mathbf{n}_1 , \mathbf{n}_2 and \mathbf{n}_3 are closed under multiplication. This constitutes the maximal case. It should be remarked that \mathbf{n}_1 , \mathbf{n}_2 and \mathbf{n}_3 in the maximal case are constructed from the single color field \boldsymbol{n} , which is sufficient for our purposes.

In all cases (maximal, intermediate and minimal), the single color field n(x) is constructed by choosing the appropriate matrix T such that

$$\boldsymbol{n}(x) = U^{\dagger}(x)TU(x) \in G/\tilde{H}.$$
(2.80)

The minimal case occurs when three of the four eigenvalues are equal, which corresponds to the largest stability subgroup $\tilde{H} = U(3)$, e.g.,

$$T_{15} = \frac{1}{2}\lambda^{15} = \frac{1}{2\sqrt{6}} \begin{pmatrix} 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & -3 \end{pmatrix}, \quad \tilde{H} = \frac{U(3) \times U(1)}{U(1)} = U(3).$$
(2.81)

The two intermediate cases are obtained for the stability subgroup $\tilde{H} = U(2) \times SU(2)$, e.g.,

$$T = \frac{1}{2\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & -1 & 0\\ 0 & 0 & 0 & -1 \end{pmatrix} = \frac{\sqrt{2}}{\sqrt{3}} \frac{\lambda_8}{2} + \frac{1}{\sqrt{3}} \frac{\lambda_{15}}{2}, \quad \tilde{H} = \frac{U(2) \times U(2)}{U(1)} = U(2) \times SU(2).$$
(2.82)

and $\tilde{H} = U(2) \times U(1)$, e.g.,

$$T_3 = \frac{1}{2}\lambda^3 = \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & 0\\ 0 & -1 & 0 & 0\\ 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \tilde{H} = \frac{U(1) \times U(1) \times U(2)}{U(1)} = U(1) \times U(2), \quad (2.83)$$

$$T_8 = \frac{1}{2}\lambda^8 = \frac{1}{2\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 & 0\\ 0 & 1 & 0 & 0\\ 0 & 0 & -2 & 0\\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \tilde{H} = \frac{U(2) \times U(1) \times U(1)}{U(1)} = U(2) \times U(1).$$
(2.84)

The maximal case occurs when all (four) eigenvalues are distinct, which corresponds to the smallest stability subgroup $\hat{H} = U(1)^3$, e.g.,

$$T' = \frac{1}{2\sqrt{5}} \begin{pmatrix} 1 & 0 & 0 & 0\\ 0 & 2 & 0 & 0\\ 0 & 0 & -1 & 0\\ 0 & 0 & 0 & -2 \end{pmatrix} = \frac{-\sqrt{5}\lambda_3}{10} + \frac{\sqrt{15}\lambda_8}{6} + \frac{2\sqrt{30}\lambda_{15}}{15} + \frac{U(1)^4}{U(1)} = U(1)^3.$$
(2.85)

Here the matrix has been chosen such that the absolute value of the elements is simple and as small as possible.

In this example, we can see the role of the three fields n_1 , n_2 , and n_3 in constructing the three cases. Note that $\dim SU(4) = 15$ and $\dim G/\tilde{H} = \dim G - \dim \tilde{H}$.

First, we find that T_{15} commutes with T_1, T_2, \dots, T_8 and T_{15} which constitute 9 generators of U(3). Therefore, n_3 spans the 15 - 9 = 6 dimensional space:

$$\boldsymbol{n}_3 = U^{\dagger} H_3 U = U^{\dagger} T_{15} U \in SU(4) / U(3) = P^3(\mathbb{C}), \qquad (2.86)$$

which generates the color field n in the minimal case of SU(4), as shown in (2.81).

Second, we find that T_8 commutes with T_1, T_2, T_3 and T_8 , which constitute 4 generators of U(2), in addition to T_{15} as a generator of U(1). Therefore, n_2 spans the 15-5=10-dimensional space:

$$\boldsymbol{n}_2 = U^{\dagger} H_2 U = U^{\dagger} T_8 U \in SU(4) / (U(2) \times U(1)), \qquad (2.87)$$

which can be used to define the color field \boldsymbol{n} in an intermediate case of SU(4) as shown in (2.84).

Third, we find that T_3 commutes with T_3 , T_8 , T_{13} , T_{14} and T_{15} . Then T_{13} , T_{14} and two linear combinations of T_8 and T_{15} (i.e., $\frac{1}{3}\sqrt{6}T_{15} - \frac{1}{3}\sqrt{3}T_8$ and $-\frac{1}{3}\sqrt{6}T_{15} - \frac{2}{3}\sqrt{3}T_8$) constitute 4 generators of U(2), in addition to T_3 as a generator of U(1). Thus, n_1 spans the 15 - 5 = 10-dimensional space:

$$\boldsymbol{n}_1 = U^{\dagger} H_1 U = U^{\dagger} T_3 U \in SU(4) / (U(2) \times U(1)), \qquad (2.88)$$

which can be adopted as the color field \boldsymbol{n} in an intermediate case of SU(4) as shown in (2.83).

The color field \boldsymbol{n} in the maximal case of SU(4) is obtained by the following linear combination of \boldsymbol{n}_1 , \boldsymbol{n}_2 and \boldsymbol{n}_3 as exemplified in (2.85):

$$\boldsymbol{n} = \frac{-\sqrt{5}}{10}\boldsymbol{n}_1 + \frac{\sqrt{15}}{6}\boldsymbol{n}_2 + \frac{2\sqrt{30}}{15}\boldsymbol{n}_3 \in SU(4)/U(1)^3 = F_3, \quad (2.89)$$

which is consistent with the above consideration based on (2.79).

For larger N, there may exist more intermediate cases, although we do not discuss such cases here.



Figure 1: The relationship between the original Yang-Mills (YM) theory and the reformulated Yang-Mills (YM') theory. A single color field \boldsymbol{n} is introduced to enlarge the original Yang-Mills theory with a gauge group G into the master Yang-Mills (M-YM) theory with the enlarged gauge symmetry $\tilde{G} = G \times G/\tilde{H}$. The reduction conditions are imposed to reduce the master Yang-Mills theory to the reformulated Yang-Mills theory with the equipollent gauge symmetry G'.

3 Reduction condition and change of variables

Our strategy for reformulating Yang-Mills theory is shown in Fig. 1. By introducing a single color field \boldsymbol{n} in the original Yang-Mills theory written in terms of the variable \mathscr{A}_{μ} with an original gauge group G = SU(N), we obtain a gauge theory called the master Yang-Mills theory with the enlarged gauge symmetry $\tilde{G} = G_{\mathscr{A}} \times [G/\tilde{H}]_{\boldsymbol{n}}$, where the degrees of freedom $[G/\tilde{H}]_{\boldsymbol{n}}$ are possessed by the color field \boldsymbol{n} . By imposing a sufficient number of constraints, say, the reduction conditions, ⁵ to eliminate the extra degrees of freedom, the master Yang-Mills theory is reduced to the gauge theory reformulated in terms of new variables with the gauge symmetry G' = SU(N), say the equipollent gauge symmetry,⁶ which is respected by the new variables.

$$G \nearrow (\text{enlargement}) \quad \tilde{G} = G_{\mathscr{A}} \times [G/\tilde{H}]_{\boldsymbol{n}} \quad \searrow (\text{reduction}) \quad G'.$$
(3.1)

The reformulated Yang-Mills theory is written in terms of new variables, i.e., the color field n(x) and the other new fields.

In the reformulated Yang-Mills theory, the color field n(x) is given as a functional of the SU(N) Yang-Mills gauge field $\mathscr{A}_{\mu}(x)$:

$$\boldsymbol{n}(x) = \boldsymbol{n}_{[\mathscr{A}]}(x). \tag{3.2}$$

Other new variables are also obtained from the original $\mathscr{A}_{\mu}(x)$ by a change of variables thanks to the existence of this color field. In fact, we shall show later that imposing the

⁵In previous papers, we called the reduction condition the new maximal Abelian gauge (new MAG). However, this is misleading for SU(N), $N \ge 3$. Therefore, we do not use this terminology in this paper.

⁶In previous papers, we called the equipollent gauge transformation the gauge transformation II.

reduction condition is a relevant prescription for obtaining $\boldsymbol{n}_{[\mathscr{A}]}(x)$. For the moment, therefore, we do not ask how this is achieved and we omit the subscript \mathscr{A} of $\boldsymbol{n}_{[\mathscr{A}]}$ to simplify the notation, keeping the above comments in mind.

We require that the original SU(N) gauge field $\mathscr{A}_{\mu}(x)$ is decomposed into two parts, $\mathscr{V}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$:

$$\mathscr{A}_{\mu}(x) = \mathscr{V}_{\mu}(x) + \mathscr{X}_{\mu}(x), \qquad (3.3)$$

such that $\mathscr{V}_{\mu}(x)$ transforms under the SU(N) gauge transformation $\Omega(x)$ identically to the original gauge field $\mathscr{A}_{\mu}(x)$, while $\mathscr{X}_{\mu}(x)$ transforms identical to an adjoint matter field:

$$\mathscr{V}_{\mu}(x) \to \mathscr{V}_{\mu}'(x) = \Omega(x)[\mathscr{V}_{\mu}(x) + ig^{-1}\partial_{\mu}]\Omega^{\dagger}(x), \qquad (3.4a)$$

$$\mathscr{X}_{\mu}(x) \to \mathscr{X}'_{\mu}(x) = \Omega(x) \mathscr{X}_{\mu}(x) \Omega^{\mathsf{T}}(x),$$
(3.4b)

by way of a single n field, which transforms according to the adjoint representation:

$$\boldsymbol{n}(x) \to \boldsymbol{n}'(x) = \Omega(x)\boldsymbol{n}(x)\Omega^{\dagger}(x).$$
 (3.4c)

The gauge transformation for the new variables is the *equipollent gauge transforma*tion, which should be compared with the original gauge transformation for $\mathscr{A}_{\mu}(x)$:

$$\mathscr{A}_{\mu}(x) \to \mathscr{A}_{\mu}'(x) = \Omega(x) [\mathscr{A}_{\mu}(x) + ig^{-1}\partial_{\mu}]\Omega^{\dagger}(x).$$
(3.5)

In the following, we use the decomposition of the Lie algebra valued function \mathscr{F} into two parts, an \tilde{H} -commutative part $\mathscr{F}_{\tilde{H}}$ and the remaining part $\mathscr{F}_{G/\tilde{H}}$:

$$\mathscr{F}(x) := \mathscr{F}_{\tilde{H}}(x) + \mathscr{F}_{G/\tilde{H}}(x) \quad \mathscr{F}_{\tilde{H}}(x) \in \mathscr{L}(\tilde{H}), \ \mathscr{F}_{G/\tilde{H}}(x) \in \mathscr{L}(G/\tilde{H}),.$$
(3.6)

where

$$[\mathscr{F}_{\tilde{H}}(x), \boldsymbol{n}(x)] = 0. \tag{3.7}$$

In this decomposition, it should be remarked that \tilde{H} does not necessarily agree with the maximal torus subgroup H for SU(N), $N \ge 3$. For SU(2), $\tilde{H} = H$ and the two parts are uniquely specified:

$$\mathscr{F} = \mathscr{F}_H + \mathscr{F}_{G/H}, \quad \mathscr{F}_H \in \mathscr{L}(H), \ \mathscr{F}_{G/H} \in \mathscr{L}(G/H).$$
 (3.8)

This is also written as

$$\mathscr{F} := \mathscr{F}_{\parallel} + \mathscr{F}_{\perp}, \tag{3.9}$$

which corresponds to the well-known decomposition of a vector F into the parallel part F_{\parallel} and perpendicular part F_{\perp} in the vector form:

$$\boldsymbol{F} = \boldsymbol{F}_{\parallel} + \boldsymbol{F}_{\perp} = \boldsymbol{n}(\boldsymbol{n} \cdot \boldsymbol{F}) + \boldsymbol{n} \times (\boldsymbol{F} \times \boldsymbol{n}), \qquad (3.10)$$

which follows from the simple identity, $\boldsymbol{n} \times (\boldsymbol{n} \times \boldsymbol{F}) = \boldsymbol{n}(\boldsymbol{n} \cdot \boldsymbol{F}) - (\boldsymbol{n} \cdot \boldsymbol{n})\boldsymbol{F}$.

3.1 Maximal case for SU(N)

In the maximal case, it is convenient to introduce a set of unit fields $n_j(x)$ $(j = 1, \dots, r)$ using the adjoint orbit representation:

$$\boldsymbol{n}_{j}(x) = U^{\dagger}(x)H_{j}U(x), \quad j \in \{1, 2, \cdots, r\},$$
(3.11)

where $r := \operatorname{rank} SU(N) = N - 1$ and H_j are the Cartan subalgebra. The fields $n_j(x)$ defined in this way are indeed unit vectors, since

$$(\boldsymbol{n}_{j}(x), \boldsymbol{n}_{k}(x)) = 2\operatorname{tr}(\boldsymbol{n}_{j}(x)\boldsymbol{n}_{k}(x)) = 2\operatorname{tr}(U^{\dagger}(x)H_{j}U(x)U^{\dagger}(x)H_{k}U(x))$$
$$= 2\operatorname{tr}(H_{j}H_{k}) = (H_{j}, H_{k}) = \delta_{jk}.$$
(3.12)

These unit vectors mutually commute,

$$[\mathbf{n}_{j}(x), \mathbf{n}_{k}(x)] = 0, \quad j, k \in \{1, 2, \cdots, r\},$$
(3.13)

since H_j are the Cartan subalgebra obeying

$$[H_j, H_k] = 0, \quad j, k \in \{1, 2, \cdots, r\}.$$
(3.14)

3.1.1 Decomposition

In the maximal case, the decomposed fields $\mathscr{V}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$ are specified by two defining equations (conditions):

(I) all $n_j(x)$ are covariant constants in the background $\mathscr{V}_{\mu}(x)$:

$$0 = D_{\mu}[\mathscr{V}]\boldsymbol{n}_{j}(x) := \partial_{\mu}\boldsymbol{n}_{j}(x) - ig[\mathscr{V}_{\mu}(x), \boldsymbol{n}_{j}(x)], \quad (j = 1, 2, \cdots, r)$$
(3.15)

(II) $\mathscr{X}^{\mu}(x)$ does not have the *H*-commutative part, i.e., $\mathscr{X}^{\mu}(x)_{H} = 0$. In other words, $\mathscr{X}_{\mu}(x)$ is orthogonal to all $n_{j}(x)$:

$$(\boldsymbol{n}_{j}(x), \mathscr{X}_{\mu}(x)) := 2 \operatorname{tr}(\boldsymbol{n}_{j}(x) \mathscr{X}_{\mu}(x)) = n_{j}^{A}(x) \mathscr{X}_{\mu}^{A}(x) = 0. \quad (j = 1, 2, \cdots, r) \quad (3.16)$$

It should be remarked that the defining equations are invariant under the gauge transformation (3.4).

First, we determine the \mathscr{X}_{μ} field by solving the defining equations. For this purpose, we take into account the following identity: Any su(N) Lie algebra valued function \mathscr{F} can be decomposed into the *H*-commutative part \mathscr{F}_{H} and the remaining G/H part $\mathscr{F}_{G/H}$ [7,21]:⁷

$$\mathscr{F} := \mathscr{F}_H + \mathscr{F}_{G/H} := \sum_{j=1}^{r=N-1} \boldsymbol{n}_j(\boldsymbol{n}_j, \mathscr{F}) + \sum_{j=1}^{r=N-1} [\boldsymbol{n}_j, [\boldsymbol{n}_j, \mathscr{F}]].$$
(3.18)

The derivation of this identity is given in Appendix B. In proving this identity, we have used the identification (3.11).

$$\delta^{AB} = n_j^A n_j^B - f^{ACD} n_j^C f^{DEB} n_j^E.$$
(3.17)

⁷This identity is equivalent to the identity [19]

We apply the identity (3.18) to \mathscr{X}_{μ} and use the second defining equation (3.16) to obtain

$$\mathscr{X}_{\mu} = \sum_{j=1}^{r} (\mathscr{X}_{\mu}, \boldsymbol{n}_{j}) \boldsymbol{n}_{j} + \sum_{j=1}^{r} [\boldsymbol{n}_{j}, [\boldsymbol{n}_{j}, \mathscr{X}_{\mu}]] = \sum_{j=1}^{r} [\boldsymbol{n}_{j}, [\boldsymbol{n}_{j}, \mathscr{X}_{\mu}]].$$
(3.19)

Then we take into account the first defining equation:

$$\mathcal{D}_{\mu}[\mathcal{A}]\boldsymbol{n}_{j} = \partial_{\mu}\boldsymbol{n}_{j} - ig[\mathcal{A}_{\mu}, \boldsymbol{n}_{j}]$$

= $\mathcal{D}_{\mu}[\mathcal{V}]\boldsymbol{n}_{j} - ig[\mathcal{X}_{\mu}, \boldsymbol{n}_{j}]$
= $-ig[\mathcal{X}_{\mu}, \boldsymbol{n}_{j}] = ig[\boldsymbol{n}_{j}, \mathcal{X}_{\mu}].$ (3.20)

Thus, $\mathscr{X}_{\mu}(x)$ is expressed in terms of $\mathscr{A}_{\mu}(x)$ and $\boldsymbol{n}_{j}(x)$ as

$$\mathscr{X}_{\mu}(x) = -ig^{-1}\sum_{j=1}^{r} [\boldsymbol{n}_{j}(x), \mathscr{D}_{\mu}[\mathscr{A}]\boldsymbol{n}_{j}(x)].$$
(3.21)

Next, the \mathscr{V}_{μ} field is expressed in terms of $\mathscr{A}_{\mu}(x)$ and $\boldsymbol{n}_{j}(x)$:

$$\mathcal{V}_{\mu}(x) = \mathscr{A}_{\mu}(x) - \mathscr{X}_{\mu}(x)$$
$$= \mathscr{A}_{\mu}(x) + ig^{-1} \sum_{j=1}^{r} [\boldsymbol{n}_{j}(x), \mathscr{D}_{\mu}[\mathscr{A}] \boldsymbol{n}_{j}(x)]$$
$$= \mathscr{A}_{\mu}(x) - \sum_{j=1}^{r} [\boldsymbol{n}_{j}(x), [\boldsymbol{n}_{j}(x), \mathscr{A}_{\mu}(x)]] + ig^{-1} \sum_{j=1}^{r} [\boldsymbol{n}_{j}(x), \partial_{\mu} \boldsymbol{n}_{j}(x)].$$
(3.22)

We now apply the identity (3.18) to \mathscr{A}_{μ} to obtain a simpler form:

$$\mathscr{V}_{\mu}(x) = \sum_{j=1}^{r} (\mathscr{A}_{\mu}(x), \boldsymbol{n}_{j}(x)) \boldsymbol{n}_{j}(x) + ig^{-1} \sum_{j=1}^{r} [\boldsymbol{n}_{j}(x), \partial_{\mu} \boldsymbol{n}_{j}(x)].$$
(3.23)

Thus, $\mathscr{V}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$ are written in terms of $\mathscr{A}_{\mu}(x)$ once $n_{j}(x)$ are given.

It should be remarked that the background field $\mathscr{V}_{\mu}(x)$ contains a part $\mathscr{C}_{\mu}(x)$ which commutes with all $n_i(x)$:

$$[\mathscr{C}_{\mu}(x), \boldsymbol{n}_{j}(x)] = 0. \quad (j = 1, 2, \cdots, r = N - 1)$$
(3.24)

Such an *H*-commutative part (or a parallel part in the vector form) $\mathscr{C}_{\mu}(x)$ in $\mathscr{V}_{\mu}(x)$ is not determined uniquely from the first defining equation (3.15) alone. However, it is determined by the second defining equation as shown above. In view of this, we further decompose $\mathscr{V}_{\mu}(x)$ into $\mathscr{C}_{\mu}(x)$ and $\mathscr{B}_{\mu}(x)$:

$$\mathscr{V}_{\mu}(x) = \mathscr{C}_{\mu}(x) + \mathscr{B}_{\mu}(x). \tag{3.25}$$

Applying the identity (3.18) to $\mathscr{C}_{\mu}(x)$ and by taking into account (3.24), we obtain

$$\mathscr{C}_{\mu}(x) = \sum_{j=1}^{r} (\mathscr{C}_{\mu}(x), \boldsymbol{n}_{j}(x)) \boldsymbol{n}_{j}(x).$$
(3.26)

If the remaining part $\mathscr{B}_{\mu}(x)$, which is not *H*-commutative, i.e., $[\mathscr{B}_{\mu}(x), \mathbf{n}_{j}(x)] \neq 0$, is perpendicular to all $\mathbf{n}_{j}(x)$:

$$(\mathscr{B}_{\mu}(x),\boldsymbol{n}_{j}(x)) = 2\mathrm{tr}(\mathscr{B}_{\mu}(x)\boldsymbol{n}_{j}(x)) = 0, \quad (j = 1, 2, \cdots, r)$$
(3.27)

then we have

$$(\mathscr{A}_{\mu}(x), \boldsymbol{n}_{j}(x)) = (\mathscr{V}_{\mu}(x), \boldsymbol{n}_{j}(x)) = (\mathscr{C}_{\mu}(x), \boldsymbol{n}_{j}(x)).$$
(3.28)

Consequently, the *H*-commutative part $\mathscr{C}_{\mu}(x)$ is

$$\mathscr{C}_{\mu}(x) = \sum_{j=1}^{r} (\mathscr{A}_{\mu}(x), \boldsymbol{n}_{j}(x)) \boldsymbol{n}_{j}(x).$$
(3.29)

and the remaining part $\mathscr{B}_{\mu}(x)$ is determined as ⁸

$$\mathscr{B}_{\mu}(x) = ig^{-1} \sum_{j=1}^{r} [\boldsymbol{n}_{j}(x), \partial_{\mu} \boldsymbol{n}_{j}(x)].$$
(3.30)

In fact, it is easy to verify that this expression indeed satisfies (3.27) and that

$$D_{\mu}[\mathscr{B}]\boldsymbol{n}_{j}(x) = \partial_{\mu}\boldsymbol{n}_{j}(x) - ig[\mathscr{B}_{\mu}(x), \boldsymbol{n}_{j}(x)] = 0 \quad (j = 1, 2, \cdots, r).$$
(3.31)

There are other ways of deriving the same result. 9

Thus, once a full set of color fields $n_j(x)$ is given, the original gauge field has the following decomposition in the Lie algebra form:

$$\mathscr{A}_{\mu}(x) = \mathscr{V}_{\mu}(x) + \mathscr{X}_{\mu}(x) = \mathscr{C}_{\mu}(x) + \mathscr{B}_{\mu}(x) + \mathscr{X}_{\mu}(x), \qquad (3.32a)$$

where each part is expressed in terms of $\mathscr{A}_{\mu}(x)$ and $\mathbf{n}_{j}(x)$ as

$$\mathscr{C}_{\mu}(x) = \sum_{j=1}^{N-1} (\mathscr{A}_{\mu}(x), \boldsymbol{n}_{j}(x)) \boldsymbol{n}_{j}(x) = \sum_{j=1}^{N-1} c_{\mu}^{j}(x) \boldsymbol{n}_{j}(x), \qquad (3.32b)$$

$$\mathscr{B}_{\mu}(x) = ig^{-1} \sum_{j=1}^{N-1} [\boldsymbol{n}_j(x), \partial_{\mu} \boldsymbol{n}_j(x)], \qquad (3.32c)$$

$$\mathscr{X}_{\mu}(x) = -ig^{-1} \sum_{j=1}^{N-1} [\boldsymbol{n}_j(x), \mathscr{D}_{\mu}[\mathscr{A}] \boldsymbol{n}_j(x)].$$
(3.32d)

In what follows, the summation over the index j should be understood when it is repeated, unless otherwise stated.

Equivalently, the decomposition (3.32) is written in the vector form as

$$\mathbf{A}_{\mu}(x) = \mathbf{V}_{\mu}(x) + \mathbf{X}_{\mu}(x) = \mathbf{C}_{\mu}(x) + \mathbf{B}_{\mu}(x) + \mathbf{X}_{\mu}(x), \qquad (3.33a)$$

⁸The SU(2) version in the vector form is expressed as $\mathbf{B}_{\mu}(x) = g^{-1}\partial_{\mu}\mathbf{n}(x) \times \mathbf{n}(x)$.

⁹For example, the same expression for \mathscr{V}_{μ} is also obtained by solving the defining equations as follows. Taking into account the commutator of the first defining equation (3.15) with \mathbf{n}_j , we have $ig^{-1}[\mathbf{n}_j(x), \partial_{\mu}\mathbf{n}_j(x)] = ig^{-1}[\mathbf{n}_j(x), ig[\mathscr{V}_{\mu}(x), \mathbf{n}_j(x)]] = [\mathbf{n}_j(x), [\mathbf{n}_j(x), \mathscr{V}_{\mu}(x)]]$. Then we obtain the relation $\mathscr{V}_{\mu}(x) = \sum_{j=1}^r (\mathscr{V}_{\mu}(x), \mathbf{n}_j(x))\mathbf{n}_j(x) + ig^{-1}\sum_{j=1}^r [\mathbf{n}_j(x), \partial_{\mu}\mathbf{n}_j(x)]$. The second defining equation (3.16) leads to $(\mathscr{A}_{\mu}(x), \mathbf{n}_j(x)) = (\mathscr{V}_{\mu}(x), \mathbf{n}_j(x))$, and hence we have (3.23).

where each part is expressed in terms of $\mathbf{A}_{\mu}(x)$ and $\mathbf{n}_{j}(x)$ as

$$\mathbf{C}_{\mu}(x) := \sum_{j=1}^{N-1} (\mathbf{A}_{\mu}(x) \cdot \mathbf{n}_{j}(x)) \mathbf{n}_{j}(x) = \sum_{j=1}^{N-1} c_{\mu}^{j}(x) \mathbf{n}_{j}(x), \qquad (3.33b)$$

$$\mathbf{B}_{\mu}(x) := g^{-1} \sum_{j=1}^{N-1} (\partial_{\mu} \mathbf{n}_j(x) \times \mathbf{n}_j(x)), \qquad (3.33c)$$

$$\mathbf{X}_{\mu}(x) = g^{-1} \sum_{j=1}^{N-1} (\mathbf{n}_j(x) \times \mathscr{D}_{\mu}[\mathbf{A}] \mathbf{n}_j(x)).$$
(3.33d)

All $n_j(x)$ fields in the maximal case are constructed from a single color field n(x). Therefore, a single color field n(x) is sufficient to specify the decomposition. See section 4.1 for the explicit form.

Finally, we point out that another equivalent expression is obtained in a slightly different way. The differentiation of $n_k(x)$ yields the relation

$$\partial_{\mu}\boldsymbol{n}_{k}(x) = [\partial_{\mu}U^{\dagger}(x)U(x), \boldsymbol{n}_{k}(x)].$$
(3.34)

If we require the covariant constantness for all $n_k(x)$ in the background of $\mathscr{V}_{\mu}(x)$:

$$D_{\mu}[\mathscr{V}]\boldsymbol{n}_{k}(x) := \partial_{\mu}\boldsymbol{n}_{k}(x) - ig[\mathscr{V}_{\mu}(x), \boldsymbol{n}_{k}(x)] = 0, \qquad (3.35)$$

the following relation must hold for all $n_k(x)$ by taking into account the relation (3.34).

$$[\mathscr{V}_{\mu}(x) + ig^{-1}\partial_{\mu}U^{\dagger}(x)U(x), \boldsymbol{n}_{k}(x)] = 0.$$
(3.36)

This shows that $\partial_{\mu} U^{\dagger}(x) U(x) - ig \mathscr{V}_{\mu}(x) \in su(N)$ commutes with all $\mathbf{n}_{k}(x)$ and that it can be written as a linear combination of all \mathbf{n}_{j} :

$$\mathscr{V}_{\mu}(x) + ig^{-1}\partial_{\mu}U^{\dagger}(x)U(x) = \sum_{j=1}^{r} c_{\mu}^{j}(x)\boldsymbol{n}_{j}(x) \in U(1)^{r}, \qquad (3.37)$$

or

$$\mathscr{V}_{\mu}(x) = \sum_{j=1}^{r} \mathbf{n}_{j}(x) c_{\mu}^{j}(x) - ig^{-1} \partial_{\mu} U^{\dagger}(x) U(x) = \sum_{j=1}^{r} \mathbf{n}_{j}(x) c_{\mu}^{j}(x) + ig^{-1} U^{\dagger}(x) \partial_{\mu} U(x).$$
(3.38)

Here \mathscr{V}_{μ} is determined up to the terms parallel to n_j . It is clear that $\mathscr{B}_{\mu}(x)$ corresponds to the pure gauge form.

3.1.2 Reduction in the maximal case

We wish to regard the new variables $\mathbf{n}^A, c^j_\mu, \mathscr{X}^A_\mu$ as those obtained by the change of variables from the original gauge field:

$$\mathscr{A}^{A}_{\mu} \Longrightarrow (\boldsymbol{n}^{A}, c^{j}_{\mu}, \mathscr{X}^{A}_{\mu}). \quad (A = 1, \cdots, N^{2} - 1; j = 1, \dots, N - 1; \mu = 1, \cdots, D)$$
(3.39)

In the maximal case, the naive counting of independent degrees of freedom is as follows.

• $\mathscr{A}_{\mu} \in \mathscr{L}(G) = su(N) \to (N^2 - 1)D$,

•
$$n \in \mathscr{L}(G/H) = su(N) - [u(1) + \dots + u(1)] \rightarrow N^2 - 1 - (N-1) = N^2 - N$$
,

•
$$c_{\mu} \in \mathscr{L}(H) = u(1) + \dots + u(1) \rightarrow (N-1)D_{2}$$

•
$$\mathscr{X}_{\mu} \in \mathscr{L}(G/H) = su(N) - u(N-1) \to (N^2 - 1)D - (N-1)D = (N^2 - N)D.$$

In the decomposition just given, therefore, there is an issue of mismatch for the independent degrees of freedom. In fact, the new variables carry $N^2 - N$ extra degrees of freedom after the decomposition. Therefore, we must eliminate $N^2 - N$ degrees of freedom. For this purpose, we intend to impose $N^2 - N$ constraints to eliminate the extra degrees of freedom.

The transformation properties of the decomposed fields $\mathbf{B}_{\mu}, \mathbf{C}_{\mu}, \mathbf{X}_{\mu}$ are uniquely determined, once we specify those for \mathbf{A}_{μ} and \mathbf{n} , as in the SU(2) case discussed in the previous paper [8]. We consider the infinitesimal version of the enlarged gauge transformation $\delta_{\omega,\theta}$, which is obtained by combining the local transformations for $\delta_{\omega}\mathbf{A}_{\mu}$ and $\delta_{\theta}\mathbf{n}$:

$$\delta_{\omega} \mathbf{A}_{\mu}(x) = D_{\mu}[\mathbf{A}] \boldsymbol{\omega}(x), \quad \delta_{\theta} \mathbf{n}(x) = g \mathbf{n}(x) \times \boldsymbol{\theta}_{\perp}(x), \tag{3.40}$$

where $\boldsymbol{\theta}_{\perp} \in \mathscr{L}(G/H)$.

We propose a constraint, which we call the reduction condition, as follows. To find the reduction condition for G = SU(N), we calculate the SU(N) version of \mathbf{X}^{μ} squared as suggested from the SU(2) case:

$$g^{2}\mathbf{X}^{\mu} \cdot \mathbf{X}_{\mu} = \{\mathbf{n}_{j} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{j}\} \cdot \{\mathbf{n}_{k} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{k}\}$$

$$= D_{\mu}[\mathbf{A}]\mathbf{n}_{k} \cdot [(\mathbf{n}_{j} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \times \mathbf{n}_{k}]$$

$$= D_{\mu}[\mathbf{A}]\mathbf{n}_{k} \cdot [(\mathbf{n}_{k} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \times \mathbf{n}_{j}]$$

$$= D_{\mu}[\mathbf{A}]\mathbf{n}_{k} \cdot [(\mathbf{n}_{j} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{k}) \times \mathbf{n}_{j}]$$

$$= D_{\mu}[\mathbf{A}]\mathbf{n}_{k} \cdot [D_{\mu}[\mathbf{A}]\mathbf{n}_{k} - (\mathbf{n}_{j} \cdot D_{\mu}[\mathbf{A}]\mathbf{n}_{k})\mathbf{n}_{j}]$$

$$= (D_{\mu}[\mathbf{A}]\mathbf{n}_{j})^{2}, \qquad (3.41)$$

where the summation is over j, k and we have used the Jacobi identity in the third and fourth equalities and $\mathbf{n}_j \cdot D_{\mu}[\mathscr{A}]\mathbf{n}_k = 0$ in the last step.

We show that a reduction condition is obtained by minimizing the functional

$$R[\mathbf{A}, \{\mathbf{n}_j\}] := \int d^D x \frac{1}{2} (D_\mu[\mathbf{A}]\mathbf{n}_j)^2, \qquad (3.42)$$

under the enlarged gauge transformation. In fact, the transformation of the integrand $(D_{\mu}[\mathbf{A}]\mathbf{n}_{j})^{2}$ under the infinitesimal enlarged gauge transformation is as follows:

$$\begin{split} \delta_{\omega,\theta} \left\{ \frac{1}{2} (D_{\mu}[\mathbf{A}]\mathbf{n}_{j})^{2} \right\} \\ &= (D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \cdot \delta_{\omega,\theta} (D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \\ &= (D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \cdot \{D_{\mu}[\mathbf{A}]\delta_{\omega,\theta}\mathbf{n}_{j} + g\delta_{\omega,\theta}\mathbf{A}_{\mu} \times \mathbf{n}_{j}\} \\ &= (D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \cdot \{D_{\mu}[\mathbf{A}](g\mathbf{n}_{j} \times \boldsymbol{\theta}_{\perp}) + gD_{\mu}[\mathbf{A}]\boldsymbol{\omega} \times \mathbf{n}_{j}\} \\ &= (D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \cdot \{g(D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \times \boldsymbol{\theta}_{\perp} + g\mathbf{n}_{j} \times D_{\mu}[\mathbf{A}]\boldsymbol{\theta}_{\perp} + gD_{\mu}[\mathbf{A}]\boldsymbol{\omega} \times \mathbf{n}_{j}\} \\ &= g(D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \cdot \{D_{\mu}[\mathbf{A}](\boldsymbol{\omega} - \boldsymbol{\theta}_{\perp}) \times \mathbf{n}_{j}\} \\ &= g(D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \cdot \{D_{\mu}[\mathbf{A}](\boldsymbol{\omega}_{\perp} - \boldsymbol{\theta}_{\perp}) \times \mathbf{n}_{j}\}, \\ &= g(\mathbf{n}_{j} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{j}) \cdot D_{\mu}[\mathbf{A}](\boldsymbol{\omega}_{\perp} - \boldsymbol{\theta}_{\perp}), \end{split}$$
(3.43)

where we have used $D_{\mu}[\mathbf{A}]\boldsymbol{\omega}_{\parallel} = D_{\mu}[\mathbf{A}](\boldsymbol{\omega}_{\parallel}\mathbf{n}) = \partial_{\mu}\boldsymbol{\omega}_{\parallel}\mathbf{n} + \boldsymbol{\omega}_{\parallel}(D_{\mu}[\mathbf{A}]\mathbf{n})$ to obtain the sixth equality.

Therefore, $(D_{\mu}[\mathbf{A}]\mathbf{n}_j)^2$ is invariant under the subset $\boldsymbol{\omega}_{\perp} = \boldsymbol{\theta}_{\perp}$ of the enlarged gauge transformation (3.40). The infinitesimal variation of the functional is

$$\delta_{\omega,\theta} R[\mathbf{A}, \{\mathbf{n}_j\}] = \int d^D x g(\mathbf{n}_j \times D_\mu[\mathbf{A}]\mathbf{n}_j) \cdot D_\mu[\mathbf{A}](\boldsymbol{\omega}_\perp - \boldsymbol{\theta}_\perp)$$

= $-\int d^D x(\boldsymbol{\omega}_\perp - \boldsymbol{\theta}_\perp) \cdot D_\mu[\mathbf{A}]g(\mathbf{n}_j \times D_\mu[\mathbf{A}]\mathbf{n}_j)$
= $-\int d^D x(\boldsymbol{\omega}_\perp - \boldsymbol{\theta}_\perp) \cdot g(\mathbf{n}_j \times D_\mu[\mathbf{A}]D_\mu[\mathbf{A}]\mathbf{n}_j).$ (3.44)

Thus, we obtain the differential form of the reduction condition:

$$\boldsymbol{\chi}[\mathbf{A},\mathbf{n}] := \mathbf{n}_j \times D_{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \mathbf{n}_j \equiv 0.$$
(3.45)

Using the Leibnitz rule for the covariant derivative $D_{\mu}[\mathbf{A}]$,

$$\boldsymbol{\chi}[\mathbf{A},\mathbf{n}] = \mathbf{n}_j \times D_{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \mathbf{n}_j = D_{\mu}[\mathbf{A}] \{\mathbf{n}_j \times D_{\mu}[\mathbf{A}] \mathbf{n}_j\}, \qquad (3.46)$$

the differential reduction condition can also be expressed in terms of \mathbf{V}_{μ} and \mathbf{X}_{μ} in the vector form:

$$\boldsymbol{\chi}[\mathbf{n}, \mathbf{C}, \mathbf{X}] := D^{\mu}[\mathbf{V}]\mathbf{X}_{\mu} \equiv 0, \qquad (3.47)$$

or in the Lie algebra form:

$$0 = \boldsymbol{\chi}[\boldsymbol{n}, \mathscr{A}_{\mu}] := D_{\mu}[\mathscr{V}]\mathscr{X}_{\mu}(x) \equiv \partial_{\mu}\mathscr{X}_{\mu}(x) - ig[\mathscr{V}_{\mu}(x), \mathscr{X}_{\mu}(x)]$$
$$= \partial_{\mu}\mathscr{X}_{\mu} - igc^{j}_{\mu}[\boldsymbol{n}_{j}, \mathscr{X}_{\mu}] - [[\partial_{\mu}\boldsymbol{n}_{j}, \boldsymbol{n}_{j}], \mathscr{X}_{\mu}].$$
(3.48)

Note that $\boldsymbol{\chi} \in SU(N)/U(1)^r$ and the number of conditions for $\boldsymbol{\chi} = (\chi^A) = 0$ $(A = 1, \dots, N^2 - 1)$ is $N^2 - 1 - (N - 1) = N^2 - N$ as expected, since $\boldsymbol{\chi}$ is subject to N - 1 orthogonality conditions:

$$(\boldsymbol{n}_j(x), \boldsymbol{\chi}(x)) = (\boldsymbol{n}_j(x), D_{\mu}[\mathcal{V}] \mathscr{X}_{\mu}(x)) = 0 \quad (j = 1, \cdots, r = N - 1).$$
 (3.49)

This follows from the defining equations (3.16) and (3.15) as

$$(\boldsymbol{n}_{j}, D_{\mu}[\mathscr{V}]\mathscr{X}_{\mu}) = (\boldsymbol{n}_{j}, \partial_{\mu}\mathscr{X}_{\mu}) - ig(\boldsymbol{n}_{j}, [\mathscr{V}_{\mu}, \mathscr{X}_{\mu}])$$

$$= \partial_{\mu}(\boldsymbol{n}_{j}, \mathscr{X}_{\mu}) - (\partial_{\mu}\boldsymbol{n}_{j}, \mathscr{X}_{\mu}) - ig(\boldsymbol{n}_{j}, [\mathscr{V}_{\mu}, \mathscr{X}_{\mu}])$$

$$= -ig([\mathscr{V}_{\mu}, \boldsymbol{n}_{j}], \mathscr{X}_{\mu}) - ig(\boldsymbol{n}_{j}, [\mathscr{V}_{\mu}, \mathscr{X}_{\mu}]) = 0.$$
(3.50)

By solving the differential reduction condition for a given $\mathbf{A}_{\mu}(x)$, the color field $\mathbf{n}(x)$ is in principle obtained, thereby, $\mathbf{n}(x)$ is obtained as a functional of the original gauge field $\mathbf{A}_{\mu}(x)$.

3.2 Minimal case for SU(N)

We now discuss general features of the minimal case for SU(N). In this case, \mathscr{A}_{μ} is decomposed into $\mathscr{V}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$, i.e., $\mathscr{A}_{\mu}(x) = \mathscr{V}_{\mu}(x) + \mathscr{X}_{\mu}(x)$, using only a single color field $\mathbf{h}(x)$ without other fields $\mathbf{n}_{j}(x)$.¹⁰ Here we require that $\mathscr{V}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$ are expressed in terms of $\mathscr{A}_{\mu}(x)$ and $\mathbf{h}(x)$ so as to obey the expected transformation property for the given gauge transformations of $\mathscr{A}_{\mu}(x)$ and $\mathbf{h}(x)$.

3.2.1 Decomposition

The respective components $\mathscr{V}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$ are specified by two defining equations (conditions):

(I) h(x) is a covariant constant in the background $\mathscr{V}_{\mu}(x)$:

$$0 = D_{\mu}[\mathscr{V}]\boldsymbol{h}(x) := \partial_{\mu}\boldsymbol{h}(x) - ig[\mathscr{V}_{\mu}(x), \boldsymbol{h}(x)]; \qquad (3.52)$$

(II) $\mathscr{X}^{\mu}(x)$ does not have the \tilde{H} -commutative part, i.e., $\mathscr{X}^{\mu}(x)_{\tilde{H}} = 0$:

$$\mathscr{X}^{\mu}(x)_{\tilde{H}} := \left(1 - 2\frac{N-1}{N}[\boldsymbol{h}, [\boldsymbol{h}, \cdot]]\right) \mathscr{X}^{\mu}(x) = 0.$$
(3.53)

Note that condition (II) is different from the orthogonality to $\boldsymbol{h}(x)$: $(\mathscr{X}_{\mu}(x), \boldsymbol{h}(x)) := 2\operatorname{tr}(\mathscr{X}_{\mu}(x)\boldsymbol{h}(x)) = \mathscr{X}_{\mu}^{A}(x)\boldsymbol{h}^{A}(x) = 0$, which is not sufficient for characterizing the \tilde{H} -commutative part, in contrast to the *H*-commutative part. This is understood from an identity used in the minimal case, see Appendix. B.2.

First, we apply the second defining equation (3.53) to $\mathscr{X}^{\mu}(x)$:

$$\mathscr{X}_{\mu}(x) = \mathscr{X}_{\mu}(x)_{\tilde{H}} + \frac{2(N-1)}{N} [\boldsymbol{h}, [\boldsymbol{h}, \mathscr{X}_{\mu}(x)]]$$
$$= \frac{2(N-1)}{N} [\boldsymbol{h}, [\boldsymbol{h}, \mathscr{X}_{\mu}(x)]].$$
(3.54)

¹⁰In the Gell-Mann representation of the generators, we adopt the last diagonal matrix $T_{N^2-1} = H_{N-1}$ for defining a single color field:

$$\boldsymbol{h}(x) := \boldsymbol{n}_r(x) = U^{\dagger}(x)H_r U(x). \tag{3.51}$$

By taking into account the first defining equation:

$$\mathcal{D}_{\mu}[\mathscr{A}]\boldsymbol{h} = \mathcal{D}_{\mu}[\mathscr{V}]\boldsymbol{h} - ig[\mathscr{X}_{\mu}, \boldsymbol{h}]$$

= $ig[\boldsymbol{h}, \mathscr{X}_{\mu}],$ (3.55)

 $\mathscr{X}_{\mu}(x)$ is expressed in terms of $\mathscr{A}_{\mu}(x)$ and $\boldsymbol{h}(x)$ as

$$\mathscr{X}_{\mu}(x) = -ig^{-1}\frac{2(N-1)}{N}[\boldsymbol{h}(x), \mathscr{D}_{\mu}[\mathscr{A}]\boldsymbol{h}(x)].$$
(3.56)

Next, the \mathscr{V}_{μ} field is expressed in terms of $\mathscr{A}_{\mu}(x)$ and $\boldsymbol{h}(x)$:

$$\mathscr{V}_{\mu}(x) = \mathscr{A}_{\mu}(x) - \mathscr{X}_{\mu}(x)$$
$$= \mathscr{A}_{\mu}(x) + ig^{-1}\frac{2(N-1)}{N}[\mathbf{h}(x), \mathscr{D}_{\mu}[\mathscr{A}]\mathbf{h}(x)].$$
(3.57)

Thus, $\mathscr{V}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$ are written in terms of $\mathscr{A}_{\mu}(x)$ once h(x) is given as a functional of $\mathscr{A}_{\mu}(x)$.

We further decompose $\mathscr{V}_{\mu}(x)$ into the \tilde{H} -commutative part $\mathscr{C}_{\mu}(x)$ and the remaining part $\mathscr{B}_{\mu}(x)$:

$$\mathscr{V}_{\mu}(x) = \mathscr{C}_{\mu}(x) + \mathscr{B}_{\mu}(x). \tag{3.58}$$

We rewrite (3.57) as

$$\mathscr{V}_{\mu}(x) = \mathscr{A}_{\mu}(x) - \frac{2(N-1)}{N} [\boldsymbol{h}(x), [\boldsymbol{h}(x), \mathscr{A}_{\mu}(x)]] + ig^{-1} \frac{2(N-1)}{N} [\boldsymbol{h}(x), \partial_{\mu} \boldsymbol{h}(x)].$$
(3.59)

The first two terms on the right-hand side of (3.59) together constitute the *H*-commutative part of $\mathscr{A}_{\mu}(x)$, i.e., $\mathscr{A}_{\mu}(x)_{\tilde{H}}$. Therefore, we obtain

$$\mathscr{C}_{\mu}(x) := \mathscr{A}_{\mu}(x)_{\tilde{H}} = \left(1 - 2\frac{N-1}{N}[\boldsymbol{h}(x), [\boldsymbol{h}(x), \cdot]]\right) \mathscr{A}_{\mu}(x)$$
$$= \mathscr{A}_{\mu}(x) - \frac{2(N-1)}{N}[\boldsymbol{h}(x), [\boldsymbol{h}(x), \mathscr{A}_{\mu}(x)]], \qquad (3.60a)$$

$$\mathscr{B}_{\mu}(x) = ig^{-1} \frac{2(N-1)}{N} [\boldsymbol{h}(x), \partial_{\mu} \boldsymbol{h}(x)].$$
(3.60b)

In fact, $\mathscr{C}_{\mu}(x)$ commutes with $\boldsymbol{h}(x)$, as we show in (3.68):

$$[\mathscr{C}_{\mu}(x), \boldsymbol{h}(x)] = 0, \quad (j = 1, 2, \cdots, r = N - 1)$$
(3.61)

and $\mathscr{B}_{\mu}(x)$ is noncommutative, $[\mathscr{B}_{\mu}(x), h(x)] \neq 0$, and is orthogonal to h(x):

$$(\mathscr{B}_{\mu}(x), \boldsymbol{h}(x)) = 2\mathrm{tr}(\mathscr{B}_{\mu}(x)\boldsymbol{h}(x)) = 0.$$
(3.62)

Thus, once a single color field h(x) is given, we have the decomposition

$$\mathscr{A}_{\mu}(x) = \mathscr{V}_{\mu}(x) + \mathscr{X}_{\mu}(x) = \mathscr{C}_{\mu}(x) + \mathscr{B}_{\mu}(x) + \mathscr{X}_{\mu}(x), \qquad (3.63a)$$

$$\mathscr{C}_{\mu}(x) = \mathscr{A}_{\mu}(x) - \frac{2(N-1)}{N} [\boldsymbol{h}(x), [\boldsymbol{h}(x), \mathscr{A}_{\mu}(x)]], \qquad (3.63b)$$

$$\mathscr{B}_{\mu}(x) = ig^{-1} \frac{2(N-1)}{N} [\boldsymbol{h}(x), \partial_{\mu} \boldsymbol{h}(x)], \qquad (3.63c)$$

$$\mathscr{X}_{\mu}(x) = -ig^{-1}\frac{2(N-1)}{N}[\mathbf{h}(x), \mathscr{D}_{\mu}[\mathscr{A}]\mathbf{h}(x)].$$
(3.63d)

Thus, all the new variables have been written in terms of h and \mathscr{A}_{μ} . It turns out that \mathscr{X}_{μ} constructed in this way belongs to the coset

$$\mathscr{X}_{\mu} \in \mathscr{G} - \tilde{\mathscr{H}} = su(N) - u(N-1), \qquad (3.64)$$

since for an arbitrary element $\tilde{h} \in \tilde{H}$

$$\begin{split} (\tilde{\boldsymbol{h}}, \mathscr{X}_{\mu}) =& 2\mathrm{tr}(\tilde{\boldsymbol{h}} \mathscr{X}_{\mu}) \\ =& -i\frac{2(N-1)}{gN} 2\mathrm{tr}(\tilde{\boldsymbol{h}}[\boldsymbol{h}, D_{\mu}[\mathscr{A}]\boldsymbol{h}]) \\ =& -i\frac{2(N-1)}{gN} 2\mathrm{tr}([\tilde{\boldsymbol{h}}, \boldsymbol{h}] D_{\mu}[\mathscr{A}]\boldsymbol{h}) \\ =& 0, \end{split}$$
(3.65)

where in the last step we have used $[\tilde{h}, h] = 0$. Similarly, it is shown that

$$\mathscr{B}_{\mu} \in \mathscr{G} - \mathscr{H} = su(N) - u(N-1).$$
(3.66)

Moreover, we can show that

$$\mathscr{C}_{\mu} \in \tilde{\mathscr{H}} = u(N-1), \tag{3.67}$$

since

$$\begin{aligned} [\boldsymbol{h}, \mathscr{C}_{\mu}] &= [\boldsymbol{h}, \mathscr{A}_{\mu}] - \frac{2(N-1)}{N} [\boldsymbol{h}, [[\mathscr{A}_{\mu}, \boldsymbol{h}], \boldsymbol{h}]] \\ &= [\boldsymbol{h}, \mathscr{A}_{\mu}] - \frac{2(N-1)}{N} \left(\frac{2}{N} [\boldsymbol{h}, \mathscr{A}_{\mu}] + \frac{2(2-N)}{\sqrt{2N(N-1)}} [\boldsymbol{h}, \{\boldsymbol{h}, \mathscr{A}_{\mu}\}] - [\boldsymbol{h}, \{\boldsymbol{h}, \{\boldsymbol{h}, \mathscr{A}_{\mu}\}\}] \right) \\ &= [\boldsymbol{h}, \mathscr{A}_{\mu}] - \frac{2(N-1)}{N} \left(\frac{2}{N} [\boldsymbol{h}, \mathscr{A}_{\mu}] + \frac{2(2-N)^{2}}{2N(N-1)} [\boldsymbol{h}, \mathscr{A}_{\mu}] - \frac{(2-N)^{2}}{2N(N-1)} [\boldsymbol{h}, \mathscr{A}_{\mu}] \right) \\ &= [\boldsymbol{h}, \mathscr{A}_{\mu}] - [\boldsymbol{h}, \mathscr{A}_{\mu}] \\ &= 0, \end{aligned}$$
(3.68)

where we have used

$$[[\mathscr{A}, \boldsymbol{h}], \boldsymbol{h}] = \{\mathscr{A}, \{\boldsymbol{h}, \boldsymbol{h}\}\} - \{\boldsymbol{h}, \{\boldsymbol{h}, \mathscr{A}\}\}$$
$$= \frac{2}{N}\mathscr{A} + \frac{2(2-N)}{\sqrt{2N(N-1)}}\{\mathscr{A}, \boldsymbol{h}\} - \{\boldsymbol{h}, \{\boldsymbol{h}, \mathscr{A}\}\},$$
(3.69)

and

$$[\boldsymbol{h}, \{\boldsymbol{h}, \mathscr{A}_{\mu}\}] = [\boldsymbol{h}\boldsymbol{h}, \mathscr{A}_{\mu}] = \frac{(2-N)}{\sqrt{2N(N-1)}} [\boldsymbol{h}, \mathscr{A}].$$
(3.70)

Thus, new variables constructed in this way indeed satisfy the desired property

$$D_{\mu}[\mathscr{V}]\boldsymbol{h} = D_{\mu}[\mathscr{B}]\boldsymbol{h} - ig[\mathscr{C}_{\mu}, \boldsymbol{h}] = 0.$$
(3.71)

It is instructive to note that the above \mathscr{C}_{μ} is written in the form

$$\mathscr{C}_{\mu} = u_{\mu}^{k} \boldsymbol{n}_{k}, \quad u_{\mu}^{k} = (\boldsymbol{n}_{k}, \mathscr{A}_{\mu}), \quad \boldsymbol{n}_{k} = U^{\dagger} T_{k} U \ (T_{k} \in su(N-1)), \tag{3.72}$$

where k runs over $k = 1, \dots, (N-1)^2$ and $N^2 - 1$. Note that these \mathbf{n}_k for $k = 1, \dots, (N-1)^2$ are not uniquely defined. This is because \mathbf{n}_k $(k = 1, \dots, (N-1)^2)$ can be changed using the rotation within $\tilde{H} = U(N-1)$ without changing \mathbf{n}_r , while \mathbf{n}_r is invariant under the $\tilde{H} = U(N-1)$ rotation. This feature is discussed for SU(3) in more detail.

3.2.2 Reduction in the minimal case

The minimal version of the reduction condition is obtained as follows. For G = SU(N), the stability group is $\tilde{H} = U(N-1)$. Therefore, the respective field variable has the following degrees of freedom at each space-time point:

• $\mathscr{A}_{\mu} \in \mathscr{L}(G) = su(N) \to (N^2 - 1)D$,

•
$$h \in \mathscr{L}(G/\tilde{H}) = su(N) - u(N-1) \to (N^2 - 1) - (N-1)^2 = 2(N-1)$$

•
$$\mathscr{C}_{\mu} \in \mathscr{L}(\tilde{H}) = u(N-1) \to (N-1)^2 D_{\mathfrak{Z}}$$

• $\mathscr{X}_{\mu} \in \mathscr{L}(G/\tilde{H}) = su(N) - u(N-1) \to 2(N-1)D.$

If we wish to regard the new variables \mathscr{C}_{μ} , \mathscr{X}_{μ} and h as those obtained from the original field variable \mathscr{A}_{μ} by the non-linear change of variables

$$\mathscr{A}_{\mu} \Longrightarrow (\boldsymbol{h}, \mathscr{C}_{\mu}, \mathscr{X}_{\mu}),$$
 (3.73)

we must give a procedure for eliminating the 2(N-1) extra degrees of freedom. Here we do not include the variable \mathscr{B}_{μ} in this counting, since it is written only in terms of **h**. For this purpose, we impose 2(N-1) reduction conditions $\chi = 0$:

•
$$\boldsymbol{\chi} \in \mathcal{G}/\mathcal{H} = su(N)/u(N-1) \rightarrow 2(N-1).$$

By introducing h in the original SU(N) Yang-Mills theory, the gauge symmetry is enlarged to $SU(N)_{\mathscr{A}} \times [SU(N)/U(N-1)]_h$. The resulting theory is called the master Yang-Mills theory. By imposing an appropriate constraint, say, the minimal version of the reduction condition, the master Yang-Mills theory is reduced to the gauge theory with the gauge symmetry SU(N), say, the equipollent gauge symmetry, which is respected by the new variables. Recall that \mathbf{X}_{μ} is transformed according to the adjoint representation under the equipollent gauge transformation. As in the SU(2) case, therefore, it is expected that such a constraint is given by minimizing the functional

$$\int d^D x \frac{1}{2} g^2 \mathbf{X}_{\mu} \cdot \mathbf{X}^{\mu} = \frac{2(N-1)^2}{N^2} \int d^D x (\boldsymbol{h} \times D_{\mu}[\mathbf{A}]\boldsymbol{h})^2$$
$$= \frac{N-1}{N} \int d^D x (D_{\mu}[\mathbf{A}]\boldsymbol{h})^2, \qquad (3.74)$$

with respect to the enlarged gauge transformation:

$$\delta \mathbf{A}_{\mu} = D_{\mu}[\mathbf{A}]\boldsymbol{\omega}, \quad \delta \boldsymbol{h} = g\boldsymbol{h} \times \boldsymbol{\theta} = g\boldsymbol{h} \times \boldsymbol{\theta}_{\perp}. \quad (\boldsymbol{\omega} \in \mathscr{L}(G), \boldsymbol{\theta}_{\perp} \in \mathscr{L}(G/\tilde{H})) \quad (3.75)$$

In fact, the enlarged gauge transformation of the functional $R[\mathbf{A}, \mathbf{h}]$:

$$R[\mathbf{A}, \boldsymbol{h}] := \int d^D x \frac{1}{2} (D_{\mu}[\mathbf{A}]\boldsymbol{h})^2, \qquad (3.76)$$

is given by

$$\delta R[\mathbf{A}, \mathbf{h}] = \int d^{D}x D_{\mu}[\mathbf{A}] \mathbf{h} \cdot \delta(D_{\mu}[\mathbf{A}] \mathbf{h})$$

$$= \int d^{D}x D_{\mu}[\mathbf{A}] \mathbf{h} \cdot (D_{\mu}[\mathbf{A}] \delta \mathbf{h} + g \delta \mathbf{A}_{\mu} \times \mathbf{h})$$

$$= \int d^{D}x D_{\mu}[\mathbf{A}] \mathbf{h} \cdot (D_{\mu}[\mathbf{A}] (g \mathbf{h} \times \boldsymbol{\theta}_{\perp}) + g D_{\mu}[\mathbf{A}] \boldsymbol{\omega} \times \mathbf{h})$$

$$= g \int d^{D}x D_{\mu}[\mathbf{A}] \mathbf{h} \cdot D_{\mu}[\mathbf{A}] \{\mathbf{h} \times (\boldsymbol{\theta}_{\perp} - \boldsymbol{\omega})\}$$

$$= -g \int d^{D}x D^{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \mathbf{h} \cdot \{\mathbf{h} \times (\boldsymbol{\theta}_{\perp} - \boldsymbol{\omega}_{\perp})\}$$

$$= g \int d^{D}x (\boldsymbol{\theta}_{\perp} - \boldsymbol{\omega}_{\perp}) \cdot (\mathbf{h} \times D^{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \mathbf{h}), \qquad (3.77)$$

where $\boldsymbol{\omega}_{\perp}$ denotes the component of $\boldsymbol{\omega}$ in the direction $\mathscr{L}(G/\tilde{H})$. Thus, we obtain the differential form of the reduction condition:

$$\boldsymbol{\chi}[\mathbf{A}, \boldsymbol{h}] := \boldsymbol{h} \times D^{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \boldsymbol{h} \equiv 0, \qquad (3.78)$$

which is also expressed in terms of \mathbf{X}_{μ} and \mathbf{V}_{μ} (\boldsymbol{h} and \mathbf{C}_{μ}):

$$\boldsymbol{\chi}[\boldsymbol{h}, \mathbf{C}, \mathbf{X}] := D^{\mu}[\mathbf{V}] \mathbf{X}_{\mu} \equiv 0.$$
(3.79)

The form of the constraint (3.78) tells us that $\boldsymbol{\chi} \in \mathscr{L}(G/\tilde{H})$, namely, $\boldsymbol{\chi}$ has no component in the direction $\mathscr{L}(\tilde{H})$. Therefore, $\boldsymbol{\chi}$ gives $(N^2-1)-(N-1)-2=2(N-1)$ independent conditions as desired.

To determine which part of the symmetry is left after imposing the minimal version of the reduction condition (3.78), we perform the gauge transformation on the enlarged gauge-fixing functional χ :

$$\delta \boldsymbol{\chi} = \delta \boldsymbol{h} \times D^{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \boldsymbol{h} + \boldsymbol{h} \times D^{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \delta \boldsymbol{h} + \boldsymbol{h} \times (g \delta \mathbf{A}_{\mu} \times D_{\mu}[\mathbf{A}] \boldsymbol{h}) + \boldsymbol{h} \times D^{\mu}[\mathbf{A}] (g \delta \mathbf{A}_{\mu} \times \boldsymbol{h})$$

$$= (gh \times \theta_{\perp}) \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h + h \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](gh \times \theta_{\perp}) + h \times (gD_{\mu}[\mathbf{A}]\boldsymbol{\omega} \times D_{\mu}[\mathbf{A}]h) + h \times D^{\mu}[\mathbf{A}](gD_{\mu}[\mathbf{A}]\boldsymbol{\omega} \times h) = g(\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) \times \theta_{\perp} + g(D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times \theta_{\perp}) \times h + gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times \theta_{\perp}) + 2gh \times (D^{\mu}[\mathbf{A}]h \times D_{\mu}[\mathbf{A}]\theta_{\perp}) + gh \times (h \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\theta_{\perp}) + gh \times (D^{\mu}[\mathbf{A}]\boldsymbol{\omega} \times D_{\mu}[\mathbf{A}]h) = g(\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\mathbf{h} \times \theta_{\perp} + 2gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) \times \theta_{\perp} + 2gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) + gh \times (h \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\theta_{\perp}) + 2gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) + gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\boldsymbol{\omega} \times h) = g(\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) \times \theta_{\perp} + 2gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) \times \theta_{\perp} + 2gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) + gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\theta_{\perp} - \boldsymbol{\omega}_{\perp})\} + 2gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) + gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\omega_{\parallel} \times h) = g(\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) + gh \times (D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\omega_{\parallel} \times h) = g(\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) \times \theta_{\perp} + 2gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h) \times \theta_{\perp} + 2gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times \omega_{\parallel}) = g(\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times \omega_{\parallel}) = g(\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times \omega_{\parallel}) = g(\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times (\theta_{\perp} + \omega_{\parallel}) + 2gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times (\theta_{\perp} + \omega_{\parallel}) + 2gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times (\theta_{\perp} + \omega_{\parallel}) + 2gh \times \{D^{\mu}[\mathbf{A}]h \times D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times (\theta_{\perp} + \omega_{\parallel}) + 2gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times (\theta_{\perp} + \omega_{\parallel}) + 2gh \times \{D^{\mu}[\mathbf{A}]h \times D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]h \times D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\theta_{\perp} - \omega_{\perp})\} + gh \times \{\mathbf{h} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\theta_$$

where $\boldsymbol{\omega}_{\parallel}$ and $\boldsymbol{\omega}_{\perp}$ denote the components of $\boldsymbol{\omega}$ in $\mathscr{L}(\tilde{H})$ and $\mathscr{L}(G/\tilde{H})$, respectively. ($\boldsymbol{\omega} = \boldsymbol{\omega}_{\parallel} + \boldsymbol{\omega}_{\perp}$). Here we have used

$$0 = D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}](\boldsymbol{\omega}_{\parallel} \times \boldsymbol{h})$$

= $D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\boldsymbol{\omega}_{\parallel} \times \boldsymbol{h} + 2D^{\mu}[\mathbf{A}]\boldsymbol{\omega}_{\parallel} \times D_{\mu}[\mathbf{A}]\boldsymbol{h} + \boldsymbol{\omega}_{\parallel} \times D^{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\boldsymbol{h}.$ (3.81)

This result shows that the minimal version of the reduction condition $\chi \equiv 0$ leaves $\theta_{\perp} = \omega_{\perp}$ intact. When $\theta_{\perp} = \omega_{\perp}$, we find from (3.80)

$$\delta \boldsymbol{\chi} = g \boldsymbol{\chi} \times \boldsymbol{\alpha}, \quad \boldsymbol{\alpha} = (\boldsymbol{\alpha}_{\parallel}, \boldsymbol{\alpha}_{\perp}) = (\boldsymbol{\omega}_{\parallel}, \boldsymbol{\omega}_{\perp} = \boldsymbol{\theta}_{\perp}). \tag{3.82}$$

4 SU(3) case

We now describe the SU(3) Yang-Mills field explicitly for practical purposes.

4.1 Maximal case for SU(3)

4.1.1 Decomposition for SU(3): maximal case

It is obvious that the maximal case of SU(3) goes in the same way as the SU(N) case given above. We introduce two fields:

$$\boldsymbol{n}_3 := U^{\dagger} T_3 U, \quad \boldsymbol{n}_8 := U^{\dagger} T_8 U, \quad U \in G = SU(3).$$
 (4.1)

As already demonstrated in the SU(N) case, the key identity for deriving the decomposition is given by

$$\mathbf{V} = \mathbf{n}_3(\mathbf{V} \cdot \mathbf{n}_3) + \mathbf{n}_8(\mathbf{V} \cdot \mathbf{n}_8) + \mathbf{n}_3 \times (\mathbf{V} \times \mathbf{n}_3) + \mathbf{n}_8 \times (\mathbf{V} \times \mathbf{n}_8).$$
(4.2)

The decomposition of the SU(3) gauge field in the vector notation is

$$\mathbf{A}_{\mu} = \mathbf{V}_{\mu} + \mathbf{X}_{\mu} = \mathbf{C}_{\mu} + \mathbf{B}_{\mu} + \mathbf{X}_{\mu},$$

$$\mathbf{C}_{\mu} := (\mathbf{n}_{3} \cdot \mathbf{A}_{\mu})\mathbf{n}_{3} + (\mathbf{n}_{8} \cdot \mathbf{A}_{\mu})\mathbf{n}_{8},$$

$$\mathbf{B}_{\mu} := g^{-1}\partial_{\mu}\mathbf{n}_{3} \times \mathbf{n}_{3} + g^{-1}\partial_{\mu}\mathbf{n}_{8} \times \mathbf{n}_{8},$$

$$\mathbf{X}_{\mu} := g^{-1}\mathbf{n}_{3} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{3} + g^{-1}\mathbf{n}_{8} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{8}.$$
(4.3)

It is important to note that n_3 and n_8 are introduced simply for convenience and that the independent dynamical degrees of freedom are provided by a single color field n. For instance, the simplest choice in the maximal case is given by

$$\mathbf{n}_3(x) = \mathbf{n}(x), \quad \mathbf{n}_8(x) = \sqrt{3}\mathbf{n}_3(x) * \mathbf{n}_3(x) = \sqrt{3}\mathbf{n}(x) * \mathbf{n}(x).$$
 (4.4)

For more details, see Appendix C.

In a similar way to the SU(2) case discussed in the previous paper [8], the transformation properties of the decomposed fields $\mathbf{B}_{\mu}, \mathbf{C}_{\mu}, \mathbf{X}_{\mu}$ are uniquely determined once we specify those for \mathbf{A}_{μ} and \mathbf{n} . For independent transformations of \mathbf{A}_{μ} and \mathbf{n} :

$$\delta_{\omega} \mathbf{A}_{\mu}(x) = D_{\mu}[\mathbf{A}]\boldsymbol{\omega}(x), \quad \delta_{\theta} \mathbf{n}(x) = g\mathbf{n}(x) \times \boldsymbol{\theta}(x) = g\mathbf{n}(x) \times \boldsymbol{\theta}_{\perp}(x), (\boldsymbol{\omega} \in SU(3), \ \boldsymbol{\theta}_{\perp} \in SU(3)/(U(1) \times U(1))),$$
(4.5)

the new variables \mathbf{V}_{μ} and \mathbf{X}_{μ} are transformed under the enlarged gauge transformation as

$$\delta_{\omega,\theta} \mathbf{V}_{\mu}(x) = D_{\mu}[\mathbf{V}] \boldsymbol{\omega}_{\parallel}(x) + D_{\mu}[\mathbf{V}] \boldsymbol{\theta}_{\perp}(x) + g \mathbf{X}_{\mu}(x) \times (\boldsymbol{\omega}_{\perp}(x) - \boldsymbol{\theta}_{\perp}(x)),$$

$$\delta_{\omega,\theta} \mathbf{X}_{\mu}(x) = g \mathbf{X}_{\mu}(x) \times (\boldsymbol{\omega}_{\parallel}(x) + \boldsymbol{\theta}_{\perp}(x)) + D_{\mu}[\mathbf{V}](\boldsymbol{\omega}_{\perp}(x) - \boldsymbol{\theta}_{\perp}(x)).$$
(4.6)

If $\boldsymbol{\omega}_{\perp}(x) = \boldsymbol{\theta}_{\perp}(x)$, in particular, \mathbf{V}_{μ} and \mathbf{X}_{μ} are transformed as

$$\delta_{\omega} \mathbf{V}_{\mu}(x) = D_{\mu}[\mathbf{V}]\boldsymbol{\omega}(x), \quad \delta_{\omega} \mathbf{X}_{\mu}(x) = g \mathbf{X}_{\mu}(x) \times \boldsymbol{\omega}(x), \tag{4.7}$$

which agrees with the equipollent gauge transformation to be obtained by the reduction condition.

In *D*-dimensional space-time, the original gauge field \mathbf{A}_{μ} for G = SU(3) has 8D components. The field \mathbf{n}_3 belongs to $G/\tilde{H} = SU(3)/(U(1) \times U(1)) = F_2$ and hence has 8-2 = 6 degrees of freedom. As already pointed out, \mathbf{n}_8 is constructed from \mathbf{n}_3 by $\mathbf{n}_8 = \sqrt{3}\mathbf{n}_3 * \mathbf{n}_3$. Note that \mathbf{B}_{μ} is uniquely determined once \mathbf{n}_3 is fixed. The maximal Abelian potential \mathbf{C}_{μ} commutes with both \mathbf{n}_3 and \mathbf{n}_8 and hence $\mathbf{C}_{\mu} \in \tilde{H} = U(1) \times U(1)$ has 2*D* components. In contrast, \mathbf{X}_{μ} has no components in the direction of \tilde{H} , i.e., $\mathbf{X}_{\mu} \in G/\tilde{H}$ has (8-2)D = 6D components. Therefore, there are 6 extra degrees of freedom after the transformation $\mathbf{A}_{\mu} \to (\mathbf{n}, \mathbf{C}_{\mu}, \mathbf{X}_{\mu})$, although the number of degrees of freedom agrees between $(\mathbf{n}, \mathbf{A}_{\mu})$ and $(\mathbf{n}, \mathbf{C}_{\mu}, \mathbf{X}_{\mu})$.

By introducing a single color fields n, the original Yang-Mills theory with the gauge symmetry SU(3) is extended into a gauge theory with the enlarged gauge

group $SU(3) \times [SU(3)/(U(1) \times U(1))]$, which we call the master Yang-Mills theory. To eliminate the 6 extra degrees of freedom, we must impose 6 independent conditions, which we call the maximal version of the reduction condition. Consequently, we can reduce the enlarged gauge symmetry to the original gauge symmetry, i.e., the equipollent gauge symmetry. Thus, we can regard

$$\mathbf{A}_{\mu} \to (\mathbf{n}, \mathbf{C}_{\mu}, \mathbf{X}_{\mu}) \tag{4.8}$$

as the change of variables and obtain a new reformulation of SU(3) Yang-Mills theory written in terms of the new variables $(\mathbf{n}, \mathbf{C}_{\mu}, \mathbf{X}_{\mu})$.

4.1.2 Reduction to SU(3): maximal case

To find the reduction condition for SU(3), we calculate the SU(3) version of the square of \mathbf{X}^{μ} , as suggested from the SU(2) case. By introducing $\mathbf{n}_s = \{\mathbf{n}_3, \mathbf{n}_8\}$, the square of \mathbf{X}^{μ} is expressed as

$$g^{2}\mathbf{X}^{\mu} \cdot \mathbf{X}_{\mu} = \{\mathbf{n}_{3} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{3}) + \mathbf{n}_{8} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{8})\}^{2} \\ = \{\mathbf{n}_{3} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{3})\}^{2} + 2\{\mathbf{n}_{3} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{3})\} \cdot \{\mathbf{n}_{8} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{8})\}^{2} \\ + \{\mathbf{n}_{8} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{8})\}^{2} \\ = \{\mathbf{n}_{3} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{3})\}^{2} + \{\mathbf{n}_{8} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{3})\}^{2} \\ + \{\mathbf{n}_{3} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{8})\}^{2} + \{\mathbf{n}_{8} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{8})\}^{2} \\ = (D_{\mu}[\mathbf{A}]\mathbf{n}_{3})^{2} + (D_{\mu}[\mathbf{A}]\mathbf{n}_{8})^{2} =: (D_{\mu}[\mathbf{A}]\mathbf{n}_{s})^{2}, \qquad (4.9)$$

where in the third equality we have used the fact that

$$\{\mathbf{n}_{3} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{3})\} \cdot \{\mathbf{n}_{8} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{8})\} = \{\mathbf{n}_{8} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{3})\}^{2} = \{\mathbf{n}_{3} \times (D_{\mu}[\mathbf{A}]\mathbf{n}_{8})\}^{2}.$$
(4.10)

From the transformation property of $(D_{\mu}[\mathbf{A}]\boldsymbol{n}_s)^2$ under the enlarged gauge transformation:

$$\delta_{\omega,\theta} \left\{ \frac{1}{2} (D_{\mu}[\mathbf{A}]\mathbf{n}_{s})^{2} \right\} = g(\mathbf{n}_{s} \times D_{\mu}[\mathbf{A}]\mathbf{n}_{s}) \cdot D_{\mu}[\mathbf{A}](\boldsymbol{\omega}_{\perp} - \boldsymbol{\theta}_{\perp}), \qquad (4.11)$$

it turns out that $(D_{\mu}[\mathbf{A}]\boldsymbol{n}_s)^2$ is invariant under the equipollent gauge transformation (4.7), i.e., the enlarged gauge transformation (4.6) when $\boldsymbol{\omega}_{\perp} = \boldsymbol{\theta}_{\perp}$. Therefore, we impose the minimizing condition of the functional R under the enlarged gauge transformation:

$$R = \int d^{D}x \frac{1}{2} (D_{\mu}[\mathbf{A}]\mathbf{n}_{s}(x))^{2}.$$
 (4.12)

From the observation that

$$0 = \delta_{\omega,\theta} R = \int d^D x \frac{1}{2} \delta_{\omega,\theta} (D_{\mu}[\mathbf{A}]\mathbf{n}_s(x))^2$$

= $-\int d^D x g(\boldsymbol{\omega}_{\perp} - \boldsymbol{\theta}_{\perp}) \cdot (\mathbf{n}_s \times D_{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\mathbf{n}_s),$ (4.13)

the minimizing condition reduces the enlarged gauge symmetry as follows:

$$\tilde{G} = SU(3)_{\omega} \times [SU(3)/(U(1) \times U(1))]_{\theta}$$

$$\rightarrow G' = SU(3)'_{\alpha}, \quad \boldsymbol{\alpha} = (\boldsymbol{\alpha}_{\parallel}, \boldsymbol{\alpha}_{\perp}) = (\boldsymbol{\omega}_{\parallel}, \boldsymbol{\omega}_{\perp} = \boldsymbol{\theta}_{\perp}).$$
(4.14)

The equipollent gauge transformation is

$$\delta_{\alpha} \mathbf{A}_{\mu} = D_{\mu} [\mathbf{A}] \boldsymbol{\alpha}, \quad \delta_{\alpha} \mathbf{n} = g \mathbf{n} \times \boldsymbol{\alpha}_{\perp}, \tag{4.15}$$

and

$$\delta_{\alpha} \mathbf{V}_{\mu}(x) = D_{\mu}[\mathbf{V}] \boldsymbol{\alpha}(x), \quad \delta_{\alpha} \mathbf{X}_{\mu}(x) = g \mathbf{X}_{\mu}(x) \times \boldsymbol{\alpha}(x).$$
(4.16)

This is simply the SU(3) version of the equipollent gauge symmetry, as discussed in the previous paper for SU(2) [8]. Note that the following quantities are also invariant under the equipollent gauge transformation (4.15):

$$(D_{\mu}[\mathbf{A}_{\mu}]\mathbf{n}_{8})^{2}, \quad (D_{\mu}[\mathbf{A}_{\mu}]\mathbf{n}_{3}) \cdot (D_{\mu}[\mathbf{A}_{\mu}]\mathbf{n}_{8}).$$
 (4.17)

This is also the case for their linear combinations.

Thus, minimizing the functional $\int d^4x (\mathbf{X}^{\mu} \cdot \mathbf{X}_{\mu})$ yields the differential form of the SU(3) version of the reduction condition:

$$\boldsymbol{\chi}_{\rm rc}[\mathbf{A}, \boldsymbol{n}_s] := \mathbf{n}_s \times D_{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \mathbf{n}_s, \qquad (4.18)$$

which is rewritten in terms of new variables as

$$\boldsymbol{\chi}_{\rm rc}[\boldsymbol{n}, \mathbf{C}, \mathbf{X}] := D^{\mu}[\mathbf{V}] \mathbf{X}_{\mu} \equiv 0.$$
(4.19)

This is called the differential reduction condition.

Moreover, we can adopt the other functional written in terms of a single color field n to write the reduction condition as

$$F[\mathbf{A}, \mathbf{n}_3] = \int d^D x \frac{1}{2} (D_{\mu}[\mathbf{A}] \mathbf{n}_3(x))^2.$$
(4.20)

This is a new option overlooked so far. We can repeat the same calculations as those given above. Then we obtain the differential form of the reduction condition:

$$\boldsymbol{\chi}_{\rm rc}[\mathbf{A},\mathbf{n}_3] = \mathbf{n}_3 \times D_{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \mathbf{n}_3. \tag{4.21}$$

However, this cannot be rewritten into such a simple form as (4.19).

4.2 Minimal case for SU(3)

4.2.1 Decomposition for SU(3): minimal case

According to the general discussion of SU(N), the decomposition in the minimal case of SU(3) is given by

$$\mathbf{A}_{\mu} = \mathbf{C}_{\mu} + \mathbf{B}_{\mu} + \mathbf{X}_{\mu},\tag{4.22}$$

$$\mathbf{C}_{\mu} := \mathbf{A}_{\mu} - \frac{4}{3}\mathbf{n}_8 \times (\mathbf{A}_{\mu} \times \mathbf{n}_8), \qquad (4.23)$$

$$\mathbf{B}_{\mu} := \frac{4}{3}g^{-1}\partial_{\mu}\mathbf{n}_{8} \times \mathbf{n}_{8}, \qquad (4.24)$$

$$\mathbf{X}_{\mu} := \frac{4}{3}g^{-1}\mathbf{n}_8 \times D_{\mu}[\mathbf{A}_{\mu}]\mathbf{n}_8, \qquad (4.25)$$

where \mathbf{n}_8 is the single color field $\mathbf{n} = \mathbf{n}_8$. In a similar way to the SU(2) case discussed in the previous paper [8], the transformation properties of the decomposed fields $\mathbf{B}_{\mu}, \mathbf{C}_{\mu}, \mathbf{X}_{\mu}$ are uniquely determined once we specify those for \mathbf{A}_{μ} and \mathbf{n}_8 . The infinitesimal enlarged gauge transformation is constructed from $\delta_{\omega} \mathbf{A}_{\mu}$ and $\delta_{\theta} \mathbf{n}_8$ as

$$\delta_{\omega} \mathbf{A}_{\mu}(x) = D_{\mu}[\mathbf{A}] \boldsymbol{\omega}(x), \quad \delta_{\theta} \mathbf{n}_{8}(x) = g \mathbf{n}_{8}(x) \times \boldsymbol{\theta}_{\perp}(x)$$
$$(\boldsymbol{\omega} \in SU(3), \boldsymbol{\theta}_{\perp} \in SU(3)/U(2)), \qquad (4.26)$$

where $\boldsymbol{\theta}_{\perp} \in \mathscr{L}(G/H)$. The new variables specified above are transformed according to

$$\delta_{\omega,\theta} \mathbf{V}_{\mu}(x) = D_{\mu}[\mathbf{V}] \boldsymbol{\omega}_{\parallel}(x) + D_{\mu}[\mathbf{V}] \boldsymbol{\theta}_{\perp}(x) + g \mathbf{X}_{\mu}(x) \times (\boldsymbol{\omega}_{\perp}(x) - \boldsymbol{\theta}_{\perp}(x)),$$

$$\delta_{\omega,\theta} \mathbf{X}_{\mu}(x) = g \mathbf{X}_{\mu}(x) \times (\boldsymbol{\omega}_{\parallel}(x) + \boldsymbol{\theta}_{\perp}(x)) + D_{\mu}[\mathbf{V}](\boldsymbol{\omega}_{\perp}(x) - \boldsymbol{\theta}_{\perp}(x)).$$
(4.27)

In particular, for $\boldsymbol{\omega}_{\perp}(x) = \boldsymbol{\theta}_{\perp}(x)$, \mathbf{V}_{μ} and \mathbf{X}_{μ} are transformed as

$$\delta_{\omega} \mathbf{V}_{\mu}(x) = D_{\mu}[\mathbf{V}]\boldsymbol{\omega}(x), \quad \delta_{\omega} \mathbf{X}_{\mu}(x) = g \mathbf{X}_{\mu}(x) \times \boldsymbol{\omega}(x).$$
(4.28)

This is the equipollent gauge transformation, which is to be obtained by imposing the reduction condition, as in the maximal case.

The original gauge field \mathbf{A}_{μ} for G = SU(3) has 8D components. The color field \mathbf{n}_8 belongs to $G/\tilde{H} = SU(3)/U(2)$ and hence has 8 - 4 = 4 degrees of freedom. The variable \mathbf{C}_{μ} commutes with \mathbf{n}_8 , and hence $\mathbf{C}_{\mu} \in \tilde{H} = U(2)$ with 4D components. Therefore, \mathbf{C}_{μ} is a non-Abelian gauge field, in sharp contrast with the maximal case, where it is the maximal Abelian gauge field. Moreover, \mathbf{X}_{μ} has no components in the direction of \tilde{H} (no \tilde{H} -commutative part), i.e., $\mathbf{X}_{\mu} \in G/\tilde{H}$ has (8 - 4)D = 4D components. Note that \mathbf{B}_{μ} is written in terms of \mathbf{n}_8 . Therefore, there are 4 extra degrees of freedom associated with the transformation $\mathbf{A}_{\mu} \to (\mathbf{n}_8, \mathbf{C}_{\mu}, \mathbf{X}_{\mu})$, although the number of degrees of freedom agrees between $(\mathbf{n}_8, \mathbf{A}_{\mu})$ and $(\mathbf{n}_8, \mathbf{C}_{\mu}, \mathbf{X}_{\mu})$.

4.2.2 Reduction to SU(3): minimal case

By introducing a single color field n_8 , the original Yang-Mills theory with gauge symmetry SU(3) is extended to the master Yang-Mills theory with the enlarged gauge group, $SU(3) \times [SU(3)/U(2)]$. Therefore, we impose a constraint which we call the minimal version of the reduction condition, to eliminate the 4 extra degrees of freedom. By imposing the minimal version of the reduction condition with the same number of independent conditions as the extra degrees of freedom, we reduce the enlarged gauge symmetry to the original gauge symmetry SU(3), which is called the equipollent gauge symmetry. Thus, we can regard

$$\mathbf{A}_{\mu} \to (\mathbf{n}_8, \mathbf{C}_{\mu}, \mathbf{X}_{\mu}) \tag{4.29}$$

$$\boldsymbol{n}_{(a)}' := W^{\dagger} \boldsymbol{n}_{(a)} W = (U^{\dagger} V^{\dagger} U) (U^{\dagger} T_a U) (U^{\dagger} V U) = U^{\dagger} (V^{\dagger} T_a V) U.$$

In particular, \boldsymbol{n}_8 is invariant, i.e., $\boldsymbol{n}'_8 = \boldsymbol{n}_8$, since $VT_8V^{\dagger} = T_8$. Therefore, $\tilde{H} = U(2)$ in G = SU(3) leaves \boldsymbol{n}_8 invariant. In other words, $\boldsymbol{n}_8 \in G/\tilde{H}$. Using the degrees of freedom of \tilde{H} , we can change $\boldsymbol{n}_{(1)}$, $\boldsymbol{n}_{(2)}$ and $\boldsymbol{n}_{(3)}$ without changing \boldsymbol{n}_8 .

¹¹Defining $\boldsymbol{n}_{(1)} := U^{\dagger}T_{1}U, \, \boldsymbol{n}_{(2)} := U^{\dagger}T_{2}U, \, \boldsymbol{n}_{(3)} := U^{\dagger}T_{3}U$, we find that $[\boldsymbol{n}_{(a)}, \boldsymbol{n}_{8}] = 0$ (a = 1, 2, 3, 8), since $[T_{a}, T_{8}] = 0$ (a = 1, 2, 3, 8). For an arbitrary element $V \in \mathcal{H} = U(2)$ generated by $\{T_{1}, T_{2}, T_{3}, T_{8}\}$, we define $W := U^{\dagger}VU$. Then we find that $\boldsymbol{n}_{(a)}$ is transformed as

as the change of variables and obtain a new reformulation of SU(3) Yang-Mills theory written in terms of the new variables $(\mathbf{n}_8, \mathbf{C}_\mu, \mathbf{X}_\mu)$.

It turns out that $\mathbf{X}^{\mu}(x) \cdot \mathbf{X}_{\mu}(x) = \frac{4}{3} (D_{\mu}[\mathbf{A}] \mathbf{n}_{8}(x))^{2}$ is invariant under the enlarged gauge transformation (4.26) when $\boldsymbol{\omega}_{\perp} = \boldsymbol{\theta}_{\perp}$. Then the minimizing functional for SU(3) in the minimal case is given by

$$F = \int d^{D}x \frac{3}{8} g^{2} \mathbf{X}^{\mu}(x) \cdot \mathbf{X}_{\mu}(x) = \int d^{D}x \frac{1}{2} (D_{\mu}[\mathbf{A}]\mathbf{n}_{8}(x))^{2}.$$
 (4.30)

We impose the reduction condition as the minimizing condition of the functional:

$$0 = \delta_{\omega,\theta} F = \int d^D x \frac{1}{2} \delta_{\omega,\theta} \{ (D_{\mu}[\mathbf{A}]\mathbf{n}_8(x)) \}^2$$

= $-\int d^D x g(\boldsymbol{\omega}_{\perp} - \boldsymbol{\theta}_{\perp}) \cdot D_{\mu}[\mathbf{A}](\mathbf{n}_8 \times D_{\mu}[\mathbf{A}]\mathbf{n}_8)$
= $-\int d^D x g(\boldsymbol{\omega}_{\perp} - \boldsymbol{\theta}_{\perp}) \cdot (\mathbf{n}_8 \times D_{\mu}[\mathbf{A}]D_{\mu}[\mathbf{A}]\mathbf{n}_8).$ (4.31)

By imposing the minimizing condition of the functional F under the enlarged gauge transformation, the enlarged gauge symmetry \tilde{G} is reduced to G':

$$G = SU(3)_{\omega} \times [SU(3)/U(2)]_{\theta}$$

$$\rightarrow G' = SU(3)'_{\alpha}, \quad \boldsymbol{\alpha} = (\boldsymbol{\alpha}_{\parallel}, \boldsymbol{\alpha}_{\perp}) = (\boldsymbol{\omega}_{\parallel}, \boldsymbol{\omega}_{\perp} = \boldsymbol{\theta}_{\perp}). \quad (4.32)$$

The equipollent gauge transformation is

$$\delta_{\alpha} \mathbf{A}_{\mu}(x) = D_{\mu}[\mathbf{A}] \boldsymbol{\alpha}(x), \quad \delta_{\alpha} \mathbf{n}_{8}(x) = g \mathbf{n}_{8}(x) \times \boldsymbol{\alpha}_{\perp}(x), \tag{4.33}$$

and

$$\delta_{\alpha} \mathbf{V}_{\mu}(x) = D_{\mu}[\mathbf{V}] \boldsymbol{\alpha}(x), \quad \delta_{\alpha} \mathbf{X}_{\mu}(x) = g \mathbf{X}_{\mu}(x) \times \boldsymbol{\alpha}(x).$$
(4.34)

Thus, minimizing the functional F yields the differential form of the SU(3) version of the reduction condition for the infinitesimal transformation:

$$\boldsymbol{\chi}_{\rm rc}[\mathbf{A}, \boldsymbol{n}_8] = \mathbf{n}_8 \times D_{\mu}[\mathbf{A}] D_{\mu}[\mathbf{A}] \mathbf{n}_8.$$
(4.35)

This is rewritten in terms of the new variables as

$$\boldsymbol{\chi}_{\rm rc}[\mathbf{n}_8, \mathbf{C}, \mathbf{X}] := D^{\mu}[\mathbf{V}] \mathbf{X}_{\mu} \equiv 0.$$
(4.36)

4.3 Interpolation between maximal and minimal cases for SU(3)

The maximal and minimal cases are interpolated by introducing a *global* parameter ϑ . This is achieved by minimizing the following functional with respect to the enlarged gauge transformation:

$$F_{\vartheta} := \text{const.} \int d^D x (D_{\mu}[\mathbf{A}]\mathbf{n}(x)) \cdot (D_{\mu}[\mathbf{A}]\mathbf{n}(x)), \qquad (4.37)$$

where $\mathbf{n}(x)$ is introduced in (2.36):

$$\mathbf{n}(x) = (\cos\vartheta(x))\mathbf{n}_3(x) + (\sin\vartheta(x))\mathbf{n}_8(x).$$
(4.38)

The minimal case is realized for $\vartheta = \frac{1}{6}\pi, \frac{1}{2}\pi, \frac{5}{6}\pi, \frac{7}{6}\pi, \frac{3}{2}\pi, \frac{11}{6}\pi$ where $\vartheta = \pi/2$ is the simplest choice. The maximal case is realized for other values of ϑ , where $\vartheta = 0$ is the simplest choice. This will be discussed in more detail elsewhere.

5 Quantization of Yang-Mills theory

To construct a quantum version of Yang-Mills theory, we adopt the functional integral quantization. We now reformulate a quantum Yang-Mills theory in terms of new variables. For this purpose, we must specify the action and the integration measure. We can immediately rewrite the Yang-Mills action $S_{\rm YM}[\mathscr{A}]$ in terms of the new variables by substituting the transformation law (3.32) or (3.63) into the original gauge field \mathscr{A} in the Yang-Mills Lagrangian $\mathscr{L}_{\rm YM}[\mathscr{A}]$. To write the integration measure explicitly in terms of the new variables, we need to know the Jacobian associated with the change of variables from the original Yang-Mills gauge field to the new variables in the reformulated Yang-Mills theory.

5.1 Jacobian for the nonlinear change of variables

We consider the Jacobian associated with the change of variables for the decomposition

$$\mathscr{A}_{\mu}(x) = \mathscr{V}_{\mu}(x) + \mathscr{X}_{\mu}(x) = \mathscr{C}_{\mu}(x) + \mathscr{B}_{\mu}(x) + \mathscr{X}_{\mu}(x).$$
(5.1)

In the following, we show that the Jacobian can be simplified by choosing appropriate bases in the group space without changing the physical content.

For G = SU(2), we have already introduced [10] the orthonormal basis ($\mathbf{e}_1(x), \mathbf{e}_2(x), \mathbf{n}(x)$) satisfying the following relations in vector form using $\mathbf{e}_a(x) = (e_a^A(x))$, $\mathbf{n}(x) = (n^A(x))$:

$$\mathbf{e}_{a}(x) \cdot \mathbf{e}_{b}(x) = \delta_{ab}, \quad \mathbf{e}_{a}(x) \cdot \mathbf{n}(x) = 0, \quad \mathbf{n}(x) \cdot \mathbf{n}(x) = 1,$$

$$\mathbf{e}_{a}(x) \times \mathbf{e}_{b}(x) = \epsilon_{ab}\mathbf{n}(x), \quad \mathbf{e}_{a}(x) \times \mathbf{n}(x) = -\epsilon_{ab}\mathbf{e}_{b}(x) \quad (a, b = 1, 2), \quad (5.2)$$

which are written in the Lie algebra form using $\boldsymbol{e}_a(x) = e^A_a(x)T^A$, $\boldsymbol{n}(x) = n^A(x)T^A$ as

$$\operatorname{tr}[\boldsymbol{e}_{a}(x)\boldsymbol{e}_{b}(x)] = \delta_{ab}/2, \quad \operatorname{tr}[\boldsymbol{e}_{a}(x)\boldsymbol{n}(x)] = 0, \quad \operatorname{tr}[\boldsymbol{n}(x)\boldsymbol{n}(x)] = 1/2, \\ [\boldsymbol{e}_{a}(x), \boldsymbol{e}_{b}(x)] = i\epsilon_{ab}\boldsymbol{n}(x), \quad [\boldsymbol{e}_{a}(x), \boldsymbol{n}(x)] = -i\epsilon_{ab}\boldsymbol{e}_{b}(x) \quad (a, b = 1, 2).$$
(5.3)

By definition, $\mathscr{C}_{\mu}(x)$ is parallel (or commutes with) $\boldsymbol{n}(x)$, i.e.,

$$[\mathscr{C}_{\mu}(x), \boldsymbol{n}(x)] = 0. \tag{5.4}$$

Hence, for SU(2), $\mathscr{C}_{\mu}(x)$ can be written as

$$\mathscr{C}_{\mu}(x) = C_{\mu}(x)\boldsymbol{n}(x), \qquad (5.5)$$

which has the components

$$\mathscr{C}^A_\mu(x) = C_\mu(x) n^A(x).$$
(5.6)

On the other hand, $\mathscr{B}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$ are orthogonal to (or noncommutative with) $\mathbf{n}(x)$,

$$\mathscr{B}_{\mu}(x) \cdot \mathbf{n}(x) = 0, \quad \mathscr{X}_{\mu}(x) \cdot \mathbf{n}(x) = 0,$$
(5.7)

and hence can be written using orthonormal bases $\{e_a(x)\}$ as

$$\mathscr{B}_{\mu}(x) = B^{a}_{\mu}(x)\boldsymbol{e}_{a}(x), \quad \mathscr{X}_{\mu}(x) = X^{a}_{\mu}(x)\boldsymbol{e}_{a}(x), \quad (5.8a)$$

which yield the components

$$\mathscr{B}^{A}_{\mu}(x) = B^{a}_{\mu}(x)e^{A}_{a}(x), \quad \mathscr{X}^{A}_{\mu}(x) = X^{a}_{\mu}(x)e^{A}_{a}(x).$$
 (5.8b)

For G = SU(N) $(N \ge 3)$, we introduce complete orthonormal bases $\{e_a, u_j\}$ for su(N) where

$$\boldsymbol{e}_a \in \mathscr{L}(G/\tilde{H}), \quad \boldsymbol{u}_j \in \mathscr{L}(\tilde{H}), \ (a = 1, \cdots, \dim G/\tilde{H}; j = 1, \cdots, \dim \tilde{H})$$
 (5.9)

which satisfy the relations in vector form using $\boldsymbol{e}_a(x) = (e_a^A(x))$, $\boldsymbol{u}_j(x) = (\boldsymbol{u}_j^A(x))$:

$$\boldsymbol{e}_{a}(x) \cdot \boldsymbol{e}_{b}(x) = \delta_{ab}, \quad \boldsymbol{e}_{a}(x) \cdot \boldsymbol{u}_{j}(x) = 0 \quad \boldsymbol{u}_{j}(x) \cdot \boldsymbol{u}_{k}(x) = \delta_{jk},$$
$$(a, b = 1, \cdots, \dim G/\tilde{H}; j, k = 1, \dots, \dim \tilde{H}), \quad (5.10)$$

which are written in the Lie algebra form using $e_a(x) = e_a^A(x)T^A$, $u_j(x) = u_j^A(x)T^A$:

$$\operatorname{tr}[\boldsymbol{e}_{a}(x)\boldsymbol{e}_{b}(x)] = \delta_{ab}/2, \quad \operatorname{tr}[\boldsymbol{e}_{a}(x)\boldsymbol{u}_{j}(x)] = 0, \quad \operatorname{tr}[\boldsymbol{u}_{j}(x)\boldsymbol{u}_{k}(x)] = 1/2\delta_{jk}.$$
(5.11)

It is instructive to note that u_j reduces to a single color field n in SU(2).

The fields $\mathscr{B}_{\mu}(x)$ and $\mathscr{X}_{\mu}(x)$ are orthogonal to (or noncommutative with) all $u_j(x)$:

$$\mathscr{B}_{\mu}(x) \cdot \boldsymbol{u}_{j}(x) = 0, \quad \mathscr{X}_{\mu}(x) \cdot \boldsymbol{u}_{j}(x) = 0 \ (j = 1, \cdots, \dim \tilde{H}).$$
 (5.12)

Hence they can be written using the bases $\{e_a(x)\}$:

$$\mathcal{B}_{\mu}(x) = B^{a}_{\mu}(x)\boldsymbol{e}_{a}(x) \in \mathcal{L}(G/\tilde{H}),$$

$$\mathcal{X}_{\mu}(x) = X^{a}_{\mu}(x)\boldsymbol{e}_{a}(x) \in \mathcal{L}(G/\tilde{H}) \ (a = 1, \cdots, \dim G/\tilde{H}).$$
(5.13)

By definition, $\mathscr{C}_{\mu}(x)$ can be written as

$$\mathscr{C}_{\mu}(x) = C^{j}_{\mu}(x)\boldsymbol{u}_{j}(x) \in \mathscr{L}(\tilde{H}) \ (j = 1, \cdots, \dim \tilde{H}).$$
(5.14)

Thus, in the minimal case, $\tilde{H} = U(N-1)$, and the decomposition is

$$\mathscr{A}^{A}_{\mu}(x) = C^{j}_{\mu}(x)\boldsymbol{u}^{A}_{j}(x) + B^{a}_{\mu}(x)\boldsymbol{e}^{A}_{a}(x) + X^{a}_{\mu}(x)\boldsymbol{e}^{A}_{a}(x)$$

(j = 1,..., dim $\tilde{H} = (N-1)^{2}$; $a = 1, \cdots, \text{dim}G/\tilde{H} = 2N-2$). (5.15)

In the maximal case, $\tilde{H} = H = U(1)^{N-1}$ and $\boldsymbol{u}_j(x)$ reduce to $\boldsymbol{n}_j(x)$ $(j = 1, \dots, \dim \tilde{H})$, which mutually commute, i.e., $[\boldsymbol{n}_j(x), \boldsymbol{n}_k(x)] = 0$ (3.13). Therefore, we obtain

$$\mathscr{A}^{A}_{\mu}(x) = C^{j}_{\mu}(x)\boldsymbol{n}^{A}_{j}(x) + B^{a}_{\mu}(x)\boldsymbol{e}^{A}_{a}(x) + X^{a}_{\mu}(x)\boldsymbol{e}^{A}_{a}(x)$$

(j = 1,..., dim $\tilde{H} = N - 1; \ a = 1, \cdots, \text{dim}G/\tilde{H} = N^{2} - N$). (5.16)

We now consider the change of variables from $\{n^{\alpha}, \mathscr{A}^{A}_{\mu}\}$ to the new variables $\{n^{\beta}, C^{k}_{\nu}, X^{b}_{\nu}\}$ defined in the su(N) bases $\{e_{b}, u_{k}\}$. Here n^{α} and n^{β} denote independent degrees of freedom after solving the constraint $n^{A}n^{A} = 1$. Then the integration measure is transformed as

$$\mathcal{D}n^{\alpha}\mathcal{D}\mathscr{A}^{A}_{\mu} = \mathcal{D}n^{\beta}\mathcal{D}C^{k}_{\nu}\mathcal{D}X^{b}_{\nu}J.$$
(5.17)

The Jacobian J is given by

$$J := \begin{vmatrix} \frac{\partial n^{\alpha}}{\partial n^{\beta}} & \frac{\partial n^{\alpha}}{\partial C_{\nu}^{k}} & \frac{\partial n^{\alpha}}{\partial X_{\nu}^{b}} \\ \frac{\partial \mathscr{A}_{\mu}}{\partial n^{\beta}} & \frac{\partial \mathscr{A}_{\mu}}{\partial C_{\nu}^{k}} & \frac{\partial \mathscr{A}_{\mu}}{\partial X_{\nu}^{b}} \end{vmatrix} = \begin{vmatrix} \delta_{\beta}^{\alpha} & 0 & 0 \\ \frac{\partial \mathscr{A}_{\mu}^{A}}{\partial n^{\beta}} & \delta_{\mu\nu} u_{k}^{A} & \delta_{\mu\nu} e_{b}^{A} \end{vmatrix}$$
$$= \begin{vmatrix} \delta_{\mu\nu} u_{k}^{A} & \delta_{\mu\nu} e_{b}^{A} \end{vmatrix} = \begin{vmatrix} u_{k}^{A} & e_{b}^{A} \end{vmatrix}^{D} = 1, \qquad (5.18)$$

where we have used the fact that $\{n^{\beta}, C_{\nu}^{k}, X_{\nu}^{b}\}$ are independent variables in the second equality, that μ and ν run from 1 to D in the fourth equality, and that $\{u_{k}^{A}, e_{b}^{A}\}$ constitute the orthonormal base in the last equality. Thus, we obtain

$$\mathcal{D}n^{\alpha}\mathcal{D}\mathscr{A}^{A}_{\mu} = \mathcal{D}n^{\beta}\mathcal{D}C^{k}_{\nu}\mathcal{D}X^{b}_{\nu}J, \quad J = 1.$$
(5.19)

For the Jacobian J to be simplified to J = 1, the fact that C_{ν}^{k}, X_{ν}^{b} are independent of n^{β} is important.

5.2 Functional integration

We now reformulate the quantum Yang-Mills theory by the functional integral method. The following steps can be clearly understood by comparing them with the corresponding ones in Fig. 1.

First, the original Yang-Mills (YM) theory in the Euclidean space is defined by the partition function (see the middle left part in Fig. 1):

$$Z_{\rm YM} = \int \mathcal{D}\mathscr{A}^A_\mu \exp(-S_{\rm YM}[\mathscr{A}]).$$
 (5.20)

Of course, to obtain a well-defined Yang-Mills theory by completely fixing the gauge, we must impose a gauge fixing condition for the SU(N) gauge symmetry, for example, the Landau gauge $\partial^{\mu} \mathscr{A}^{A}_{\mu}(x) = 0$, which is called the overall gauge fixing condition (see the down arrows in Fig. 1). For a while, we disregard this procedure, as it can be carried out following the standard method.

Second, the color field n(x) is introduced to extend the original YM theory (see the up arrow in Fig. 1) to the enlarged Yang-Mills theory called the master Yang-Mills (M-YM) theory, which is defined by a partition function written in terms of both $n^{\alpha}(x)$ and $\mathscr{A}^{A}_{\mu}(x)$ (see the top part in Fig. 1),

$$\tilde{Z}_{\rm YM} = \int \mathcal{D}n^{\alpha} \int \mathcal{D}\mathscr{A}_{\mu}^{A} \exp(-S_{\rm YM}[\mathscr{A}]).$$
(5.21)

Third, we regard (5.21) as

$$\tilde{Z}_{\rm YM} = \int \mathcal{D}n^{\beta} \int \mathcal{D}C_{\nu}^{k} \int \mathcal{D}X_{\nu}^{b} J \exp(-\tilde{S}_{\rm YM}[\boldsymbol{n},\mathscr{C},\mathscr{X}]), \qquad (5.22)$$

where J is the Jacobian associated with the change of variables, and the action $\tilde{S}_{\text{YM}}[\boldsymbol{n}, \mathscr{C}, \mathscr{X}]$ is obtained by substituting the decomposition (5.15) of \mathscr{A}^A_{μ} into $S_{\text{YM}}[\mathscr{A}]$:

$$\tilde{S}_{\text{YM}}[\boldsymbol{n}, \mathscr{C}, \mathscr{X}] = S_{\text{YM}}[\mathscr{A}].$$
 (5.23)

Starting with this form, we wish to obtain a new Yang-Mills (YM') theory written in terms of new variables $(n^{\beta}, C_{\nu}^{k}, X_{\nu}^{b})$ (see the middle right part in Fig. 1), which is equipollent to the original Yang-Mills theory. Here, as already discussed in the previous subsection,

$$(n^{\alpha}(x), \mathscr{A}^{A}_{\mu}(x)) \to (n^{\beta}(x), C^{k}_{\nu}(x), X^{b}_{\nu}(x))$$
 (5.24)

should be considered as a change of variables in the sense that $n^{\beta}(x), C_{\nu}^{k}(x), X_{\nu}^{b}(x)$ become *independent* field variables in the new YM' theory.

However, the M-YM has enlarged gauge symmetry, which is larger than the original gauge symmetry G = SU(N). Therefore, to obtain a YM' theory that is equipollent to the original YM theory, the extra gauge degrees of freedom must be eliminated. To fix the enlarged gauge symmetry in the M-YM theory and retain only the same gauge symmetry as that in the original YM, therefore, we impose the constraint $\boldsymbol{\chi}[\mathscr{A}, \boldsymbol{n}] = 0$ (the reduction condition), which is also written in terms of the new variables as $\boldsymbol{\tilde{\chi}} := \boldsymbol{\tilde{\chi}}[\boldsymbol{n}, \mathscr{C}, \mathscr{X}] = 0$. (see the diagonal down arrow in Fig. 1).

The constraint $\boldsymbol{\chi}[\mathscr{A}, \boldsymbol{n}] = 0$ is introduced in the functional integral as follows. We write unity in the form

$$1 = \int \mathcal{D}\boldsymbol{\chi}^{\theta} \delta(\boldsymbol{\chi}^{\theta}) = \int \mathcal{D}\boldsymbol{\theta} \delta(\boldsymbol{\chi}^{\theta}) \det\left(\frac{\delta\boldsymbol{\chi}^{\theta}}{\delta\boldsymbol{\theta}}\right), \qquad (5.25)$$

where χ^{θ} is the constraint written in terms of the gauge-transformed variables, i.e., $\chi^{\theta} := \chi[\mathscr{A}, n^{\theta}]$. As an intermediate step for moving from M-YM to YM', we insert this into the functional integral (5.22), and cast the partition function of the M-YM theory into an intermediate form (see the middle right part in Fig. 1)

$$\tilde{Z}_{\rm YM} = \int \mathcal{D}n^{\beta} \int \mathcal{D}C_{\nu}^{k} \int \mathcal{D}X_{\nu}^{b} J$$

$$\times \int \mathcal{D}\boldsymbol{\theta}\delta(\boldsymbol{\chi}^{\theta}) \det\left(\frac{\delta\boldsymbol{\chi}^{\theta}}{\delta\boldsymbol{\theta}}\right) \exp(-\tilde{S}_{\rm YM}[\boldsymbol{n},\mathscr{C},\mathscr{X}]). \tag{5.26}$$

We next perform the change of variables $\mathbf{n} \to \mathbf{n}^{\theta}$ obtained through a local rotation by angle θ and the corresponding gauge transformations for the other new variables \mathscr{C}_{μ} and $\mathscr{X}_{\mu}: \mathscr{C}_{\mu}, \mathscr{X}_{\mu} \to \mathscr{C}_{\mu}^{\theta}, \mathscr{X}_{\mu}^{\theta}$. From the gauge invariance of the action $\tilde{S}_{\text{YM}}[\mathbf{n}, \mathscr{C}, \mathscr{X}]$ and the measure $\mathcal{D}n^{\beta}\mathcal{D}C_{\nu}^{k}\mathcal{D}X_{\nu}^{b}$, we can rename the dummy integration variables $\mathbf{n}^{\theta}, \mathscr{C}_{\mu}^{\theta}, \mathscr{X}_{\mu}^{\theta}$ as $\mathbf{n}, \mathscr{C}_{\mu}, \mathscr{X}_{\mu}$, respectively. Thus, the integrand does not depend on θ , and the gauge volume $\int \mathcal{D}\boldsymbol{\theta}$ can be removed:

$$\tilde{Z}_{\rm YM} = \int \mathcal{D}\boldsymbol{\theta} \int \mathcal{D}n^{\beta} \int \mathcal{D}C_{\nu}^{k} \int \mathcal{D}X_{\nu}^{b}J \\ \times \delta(\boldsymbol{\chi}) \det\left(\frac{\delta\boldsymbol{\chi}}{\delta\boldsymbol{\theta}}\right) \exp(-\tilde{S}_{\rm YM}[\boldsymbol{n},\mathscr{C},\mathscr{X}]).$$
(5.27)

Thus, we have arrived at the reformulated YM' theory with the partition function:

$$Z'_{\rm YM} = \int \mathcal{D}n^{\beta} \int \mathcal{D}C^{k}_{\nu} \int \mathcal{D}X^{b}_{\nu} J\delta(\tilde{\boldsymbol{\chi}}) \Delta^{rc}_{FP} \exp(-\tilde{S}_{\rm YM}[\boldsymbol{n},\mathscr{C},\mathscr{X}]), \qquad (5.28)$$

where the reduction condition is written in terms of the new variables:

$$\tilde{\boldsymbol{\chi}} := \tilde{\boldsymbol{\chi}}[\boldsymbol{n}, \mathscr{C}, \mathscr{X}] := D^{\mu}[\mathscr{V}]\mathscr{X}_{\mu}, \qquad (5.29)$$

and $\Delta_{FP}^{rc} := \det \left(\frac{\delta \chi}{\delta \theta}\right)_{\chi=0} = \det \left(\frac{\delta \chi}{\delta n^{\theta}}\right)_{\chi=0}$ is the Faddeev-Popov determinant associated with the reduction condition. It is important to note that the independent variables are regarded as $n^{\beta}(x)$, $C_{\nu}^{k}(x)$ and $X_{\nu}^{b}(x)$ in the reformulated Yang-Mills theory to simplify the Jacobian.

As already mentioned above, in order to obtain a well-defined Yang-Mills theory by completely fixing the gauge, we must impose the overall gauge fixing condition (see the down arrows in Fig. 1) in addition to the reduction condition. However, we have omitted to write this procedure while simplifying the presentation, since the overall gauge fixing can be performed according to standard procedures. Moreover, a systematic treatment of gauge fixing and the associated Faddeev-Popov determinant or the introduction of ghost fields can be carried out using the BRST symmetry of the new theory. Although the BRST treatment can also be performed for the SU(N)case following the method already performed for SU(2) [8], such a treatment is rather involved. Therefore, we have given a heuristic explanation based on the Faddeev-Popov trick for SU(N) in the above, even though it is possible to develop the BRST approach. Some applications of this reformulation will be given in a subsequent paper.

6 Conclusion and discussion

In this paper we have proposed a new version of the SU(N) Yang-Mills theory reformulated in terms of new field variables, which are obtained by a nonlinear change of variables from the original Yang-Mills gauge field. The basic idea of our reformulation originates from the Cho-Faddeev-Niemi decomposition of the gauge potential. However, our reformulation differs from other proposals [18, 19] for the gauge group $SU(N), N \geq 3$, including the SU(3) case, which is most interesting from the physical viewpoint, although it agrees with the conventional approach for the SU(2) gauge group. In particular, we have mainly studied the maximal and minimal cases.

This reformulation of Yang-Mills theory has already been applied to a few physical problems. It enables us to study the low-energy dynamics of Yang-Mills theory by explicitly extracting the topological degrees of freedom such as magnetic monopoles and vortices [25]. Such topological configurations are believed to provide the dominant degrees of freedom responsible for quark confinement from the viewpoint of dual superconductivity. Using the reformulation, the dual superconductivity in Yang-Mills theory can indeed be understood in a gauge-invariant way, as demonstrated recently by a non-Abelian Stokes theorem for the Wilson loop operator [15]. The non-Abelian Stokes theorem has played a crucial role in understanding the gauge-invariant meaning of the Abelian dominance in the Wilson loop operator [26], which was confirmed using the Abelian projection approach. From this viewpoint, the reformulation given in this paper will give a useful framework for other applications, which will be discussed in subsequent papers.

A SU(N) group

The group G = SU(N) has rank N - 1, and the Cartan subalgebra is constructed from N-1 diagonal generators H_k . Hence, there must be N(N-1) off-diagonal shift operators E_{α} , since dim $SU(N) := N^2 - 1 = (N-1) + N(N-1)$. Therefore, the total number of roots is N(N-1), of which there are N-1 simple roots, and the other roots are constructed as linear combinations of the simple roots.

An element of SU(N) in the fundamental representation is expressed by the $N \times N$ unitary matrices with determinant 1 that are generated by traceless Hermitian matrices, i.e., $N^2 - 1$ linearly independent generators $T^A(A = 1, \dots, N^2 - 1)$. The generators are normalized as $tr(T_A T_B) = \frac{1}{2}\delta_{AB}$. The diagonal generator H_m is defined by

$$(H_m)_{ab} = \frac{1}{\sqrt{2m(m+1)}} (\sum_{k=1}^m \delta_{ak} \delta_{bk} - m \delta_{a,m+1} \delta_{b,m+1})$$
(A.1)

$$= \frac{1}{\sqrt{2m(m+1)}} \operatorname{diag}(1, \cdots, 1, -m, 0, \cdots, 0), \qquad (A.2)$$

where for m = 1 to N - 1, the first m diagonal elements (beginning from the upper left-hand corner) of H_m are 1, the next element is -m, and the rest of the diagonal elements are 0. Thus, H_m is traceless. Each off-diagonal generator E_{α} has a single nonzero element $1/\sqrt{2}$.

The weight vector ν_j is defined as the eigenvector of H_j , i.e.,

$$H_j|\nu\rangle = \nu^j|\nu\rangle. \tag{A.3}$$

We have N weight vectors in the fundamental representation N (N-dimensional irreducible representation of SU(N)) given by

$$\nu^{1} = \left(\frac{1}{2}, \frac{1}{2\sqrt{3}}, \cdots, \frac{1}{\sqrt{2m(m+1)}}, \cdots, \frac{1}{\sqrt{2(N-1)N}}\right),$$

$$\nu^{2} = \left(-\frac{1}{2}, \frac{1}{2\sqrt{3}}, \cdots, \frac{1}{\sqrt{2m(m+1)}}, \cdots, \frac{1}{\sqrt{2(N-1)N}}\right),$$

$$\nu^{3} = \left(0, -\frac{1}{\sqrt{3}}, \frac{1}{2\sqrt{6}}, \cdots, \frac{1}{\sqrt{2(N-1)N}}\right),$$

$$\vdots$$

$$\nu^{m+1} = \left(0, 0, \cdots, 0, -\frac{m}{\sqrt{2m(m+1)}}, \cdots, \frac{1}{\sqrt{2(N-1)N}}\right),$$

$$\vdots$$

$$\nu^{N} = \left(0, 0, \cdots, 0, \frac{-N+1}{\sqrt{2(N-1)N}}\right).$$
(A.4)

All the weight vectors have the same length, and the angles between different weights are the same:

$$\nu^{i} \cdot \nu^{i} = \frac{N-1}{2N}, \quad \nu^{i} \cdot \nu^{j} = -\frac{1}{2N}, \quad (\text{for } i \neq j)$$
(A.5)

The weights constitute a polygon in the (N-1)-dimensional space. This implies that any weight can be used as the highest weight. A weight is called **positive** if its *last* nonzero component is positive. With this definition, the weights satisfy

$$\nu^1 > \nu^2 > \dots > \nu^N. \tag{A.6}$$

The simple roots are given by

$$\alpha^{i} = \nu^{i} - \nu^{i+1}, \quad (i = 1, \cdots, N-1)$$
 (A.7)

and are written explicitly as

$$\begin{aligned}
\alpha^{1} &= (1, 0, \cdots, 0), \\
\alpha^{2} &= \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}, 0, \cdots, 0\right), \\
\alpha^{3} &= \left(0, -\frac{1}{\sqrt{3}}, \sqrt{\frac{2}{3}}, 0, \cdots, 0\right), \\
\vdots \\
\alpha^{m} &= \left(0, 0, \cdots, -\sqrt{\frac{m-1}{2m}}, \sqrt{\frac{m+1}{2m}}, 0, \cdots, 0\right), \\
\vdots \\
\alpha^{N-1} &= \left(0, 0, \cdots, -\sqrt{\frac{N-2}{2(N-1)}}, \sqrt{\frac{N}{2(N-1)}}\right).
\end{aligned}$$
(A.8)

As can be shown from (A.5), all these roots have length 1, the angles between successive roots are the same, $2\pi/3$, and other pairs of roots are orthogonal:

$$\alpha^{j} \cdot \alpha^{j} = 1, \quad \alpha^{i} \cdot \alpha^{j} = -\frac{1}{2}, \quad (j = i \pm 1)$$
$$\alpha^{i} \cdot \alpha^{j} = 0, \quad (j \neq i, i \pm 1)$$
(A.9)

If we choose ν^1 as the highest-weight $\vec{\Lambda}$ of the fundamental representation **N**, some of the roots are orthogonal to ν^1 . From the above construction, it is easy to see that only one simple root α_1 is non-orthogonal to ν^1 , and that all the other simple roots are orthogonal:

$$\nu^{1} \cdot \alpha^{1} \neq 0, \quad \nu^{1} \cdot \alpha^{2} = \nu^{1} \cdot \alpha^{3} = \dots = \nu^{1} \cdot \alpha^{N-1} = 0.$$
 (A.10)

Therefore, all the linear combinations constructed from $\alpha^2, \dots, \alpha^{N-1}$ are also orthogonal to ν^1 . Nonorthogonal roots are only obtained when α^1 is included in the linear combinations. It is not difficult to show that the total number of non-orthogonal roots is 2(N-1), and hence there are N(N-1) - 2(N-1) = (N-2)(N-1) orthogonal roots. The (N-2)(N-1) shift operators E_{α} corresponding to these orthogonal roots together with the N-1 diagonal generators of the Cartan subalgebra H_k constitute the maximal stability subgroup $\tilde{H} = U(N-1)$, since (N-2)(N-1) + (N-1) = $(N-1)^2 = \dim U(N-1).$ Thus, for the fundamental representation, the stability subgroup \tilde{H} of SU(N) is given by

$$\tilde{H} = U(N-1). \tag{A.11}$$

To describe the coset space G/\tilde{H} , we need only N-1 complex numbers, since

$$G/\tilde{H} = SU(N)/U(N-1) = CP^{N-1}$$
 (A.12)

has dim $G/\tilde{H} = 2(N-1)$, where $CP^{N-1} = P^{N-1}(\mathbb{C})$ is the (N-1)-dimensional complex projective space, a submanifold of the flag manifold F_{N-1} .

B Derivation of identities

B.1 Maximal case

The identity (3.18) is proved as follows. We consider any su(N) Lie algebra valued function \mathscr{V} . Using the adjoint rotation, $\mathscr{V}' = U\mathscr{V}U^{\dagger}$, we have only to prove

$$\mathscr{V}' = \sum_{j=1}^{N-1} H_j(H_j, \mathscr{V}') + \sum_{j=1}^{N-1} [H_j, [H_j, \mathscr{V}']].$$
(B.1)

The Cartan decomposition for \mathscr{V}' is

$$\mathscr{V}' = \sum_{k=1}^{N-1} V^k H_k + \sum_{\alpha=1}^{(N^2 - N)/2} (W^{*\alpha} \tilde{E}_{\alpha} + W^{\alpha} \tilde{E}_{-\alpha}),$$
(B.2)

where the Cartan basis is given by

$$\vec{H} = (H_1, H_2, H_3, \cdots, H_{N-1}) = (T^3, T^8, T^{15}, \cdots, T^{N^2 - 1}),$$

$$\tilde{E}_{\pm 1} = \frac{1}{\sqrt{2}} (T^1 \pm iT^2), \quad \tilde{E}_{\pm 2} = \frac{1}{\sqrt{2}} (T^4 \pm iT^5),$$

$$\cdots, \quad \tilde{E}_{\pm (N^2 - N)/2} = \frac{1}{\sqrt{2}} (T^{N^2 - 3} \pm iT^{N^2 - 2}),$$
(B.3)

and the complex field is defined by

$$W^{1} = \frac{1}{\sqrt{2}} (V^{1} + iV^{2}), \quad W^{2} = \frac{1}{\sqrt{2}} (V^{4} + iV^{5}),$$

$$\cdots, \quad W^{(N^{2} - N)/2} = \frac{1}{\sqrt{2}} (V^{N^{2} - 3} + iV^{N^{2} - 2}).$$
(B.4)

We now calculate the double commutator as

$$[H_{j}, [H_{j}, \mathscr{V}']] = \sum_{j=1}^{N-1} V_{k}[H_{j}, [H_{j}, H_{k}]] + \sum_{\alpha=1}^{(N^{2}-N)/2} (W^{*\alpha}[H_{j}, [H_{j}, \tilde{E}_{\alpha}]] + W^{\alpha}[H_{j}, [H_{j}, \tilde{E}_{-\alpha}]]) = \sum_{\alpha=1}^{(N^{2}-N)/2} (W^{*\alpha}[H_{j}, \alpha_{j}\tilde{E}_{\alpha}] + W^{\alpha}[H_{j}, -\alpha_{j}\tilde{E}_{-\alpha}]) = \alpha_{j}\alpha_{j} \sum_{\alpha=1}^{(N^{2}-N)/2} (W^{*\alpha}\tilde{E}_{\alpha} + W^{\alpha}\tilde{E}_{-\alpha}).$$
(B.5)

On the other hand, we have

$$(H_{j}, \mathscr{V}') = \sum_{k=1}^{N-1} V_{k}(H_{j}, H_{k}) + \sum_{\alpha=1}^{(N^{2}-N)/2} (W^{*\alpha}(H_{j}, \tilde{E}_{\alpha}) + W^{\alpha}(H_{j}, \tilde{E}_{-\alpha})) = V^{j},$$
(B.6)

since $(H_j, \tilde{E}_\alpha) = \operatorname{tr}(H_j \tilde{E}_\alpha) = 0$. Thus the RHS of (B.1) reduces to

$$\sum_{j=1}^{N-1} V_j H_j + \sum_{j=1}^{N-1} \alpha_j \alpha_j \sum_{\alpha=1}^{(N^2 - N)/2} (W^{*\alpha} \tilde{E}_{\alpha} + W^{\alpha} \tilde{E}_{-\alpha}).$$
(B.7)

This is equal to the Cartan decomposition of $\mathscr V$ itself, since

$$\sum_{j=1}^{N-1} \alpha_j \alpha_j = 1. \tag{B.8}$$

B.2 Minimal case

Any su(N) Lie algebra valued function $\mathscr{F}(x)$ is decomposed into the \tilde{H} -commutative part $\mathscr{F}_{\tilde{H}}$ and the remaining part $\mathscr{F}_{G/\tilde{H}}$ as

$$\mathscr{F} = \mathscr{F}_{\tilde{H}} + \mathscr{F}_{G/\tilde{H}}, \quad \mathscr{F}_{\tilde{H}} = \tilde{\mathscr{F}} + \boldsymbol{h}(\boldsymbol{h}, \mathscr{F}), \quad \mathscr{F}_{G/\tilde{H}} = 2\frac{N-1}{N}[\boldsymbol{h}, [\boldsymbol{h}, \mathscr{F}]], \quad (B.9)$$

where $h(h, \mathscr{F}) = 2 \operatorname{tr}(\mathscr{F}h)h$ and we have defined the matrix $\tilde{\mathscr{F}}$ in which all the elements in both the last column and the last row are zero. It is important to remark that

$$[\tilde{\mathscr{F}}, \boldsymbol{h}] = 0. \tag{B.10}$$

This identity shows that the \tilde{H} -commutative part is not necessarily written in a form that is proportional to h and there are additional contributions of $\tilde{\mathscr{F}}$ to the \tilde{H} -commutative part. This forces us to use (3.53) for the \tilde{H} -commutative part.

For any su(N) Lie algebra valued function $\mathcal{M}(x)$, the following identity holds:

$$\mathscr{V} = \sum_{A=1}^{N^2 - 1} \mathscr{V}^A T_A = \tilde{\mathscr{V}} + \boldsymbol{h}(\boldsymbol{h}, \mathscr{V}) + 2 \frac{N - 1}{N} [\boldsymbol{h}, [\boldsymbol{h}, \mathscr{V}]], \quad (B.11)$$

where $(\mathbf{h}, \mathscr{V}) := 2 \operatorname{tr}(\mathscr{V}\mathbf{h})$ and we have defined the matrix $\widetilde{\mathscr{V}}$ in which all the elements in the last column and the last row are zero:

$$\tilde{\mathscr{V}} = \sum_{A=1}^{(N-1)^2 - 1} \mathscr{V}^A T_A = \sum_{A=1}^{(N-1)^2 - 1} (\mathscr{V}, T_A) T_A = \sum_{A=1}^{(N-1)^2 - 1} 2 \operatorname{tr}(\mathscr{V}T_A) T_A.$$
(B.12)

Note that $\tilde{\mathscr{V}} + \boldsymbol{h}(\boldsymbol{h}, \mathscr{V})$ commutes with \boldsymbol{n} , since $[\tilde{\mathscr{V}}, \boldsymbol{h}] = 0$.

The Cartan decomposition for $\mathscr{V}_N \in su(N)$ is

$$\mathcal{V}_{N} = \sum_{k=1}^{N-1} V_{k} H_{k} + \sum_{\alpha=1}^{(N^{2}-N)/2} (W^{*\alpha} \tilde{E}_{\alpha} + W^{\alpha} \tilde{E}_{-\alpha})$$

$$= \tilde{\mathcal{V}}_{N} + M_{N-1} H_{N-1} + \sum_{\alpha=[(N-1)^{2}-(N-1)]/2+1}^{(N^{2}-N)/2} (W^{*\alpha} \tilde{E}_{\alpha} + W^{\alpha} \tilde{E}_{-\alpha}).$$
(B.13)

We now calculate the double commutator (r = N - 1) as

$$[H_{r}, [H_{r}, \mathscr{V}_{N}]]$$

$$= \sum_{j=1}^{N-1} V_{k}[H_{r}, [H_{r}, H_{k}]] + \sum_{\alpha=1}^{(N^{2}-N)/2} (W^{*\alpha}[H_{r}, [H_{r}, \tilde{E}_{\alpha}]] + W^{\alpha}[H_{r}, [H_{r}, \tilde{E}_{-\alpha}]])$$

$$= \sum_{\alpha=[(N-1)^{2}-(N-1)]/2+1}^{(N^{2}-N)/2} (W^{*\alpha}[H_{r}, \alpha_{r}\tilde{E}_{\alpha}] + W^{\alpha}[H_{r}, -\alpha_{r}\tilde{E}_{-\alpha}])$$

$$= \alpha_{r}\alpha_{r} \sum_{\alpha=[(N-1)^{2}-(N-1)]/2+1}^{(N^{2}-N)/2} (W^{*\alpha}\tilde{E}_{\alpha} + W^{\alpha}\tilde{E}_{-\alpha}). \qquad (B.14)$$

On the other hand, we have

$$(H_r, \mathscr{V}_N) = \sum_{k=1}^{N-1} V_k(H_r, H_k) + \sum_{\alpha=1}^{(N^2 - N)/2} (W^{*\alpha}(H_r, \tilde{E}_{\alpha}) + W^{\alpha}(H_r, \tilde{E}_{-\alpha})) = V_r,$$
(B.15)

since $(H_j, \tilde{E}_\alpha) = \operatorname{tr}(H_j \tilde{E}_\alpha) = 0.$

For any Lie algebra valued function $\mathscr{V}_N(x)$, we obtain the identity

$$\begin{aligned} \mathscr{V}_N &= \sum_{A=1}^{N^2 - 1} \mathscr{V}_A T_A \\ &= \widetilde{\mathscr{V}}_N + (\mathscr{V}_N, H_r) H_r + \frac{1}{\alpha_r^2} [H_r, [H_r, \mathscr{V}_N]] \\ &= \widetilde{\mathscr{V}}_N + (\mathscr{V}_N, H_r) H_r + \frac{2(N-1)}{N} [H_r, [H_r, \mathscr{V}_N]]. \end{aligned}$$
(B.16)

C More SU(3) cases

Recall that \mathbf{n}_3 and \mathbf{n}_8 are representatives of the maximal case and minimal case, respectively, which are distinguished by the value of ϑ . To take into account all the degrees of freedom of \mathbf{n} , it is necessary to introduce another commutative and orthogonal unit vector \mathbf{n}' in addition to \mathbf{n} , i.e., $\mathbf{n} \cdot \mathbf{n}' = 0$, since $\{\mathbf{n}, \mathbf{n}'\}$ constitutes the complete set for bases of the maximal Abelian subgroup $U(1) \times U(1)$.¹² Such an

¹²This does not necessarily mean that the analogous relationship for $\{n_3, n_8\}$ also holds for $\{n, n'\}$.

example of \mathbf{n}' for \mathbf{n} is given by

$$\mathbf{n} = \cos \vartheta \mathbf{n}_3 + \sin \vartheta \mathbf{n}_8,\tag{C.1}$$

$$\mathbf{n}' = \cos(\vartheta + \pi/2)\mathbf{n}_3 + \sin(\vartheta + \pi/2)\mathbf{n}_8$$

= $-\sin\vartheta\mathbf{n}_3 + \cos\vartheta\mathbf{n}_8$, (C.2)

which has the matrix representation

$$\begin{pmatrix} \mathbf{n} \\ \mathbf{n}' \end{pmatrix} = \begin{pmatrix} \cos\vartheta & \sin\vartheta \\ -\sin\vartheta & \cos\vartheta \end{pmatrix} \begin{pmatrix} \mathbf{n}_3 \\ \mathbf{n}_8 \end{pmatrix}.$$
(C.3)

In fact, the set $\{\mathbf{n}, \mathbf{n}'\}$ satisfies the key relation

$$\mathbf{V} = \mathbf{n}(\mathbf{V} \cdot \mathbf{n}) + \mathbf{n}'(\mathbf{V} \cdot \mathbf{n}') + \mathbf{n} \times (\mathbf{V} \times \mathbf{n}) + \mathbf{n}' \times (\mathbf{V} \times \mathbf{n}'), \quad (C.4)$$

which is derived by combining

$$\mathbf{n} \times (\mathbf{V} \times \mathbf{n}) + \mathbf{n}' \times \mathbf{V} \times \mathbf{n}') = \mathbf{n}_3 \times (\mathbf{V} \times \mathbf{n}_3) + \mathbf{n}_8 \times (\mathbf{V} \times \mathbf{n}_8), \qquad (C.5)$$

and

$$(\mathbf{V} \cdot \mathbf{n})\mathbf{n} + (\mathbf{V} \cdot \mathbf{n}')\mathbf{n}' = (\mathbf{V} \cdot \mathbf{n}_3)\mathbf{n}_3 + (\mathbf{V} \cdot \mathbf{n}_8)\mathbf{n}_8.$$
(C.6)

Thus, we obtain the decomposition in the vector form as

$$\mathbf{A}_{\mu} = \mathbf{V}_{\mu} + \mathbf{X}_{\mu} = \mathbf{C}_{\mu} + \mathbf{B}_{\mu} + \mathbf{X}_{\mu},$$

$$\mathbf{C}_{\mu} = (\mathbf{n} \cdot \mathbf{A}_{\mu})\mathbf{n} + (\mathbf{n}' \cdot \mathbf{A}_{\mu})\mathbf{n}',$$

$$\mathbf{B}_{\mu} = g^{-1}\partial_{\mu}\mathbf{n} \times \mathbf{n} + g^{-1}\partial_{\mu}\mathbf{n}' \times \mathbf{n}',$$

$$\mathbf{X}_{\mu} = g^{-1}\mathbf{n} \times D_{\mu}[\mathbf{A}]\mathbf{n} + g^{-1}\mathbf{n}' \times D_{\mu}[\mathbf{A}]\mathbf{n}'.$$
 (C.7)

This decomposition is obtained by solving the following defining equations: (a) the covariant constantness of \mathbf{n} and \mathbf{n}' in the background \mathbf{V}_{μ} :

$$0 = D_{\mu}[\mathbf{V}]\boldsymbol{n}_{s}(x) := \partial_{\mu}\mathbf{n}_{s}(x) + g\mathbf{V}_{\mu}(x) \times \mathbf{n}_{s}(x) \quad (\mathbf{n}_{s} = \mathbf{n}, \mathbf{n}'), \quad (C.8)$$

and (b) the orthogonality of \boldsymbol{n} and \boldsymbol{n}' to \mathbf{X}_{μ} :

$$0 = \mathbf{n}_s(x) \cdot \mathbf{X}_\mu(x) := \mathbf{n}_s^A(x) X_\mu^A(x) \quad (\mathbf{n}_s = \mathbf{n}, \mathbf{n}').$$
(C.9)

These defining equations can be solved using the completeness identity (C.4). It is easy to see that we can obtain the same reduction condition as that given in the text.

We consider the determinant of the color field \boldsymbol{n} as a function of ϑ , which is invariant under the similarity transformation (adjoint rotation) $\boldsymbol{n} \to U^{\dagger} \boldsymbol{n} U$. Therefore, it can be a character for classification. The minimal case, i.e., $\vartheta = \frac{1}{6}\pi, \frac{1}{2}\pi, \frac{5}{6}\pi, \frac{7}{6}\pi, \frac{3}{2}\pi, \frac{11}{6}\pi$ is realized if and only if $|\det(\boldsymbol{n})|$ takes the maximum value, i.e., $\det(\boldsymbol{n}) = \pm 1$:

$$\det(\boldsymbol{n}) = -\frac{1}{3\sqrt{3}}\sin\vartheta\left(\sin\vartheta - \frac{\sqrt{3}}{2}\right)\left(\sin\vartheta + \frac{\sqrt{3}}{2}\right) = \frac{1}{12\sqrt{3}}\sin3\vartheta, \qquad (C.10)$$

or equivalently, if and only if $det(\mathbf{n}') = 0$, since

$$\det(\mathbf{n}') = -\frac{1}{3\sqrt{3}}\cos\vartheta\left(\cos\vartheta - \frac{\sqrt{3}}{2}\right)\left(\cos\vartheta + \frac{\sqrt{3}}{2}\right) = \frac{1}{12\sqrt{3}}\cos3\vartheta.$$
(C.11)

On the other hand, $\det(\mathbf{n}) = 0$ is obtained for $\vartheta = 0, \pi/3, 2\pi/3, \pi, 4\pi/3, 5\pi/3$. It turns out that these cases represent the conventional approach. In fact, we find that $\det(\mathbf{n}_3) = 0$.

Acknowledgments

This work is financially supported by a Grant-in-Aid for Scientific Research (C) No. 18540251 from the Japan Society for the Promotion of Science (JSPS).

References

- [1] C.N. Yang and R.L. Mills, Phys. Rev. **96**, 191(1954).
- [2] G. 't Hooft, Nucl.Phys. B **190** [FS3], 455(1981).
- [3] A. Kronfeld, M. Laursen, G. Schierholz and U.-J. Wiese, Phys.Lett. B 198, 516(1987).
- [4] L. Del Debbio, M. Faber, J. Giedt, J. Greensite and S. Olejnik, [hep-lat/9801027], Phys. Rev. D58, 094501 (1998).
 L. Del Debbio, M. Faber, J. Greensite and S. Olejnik, [hep-lat/9610005], Phys. Rev. D55, 2298(1997).
- [5] Y.M. Cho, Phys. Rev. D 21, 1080(1980).
 Y.M. Cho, Phys. Rev. D 23, 2415(1981).
- [6] L. Faddeev and A.J. Niemi, [hep-th/9807069], Phys. Rev. Lett. 82, 1624(1999).
 L.D. Faddeev and A.J. Niemi, [hep-th/0608111], Nucl. Phys. B776, 38(2007).
- S.V. Shabanov, [hep-th/9903223], Phys. Lett. B 458, 322(1999).
 S.V. Shabanov, [hep-th/9907182], Phys. Lett. B 463, 263(1999).
- [8] K.-I. Kondo, T. Murakami and T. Shinohara, [hep-th/0504107], Prog. Theor. Phys. 115, 201(2006).
 K.-I. Kondo, T. Murakami and T. Shinohara, [hep-th/0504198], Eur. Phys. J. C 42, 475(2005).
- [9] K.-I. Kondo, [hep-th/0404252], Phys.Lett. B600, 287(2004).
- [10] K.-I. Kondo, [hep-th/0609166], Phys. Rev. D 74, 125003 (2006).
- [11] S. Kato, K.-I. Kondo, T. Murakami, A. Shibata, T. Shinohara and S. Ito, [heplat/0509069], Phys. Lett. B 632, 326(2006).
- [12] S. Ito, S. Kato, K.-I. Kondo, T. Murakami, A. Shibata and T. Shinohara, [heplat/0604016], Phys. Lett. B 645, 67(2007).
- [13] A. Shibata, S. Kato, K.-I. Kondo, T. Murakami, T. Shinohara and S. Ito, e-Print: arXiv:0706.2529 [hep-lat], Phys.Lett. B653, 101(2007).
- [14] K. Wilson, Phys. Rev. D 10, 2445(1974).
- [15] K.-I. Kondo, arXiv:0801.1274 [hep-th], Phys. Rev. D 77, 085029 (2008).
- [16] K.-I. Kondo, [hep-th/9805153], Phys. Rev. D 58, 105016 (1998).
- [17] K.-I. Kondo and Y. Taira, [hep-th/9906129], Mod. Phys. Lett. A 15, 367(2000);
 K.-I. Kondo and Y. Taira, [hep-th/9911242], Prog. Theor. Phys. 104, 1189(2000).
 K.-I. Kondo and Y. Taira, Nucl. Phys. Proc. Suppl. 83, 497(2000).

- [18] Y.M. Cho, MPI-PAE/PTh 14/80 (1980).
 Y.M. Cho, Phys. Rev. Lett. 44, 1115(1980).
 Y.D. Kim, I.G. Koh and Y.J. Park, Phys. Rev. D 25, 587(1982).
 W.S. L'Yi, Y.J. Park, I.G. Koh, Y. Kim, Phys. Rev. D 25, 3308(1982).
- [19] L. Faddeev and A.J. Niemi, [hep-th/9812090], Phys. Lett.B 449, 214(1999).
- [20] L. Faddeev and A.J. Niemi, [hep-th/9907180], Phys. Lett.B 464, 90(1999).
- [21] S.V. Shabanov, [hep-th/0202146], J. Math. Phys. 43, 4127(2002).
- [22] E.J. Weinberg and P. Yi, [hep-th/0609055], Phys. Rept. 438, 65(2007).
- [23] A. Shibata, S. Kato, K.-I. Kondo, T. Murakami, T. Shinohara, and S. Ito, e-Print: arXiv:0710.3221 [hep-lat], POS(LATTICE-2007) 331, Oct 2007, 7pp, Talk given at 25th International Symposium on Lattice Field Theory, Regensburg, Germany, 30 Jul - 4 Aug 2007.
- [24] K.-I. Kondo, A. Shibata, T. Shinohara, T. Murakami, S. Kato and S. Ito, Chiba Univ. Preprint, CHIBA-EP-168, e-Print: arXiv:0803.2451 [hep-lat].
- [25] K.-I. Kondo, Chiba Univ. Preprint, CHIBA-EP-171, arXiv:0802.3829 [hep-th],
 J. Phys. G: Nucl. Part. Phys. 35, 085001 (2008).
- [26] K.-I. Kondo and A. Shibata, Chiba Univ. Preprint, CHIBA-EP-170, arXiv:0801.4203 [hep-th].