

Equilibrium boundary conditions, dynamic vacuum energy, and the Big Bang

F.R. Klinkhamer*

*Institute for Theoretical Physics,
University of Karlsruhe (TH),
76128 Karlsruhe, Germany*

Abstract

The smallness of the cosmological constant Λ in an equilibrium context may be due to the existence of a self-tuning vacuum variable q . Here, a nonequilibrium context is considered with a corresponding time-dependent cosmological parameter $\Lambda(t)$ or vacuum energy density $\rho_V(t)$. A specific model of a closed Friedmann–Robertson–Walker universe is presented, which is determined by equilibrium boundary conditions at one particular instant of time and an appropriate form of vacuum-energy dynamics. The model has a standard Big Bang phase at early times and reproduces the main characteristics of the present universe.

PACS numbers: 11.30.Cp, 04.20.Cv, 98.80.Jk, 95.36.+x

Keywords: Lorentz invariance, general relativity, cosmology, dark energy

*Electronic address: frans.klinkhamer@physik.uni-karlsruhe.de

I. INTRODUCTION

It has been argued that the effective cosmological constant Λ [1, 2, 3, 4, 5] (or gravitating vacuum energy density ρ_V) vanishes in a perfect quantum vacuum, provided that this vacuum can be considered to be a self-sustained medium at zero external pressure and that there exists a new type of conserved charge q which self-adjusts so as to give vanishing internal pressure [6]. As the perfect quantum vacuum is taken to be Lorentz invariant (see, e.g., Refs. [7, 8] for bounds on Lorentz violation), this vacuum variable q must be of an entirely new type, different from known conserved charges such as baryon minus lepton number, $B - L$. The detailed microscopic theory is, of course, unknown, but two examples of possible theories with such a vacuum variable q have been given in Ref. [6].

For the perfect Lorentz-invariant quantum vacuum, the vacuum variable q would be constant over the whole of spacetime. The previous discussion then applies to an equilibrium situation and describes what may be called the “statics of dark energy.” The outstanding questions are how the equilibrium argument relates to the observed *expanding* universe and which physical principle governs the “dynamics of dark energy.” Obviously, these are profound questions and the present article can only hope to provide a small step towards a possible solution. In fact, the first question is temporarily replaced by the following restricted question: *is it possible at all to relate equilibrium boundary conditions for $\rho_V(0)$ to an expanding universe which matches the observations, even if we are free to choose the type of vacuum-energy dynamics, $d\rho_V/dt \neq 0$?* In mathematical terms, we are after an “existence proof” for this type of universe, with equilibrium boundary conditions setting the numerical value of the vacuum energy density ρ_V at one moment in time.

It turns out to be rather difficult to construct such an “existence proof,” but, in the end, we have been able to find one suitable universe. The main lesson we will learn from this exercise is the necessity of some type of metastability of the imperfect quantum vacuum (Lorentz invariance being perturbed by the presence of matter) and we will get some idea of what type of metastability would be required to reproduce our known universe [2, 3, 4, 5]. In a way, we are seeking the “Kepler laws” of our accelerating universe [5] and leave the underlying physics to future generations.

II. DYNAMIC VACUUM ENERGY DENSITY

The crucial issue is the exchange of energy between the deep vacuum (as described by the variable q , for example) and the low-energy degrees of freedom corresponding to the physics of the standard model and general relativity. The detailed microscopic theory is unknown, but we can try to seek guidance from the concrete four-form theory discussed in Ref. [6].

This particular theory, coupled to low-energy matter, is given by the action [6, 9, 10]

$$S = \int_{\mathbb{R}^4} d^4x \sqrt{-g} \left(\frac{R}{16\pi G} - \frac{1}{2} \partial_\mu \Phi \partial_\nu \Phi g^{\mu\nu} - \epsilon(F) \left(1 - \frac{1}{2} \Phi^2/M^2 \right) \right). \quad (1a)$$

$$F^2 \equiv -\frac{1}{24} F_{\mu\nu\rho\sigma} F_{\alpha\beta\gamma\delta} g^{\alpha\mu} g^{\beta\nu} g^{\gamma\rho} g^{\delta\sigma}, \quad (1b)$$

where $R(x)$ is the Ricci curvature scalar from the metric $g_{\mu\nu}(x)$, $F_{\mu\nu\rho\sigma}(x)$ the four-form field strength of a three-form gauge field $A_{\nu\rho\sigma}(x)$, $\epsilon(F)$ an arbitrary even function of F , and $\Phi(x)$ a real scalar field with coupling constant $1/M^2$ to the deep-vacuum energy density $\epsilon(F)$. As will become clear, a strong hierarchy $M^4 \gg \epsilon(F)$ appears to be required in order to produce the present universe. Here, and in the rest of this section, we use natural units with $\hbar = c = 1$.

The variational principle applied to action (1a) results in three field equations, a generalized Maxwell equation for the $F_{\mu\nu\rho\sigma}$ field, a generalized Klein–Gordon equation for the Φ field, and the Einstein equation for the $g_{\mu\nu}$ field with an energy-momentum tensor from both $F_{\mu\nu\rho\sigma}$ and Φ fields.

Now, take the following *Ansatz*: a spatially-flat ($k = 0$) Friedmann–Robertson–Walker (FRW) metric, a Levi–Civita-type four-form field, and a homogenous scalar field. Specifically, the *Ansatz* fields are given by

$$g_{\mu\nu}(x) = \text{diag}(+1, -a^2(t), -a^2(t), -a^2(t)), \quad (2a)$$

$$F_{\mu\nu\rho\sigma}(x) = q(t) e_{\mu\nu\rho\sigma}, \quad (2b)$$

$$\Phi(x) = \Phi(t), \quad (2c)$$

with scale factor $a(t)$ and totally antisymmetric Levi–Civita symbol $e_{\mu\nu\rho\sigma}$. In terms of the vacuum compressibility [6] $\chi \equiv 1/(q^2 d^2\epsilon/dq^2)$, the generalized Maxwell equation reduces to

$$\frac{\dot{q}}{q} = \chi q \epsilon' \frac{\Phi \dot{\Phi}}{M^2 - \frac{1}{2} \Phi^2}, \quad (3)$$

where the overdot stands for differentiation with respect to the time coordinate t and the prime for differentiation with respect to the vacuum variable q . As a first approximation, consider the case of an incompressible fluid, $\chi = 0$, so that the vacuum variable does not change with time, $q = q_0 = \text{const}$.

For the *Ansatz* fields, the vacuum energy density in the Einstein field equation is given by

$$\rho_V = \left(\epsilon(q) - q \frac{d\epsilon}{dq} \right) \left(1 - \frac{1}{2} \Phi^2/M^2 \right) \equiv \tilde{\epsilon}(q) \left(1 - \frac{1}{2} \Phi^2/M^2 \right), \quad (4)$$

which equals the previous result $\tilde{\epsilon}(q)$ from Ref. [6] multiplied by an overall factor containing Φ^2/M^2 . [The reason for obtaining this simple result is that the metric does not enter the coupling of Φ^2 and $\epsilon(F)$ in (1a).] The time derivative of this vacuum energy density is simply

$$\dot{\rho}_V = -\tilde{\epsilon} \Phi \dot{\Phi}/M^2, \quad (5)$$

for vanishing \dot{q} due to the assumed vacuum incompressibility, $\chi = 0$.

At this moment, it is useful to define

$$\mu_0^2 \equiv \tilde{\epsilon}(q_0)/M^2, \quad (6)$$

for the constant equilibrium value q_0 (assuming $\mu_0^2 \geq 0$), and to recall that the energy-momentum tensor of the perfect fluid corresponding to the homogeneous scalar field $\Phi(t)$ has the following energy density and pressure [3, 4, 5]:

$$\rho_M = \frac{1}{2} \dot{\Phi}^2 + \frac{1}{2} \mu_0^2 \Phi^2, \quad P_M = \frac{1}{2} \dot{\Phi}^2 - \frac{1}{2} \mu_0^2 \Phi^2. \quad (7)$$

Also introduce the matter and vacuum equations of state

$$P_M = w \rho_M, \quad P_V = w_V \rho_V = -\rho_V. \quad (8)$$

Relation (5) can then be written solely in terms of μ_0 , $\rho_M(t)$, and $w(t)$:

$$\dot{\rho}_V = -\text{sgn}(\mu_0^2 \Phi \dot{\Phi}) |\mu_0| \sqrt{1 - w^2(t)} \rho_M(t), \quad (9)$$

which allows us to study simple cosmological models. However, (9) holds only for the case of $\dot{q} = 0$ (having assumed $\chi = 0$), for matter described by a single scalar field Φ , and for the flat $k = 0$ FRW universe. Therefore, we keep, in the next section, the equation for $\dot{\rho}_V$ as general as possible, but still proportional to ρ_M .

III. CLOSED FRW UNIVERSE

The spatially flat ($k = 0$) FRW universe does not have an obvious time where to apply equilibrium boundary conditions, apart from the limiting case $a(t) \rightarrow \infty$ for $t \rightarrow \infty$. But, for $t \rightarrow \infty$, the matter density can be expected to vanish, which complicates the discussion. For this reason, we turn to the closed ($k = 1$) FRW universe with metric [2],

$$g_{00}(x) = 1, \quad g_{m0}(x) = 0, \quad g_{mn}(x) = -a^2(t) \tilde{g}_{mn}(x), \quad (10)$$

in terms of the standard metric \tilde{g}_{mn} of a unit 3-sphere for spatial indices $m, n = 1, 2, 3$. The scale factor $a(t)$ now corresponds to the radius of the universe. Henceforth, we discuss only classical relativity and use units with $c = 8\pi G/3 = 1$, unless stated otherwise.

The closed ($k = 1$) FRW universe can have an equilibrium point $t_{\text{eq}} \equiv 0$ where the expansion momentarily stops ($\dot{a}/a = 0$), provided the following condition holds:

$$(8\pi G/3) \left(\rho_V(t_{\text{eq}}) + \rho_M(t_{\text{eq}}) \right) = k a(t_{\text{eq}})^{-2} \Big|_{k=1}, \quad (11)$$

with the dimensionless curvature parameter k shown explicitly. A thermodynamical argument suggests a further Gibbs–Duhem-like condition at the equilibrium point:

$$\rho_V(t_{\text{eq}}) = w \rho_M(t_{\text{eq}}) + \frac{1}{2} (1 + w) \rho_M(t_{\text{eq}}). \quad (12)$$

Here, the first term on the right-hand side corresponds to the flat-spacetime condition $\rho_V = P_M = w \rho_M$ from pressure equilibrium $P_V + P_M = P_{\text{ext}} = 0$ and the vacuum equation of state $P_V = -\rho_V$; see Ref. [6] and references therein. The second term on the right-hand side of (12) describes the gravitational effects, even though Newton’s gravitational constant G does not appear explicitly. Specifically, the complete relation (12) follows from the conditions $P_V + P_M + P_{\text{grav}} = 0$ and $\rho_V + \rho_M + \rho_{\text{grav}} = 0$ for a gravitational equation of state $P_{\text{grav}} = -(1/3) \rho_{\text{grav}}$; see Sec. 7 of Ref. [11] for further discussion. The conditions (11) and (12) will be seen to nullify the respective right-hand sides of the differential equations (16) and (13) below.

Consider, then, the dynamics of this closed ($k = 1$) FRW universe, which is governed by the 00–component of the Einstein equation,

$$\ddot{a}/a = -(4\pi G/3) \left(\rho_{\text{total}} + 3 P_{\text{total}} \right) = (8\pi G/3) \left(\rho_V - \frac{1}{2} (1 + 3w) \rho_M \right), \quad (13)$$

the energy-conservation equation,

$$(\dot{\rho}_V + \dot{\rho}_M) = -3(\dot{a}/a) (1 + w) \rho_M, \quad (14)$$

and a dynamical vacuum-energy equation which generalizes (9). In addition, there are the equations of state (8) for matter and vacuum with, for simplicity, a time-independent value of the matter parameter w .

Given boundary conditions (11) and (12), the problem is to get away from the *static* Einstein universe [1]. It appears that the only way to achieve this is to consider either a modification of gravity (e.g., modified Einstein field equations as discussed in Ref. [12]) or a new type of metastability of the perturbed quantum vacuum as will be discussed here. Specifically, we assume the time variation of the vacuum energy to be described by the following *Ansatz*:

$$\dot{\rho}_V = (1/\tau) \gamma \rho_M, \quad (15)$$

where τ is a fundamental time constant and $\gamma = \gamma[a(t)]$ a dimensionless functional with appropriate normalization. As mentioned before, the origin of (15) needs to be explained

by the detailed microphysics [perhaps similar to the discussion in Sec. II leading up to (9)], but, here, we take a purely phenomenological (“Keplerian”) approach and simply assume a particular form for $\dot{\rho}_V$.

Several remarks on (15) are in order:

1. $\dot{\rho}_V$ vanishes if $\rho_M = 0$, so that the perfect Lorentz-invariant quantum vacuum without matter remains stable;
2. $\dot{\rho}_V$ does not necessarily vanish if $\dot{a}/a = 0$ and, in particular, $\dot{\rho}_V$ does not vanish at $t = t_{\text{eq}}$, so that the universe can get away from the static case;
3. time-reversal invariance around t_{eq} is broken, as long as γ in (15) does not depend on t explicitly.

In fact, our *Ansatz* (15) resembles Eq. (8) of Ref. [13], but differs by point 1 in particular. From point 2 and assuming $\gamma[a(t_{\text{eq}})]/\tau > 0$, there is the possibility of having a “Big Bang” with $a(t_{\text{BB}}) = 0$ at $t_{\text{BB}} < t_{\text{eq}}$. From point 3, there is also the possibility that the universe does not return to vanishing 3-volume for $t > t_{\text{eq}}$ but keeps expanding forever. With nonnegative $\gamma[a(t)]/\tau$ for $t > t_{\text{eq}}$, the universe may approach a de-Sitter-like universe [2, 3, 4], if $\rho_M \rightarrow 0$ and $\rho_V \rightarrow \text{const}$ for $t \rightarrow \infty$.

Equations (13), (14), and (15) with boundary conditions (11) and (12) can now be solved numerically to give $a(t)$, $\rho_M(t)$, and $\rho_V(t)$. As we intend to take equilibrium-point boundary conditions also for the standard case with $\rho_V(t) = \gamma[a(t)]/\tau = 0$, we use the second-order Einstein equation (13) instead of the first-order Friedmann equation,

$$(\dot{a}/a)^2 = (8\pi G/3) (\rho_V + \rho_M) - k/a^2 \Big|_{k=1}. \quad (16)$$

It is well known [2] that, with appropriate boundary conditions, the differential equations (13) and (16) are equivalent when combined with the energy-conservation equation (14). Incidentally, the 11-component of the Einstein equation is also satisfied, as are the 22 and 33 components by isotropy.

In the next section, we will make a specific choice for $\gamma[a]$ and will present a particular model universe with dynamic vacuum energy density. For comparison, we give here a standard universe with the same extremal radius at $t = t_{\text{max}} \equiv 0$ (in units with $8\pi G/3 = c = 1$),

$$\begin{pmatrix} a(0) \\ \rho_M(0) \\ \rho_V(0) \\ w \\ 1/\tau \end{pmatrix} = \begin{pmatrix} 10 \\ 1/100 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad (17)$$

closed FRW universe with pressureless matter




FIG. 1: Closed FRW universe with pressureless matter ($w = 0$) and boundary conditions (17) for units with $8\pi G/3 = c = 1$. On the first row are shown the scale factor $a(t)$ and various derivatives, \dot{a}/a and \ddot{a}/a . On the second row are shown $a(t)$ scaled by a power of the elapsed time since $t_{\text{BB}} = -5\pi \approx -15.71$ where $a(t)$ vanishes, the matter energy density ρ_M multiplied by a^3 , and the matter density parameter $\Omega_M \equiv \rho_M/\rho_c$ defined in terms of the critical density $\rho_c \equiv (\dot{a}/a)^2$. The two boundary conditions on a and ρ_M at $t = t_{\text{max}} \equiv 0$ are indicated by the heavy dots.

so that (15) becomes trivial. Note that boundary conditions (17) imply $\dot{a}/a = 0$ at $t = 0$ by the Friedmann equation (16). The corresponding numerical solution is displayed in Fig. 1 and the analytic solution is given by [2]

$$\rho_M(a) = a_{\text{max}}/a^3, \quad a = a_{\text{max}} \sin^2(\theta/2), \quad (18a)$$

$$\rho_V(a) = 0, \quad t = (\theta - \sin \theta - \pi) a_{\text{max}}/2, \quad (18b)$$

for an auxiliary angle $\theta \in [0, 2\pi]$. The time-symmetric solution (18ab) has Big Bang coordinate time $t_{\text{BB}} = -\pi a_{\text{max}}/2$ and Big Crunch coordinate time $t_{\text{BC}} = +\pi a_{\text{max}}/2$. For $t \downarrow t_{\text{BB}}$, the behavior of $a(t)$ approaches that of the flat ($k = 0$) FRW universe, $a(t) \propto (t - t_{\text{BB}})^{2/3}$.

IV. MODEL UNIVERSE FROM $\gamma[a]$ ANSATZ

As explained in the Introduction, our goal is relatively modest: to find at least one functional $\gamma[a]$ so that equations (13), (14) and (15), with boundary conditions (11) and (12) can produce a solution which more or less reproduces our known universe (see, e.g., Refs. [14, 15, 16] and references therein), which is spatially flat to a high degree of precision and approximately consists of 75% “dark energy” and 25% matter (primarily nonbaryonic “cold dark matter”). With three coupled nonlinear ordinary differential equations (ODEs), this modest goal is surprisingly difficult to reach. Still, we have been successful by first

considering the inverse problem: (1) to find, given a more or less reasonable $a_{\text{designer}}(t)$, which densities $\rho_M(t)$ and $\rho_V(t)$ are required; (2) to determine, by differentiation of the $\rho_V(t)$ from the first step, the required $\gamma[a(t)]/\tau$ from (15).

Inspired by these designer-universe results, we make the following *Ansatz* for the (dimensionless) vacuum dynamics functional:

$$\begin{aligned} \gamma[a] = N \frac{a}{a_{\text{eq}}} & \left(\frac{(a_{\text{eq}} - a)^6}{(c_1 a_{\text{eq}})^6 + (a_{\text{eq}} - a)^6} \frac{a}{a_{\text{eq}}} \sin \left(c_2 \pi \frac{a}{a_{\text{eq}}} \right) \right. \\ & \left. + \frac{a^6}{(c_1 a_{\text{eq}})^6 + a^6} \left(\frac{(c_3 a_{\text{eq}})^{1/3}}{(c_3 a_{\text{eq}})^{1/3} + |a_{\text{eq}} - a|^{1/3}} \right)^4 \right), \end{aligned} \quad (19)$$

with numerical coefficients c_n and normalization factor $N \equiv 1 + (c_1)^6$, so that $\gamma[a_{\text{eq}}] = 1$. Roughly speaking, this *Ansatz* for $\gamma[a]$ consists of a term $a^2 \sin a$ modulated to be effective near $a = 0$ and a sharply-peaked positive term of the form $(1-t)^{-4}$ modulated to be effective just below $a = a_{\text{eq}}$ for $t \leq 0$. The behavior $\gamma[a] \propto a^3$ near $a = 0$ will be seen to allow for a finite limiting value of $\rho_V(a)$ by compensating the divergent behavior of $\rho_M \propto 1/a^3$ on the right-hand side of (15).

Now take the following numerical values (in units with $8\pi G/3 = c = 1$) for the boundary conditions (11)–(12), the vacuum decay constant $1/\tau$, and the *Ansatz* coefficients c_n :

$$\begin{pmatrix} a(0) \\ \rho_M(0) \\ \rho_V(0) \\ w \\ 1/\tau \\ c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 10 \\ 2/300 \\ 1/300 \\ 0 \\ 50 \\ 1/5 \\ 9/4 \\ 1/15 \end{pmatrix}, \quad (20)$$

with the implicit equilibrium condition $\dot{a}/a = 0$ at $t = 0$ from the Friedmann equation (16). The numerical solution of the coupled ODEs (13), (14), and (15) with these boundary conditions is given in Fig. 2. Observe that $\gamma[a]/\tau = 0$ would give a static universe with $a(t) = a(0)$, $\rho_V(t) = \rho_V(0)$, and $\rho_M(t) = \rho_M(0)$ at the values indicated by the heavy dots in Fig. 2. As explained in the previous section, we appeal to a new type of metastability of the perturbed quantum vacuum ($\gamma/\tau \neq 0$) in order to get away from this static universe.

Turning to the detailed model results of Fig. 2, the “Big Bang” would occur at coordinate time $t = t_{\text{BB}} = -0.91636$, which differs by one order of magnitude from the result without vacuum energy in Fig. 1. The “present universe” [14, 15, 16] with density ratio $\rho_V/\rho_M \approx 3$ and total density parameter $\Omega_V + \Omega_M \approx 1$ (see below for its definition) would approximately correspond to the time $t = t_0 = -0.5842$ in Fig. 2 (choosing the latest time of two possible

closed FRW universe with pressureless matter and dynamic vacuum energy




FIG. 2: Closed FRW universe with pressureless matter ($w = 0$) and dynamic vacuum energy ($w_V = -1$) for boundary conditions (20) and units with $8\pi G/3 = c = 1$. The assumed behavior of the vacuum-energy dynamics is given by (15) with the functional $\gamma[a(t)]$ from (19). The three nonzero equilibrium boundary conditions on a , ρ_M , and ρ_V at $t = t_{\text{eq}} \equiv 0$ are indicated by the heavy dots (only shown if clearly different from zero) and the assumed functional $\gamma[a]$ (multiplied by τ^{-1} and divided by a) is indicated by the heavy curve in the left-most panel on the second row from the top. The expansion of the universe is accelerated ($\ddot{a}/a > 0$) if $\rho_V/\rho_M > 1/2$, as indicated by the dashed curve in the right-most panel of the second row from the top. The scale factor a vanishes at $t = t_{\text{BB}} = -0.91636$ and the matter energy density ρ_M diverges as $1/a^3$ with the vacuum energy density ρ_V staying finite (see, in particular, the right-most panel of the third row from the top and the three panels on the fourth row, where Ω_{VM} stands for $\Omega_V + \Omega_M$).

times, which are both close to the maximum of ρ_V/ρ_M). The model values of the “present universe” are then

$$\begin{pmatrix} t_0 \\ t_0 - t_{\text{BB}} \\ a \\ \dot{a}/a \\ \rho_M \\ \rho_V \\ \rho_V/\rho_M \\ \Omega_V + \Omega_M \end{pmatrix} = \begin{pmatrix} -0.5842 \\ 0.3322 \\ 5.582 \\ 2.985 \\ 2.384 \\ 6.557 \\ 2.750 \\ 1.004 \end{pmatrix}, \quad (21)$$

where Ω_X is the energy density ρ_X relative to the critical density $\rho_c \equiv (\dot{a}/a)^2$ in units with $8\pi G/3 = c = 1$

By identifying the calculated value $\dot{a}/a = 2.985$ with the measured value [17] of the Hubble constant $H_0 = h/(9.78 \cdot 10^9 \text{ yr}) \approx 0.70/(9.78 \text{ Gyr})$, the present age of the model universe $t_0 - t_{\text{BB}} \approx 0.3322$ becomes

$$\tau_0 \approx 13.85 (0.70/h) \text{ Gyr}. \quad (22)$$

Similarly, the present radius of the model universe $a_0 \approx 5.582$ becomes of the order of $2 \times 10^{11} \text{ lyr}$, significantly larger than the present particle horizon. It is far from trivial that more or less reasonable values for ρ_{V0}/ρ_{M0} , $\Omega_{V0} + \Omega_{M0}$, and τ_0 can be produced at all in our approach.

In the same way, the equilibrium time $t_{\text{eq}} - t_{\text{BB}} \approx 0.91636$ would correspond to $\tau_{\text{eq}} \approx 38.22 \text{ Gyr}$, but there need not be a Big Crunch at even later times because of the lack of time-reversal invariance, as mentioned in the previous section. With the measured photon temperature $T_{\gamma 0} \approx 3 \text{ K}$, the matter equation-of-state parameter w would need to change to a value $1/3$ (relativistic matter for $T \gtrsim 3000 \text{ K}$) for $0 \leq a \lesssim 6 \times 10^{-3}$, in order to recover the standard nucleosynthesis of the very early universe [2, 4].

As to the phenomenology of $\gamma[a]$, we clearly recognize three phases in Fig. 2, where $\gamma[a]$ is positive, negative, and again positive as the time coordinate t moves away from $t_{\text{eq}} = 0$ in the negative direction. (Other structures of $\gamma[a]$ are not excluded *a priori*, but the one found suffices for the present discussion.) The resulting behavior of $\rho_V(t)$ from (15) is shown in the figure panel under the one of $\tau^{-1} \gamma[a]/a$. Note that $\rho_V(t_{\text{BB}})$ need not be negative, as different $1/\tau$ values and coefficients c_n in (19) can give positive $\rho_V(t_{\text{BB}})$ values of order 1 or perhaps even $\rho_V(t_{\text{BB}}) = 0$. Different $1/\tau$ values and coefficients c_n can also give a ρ_V/ρ_M peak above 3, but it may be difficult to keep the “present age” of the universe at the value (22) and to prevent it from dropping to significantly lower values.

From the approximate linearity of $a(t)$ up to the “present value” $t_0 \approx -0.5842$ in Fig. 2, it is possible to relate the time coordinate t just below t_0 to the redshift z used by observational

cosmology through the approximate relation $1+z \approx (t_0 - t_{\text{BB}})/(t - t_{\text{BB}})$. Then, a coordinate time $t = -0.75$ would correspond to a redshift $z \approx 1$. If future observations can measure $\rho_V(z)$ and $\rho_M(z)$ up to $z \approx 10$ (see, e.g., Ref. [18] for theoretical considerations), this would indirectly constrain the behavior of $\gamma[a(t)]/\tau$ for $t \in [t_{\text{BB}}, t_0]$. But these observations can perhaps also constrain $\gamma[a(t)]/\tau$ over the *whole* range $[t_{\text{BB}}, t_{\text{eq}}]$ if there are effective two-boundary conditions such as $\rho_V(t_{\text{BB}}) = 0$ and $\rho_V(t_{\text{eq}}) \neq 0$ from the underlying physics.

By way of summary, we list the following features of the model universe from Fig. 2:

1. Gibbs–Duhem-like boundary condition (12) at $t = t_{\text{eq}} \equiv 0$ with $\rho_V(t_{\text{eq}}) = \frac{1}{2} \rho_M(t_{\text{eq}})$ for $w = 0$, which may result from the self-tuning of a vacuum variable q to an equilibrium value q_0 ;
2. finite $|\rho_V(t)|$ within a factor of order 10^4 from the value set at $t = t_{\text{eq}}$;
3. a standard Big Bang phase having $a(t) \propto (t - t_{\text{BB}})^{2/3}$ for $w = 0$, matter energy density $\rho_M \propto a^{-3}$, and suppressed vacuum energy density with $\rho_V/\rho_M \rightarrow 0$ for $t \downarrow t_{\text{BB}}$;
4. an accelerating universe for “present times,” with ρ_V/ρ_M of order 1 and an approximately flat 3–geometry.

Features 1 and 2 suggest that a nonvanishing vacuum energy density ρ_V (or cosmological parameter Λ) does not require fine-tuning by factors of order $(E_{\text{Planck}}/10^{-3} \text{ eV})^4 \sim 10^{124}$ [6]. Still, it remains to be explained theoretically that the fundamental vacuum-dynamics constant $c\tau$ is of the order of the length scale $a(t_{\text{eq}}) \approx 4 \times 10^{10} \text{ yr} \approx \hbar c / (5 \times 10^{-34} \text{ eV})$. [Note that $a(t_{\text{eq}})$ also determines the two equilibrium energy densities from conditions (11)–(12) for a given value of w .] The theoretical explanation of this very small energy scale \hbar/τ would, most likely, trace back to the detailed microphysics, perhaps along the lines of the toy model discussed in Sec. II with mass $|\mu_0|$ from (6) setting the scale in the vacuum-decay equation (9). Inversely, there is the possibility that observational cosmology, by measuring the behavior of the vacuum energy density as mentioned above, can provide information on the microscopic structure of the quantum vacuum.

ACKNOWLEDGMENTS

It is a pleasure to thank G.E. Volovik for valuable discussions.

[1] A. Einstein, “Kosmologische Betrachtungen zur allgemeinen Relativitätstheorie,” Sitzungsber. Preuss. Akad. Wiss. 1917, p. 142; translated as: “Cosmological considerations on the general

- theory of relativity,” in: *The Principle of Relativity*, edited by H.A. Lorentz *et al.* (Dover Publ., New York, 1952), Chap. 9.
- [2] S. Weinberg, *Gravitation and Cosmology* (Wiley, New York, 1972).
- [3] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cambridge, England, 1973).
- [4] V. Mukhanov, *Physical Foundations of Cosmology* (Cambridge University Press, Cambridge, England, 2005).
- [5] S. Weinberg, *Cosmology* (Oxford University Press, Oxford, England, 2008).
- [6] F.R. Klinkhamer and G.E. Volovik, “Self-tuning vacuum variable and cosmological constant,” to appear in PRD, arXiv:0711.3170 [gr-qc].
- [7] A. Kostelecký and M. Mewes, “Signals for Lorentz violation in electrodynamics,” Phys. Rev. D **66**, 056005 (2002), arXiv:hep-ph/0205211.
- [8] F.R. Klinkhamer and M. Risse, “Ultrahigh-energy cosmic-ray bounds on nonbirefringent modified-Maxwell theory,” Phys. Rev. D **77**, 016002 (2008), arXiv:0709.2502 [hep-ph].
- [9] M.J. Duff and P. van Nieuwenhuizen, “Quantum inequivalence of different field representations,” Phys. Lett. B **94**, 179 (1980).
- [10] A. Aurilia, H. Nicolai, and P.K. Townsend, “Hidden constants: The theta parameter of QCD and the cosmological constant of N=8 supergravity,” Nucl. Phys. B **176**, 509 (1980).
- [11] G.E. Volovik, “Cosmological constant and vacuum energy,” Annalen Phys. (Leipzig) **14**, 165 (2005), arXiv:gr-qc/0405012.
- [12] G.E. Volovik, “Evolution of cosmological constant in effective gravity,” JETP Lett. **77**, 339 (2003), arXiv:gr-qc/0302069.
- [13] C. Barcelo, “Cosmology as a search for overall equilibrium,” JETP Lett. **84**, 635 (2007), arXiv:gr-qc/0611090.
- [14] D.J. Eisenstein *et al.* [SDSS Collaboration], “Detection of the baryon acoustic peak in the large-scale correlation function of SDSS luminous red galaxies,” Astrophys. J. **633**, 560 (2005), arXiv:astro-ph/0501171.
- [15] A.G. Riess *et al.*, “New Hubble Space Telescope discoveries of type Ia supernovae at $z > 1$: Narrowing constraints on the early behavior of dark energy,” Astrophys. J. **659**, 98 (2007), astro-ph/0611572.
- [16] E. Komatsu *et al.*, “Five-year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Cosmological interpretation,” arXiv:0803.0547v1 [astro-ph].
- [17] W.L. Freedman *et al.* [Hubble Space Telescope Collaboration], “Final results from the Hubble Space Telescope Key Project to measure the Hubble constant,” Astrophys. J. **553**, 47 (2001), arXiv:astro-ph/0012376.
- [18] V. Sahni and A. Starobinsky, “Reconstructing dark energy,” Int. J. Mod. Phys. D **15**, 2105 (2006), arXiv:astro-ph/0610026.