

# Cartan's structure of symmetry pseudo-group and coverings for the $r$ -th modified dispersionless Kadomtsev–Petviashvili equation

**Oleg I. Morozov**

Department of Mathematics, Moscow State Technical University  
of Civil Aviation, Kronshtadtskiy Blvd 20, Moscow 125993, Russia  
oim@foxcub.org

**Abstract.** We derive two non-equivalent coverings for the  $r$ -th mdKP equation from Maurer–Cartan forms of its symmetry pseudo-group. Also we find Bäcklund transformations between the covering equations.

AMS classification scheme numbers: 58H05, 58J70, 35A30

## 1. Introduction

The role of coverings in studying nonlinear differential equations (DEs) is well-known, [20, 21, 22, 23]. They lead to a number of useful techniques such as inverse scattering transformations, Bäcklund transformations, recursion operators, nonlocal symmetries and nonlocal conservation laws. For a given DE, a problem of constructing a covering is very difficult, see, e.g., [41, 8, 6, 14, 29, 30, 38, 15, 35, 36, 42, 40, 28, 12, 13]. One of the possible approaches to solution lies in the framework of Élie Cartan's structure theory of Lie pseudo-groups, [27, 3, 33, 34].

In the present paper we apply the method of [33, 34] to the  $r$ -th modified dispersionless Kadomtsev–Petviashvili equation ( $r$ -mdKP), [1]. We use Élie Cartan's method of equivalence, [4, 11, 17, 37], to compute Maurer–Cartan (MC) forms of the pseudo-group of contact symmetries of  $r$ -mdKP, and then find two linear combinations of these forms, whose horizontalizations provide covering equations of  $r$ -mdKP. Previously this approach was applied in [34] to a particular case of  $r$ -mdKP – modified dispersionless Kadomtsev–Petviashvili equation (mdKP), or modified Khokhlov–Zabolotskaya equation, [27, 25, 26]. Coverings for particular cases of  $r$ -mdKP were found in [10, 7, 18, 5] via other methods.

## 2. Preliminaries

### 2.1. Coverings of DEs

Let  $\pi_\infty : J^\infty(\pi) \rightarrow \mathbb{R}^n$  be the infinite jet bundle of local sections of the bundle  $\pi : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ . The coordinates on  $J^\infty(\pi)$  are  $(x^i, u_I)$ , where  $I = (i_1, \dots, i_k)$  are symmetric multi-indices,  $i_1, \dots, i_k \in \{1, \dots, n\}$ ,  $u_\emptyset = u$ , and for any local section  $f$  of  $\pi$  there exists a section  $j_\infty(f) : \mathbb{R}^n \rightarrow J^\infty(\pi)$  such that  $u_I(j_\infty(f)) = \partial^{\#I}(f)/\partial x^{i_1} \dots \partial x^{i_k}$ ,  $\#I = \#(i_1, \dots, i_k) = k$ . The *total derivatives* on  $J^\infty(\pi)$  are defined in the local coordinates as

$$D_i = \frac{\partial}{\partial x^i} + \sum_{\#I \geq 0} u_{Ii} \frac{\partial}{\partial u_I}.$$

We have  $[D_i, D_j] = 0$  for  $i, j \in \{1, \dots, n\}$ . A DE  $F(x^i, u_K) = 0$  defines a submanifold  $\mathcal{E}^\infty = \{D_I(F) = 0 \mid \#I \geq 0\} \subset J^\infty(\pi)$ , where  $D_I = D_{i_1} \circ \dots \circ D_{i_k}$  for  $I = (i_1, \dots, i_k)$ . We denote restrictions of  $D_i$  on  $\mathcal{E}^\infty$  as  $\overline{D}_i$ .

In local coordinates, a *covering* over  $\mathcal{E}^\infty$  is a bundle  $\widetilde{\mathcal{E}}^\infty = \mathcal{E}^\infty \times \mathcal{Q} \rightarrow \mathcal{E}^\infty$  with fibre coordinates  $q^\alpha$ ,  $\alpha \in \{1, \dots, N\}$  or  $\alpha \in \mathbb{N}$ , equipped with *extended total derivatives*

$$\widetilde{D}_i = \overline{D}_i + \sum_{\alpha} T_i^\alpha(x^j, u_I, q^\beta) \frac{\partial}{\partial q^\alpha}$$

such that  $[\widetilde{D}_i, \widetilde{D}_j] = 0$  whenever  $(x^i, u_I) \in \mathcal{E}^\infty$ .

Dually, the covering is defined by the following differential 1-forms, [41],

$$\omega^\alpha = dq^\alpha - T_i^\alpha(x^j, u_I, q^\beta) dx^i$$

such that  $d\omega^\alpha \equiv 0 \pmod{\omega^\beta, \bar{\vartheta}_I}$  iff  $(x^i, u_I) \in \mathcal{E}^\infty$ , where  $\bar{\vartheta}_I$  are restrictions of the contact forms  $\vartheta_I = du_I - u_{I,k} dx^k$  on  $\mathcal{E}^\infty$ .

## 2.2. Cartan's structure theory of contact symmetry pseudo-groups of DEs

A *pseudo-group* on a manifold  $M$  is a collection of local diffeomorphisms of  $M$ , which is closed under composition *when defined*, contains an identity and is closed under inverse. A *Lie pseudo-group* is a pseudo-group whose diffeomorphisms are local analytic solutions of an involutive system of partial differential equations. Élie Cartan's approach to Lie pseudo-groups is based on a possibility to characterize transformations from a pseudo-group in terms of a set of invariant differential 1-forms called *Maurer–Cartan forms*. The MC forms for a Lie pseudo-group can be computed by means of algebraic operations and differentiation. Expressions of differentials of the MC forms in terms of themselves give *structure equations* of the pseudo-group. The structure equations contain the full information about their pseudo-group.

EXAMPLE 1. Consider the bundle  $J^2(\pi)$  of jets of the second order of the bundle  $\pi$ . A differential 1-form  $\vartheta$  on  $J^2(\pi)$  is called a *contact form* if it is annihilated by all 2-jets of local sections:  $j_2(f)^*\vartheta = 0$ . In the local coordinates every contact 1-form is a linear combination of the forms  $\vartheta_0 = du - u_i dx^i$ ,  $\vartheta_i = du_i - u_{ij} dx^j$ ,  $i, j \in \{1, \dots, n\}$ ,  $u_{ji} = u_{ij}$ . A local diffeomorphism  $\Delta : J^2(\pi) \rightarrow J^2(\pi)$ ,  $\Delta : (x^i, u, u_i, u_{ij}) \mapsto (\bar{x}^i, \bar{u}, \bar{u}_i, \bar{u}_{ij})$ , is called a *contact transformation* if for every contact 1-form  $\bar{\vartheta}$  the form  $\Delta^*\bar{\vartheta}$  is also contact. We denote by  $\text{Cont}(J^2(\pi))$  the pseudo-group of contact transformations on  $J^2(\pi)$ . Consider the following 1-forms

$$\begin{aligned} \Theta_0 &= a \vartheta_0, & \Theta_i &= g_i \Theta_0 + a B_i^k \vartheta_k, & \Xi^i &= c^i \Theta_0 + f^{ik} \Theta_k + b_k^i dx^k, \\ \Theta_{ij} &= a B_k^i B_l^j (du_{kl} - u_{klm} dx^m) + s_{ij} \Theta_0 + w_{ij}^k \Theta_k + z_{ijk} \Xi^k, \end{aligned} \quad (1)$$

defined on  $J^2(\pi) \times \mathcal{H}$ , where  $\mathcal{H}$  is an open subset of  $\mathbb{R}^{(2n+1)(n+3)(n+1)/3}$  with local coordinates  $(a, b_k^i, c^i, f^{ik}, g_i, s_{ij}, w_{ij}^k, u_{ijk})$ ,  $i, j, k \in \{1, \dots, n\}$ ,  $i \leq j$ , such that  $a \neq 0$ ,  $\det(b_k^i) \neq 0$ ,  $f^{ik} = f^{ki}$ ,  $u_{ijk} = u_{ikj} = u_{jik}$ , while  $(B_k^i)$  is the inverse matrix for the matrix  $(b_k^i)$ . As it is shown in [32], the forms (1) are MC forms for  $\text{Cont}(J^2(\pi))$ , that is, a local diffeomorphism  $\widehat{\Delta} : J^2(\pi) \times \mathcal{H} \rightarrow J^2(\pi) \times \mathcal{H}$  satisfies the conditions  $\widehat{\Delta}^* \bar{\Theta}_0 = \Theta_0$ ,  $\widehat{\Delta}^* \bar{\Theta}_i = \Theta_i$ ,  $\widehat{\Delta}^* \bar{\Xi}^i = \Xi^i$ , and  $\widehat{\Delta}^* \bar{\Theta}_{ij} = \Theta_{ij}$  if and only if it is projectable on  $J^2(\pi)$ , and its projection  $\Delta : J^2(\pi) \rightarrow J^2(\pi)$  is a contact transformation. The structure equations for  $\text{Cont}(J^2(\pi))$  have the form

$$\begin{aligned} d\Theta_0 &= \Phi_0^0 \wedge \Theta_0 + \Xi^i \wedge \Theta_i, \\ d\Theta_i &= \Phi_i^0 \wedge \Theta_0 + \Phi_i^k \wedge \Theta_k + \Xi^k \wedge \Theta_{ik}, \\ d\Xi^i &= \Phi_0^i \wedge \Xi^i - \Phi_k^i \wedge \Xi^k + \Psi^{i0} \wedge \Theta_0 + \Psi^{ik} \wedge \Theta_k, \\ d\Theta_{ij} &= \Phi_i^k \wedge \Theta_{kj} + \Phi_j^k \wedge \Theta_{ki} - \Phi_0^0 \wedge \Theta_{ij} + \Upsilon_{ij}^0 \wedge \Theta_0 + \Upsilon_{ij}^k \wedge \Theta_k + \Lambda_{ijk} \wedge \Xi^k, \end{aligned}$$

where the additional forms  $\Phi_0^0$ ,  $\Phi_i^0$ ,  $\Phi_i^k$ ,  $\Psi^{i0}$ ,  $\Psi^{ij}$ ,  $\Upsilon_{ij}^0$ ,  $\Upsilon_{ij}^k$ , and  $\Lambda_{ijk}$  depend on differentials of the coordinates of  $\mathcal{H}$ .

EXAMPLE 2. Suppose  $\mathcal{E}$  is a second-order differential equation in one dependent and  $n$  independent variables. We consider  $\mathcal{E}$  as a submanifold in  $J^2(\pi)$ . Let  $\text{Cont}(\mathcal{E})$  be the group of contact symmetries for  $\mathcal{E}$ . It consists of all the contact transformations on  $J^2(\pi)$  mapping  $\mathcal{E}$  to itself. Let  $\iota_0 : \mathcal{E} \rightarrow J^2(\pi)$  be an embedding, and  $\iota = \iota_0 \times \text{id} : \mathcal{E} \times \mathcal{H} \rightarrow J^2(\pi) \times \mathcal{H}$ . The MC forms of  $\text{Cont}(\mathcal{E})$  can be derived from the forms  $\theta_0 = \iota^* \Theta_0$ ,  $\theta_i = \iota^* \Theta_i$ ,  $\xi^i = \iota^* \Xi^i$ , and  $\theta_{ij} = \iota^* \Theta_{ij}$  by means of Cartan's method of equivalence, see details and examples in [9, 31, 32].

### 3. Cartan's structure of the contact symmetry pseudo-group for $r$ -mdKP

The  $r$ -th mdKP

$$u_{tx} = -\frac{(3-r)(1-r)}{2} u_x^2 u_{xx} + \frac{r(3-r)}{2-r} u_x u_{xy} + \frac{3-r}{(2-r)^2} u_{yy} + \frac{(3-r)(1-r)}{2-r} u_y u_{xx},$$

$r \in \mathbb{Z} \setminus \{2\}$ , was derived in [1]. For a convenience of computations we use the following change of variables:

$$\tilde{t} = (3-r)t, \quad \tilde{x} = x, \quad \tilde{y} = (2-r)y, \quad \tilde{u} = -(1-r)u,$$

where  $r \notin \{1, 2, 3\}$ . Then we have

$$\tilde{u}_{\tilde{y}\tilde{y}} = \tilde{u}_{\tilde{x}\tilde{x}} + \left( \frac{1}{2(1-r)} \tilde{u}_{\tilde{x}}^2 + \tilde{u}_{\tilde{y}} \right) \tilde{u}_{\tilde{x}\tilde{x}} + \frac{r}{1-r} \tilde{u}_{\tilde{x}} \tilde{u}_{\tilde{x}\tilde{y}}.$$

We drop tildes and denote  $\kappa = \frac{r}{1-r}$ ; this yields

$$u_{yy} = u_{tx} + \left( \frac{\kappa+1}{2} u_x^2 + u_y \right) u_{xx} + \kappa u_x u_{xy}. \quad (2)$$

The exceptional cases  $r = 2$  and  $r = 3$  correspond to  $\kappa = -2$  and  $\kappa = -\frac{3}{2}$ , respectively. We will not consider the case of  $r = 1$ . The case of  $\kappa = -1$  is exceptional, too, since  $r \rightarrow \infty$  when  $\kappa \rightarrow -1$ . In the cases of  $\kappa = 0$ ,  $\kappa = 1$ , and  $\kappa = -1$  Eq. (2) gets the forms of the mdKP equation, [27, 25, 26],

$$u_{yy} = u_{tx} + \left( \frac{1}{2} u_x^2 + u_y \right) u_{xx}, \quad (3)$$

the dBKP equation, [39, 18],

$$u_{yy} = u_{tx} + (u_x^2 + u_y) u_{xx} + u_x u_{xy}. \quad (4)$$

and the equation describing Lorentzian hyper-CR Einstein–Weil structures, [7, 10],

$$u_{yy} = u_{tx} + u_y u_{xx} - u_x u_{xy}. \quad (5)$$

We use the method outlined in the previous section to compute MC forms and structure equations of the pseudo-group of contact symmetries for Eq. (2). The results depend on  $\kappa$ .

When  $\kappa \notin \{-2, -1\}$ , the structure equations have the form

$$\begin{aligned}
d\theta_0 &= (\eta_1 + \xi^2 - \frac{\kappa-4}{8}\xi^3 - \frac{\kappa}{\kappa+2}\theta_{22}) \wedge \theta_0 + \xi^1 \wedge \theta_1 + \xi^2 \wedge \theta_2 + \xi^3 \wedge \theta_3, \\
d\theta_1 &= \left(\frac{3}{2}\eta_1 - \frac{\kappa-4}{8}\xi^3 - \frac{3(\kappa+1)}{\kappa+2}\theta_{22}\right) \wedge \theta_1 + ((\kappa+1)\theta_2 + (\kappa+2)\xi^2) \wedge \theta_3 + \xi^1 \wedge \theta_{11} \\
&\quad + \left(\frac{2\kappa^2+15\kappa+4}{8}\theta_2 - \frac{\kappa^2-10\kappa+8}{8}\xi^2 + \kappa\theta_{23}\right) \wedge \theta_0 + \xi^2 \wedge \theta_{12} + \xi^3 \wedge \theta_{13}, \\
d\theta_2 &= \left(\frac{1}{2}\eta_1 - \frac{\kappa+1}{\kappa+2}\theta_{22} + \frac{3\kappa+4}{8}\xi^3\right) \wedge \theta_2 + \xi^1 \wedge \theta_{12} + \xi^2 \wedge \theta_{22} + \xi^3 \wedge \theta_{23}, \\
d\theta_3 &= \left(\eta_1 - \frac{2(\kappa+1)}{\kappa+2}\theta_{22} + \frac{\kappa+4}{8}\xi^3\right) \wedge \theta_3 + \left(\frac{\kappa-4}{8}\theta_{22} - \frac{\kappa^2-16}{64}\xi^3\right) \wedge \theta_0 + \frac{\kappa+2}{2}\xi^2 \wedge \theta_2 \\
&\quad + \xi^1 \wedge \theta_{13} + \xi^2 \wedge \theta_{23} + \xi^3 \wedge \theta_{12}, \\
d\xi^1 &= -\left(\frac{1}{2}\eta_1 - \frac{2\kappa+3}{\kappa+2}\theta_{22}\right) \wedge \xi^1, \\
d\xi^2 &= \left(\frac{1}{2}\eta_1 + \frac{1}{\kappa+2}\theta_{22} + \xi^3\right) \wedge \xi^2 + \left(\frac{\kappa-4}{8}\theta_0 - \theta_3\right) \wedge \xi^1 - \theta_2 \wedge \xi^3, \\
d\xi^3 &= -(\kappa+2)(\theta_2 + \xi^2) \wedge \xi^1 + \theta_{22} \wedge \xi^3, \\
d\theta_{11} &= 2\eta_1 \wedge \theta_{11} - \kappa\eta_2 \wedge \theta_0 + \eta_3 \wedge \xi^2 + \eta_4 \wedge \xi^3 + \eta_5 \wedge \xi^1 + \kappa(\theta_3 + \frac{\kappa+8}{4}\theta_{12}) \wedge \theta_0 \\
&\quad + \left((4\kappa+3)\theta_{23} - \frac{\kappa^2-18\kappa+20}{4}\xi^2\right) \wedge \theta_1 - \left((2\kappa+3)\theta_{13} + \frac{5\kappa^2+31\kappa+24}{4}\theta_1\right) \wedge \theta_2 \\
&\quad - \kappa\theta_3 \wedge \theta_{12} - \left(\frac{5\kappa+6}{\kappa+2}\theta_{22} + \frac{\kappa-4}{8}\xi^3\right) \wedge \theta_{11} + (2\kappa+4)\xi^2 \wedge \theta_{13}, \\
d\theta_{12} &= \eta_1 \wedge \theta_{12} + \eta_2 \wedge \xi^3 + \eta_3 \wedge \xi^1 + \theta_2 \wedge (\theta_{23} - \xi^2) + \left(\theta_3 - \frac{\kappa-4}{8}\theta_0 + \frac{3\kappa+4}{\kappa+2}\theta_{12}\right) \wedge \theta_{22} \\
&\quad + \frac{3\kappa+4}{8}\xi^3 \wedge \theta_{12}, \\
d\theta_{13} &= \frac{3}{2}\eta_1 \wedge \theta_{13} + \eta_2 \wedge \xi^2 + \eta_3 \wedge \xi^3 + \eta_4 \wedge \xi^1 + \frac{\kappa^3+10\kappa^2-32\kappa-96}{64}\theta_0 \wedge \xi^2 \\
&\quad - \left(\frac{\kappa^3+\kappa^2-14\kappa-24}{16}\theta_2 + \frac{\kappa^2-3\kappa-4}{8}\theta_{23}\right) \wedge \theta_0 + \left(\frac{\kappa-4}{8}\theta_{22} - \frac{\kappa^2-16}{64}\xi^3\right) \wedge \theta_1 \\
&\quad - \left(\frac{7\kappa^2+32\kappa+24}{8}\theta_3 + (\kappa+2)\theta_{12}\right) \wedge \theta_2 + \left((2\kappa+1)\theta_{23} + \frac{\kappa^2+22\kappa+24}{8}\xi^2\right) \wedge \theta_3 \\
&\quad + \frac{3(\kappa+2)}{2}\xi^2 \wedge \theta_{12} - \left(\frac{4\kappa+5}{\kappa+2}\theta_{22} - \frac{\kappa+4}{8}\xi^3\right) \wedge \theta_{13}, \\
d\theta_{22} &= \left(\frac{3\kappa^2+18\kappa+24}{8}\theta_2 + (\kappa+2)(\theta_{23} + \xi^2)\right) \wedge \xi^1 \\
d\theta_{23} &= \frac{1}{2}\eta_1 \wedge \theta_{23} + \eta_2 \wedge \xi^1 - \left(\frac{3(\kappa+4)}{8}\theta_{22} + \frac{9\kappa^2+48\kappa+112}{64}\xi^3\right) \wedge \theta_2 + \left(\frac{3(\kappa+4)}{8}\theta_{23} - \xi^2\right) \wedge \xi^3 \\
&\quad + \left(\frac{2\kappa+3}{\kappa+2}\theta_{23} - \frac{3\kappa-4}{8}\xi^2\right) \wedge \theta_{22}, \\
d\eta_1 &= 0, \\
d\eta_2 &= \eta_6 \wedge \xi^1 + \left(\eta_1 + \frac{2(2\kappa+3)}{\kappa+2}\theta_{22} - \frac{3(\kappa+4)}{8}\xi^3\right) \wedge \eta_2 + \theta_3 \wedge \left(\frac{3(\kappa-4)}{8}\theta_{22} - \xi^3\right) \\
&\quad + \left(\frac{3\kappa^2-16\kappa+16}{64}\theta_{22} - \frac{\kappa-4}{8}\xi^3\right) \wedge \theta_0 + \left(\frac{3(\kappa+4)}{8}\theta_2 + \xi^2\right) \wedge \theta_{23} \\
&\quad + \left(\frac{3(\kappa+4)}{8}\theta_{22} + \frac{9\kappa^2+48\kappa+112}{64}\xi^3\right) \wedge \theta_{12}, \\
d\eta_3 &= \eta_6 \wedge \xi^3 + \eta_7 \wedge \xi^1 + ((\kappa+3)\theta_2 + (\kappa+2)\xi^2) \wedge \eta_2 + \left(\frac{\kappa-4}{8}\theta_1 - \theta_{13}\right) \wedge \theta_{22} \\
&\quad + \left(\frac{3}{2}\eta_1 + \frac{5\kappa+7}{\kappa+2}\theta_{22} + \frac{3\kappa+4}{8}\xi^3\right) \wedge \eta_3 + (\kappa+2)\theta_3 \wedge (\theta_{23} + \xi^2) \\
&\quad + \frac{\kappa-4}{64}((3\kappa^2 + 18\kappa + 32)\theta_2 + 8(\kappa+2)(\theta_{23} + \xi^2)) \wedge \theta_0 \\
&\quad + \left(\frac{3\kappa^2+18\kappa+32}{8}\theta_3 + \frac{6\kappa^2+29\kappa+28}{4}\theta_{12}\right) \wedge \theta_2 - \left(3(\kappa+1)\theta_{23} - \frac{3\kappa^2+34\kappa+32}{8}\xi^2\right) \wedge \theta_{12}, \\
d\eta_4 &= \eta_6 \wedge \xi^2 + \eta_7 \wedge \xi^3 + \eta_8 \wedge \xi^1 + \left(2\eta_1 - \frac{2(3\kappa+4)}{\kappa+2}\theta_{22} + \frac{\kappa+4}{8}\xi^3\right) \wedge \eta_4
\end{aligned}$$

$$\begin{aligned}
& + \kappa \left( \frac{\kappa-4}{8} \theta_0 - 2 \theta_3 \right) \wedge \eta_2 + \left( (2\kappa + 4) \theta_2 + \frac{5(\kappa+2)}{2} \xi^2 \right) \wedge \eta_3 \\
& - \frac{\kappa}{16} \left( 2(\kappa - 4) \theta_3 - (\kappa^2 - 2\kappa - 4) \theta_{12} \right) \wedge \theta_0 + \frac{\kappa}{8} (7\kappa + 5) \theta_3 \wedge \theta_{12} \\
& + \frac{1}{16} \left( (2\kappa^3 + 3\kappa^2 - 30\kappa - 112) \theta_2 - 2(2\kappa^2 - 5\kappa + 4) \theta_{23} \right) \wedge \theta_1 \\
& - \frac{1}{32} (\kappa^3 + 10\kappa^2 - 32\kappa - 96) \theta_1 \wedge \xi^2 \\
& - \left( 6(\kappa + 1) \theta_{23} + \frac{1}{4} (\kappa^2 + 30\kappa + 36) \xi^2 \right) \wedge \theta_{13} \\
& - \frac{1}{8} (20\kappa^2 + 101\kappa + 92) \theta_2 \wedge \theta_{13} - \frac{1}{64} (8(\kappa - 4) \theta_{22} - (\kappa^2 - 16) \xi^3) \wedge \theta_{11}, \\
d\eta_5 = & - \kappa \eta_6 \wedge \theta_0 + \eta_7 \wedge \xi^2 + \eta_8 \wedge \xi^3 + \eta_9 \wedge \xi^1 + \left( \frac{5}{2} \eta_1 - \frac{7\kappa+9}{\kappa+2} \theta_{22} - \frac{\kappa-4}{8} \xi^3 \right) \wedge \eta_5 \\
& + (5\kappa + 3) \eta_2 \wedge \theta_1 + \eta_3 \wedge \left( \frac{2\kappa^2+17\kappa-4}{8} \theta_0 + (\kappa - 1) \theta_3 \right) - 3(\kappa + 1) \theta_{12} \wedge \theta_{13}. \\
& + \left( (3\kappa + 5) \theta_2 + 3(\kappa + 2) \xi^2 \right) \wedge \eta_4 - \left( 9(\kappa + 1) \theta_{23} - \frac{3\kappa^2-78\kappa-96}{8} \xi^2 \right) \wedge \theta_{11} \\
& + \left( \frac{\kappa^3-22\kappa^2+52\kappa+90}{32} \theta_1 - \frac{\kappa^2+2\kappa-8}{4} \theta_{13} \right) \wedge \theta_0 - 2(\kappa + 2) \theta_3 \wedge \theta_{13} \\
& + \left( \frac{\kappa^2-22\kappa-20}{4} \theta_3 - \frac{6\kappa^2+39\kappa+24}{4} \theta_{12} \right) \wedge \theta_1 - \frac{24\kappa^2+71\kappa+64}{4} \theta_2 \wedge \theta_{11},
\end{aligned}$$

where

$$\begin{aligned}
\xi^1 &= q^{-1} dt, \\
\xi^2 &= q u_{xx}^2 \left( \left( \frac{\kappa+1}{2} u_x^2 - u_y \right) dt + dx - u_x dy \right), \\
\xi^3 &= u_{xx} (dy - (\kappa + 2) u_x dt), \\
\eta_1 &= 2 \frac{dq}{q} + \frac{2(2\kappa + 3)}{\kappa + 2} \frac{du_{xx}}{u_{xx}},
\end{aligned} \tag{6}$$

and  $q = B_1^1 \neq 0$ . We need not explicit expressions for the other MC forms in the sequel.

In the case of  $\kappa = -1$  the contact symmetry pseudo-group of Eq. (5) has the following structure equations

$$\begin{aligned}
d\theta_0 &= (\eta_1 + \frac{5}{8} \xi^3 + \theta_{22}) \wedge \theta_0 + \xi^1 \wedge \theta_1 + \xi^2 \wedge \theta_2 + \xi^3 \wedge \theta_3, \\
d\theta_1 &= \frac{3}{2} \eta_1 \wedge \theta_1 + \eta_2 \wedge \left( \frac{1}{8} \theta_0 + \theta_3 \right) - \left( \frac{9}{8} \theta_2 + \theta_{23} - \frac{1}{2} \xi^2 \right) \wedge \theta_0 - \frac{5}{8} \theta_1 \wedge \xi^3 + \xi^1 \wedge \theta_{11} \\
&+ \xi^2 \wedge \theta_{12} + \xi^3 \wedge \theta_{13}, \\
d\theta_2 &= \frac{1}{8} (4\eta_1 + \xi^3) \wedge \theta_2 + \xi^1 \wedge \theta_{12} + \xi^2 \wedge \theta_{22} + \xi^3 \wedge \theta_{23}, \\
d\theta_3 &= \eta_1 \wedge \theta_3 + \frac{1}{2} \eta_2 \wedge \theta_2 - \frac{5}{64} (8\theta_{22} + 3\xi^3) \wedge \theta_0 + \frac{3}{8} \xi^3 \wedge \theta_3 + \xi^1 \wedge \theta_{13} + \xi^2 \wedge \theta_{23} \\
&+ \xi^3 \wedge \theta_{12}, \\
d\xi^1 &= -\frac{1}{2} (\eta_1 - 2\theta_{22}) \wedge \xi^1, \\
d\xi^2 &= -\frac{1}{8} (5\theta_0 + 8\theta_3) \wedge \xi^1 + \frac{1}{2} (\eta_1 + 2\theta_{22} + \xi^3) \wedge \xi^2 - \frac{1}{2} (\eta_2 + \theta_2) \wedge \xi^3, \\
d\xi^3 &= -(\eta_2 + \theta_2) \wedge \xi^1 + \theta_{22} \wedge \xi^3, \\
d\theta_{11} &= 2\eta_1 \wedge \theta_{11} + \eta_2 \wedge \left( \frac{3}{4} \theta_1 + 2\theta_{13} \right) + \eta_4 \wedge \xi^2 + \eta_5 \wedge \xi^3 + \eta_6 \wedge \xi^1 - \left( \theta_{22} - \frac{5}{8} \xi^3 \right) \wedge \theta_{11} \\
&+ \left( \eta_3 - \theta_3 - \frac{7}{4} \theta_{12} \right) \wedge \theta_0 - \left( \frac{1}{2} \theta_2 - \theta_{23} - \frac{1}{2} \xi^2 \right) \wedge \theta_1 + \theta_2 \wedge \theta_{13} + \theta_3 \wedge \theta_{12}, \\
d\theta_{12} &= \eta_1 \wedge \theta_{12} + \eta_2 \wedge \left( \frac{1}{8} \theta_2 + \theta_{23} + \xi^2 \right) + \eta_3 \wedge \xi^3 + \eta_4 \wedge \xi^1 + \frac{5}{8} \theta_0 \wedge \theta_{22} - \xi^2 \wedge \theta_{23} \\
&- \left( \theta_{23} - \frac{9}{8} \xi^2 \right) \wedge \theta_2 + (\theta_3 + \theta_{12}) \wedge \theta_{22} + \frac{1}{8} \xi^3 \wedge \theta_{12}, \\
d\theta_{13} &= \frac{3}{2} \eta_1 \wedge \theta_{13} + \frac{1}{64} \eta_2 \wedge (35\theta_0 + 24\theta_3 + 96\theta_{12}) + \eta_3 \wedge \xi^2 + \eta_4 \wedge \xi^3 + \eta_5 \wedge \xi^1
\end{aligned}$$

$$\begin{aligned}
& + \frac{5}{16} (2\theta_2 + \xi^2) \wedge \theta_0 - \frac{5}{64} (8\theta_{22} - 3\xi^3) \wedge \theta_1 + \left(\frac{1}{8}\theta_3 + \theta_{12}\right) \wedge \theta_2 + \theta_3 \wedge \theta_{23} \\
& - \left(\theta_{22} - \frac{3}{8}\xi^3\right) \wedge \theta_{13}, \\
d\theta_{22} &= \frac{1}{8} (4\eta_2 + 9\theta_2 + 4\xi^2 + 8\theta_{23}) \wedge \xi^1, \\
d\theta_{23} &= \frac{1}{2} \eta_1 \wedge \theta_{23} + \frac{1}{2} \eta_2 \wedge (\theta_{22} + \xi^3) + \eta_3 \wedge \xi^1 + \frac{1}{64} \theta_2 \wedge (72\theta_{22} - 73\xi^3) \\
& + (\theta_{23} + \frac{3}{8}\xi^2) \wedge \theta_{22} + \frac{1}{8} (9\theta_{23} + 4\xi^2) \wedge \xi^3, \\
d\eta_1 &= 0, \\
d\eta_2 &= \frac{1}{2} (\eta_1 + \xi^3) \wedge \eta_2 - \frac{5}{8} \theta_0 \wedge \xi^1 - (\theta_2 + \frac{1}{2}\xi^2) \wedge \xi^3 - \theta_3 \wedge \xi^1 - \theta_{22} \wedge \xi^2, \\
d\eta_3 &= \eta_7 \wedge \xi^1 + \eta_1 \wedge \eta_3 + \frac{1}{64} \eta_2 \wedge (41\theta_2 + 72\theta_{23} + 36\xi^3) + \frac{1}{8} \eta_3 \wedge (16\theta_{22} + 9\xi^3) \\
& + \frac{5}{64} (7\theta_{22} + 8\xi^3) \wedge \theta_0 - \frac{1}{64} (72\theta_{23} + 41\xi^2) \wedge \theta_2 - \left(\frac{1}{8}\theta_{22} + \xi^3\right) \wedge \theta_3 \\
& + \frac{1}{64} (72\theta_{22} + 73\xi^3) \wedge \theta_{12} + \frac{1}{8} \theta_{23} \wedge \xi^2, \\
d\eta_4 &= \eta_7 \wedge \xi^3 + \eta_8 \wedge \xi^1 + \frac{1}{8} (5\theta_0 + 8\theta_3 + 6\theta_{12}) \wedge \eta_2 + (2\eta_2 + 2\theta_2 - \xi^2) \wedge \eta_3 \\
& + \frac{1}{8} (12\eta_1 - 16\theta_{22} + \xi^3) \wedge \eta_4 + \frac{5}{64} \theta_0 \wedge (17\theta_2 + 8\theta_{23}) - \frac{5}{8} \theta_1 \wedge \theta_{22} \\
& + \frac{1}{8} (17\theta_3 + 10\theta_{12}) \wedge \theta_2 + \theta_3 \wedge \theta_{23} - \frac{5}{8} \theta_{12} \wedge \xi^2 - \theta_{13} \wedge \theta_{22}, \\
d\eta_5 &= \eta_7 \wedge \xi^2 + \eta_8 \wedge \xi^3 + \eta_9 \wedge \xi^1 - \frac{5}{32} \eta_2 \wedge (7\theta_1 + 8\theta_{13}) + \left(\frac{5}{8}\theta_0 + 2\theta_3\right) \wedge \eta_3 \\
& + \left(\frac{5}{2}\eta_2 + 2\theta_2\right) \wedge \eta_4 + \left(2\eta_1 - 2\theta_{22} + \frac{3}{8}\xi^3\right) \wedge \eta_5 + \frac{5}{16} (\theta_{12} - 2\theta_3) \wedge \theta_0 \\
& - \frac{5}{16} (5\theta_2 + 2\theta_{23} + 2\xi^2) \wedge \theta_1 + \frac{13}{8} \theta_{12} \wedge \theta_3 - \frac{5}{64} (8\theta_{22} + 3\xi^3) \wedge \theta_{11} \\
& - \frac{1}{8} (11\theta_2 + 4\xi^2) \wedge \theta_{13}, \\
d\eta_6 &= \eta_7 \wedge \theta_0 + \eta_8 \wedge \xi^2 + \eta_9 \wedge \xi^3 + \eta_{10} \wedge \xi^1 + \eta_2 \wedge (\eta_5 - \frac{15}{8}\theta_{11}) - 2\eta_3 \wedge \theta_1 \\
& - \eta_4 \wedge \left(\frac{19}{8}\theta_0 + 2\theta_3\right) - 2\eta_5 \wedge \theta_2 - \frac{1}{8} \eta_6 \wedge (20\eta_1 + 16\theta_{22} - 5\xi^3) \\
& + \frac{1}{32} (5\theta_1 + 72\theta_{13}) \wedge \theta_0 + \frac{3}{4} (\theta_3 + 3\theta_{12}) \wedge \theta_1 + \frac{5}{4} \theta_{11} \wedge \theta_2 + 2\theta_{13} \wedge \theta_3
\end{aligned}$$

with

$$\begin{aligned}
\xi^1 &= q^{-1} dt, \\
\xi^2 &= q u_{xx}^2 \left( (s^2 u_{xx}^2 - s u_x u_{xx} - u_y) dt + dx - s u_{xx} dy \right), \\
\xi^3 &= u_{xx} \left( (u_x - 2s u_{xx}) dt + dy \right), \\
\eta_1 &= 2 \frac{dq}{q} + 2 \frac{du_{xx}}{u_{xx}},
\end{aligned} \tag{7}$$

where  $s = B_3^2 \in \mathbb{R}$ .

#### 4. Coverings of r-mdKP

Following [33, 34], we find linear combinations of the MC forms (6) and (7), which provide coverings of Eq. (2) in the cases of  $\kappa \notin \{-2, -\frac{3}{2}, -1\}$ ,  $\kappa = -\frac{3}{2}$ , and  $\kappa = -1$ , respectively.

##### 4.1. General case

When  $\kappa \notin \{-2, -\frac{3}{2}, -1\}$ , we take the following linear combination of the MC forms (6)

$$\omega = \eta_1 - \lambda_1 \xi^1 - \lambda_2 \xi^2 - \lambda_3 \xi^3$$

$$\begin{aligned}
&= 2 \frac{dq}{q} + \frac{2(2\kappa+3)}{\kappa+2} \frac{du_{xx}}{u_{xx}} - \lambda_2 q u_{xx} dx + (\lambda_2 q u_x u_{xx} - \lambda_3 u_{xx}) dy \\
&+ \left( \lambda_2 q \left( u_y - \frac{\kappa+1}{2} u_x^2 \right) u_{xx}^2 + \lambda_3 (\kappa+2) u_x u_{xx} - \lambda_1 q^{-1} \right) dt,
\end{aligned}$$

with  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ , and put

$$q = -\frac{(\kappa+1)v}{(\kappa+2)v_1^{2\kappa+3}}, \quad u_{xx} = \frac{v_1^{\kappa+2}}{v},$$

where  $v$  and  $v_1$  are new independent variables. This gives

$$\omega = -\frac{2(\kappa+1)}{(\kappa+2)v} (dv - A v_1 dt - v_1 dx - B v_1 dy) \quad (8)$$

with

$$\begin{aligned}
A &= -\frac{(\kappa+2)^2}{4(\kappa+1)} \left( \lambda_1 \lambda_2 v_1^{2(\kappa+1)} + 2\lambda_3 (\kappa+1) u_x v_1^{\kappa+1} \right) + \frac{\kappa+1}{2} u_x^2 - 2u_y, \\
B &= -u_x - \frac{\lambda_3 (\kappa+2)}{2(\kappa+1)} v_1^{\kappa+1}.
\end{aligned}$$

The form (8) is equal to zero whenever  $v_1 = v_x$  and

$$\begin{aligned}
v_t &= \left( \frac{\lambda_1 \lambda_2 (\kappa+2)^2}{4(\kappa+1)^2} v_x^{2(\kappa+1)} + \frac{\lambda_3 (\kappa+2)^2}{2(\kappa+1)} v_x^{\kappa+1} + \frac{\kappa+1}{2} u_x^2 - u_y \right) v_x, \\
v_y &= -\left( \frac{\lambda_3 (\kappa+2)}{\kappa+1} v_x^{\kappa+1} + u_x \right) v_x.
\end{aligned}$$

This system is compatible, i.e.,  $(v_t)_y = (v_y)_t$ , whenever

$$\begin{aligned}
u_{yy} - u_{tx} - \left( \frac{\kappa+1}{2} u_x^2 + u_y \right) u_{xx} - \kappa u_x u_{xy} \\
+ \frac{(\kappa+2)^2}{4(\kappa+1)^2} v_x^{2(\kappa+1)} (\lambda_3^2 (\kappa+2)^2 - \lambda_1 \lambda_2 (2\kappa+3)) = 0.
\end{aligned}$$

This equation coincides with Eq. (2) iff  $\lambda_3^2 (\kappa+2)^2 - \lambda_1 \lambda_2 (2\kappa+3) = 0$ . So we put

$$\lambda_1 = \frac{\lambda_3^2 (\kappa+2)^2}{\lambda_2 (2\kappa+3)}.$$

This yields

$$\begin{aligned}
v_t &= \left( \frac{\lambda_3^2 (\kappa+2)^4}{(2\kappa+3)(\kappa+1)^2} v_x^{2(\kappa+1)} + \frac{\lambda_3 (\kappa+2)^2}{2(\kappa+1)} u_x v_x^{\kappa+1} + \frac{\kappa+1}{2} u_x^2 - u_y \right) v_x, \\
v_y &= -\left( \frac{\lambda_3 (\kappa+2)}{\kappa+1} v_x^{\kappa+1} + u_x \right) v_x.
\end{aligned}$$

When  $\lambda_3 = 0$ , we have

$$v_t = \left( \frac{\kappa+1}{2} u_x^2 - u_y \right) v_x, \quad v_y = -u_x v_x. \quad (9)$$

In the case of  $\kappa = 0$  this covering for Eq. (3) was obtained in [34]. When  $\lambda_3 \neq 0$ , we put  $v = \left( \frac{\lambda_3 (\kappa+2)}{2(\kappa+1)} \right)^{1/(\kappa+1)} w$ . Then

$$\begin{aligned}
w_t &= \left( \frac{(\kappa+2)^2}{2\kappa+3} w_x^{2(\kappa+1)} + (\kappa+2) u_x w_x^{\kappa+1} + \frac{\kappa+1}{2} u_x^2 - u_y \right) w_x, \\
w_y &= -\left( w_x^{\kappa+1} + u_x \right) w_x.
\end{aligned} \quad (10)$$



For  $\kappa = 0$  this covering of (3) was found in [5] by means of another technique and in [34] via the method described above. For  $\kappa = 1$  the covering (10) of Eq. (4) was obtained in [18].

From (9) we have

$$u_x = -\frac{v_y}{v_x}, \quad u_y = \frac{\kappa + 1}{2} \left( \frac{v_y}{v_x} \right)^2 - \frac{v_t}{v_x}. \quad (11)$$

The integrability condition  $(u_x)_y = (u_y)_x$  of this system gives

$$v_{yy} = v_{tx} + \left( \frac{(\kappa + 1) v_y^2}{v_x^2} - \frac{v_t}{v_x} \right) v_{xx} - \frac{\kappa v_y}{v_x} v_{xy}. \quad (12)$$

For  $\kappa = 0$  this equation was obtained in [2]. Also, from (10) we get

$$u_x = -\frac{w_y}{w_x} - w_x^{\kappa+1}, \quad u_y = -\frac{w_t}{w_x} + \frac{(\kappa + 1) w_y^2}{2 w_x^2} - w_x^\kappa w_y - \frac{(\kappa + 1) w_x^{2(\kappa+1)}}{2(2\kappa + 3)} \quad (13)$$

This system yields

$$\begin{aligned} w_{yy} = w_{tx} + & \left( (\kappa + 1) \frac{w_y^2}{w_x^2} - \frac{w_t}{w_x} + \kappa w_x^\kappa w_y + \frac{(\kappa+1)^2}{2\kappa+3} w_x^{2(\kappa+1)} \right) w_{xx} \\ & - \kappa \left( \frac{w_y}{w_x} + w_x^\kappa \right) w_{xy}. \end{aligned} \quad (14)$$

Substitution for (11) in (10) gives a Bäcklund transformation

$$w_t = \frac{(\kappa + 2)^2}{2\kappa + 3} w_x^{2\kappa+3} - \frac{(\kappa + 2)v_y}{v_x} w_x^{\kappa+2} + \frac{v_t}{v_x} w_x, \quad w_y = -w_x^{\kappa+2} + \frac{v_y}{v_x} w_x$$

from Eq. (12) to Eq. (14). The inverse Bäcklund transformation appears from substitution for (13) in (9).

#### 4.2. Case of $\kappa = -\frac{3}{2}$

In the case of  $\kappa = -\frac{3}{2}$  we take the following combination of the MC forms (6)

$$\omega = \eta_1 - \lambda_1 \xi^1 - 4\xi^2 = 2 \frac{dq}{q} + (q u_{xx}^2 (u_x^2 + 4u_y) - \lambda_1 q^{-1}) dt - 4q u_{xx} (dx - u_x dy)$$

and the following change of variables:  $q = -v^{-2}$ ,  $u_{xx} = (v v_1)^{1/2}$ . Then we have

$$\omega = -4 \frac{dv}{v} - \left( \frac{(u_x^2 + 4u_y) v_1}{v} - \lambda_1 v^2 \right) dt + \frac{4v_1}{v} dx - \frac{4u_x v_1}{v} dy.$$

This form is equal to zero whenever  $v_1 = v_x$  and

$$v_t = \frac{1}{4} \lambda_1 v^3 - \left( \frac{1}{4} u_x^2 + u_y \right) v_x, \quad v_y = -u_x v_x.$$

This system is compatible for every value of  $\lambda_1$  whenever Eq. (2) with  $\kappa = -\frac{3}{2}$  is satisfied. When  $\lambda_1 = 0$ , we have Eqs. (9) with  $\kappa = -\frac{3}{2}$ :

$$v_t = -\left( \frac{1}{4} u_x^2 + u_y \right) v_x, \quad v_y = -u_x v_x. \quad (15)$$

When  $\lambda_1 \neq 0$ , we put  $v = 2 \lambda_1^{-1/2} w$ . Then we get

$$w_t = w^3 - \left( \frac{1}{4} u_x^2 + u_y \right) w_x, \quad w_y = -u_x w_x. \quad (16)$$

Exclusion of  $u_x$  and  $u_y$  from Eqs. (15) and (16) gives equations

$$v_{yy} = v_{tx} - \left( \frac{v_y^2}{2v_x^2} + \frac{v_t}{v_x} \right) v_{xx} + \frac{3v_y}{2v_x} v_{xy}, \quad (17)$$

$$w_{yy} = w_{tx} - \left( \frac{w_y^2}{2w_x^2} + \frac{w_t}{w_x} - \frac{w^3}{w_x} \right) w_{xx} + \frac{3w_y}{2w_x} w_{xy} - 3w^2 w_x, \quad (18)$$

and a Bäcklund transformation from (17) to (18):

$$w_t = w^3 + \frac{v_t}{v_x} w_x, \quad w_y = \frac{v_y}{v_x} w_x.$$

#### 4.3. Case of $\kappa = -1$

When  $\kappa = -1$ , we take the following combination of the MC forms (7):

$$\begin{aligned} \omega = \eta_1 + \frac{1}{2} \lambda_3^2 \xi^1 + 2\xi^2 - \lambda_3 \xi^3 = 2q u_{xx}^2 dx - u_{xx} (2q s u_{xx}^2 + \lambda_3) dy + 2 \frac{dq}{q} + 2 \frac{du_{xx}}{u_{xx}} \\ + \frac{1}{2q} \left( (\lambda_3 + q u_{xx} (2s u_{xx} - u_x))^2 - q^2 u_{xx}^2 (u_x^2 + 4u_y) \right) dt. \end{aligned}$$

Then we substitute for  $q = (v v_1)^{-1}$ ,  $s = u_x v_1^{-1}$ ,  $u_{xx} = v_1$  and obtain

$$\omega = -\frac{1}{2v} (4dv - (\lambda_3^2 v^2 + 2\lambda_3 u_x v - 4u_y) v_1 dt - 4v_1 dx + 2(\lambda_3 v + 2u_x) v_1 dy).$$

This form is equal to zero whenever  $v_1 = v_x$  and

$$v_t = \left( \frac{\lambda_3^2}{4} v^2 + \frac{1}{2} u_x v - u_y \right) v_x, \quad v_y = -\left( \frac{1}{2} \lambda_3 v + u_x \right) v_x.$$

This system is compatible for every value of  $\lambda_3$  whenever Eq. (5) is satisfied. For  $\lambda_3 = 0$  we have

$$v_t = -u_y v_x, \quad v_y = -u_x v_x. \quad (19)$$

When  $\lambda_3 \neq 0$ , we put  $v = 2\lambda_3^{-1} w$ . Then we have

$$w_t = (w^2 + u_x w - u_y) w_x, \quad w_y = -(w + u_x) w_x. \quad (20)$$

Excluding  $u_x$  and  $u_y$  from systems (19) and (20), we get equations

$$v_{yy} = v_{tx} - \frac{v_t}{v_x} v_{xx} + \frac{v_y}{v_x} v_{xy} \quad (21)$$

and

$$w_{yy} = w_{tx} - \frac{w_t + w w_x}{w_x} w_{xx} + \frac{w_y + w w_x}{w_x} w_{xy}, \quad (22)$$

correspondingly. The Bäcklund transformation between Eqs. (21) and (22) has the form

$$w_t = \left( w^2 + \frac{v_t - w v_y}{v_x} \right) w_x, \quad w_y = \left( w - \frac{v_y}{v_x} \right) w_x.$$

REMARK 1. A one-parametric family of coverings with a nonremovable parameter

$$v_t = -(u_y - \lambda u_x - \lambda^2) v_x, \quad v_y = -(u_x + \lambda) v_x, \quad (23)$$

for Eq. (5) is presented in [7]. This family can be obtained from (19) by means the following technique, [22, § 3.6] [19, ?, 30, 16]. Eq. (5) has the infinitesimal symmetry

$$X = y \frac{\partial}{\partial x} + 2x \frac{\partial}{\partial u},$$

which can't be lifted into a symmetry of the covering (19). Then the deformation  $e^{\lambda X}$  transforms the covering (19) into (23). Indeed, we have

$$\begin{aligned} \tilde{t} &= e^{\lambda X}(t) = t, & \tilde{x} &= e^{\lambda X}(x) = x + \lambda y, & \tilde{y} &= e^{\lambda X}(y) = y, \\ \tilde{u} &= e^{\lambda X}(u) = u + 2\lambda x + \lambda^2 y, \end{aligned}$$

and therefore

$$\tilde{u}_{\tilde{t}} = e^{\lambda X}(u_t) = u_t, \quad \tilde{u}_{\tilde{x}} = e^{\lambda X}(u_x) = u_x + 2\lambda, \quad \tilde{u}_{\tilde{y}} = e^{\lambda X}(u_y) = u_y - \lambda u_x - \lambda^2.$$

Since  $\tilde{v} = e^{\lambda X}(v) = v$ , for the form

$$\tilde{\omega}_1 = d\tilde{v} + \tilde{u}_{\tilde{y}} \tilde{v}_{\tilde{x}} d\tilde{t} - \tilde{v}_{\tilde{x}} d\tilde{x} + \tilde{u}_{\tilde{x}} \tilde{v}_{\tilde{x}} d\tilde{y},$$

which defines the covering (19) in the tilded variables, we have

$$(e^{\lambda X})^* \tilde{\omega}_1 = dv + (u_y - \lambda u_x - \lambda^2) v_x dt - v_x dx + (u_x + \lambda) v_x dy.$$

This form defines the family of coverings (23).

Similarly, we derive a new one-parametric family of coverings with a nonremovable parameter from the covering (20). We have  $\tilde{w} = e^{\lambda X}(w) = w$ , so the form

$$\tilde{\omega}_2 = d\tilde{w} - (\tilde{w}^2 + \tilde{u}_{\tilde{x}} \tilde{w} - \tilde{u}_{\tilde{y}}) \tilde{w}_{\tilde{x}} d\tilde{t} - \tilde{w}_{\tilde{x}} d\tilde{x} + (\tilde{w} + \tilde{u}_{\tilde{x}}) \tilde{w}_{\tilde{x}} d\tilde{y},$$

which defines the covering (20) in the tilded variables, provides

$$\begin{aligned} (e^{\lambda X})^* \tilde{\omega}_2 &= dw - (w^2 + (u_x + 2\lambda)w - u_y + \lambda u_x + \lambda^2) w_x dt - w_x dx \\ &\quad + (w + u_x + \lambda) w_x dy. \end{aligned}$$

This form defines a family of coverings

$$w_t = (w^2 + (u_x + 2\lambda)w - u_y + \lambda u_x + \lambda^2) w_x, \quad w_y = -(w + u_x + \lambda) w_x.$$

REMARK 2. In the case of  $\kappa = -2$  Eqs. (9) define a covering for Eq. (2), too, while we can't obtain this result by the method described above.

## References

- [1] Błaszak, M.: Classical R-matrices on Poisson algebras and related dispersionless systems, *Phys. Lett. A* **297**, 191–195 (2002)
- [2] Bogdanov, L.V., Konopelchenko B.G., Nonlinear Beltrami equations and  $\tau$ -functions for dispersionless hierarchies, *Phys. Lett. A* **322**, 330–337 (2004)
- [3] Bryant, R.L., Griffiths, Ph.A.: Characteristic cohomology of differential systems (II): conservation laws for a class of parabolic equations, *Duke Math. J.* **78**, 531–676 (1995)
- [4] Cartan, É.: *Œuvres Complètes*, **2**, Part II, Gauthier - Villars, Paris (1953)
- [5] Chang, J.-H., Tu, M.-H.: On the Miura map between the dispersionless KP and dispersionless modified KP hierarchies. *J. Math. Phys.*, **41**, 5391–5406 (2000).
- [6] Dodd, R., Fordy, A.: The prolongation structures of quasi-polynomial flows. *Proc. Roy. Soc. London, A*, **385**, 389–429 (1983)

- [7] Dunajski, M.: A class of Einstein–Weil spaces associated to an integrable system of hydrodynamic type, *J. Geom. Phys.* **51**, 126–137 (2004)
- [8] Estabrook F.B.: Moving frames and prolongation algebras. *J. Math. Phys.* **23**, 2071–2076 (1982)
- [9] Fels, M. and Olver, P.J.: Moving coframes. I. A practical algorithm. *Acta. Appl. Math.* **51**, 161–213 (1998)
- [10] Ferapontov, E.V., Khusnutdinova, K.R.: The characterization of two-component (2+1)-dimensional integrable systems of hydrodynamic type, *J. Phys. A.: Math. Gen.* **37** 2949–2963 (2004)
- [11] Gardner, R.B.: The method of equivalence and its applications. CBMS–NSF regional conference series in applied math., SIAM, Philadelphia (1989)
- [12] Harrison, B.K.: On methods of finding Bäcklund transformations in systems with more than two independent variables. *J. Nonlinear Math. Phys.*, **2**, 201–215 (1995)
- [13] Harrison, B.K.: Matrix methods of searching for Lax pairs and a paper by Estévez. *Proc. Inst. Math. NAS Ukraine*, **30**, Part 1, 17–24 (2000)
- [14] Hoenselaers C.: More prolongation structures. *Prog. Theor. Phys.* **75**, 1014–1029 (1986)
- [15] Igonin, S.: Coverings and the fundamental group for partial differential equations. *J. Geom. Phys.*, **56**, 939–998 (2006).
- [16] Igonin, S., Kersten, P., Krasil’shchik, I.: On symmetries and cohomological invariants of equations possessing flat representations. Preprint DIPS-07, The Diffiety Institute, Pereslavl-Zalessky (2002)
- [17] Kamran, N.: Contributions to the study of the equivalence problem of Élie Cartan and its applications to partial and ordinary differential equations. *Mem. Cl. Sci. Acad. Roy. Belg.*, **45**, Fac. 7 (1989)
- [18] Konopelchenko, B., Martínez Alonso L. Dispersionless scalar hierarchies, Whitham hierarchy and the quasi-classical  $\bar{\partial}$ -method, *J. Math. Phys.* **43** 3807–3823 (2003)
- [19] Krasil’shchik, I.S.: On one-parametric families of Bäcklund transformations. Preprint DIPS-1/2000, The Diffiety Institute, Pereslavl-Zalessky (2000)
- [20] Krasil’shchik, I.S. and Vinogradov, A.M.: Nonlocal symmetries and the theory of coverings. *Acta Appl. Math.*, **2**, 79–86 (1984)
- [21] Krasil’shchik, I.S., Lychagin, V.V., Vinogradov, A.M.: *Geometry of jet spaces and nonlinear partial differential equations*. Gordon and Breach, New York (1986)
- [22] Krasil’shchik, I.S., Vinogradov, A.M.: Nonlocal trends in the geometry of differential equations: symmetries, conservation laws, and Bäcklund transformations. *Acta Appl. Math.*, **15**, 161–209 (1989)
- [23] Krasil’shchik, I.S., Vinogradov, A.M. (eds.): *Symmetries and conservation laws for differential equations of mathematical physics*. Transl. Math. Monographs 182, Amer. Math. Soc., Providence (1999).
- [24] Krasil’shchik I.S. and Igonin S.A., On one-parametric families of Bäcklund transformations. *Lie Groups, Geometric Structures and Differential Equations—One Hundred Years After Sophus Lie* (T. Morimoto, H. Sato, and K. Yamaguchi, eds.), *Advanced Studies in Pure Mathematics*, vol. 37, Math. Soc. of Japan, 2002, 99–114, [arXiv:nlin.SI/0010040](https://arxiv.org/abs/nlin.SI/0010040)
- [25] Krichever, I.M.: The averaging method for two-dimensional ”integrable” equations. *Funct. Anal. Appl.* **22**, 200–213 (1988)
- [26] Kupershmidt, B.A.: The quasiclassical limit of the modified KP hierarchy. *J. Phys. A Math. Gen.* **23**, 871–886 (1990)
- [27] Kuz’mina, G.M.: On a possibility to reduce a system of two first-order partial differential equations to a single equation of the second order. *Proc. Moscow State Pedagog. Inst.* **271**, 67–76 (1967) (in Russian)
- [28] Marvan, M.: On zero-curvature representations of partial differential equations. *Proc. Conf. on Diff. Geom. and Its Appl., Opava (Czech Republic)*, 103–122 (1992)
- [29] Marvan, M.: A direct procedure to compute zero-curvature representations. The case  $\mathfrak{sl}_2$ . In: *Proc. Int. Conf. on Secondary Calculus and Cohomological Physics, Moscow, Russia, August 24–31,*

1997. Available via the Internet at ELibEMS, <http://www.emis.de/proceedings>.
- [30] Marvan, M.: On the horizontal gauge cohomology and nonremovability of the spectral parameter. *Acta Appl. Math* **72**, 51–65 (2002)
  - [31] Morozov, O.I.: Moving coframes and symmetries of differential equations. *J. Phys. A, Math. Gen.*, **35**, 2965–2977 (2002)
  - [32] Morozov, O.I.: Contact-equivalence problem for linear hyperbolic equations. *J. Math. Sci.*, **135**, 2680–2694 (2006)
  - [33] Morozov, O.I.: Coverings of differential equations and Cartan’s structure theory of Lie pseudo-groups. *Acta Appl. Math.* **99**, 309–319 (2007)
  - [34] Morozov, O.I.: Cartan’s structure theory of symmetry pseudo-groups, coverings and multi-valued solutions for the Khokhlov–Zabolotskaya equation, *Acta Appl. Math.* **101**, 231–241 (2008)
  - [35] Morris, H.C.: Prolongation structures and nonlinear evolution equations in two spatial dimensions. *J. Math. Phys.*, **17**, 1870–1872 (1976)
  - [36] Morris, H.C.: Prolongation structures and nonlinear evolution equations in two spatial dimensions: a general class of equations. *J. Phys. A, Math. Gen.*, **12**, 261–267 (1979)
  - [37] Olver, P.J.: *Equivalence, invariants, and symmetry*. Cambridge, Cambridge University Press (1995)
  - [38] Sakovich, S.Yu.: On zero-curvature representations of evolution equations. *J. Phys. A, Math. Gen.*, **28**, 2861–2869 (1995)
  - [39] Takasaki, K.: Quasi-classical limit of BKP hierarchy and  $W$ -infinity symmetries, *Lett. Math. Phys.* **28**, 177–185 (1993)
  - [40] Tondo, G.S.: The eigenvalue problem for the three-wave resonant interaction in  $(2+1)$  dimensions via the prolongation structure. *Lett. Nuovo Cimento*, **44**, 297–302 (1985)
  - [41] Wahlquist, H.D., Estabrook F.B.: Prolongation structures of nonlinear evolution equations. *J. Math. Phys.*, **16**, 1–7 (1975)
  - [42] Zakharov, V.E.: Integrable systems in multidimensional spaces. *Lect. Notes Phys.*, **153**, 190–216 (1982)