## Poincaré Recurrence, Zermelo's Second Law Paradox, and Probabilistic Origin in Statistical Mechanics

P. D. Gujrati\*

The Department of Physics, The University of Akron, Akron, OH 44325 (Dated: September 11, 2021)

Abstract

We show that Poincaré recurrence does not mean that the entropy will eventually decrease, contrary to the claim of Zermelo, and that the probabilitistic origin in statistical physics must lie in the external noise, and not the preparation of the system.

As first pointed out by Kröning [1], and later developed by Boltzmann [2], any deep understanding of the second law of thermodynamics in terms of entropy due to Clausius [3] must involve a probabilistic approach; see [4, 5] for two of the excellent reviews. This was the first approach in physics to establish that fundamental laws of Nature need not be strictly deterministic. However, as many phenomena at the microscopic level such as nuclear decay also require a probabilistic approach for their understanding, the probabilistic interpretation is not just a consequence of a macroscopic system; yet it has to be exploited in statistical physics. To appreciate this probabilistic approach in statistical physics, we note that in the Gibbs formulation, the entropy is given by the average of the degree of uncertainty  $u_i(t) = -\ln p_i(t)$  of the *i*th microstate:

$$S(t) = \sum p_i(t)u_i(t) \ge 0, \quad \sum p_i(t) = 1;$$
 (1)

here  $p_i(t)$  is the probability for the *i*th microstate at time t, and the sum is over all distinct microstates W. We allow the possibility that some of the probabilities may be zero. Thus, one envisions the system to be in different microstates with certain probabilities. To introduce the concept of  $p_i(t)$ , we construct a Gibbs ensemble as containing  $\mathcal{N}$  replicas of the system,  $\mathcal{N}_i$  of which are in microstate *i*. Then,  $p_i(t) \equiv \mathcal{N}_i/\mathcal{N}$  in the limit  $\mathcal{N} \to \infty$ ; the limit will always be implicit. If there is only one microstate i = 0 possible [6] so that  $\mathcal{N}_0 = \mathcal{N}$ , the entropy is identically zero as the system is in i = 0 with certainty. According to the second law, the entropy of an isolated system cannot decrease. The equilibrium is attained when the entropy becomes maximum, which occurs when all microstates have the same probability:

$$p_i(t) \to 1/W. \tag{2}$$

Once the equilibrium is achieved, the entropy cannot decrease if the system is left undisturbed.

However, the application of the Poincaré recurrence theorem [7], see below, gave rise to Zermelo's [8] paradox, which has not been resolved to everyone's satisfaction yet, and is the subject matter of this work. The recurrence theorem is valid for a classical system and basically states that provided an isolated mechanical system, in which the forces do not depend on the velocities of the particles, remains in a *finite* part of the phase space during its evolution, then the uniqueness of classical trajectories implies that a given initial state must come arbitrary close to itself infinitely many times. Zermelo [8] argued that since the entropy is determined by the phase point, then it must also return to its original value according to the recurrence theorem. Thus, if the entropy increases during a part of the time, it must decrease during another and this increase and decrease in the entropy must occur infinitely many times, thereby violating the second law. In addition, it is not just the microstate itself, but its probability of occurrence that determines the entropy. As we demonstrate here, not appreciating this fact has given rise to the paradox due to Zermelo. We will also show that the origin of this probabilistic behavior in not in the method of preparation of the system, which leaves the system deterministic; rather it lies in the stochastic interaction with the environment, no matter how weak, for the second law to work.

According to Boltzmann [9], recurrences are not inconsistent with the statistical viewpoint: they are merely statistical fluctuations, which are almost certain to occur. Indeed, Boltzmann [9], Smoluchowski [10], and others recognized that the period of a Poincaré cycle is so much larger for a macroscopic system to be almost infinitely large so that the *violation* of the second law (decrease in entropy of an isolated system) would be almost impossible to occur in our life. The period of the cycle will be many orders of magnitude larger than the present age of the universe [12]. While an appealing argument, it is hard to understand its relevance as the argument compares presumably a system-intrinsic time, the recurrence time, with a system-extrinsic time, the time of observation or the age of the universe. Moreover, the recurrence theorem is valid for a *deterministic system* as will be detailed below, while the second law is valid for a stochastic system requiring a probabilistic approach which necessiates exploiting an ensemble . In particular, Poincaré's recurrence theorem states that the mechanical system will revisit the neighborhood of its initial state with *cer*tainty (with probability p = 1), while for a statistical system, the probability of revisit is extremely small (indeed  $p \simeq 1/W$  for a macroscopically large system) as we discuss below.




FIG. 1: Schematic evolution of an initial microstate 0 as a function of time in the phase space. A cell shows a microstate and the numbers show the sequential emrgence of microstates in time. (a) shows not only the unique deterministic evolution, but also one of many possible stochastic evolution. Being a deterministic evolution, no microstate recurs except the initial one at time  $t_{\rm R}$ . (b) shows one of the many possible stochastic evolution at time  $t_{\rm R}$ .

We first discuss the recurrence theorem for completeness and then its relevance for the second law.

**Theorem 1** Poincaré Recurrence A microstate of a finite classical system evolving deterministically and confined to a finite region of the phase space during its evolution recurs infinitely many times.

**Proof.** We consider a classical system consisting of N particles, which we take to be point-like for simplicity, with energy E in a volume V. We restrict N, E and V to be finite to ensure that the system moves in a *finite* region of size  $|\Gamma_0|$  in the phase space; see the shaded region in Fig. 1. A microstate is commonly defined not by a point in the phase space, but by a small volume (shaded cells in Fig. 1) of the size  $\tau_0 \equiv h^{2r}$ ; h is Planck's constant and r = 3N [13]. The number of distinct microstates W is

$$W \equiv \left| \Gamma_0 \right| / \tau_0,$$

which is exponentially large of the order of  $c^N$ , with  $c \geq 1$ some constant. We assume, for simplicity, that the time required for a microstate to evolve into a different microstate is some constant  $\Delta$ , and observe the system at times  $t = t_j = j\Delta, j = 1, 2, \cdots$ , to determine the microstates. The dynamics of an isolated system with a given Hamiltonian is completely deterministic [14]: an initial microstate i evolves in a *unique* fashion into a microstate  $i_j$  at time  $t = t_j$ , which we represent by the one-to-one mapping  $i \rightarrow i_j$ . Due to the unique evolution, the system visits each of the W microstates in time without *repeating* until it has visited all of them. It will then revisit the initial microstate (it cannot visit any other microstate because of the unique evolution) and then repeat the entire sequence  $\{i_i\}$  exactly in the same order over and over. We show in Fig. 1a the deterministic evolution of the initial microstate i = 0 through microstates  $0_j$  at  $t = t_j$ , shown schematically by j = $1, 2, \cdots, W' \equiv W - 1$ . The next microstate at j = Wwill be 0 [15], and the entire ordered sequence  $\{0_i\}$  will be visited during the next cycle. The recurrence time  $t_{\rm R}$ , also known as the Poincaré cycle, is given by  $t_{\rm R} \equiv W\Delta$ . Each microstate will be revisited several times in a time  $t \gg t_{\rm R}$ . This proves the recurrence theorem [15].

A more general proof can be found in [12].

**Theorem 2** The entropy in a Poincaré cycle remains constant so that the second law is never violated.

**Proof.** Consider Fig. 1a. Since the system is with certainty in only one microstate  $j \pmod{W}$  at instant  $t = t_j$ , its entropy  $S(t_j) = 0$  identically for all j. As  $S(t_j)$  can never decrease, the phenomenon of recurrence does not violate the second law.

The same conclusion is also obtained in the ensemble approach. We prepare each replica in the same microstate 0 initially and follow its evolution in time. Because the evolution is deterministic, each replica in the ensemble will be in the same microstate  $0_j$  at  $t = t_j$ . Thus,  $p_i(t_j) = \delta_{ij}$ , which again gives  $S(t_j) = 0$ . Let us now consider the system to be initially in a "macrostate" consisting of two possible microstates 0 and 1 with probabilities  $p_0$  and  $p_1 \equiv 1 - p_0$ , respectively. This can also be done for a quantum system. There are  $\mathcal{N}p_0$  replicas in the microstate 0, and  $\mathcal{N}p_1$  in the microstate 1, and the initial entropy is  $S(0) = -p_0 \ln p_0 - p_1 \ln p_1$ . Since the evolution  $(0 \rightarrow 0_i, 1 \rightarrow 1_i)$  at some later time  $t_i$  is deterministic, all the  $\mathcal{N}p_0$  replicas are in microstate  $0_i$ , and the remaining  $\mathcal{N}p_1$  in microstate  $1_i$ , so that  $\Pr(0_i) = p_0$ , and  $Pr(1_i) = p_1$ . Consequently,  $S(t_i) = S(0)$  so the entropy remains constant. It is easy to extend the calculation to an initial "macrostate" consisting of any number of microstates i, in particular all microstates W, with probabilities  $p_i$  with the same conclusion that the entropy given by (1) remains constant during the Poincaré cycle. This completes the proof.  $\blacksquare$ 

The above conclusion is consistent with *time-reversal invariance* in a deterministic dynamics. As the evolution  $i \rightarrow i_j$  is one-to-one, it can be inverted at any time. Thus, the forward evolution  $i \rightarrow i_1 \rightarrow i_2 \rightarrow \cdots \rightarrow i_j$  of an initial microstate can be uniquely inverted to give  $i_j \rightarrow$  $i_{j-1} \rightarrow i_{j-2} \rightarrow \cdots \rightarrow i_1 \rightarrow i$ , and we recover the initial microstate in this reversal. The entropy in this reversal remains constant to ensure time-reversal invariance.

It should be commented that the non-zero initial entropy for a deterministic system (classical or quantum) considered above is due to our mode of preparation. It is due to our ignorance about the system and does not represent an intrinsic property of the system. The notion of probability here is brought into the discussion due to the preparation of the system, and we have total control to change its probabilistic nature and to change the entropy so that the latter is not an intrinsic characteristic of the system. This entropy of the deterministic system can be readily changed to zero by making a precise measurement to determine which microstate the system is in [6]. This "collapse" of the "macrostate" means that once the system is known to be in a particular microstate with certainty after measurement, its entropy will remain zero as shown above for ever, even though the system is not in equilibrium. This is not how we expect the thermodynamic entropy to behave. Moreover, being a constant, this entropy will never become the maximum possible equilibrium entropy  $\ln W$ , unless it is already at the maximum. This gives us the following

## **Corollary 3** A deterministic system will never equilibrate if it was not in equilibrium initially.

Ideal gases confined by idealized walls to form isolated systems have no mechanism to achieve equilibrium, and will remain in non-equilibrium states for ever if they were so initially; see footnote 6 in [4]. For the concept of entropy to be useful requires it to be an intrinsic property of the system which should not be affected by the measurements if we wait long enough after the measurements. Thus, the concept of entropy requires a particular kind of probabilistic approach in which the evolution must not be deterministic; rather, it must be *stochastic*. Even an isolated system is not truly deterministic in Nature. A real system must be confined by a real container, which cannot be a perfect insulator. Even the container will introduce environmental noise in the system. Thus, there are always stochastic processes going on in a real system, which cannot be eliminated, though they can be minimized. In the case the external noise is too strong, then there is no sense in not considering the environment as part of the system for its thermodynamic investigation. It is the limit in which the external noise is too weak that is relevant for a sensible thermodynamical description of a system, so that the external noise will not alter the average properties such as the average energy of the system [11]. For quantum systems, this requires considering the Landau-von Neumann density matrix, rather than eigenstates; see, for example, [12]. The derivation in [12] clearly shows the uncertainty introduced by the presence of the environment. The latter is not part of the system, just as in thermodynamics. It is only in this case that the entropy will vary as the probabilities of various microstates change in time, as we describe below.

The stochasticity introduces a new time scale  $\Delta'$  over which the system evolves deterministically as above. Over this time-period, the the mapping  $i \to i_i$  is oneto-one and can be inverted to study time-reversal. The entropy remains constant during this period. At the end of each time period  $\Delta'$ , i.e. at time  $t = t'_k \equiv k$  $\Delta', k = 1, 2, \cdots$ , the current microstate *i* will undergo a stochastic "jump" (shown by the double arrow  $i \rightarrow j$ ) to any of the W microstates i [16] brought about by the environmental noise. We take these "jumps" to occur instantaneously just for simplicity. The "jump" may create a new microstate not generated so far, or bring it back to a previously visited microstate, including the initial microstate, generated during its deterministic evolution. Such a jump to a previously generated microstate (not the initial microstate) would have been forbidden in a deterministic evolution alone as noted above. Many such stochastic "jumps" are needed to bring the system to equilibrium, which requires a time interval  $t_{eq}$ , so that  $\Delta' < t_{\rm eq}$ . The presence of stochastic "jumps" give rise to a probabilistic nature to the microstates, their probability of occurrence changing with time. This in turn changes the entropy with time whenever "jumps" occur.

How  $\Delta'$  relates to the timescale  $\Delta$  depends on the strength of the noise; here we will assume  $\Delta' \gtrsim \Delta$ , which can be reversed without affecting our conclusions. For an isolated system,  $\Delta' \rightarrow \infty$ , which is consistent with our Corollary that the deterministic evolution cannot bring about equilibration (entropy maximization). The external noise causes the entropy to increase with time if the initial state was out of equilibrium as shown elsewhere [11]. Therefore, we now turn to the stochastic evolution to make contact with the second law.

We do not have to consider the actual nature of the noise; all that is required is its presence. The actual nature will only determine the value of  $\Delta'$ , but not the final equilibrium state, which remains oblivious to the actual noise. This is what allows the statistical mechanical approach to make predictions about the equilibrium state. We consider an ensemble of  $\mathcal{N}$  replicas, each replica being identically prepared in the same microstate 0, so that  $p_i(0) = \delta_{i0}$ . Consequently, S(0) = 0. This obviously represents an extreme non-equilibrium situation. However, since the evolution is stochastic, a microstate *i* makes a "jump" to another microstate *j* (*i*  $\rightarrow$  *j*) or remain the same (*i*  $\rightarrow$  *i*) caused by the noise. It is also possible to have *i*  $\rightarrow$  *j*  $\rightarrow$  *i*, as shown in Fig. 1b, where we show that the system leaves the original microstate 0 but

comes back to it at  $t_k$ . Thus, the recurrence can happen at any time  $t \ge \Delta'$ , albeit without certainty (probability  $p_0(t) < 1$ ) and has no particular significance or relevance for the Poincaré cycle for a finite system, where recurrence occurs with certainty. This distinction in the probability of recurrence is very important, as the entropy is determined by the probability. Just because the initial microstate has recurred does not necessarily mean that the entropy has reversed to its initial value S(0) = 0, contrary to the claim by Zermelo. One needs to consider its probability also. To establish this, we proceed as follows.

At  $t'_1$ , there will be  $\mathcal{N}p_i(t'_1)$  replicas in the *i*th microstates. In particular, there is a non-zero probability  $p_0(t'_1) < 1$  that the system will be back in its original microstate 0. However, this in no way means that the average degree of uncertainty  $S(t'_1)$  has reduced to zero, as it is obtained by (1), which requires a sum over *all* microstates that are present in the ensemble. The degree of uncertainty of the initial microstate

$$u_0(t_1') = -\ln p_0(t_1') > 0,$$

so that  $S(t'_1) > p_0(t'_1)u_0(t'_1) > 0$ . The entropy has increased. For  $t'_1 < t < t'_2$ , each replica evolves deterministically so that the entropy remains constant, as follows from Theorem 2. This is true during each of the intervals  $t'_k < t < t'_{k+1}$ , with the entropy changing at  $t'_k$  as the probabilities  $p_i(t'_k)$  change. During all this time, the system has a non-zero probability  $p_0(t) < 1$  to be in the initial microstate 0. Eventually, the system equilibrates when (2) holds so that

$$u_i(t) \rightarrow \ln W, \quad i = 0, 1, \cdots, W',$$

which is exactly the entropy  $S(t) = \ln W$  of the system obtained by summing over all microstates; see (1). We observe that in equilibrium, the entropy is exactly the degree of uncertainty of any microstate and, in particular, the initial state.

Thus, we come to the following theorem:

**Theorem 4** Even for a stochastic evolution, which is needed for a statistical system, the recurrence of the initial microstate does not violate the second law.

The entropy remains constant after equilibrium is reached. The recurrence of the initial microstate 0 (but with  $p_0(t) = 1/W$ ) in the stochastic case does not mean that the entropy reverts to the initial entropy S(0). On the other hand, the *true recurrence* of the initial microstate 0  $[p_0(0) = 1, S(0) = 0]$  requires  $p_0(t) = 1$ , for which all replicas must be in the microstate 0 simultaneously. This can occur in only one way. However, such a true recurrence is impossible in stochastic systems. To show this, we consider the situation in equilibrium; see (2). The number of possible ways the replicas can be arranged at time t, consistent with microstate probabilities  $p_i(t)$ , is

$$\mathcal{N}!/\prod [\mathcal{N}p_i(t)]! \approx e^{\mathcal{N}S(t)} = W^{\mathcal{N}}$$

one of which is the true recurrent state. Hence, the probability for the initial microstate to truly recur is  $W^{-\mathcal{N}} \to 0$  as  $\mathcal{N} \to \infty$ . The recurrence of 0 occurs [5, 12] several times, but with  $p_0(t) < 1$ , so that other microstates have to be considered to determine S(t).

In conclusion, we have shown that the entropy of an isolated deterministic system in a Poincaré cycle remains constant, so there is no violation of the second law. Furthermore, we have also shown that the second law requires that the probabilistic nature of microstates must be caused by external noise, and not the mode of preparation alone. Once the entropy reaches its maximum value, it remains constant. It never decreases. Thus, the second law is never violated.

- \* Electronic address: pdg@arjun.physics.uakron.edu
- [1] A. Kröning, Ann. Physik, 99, 315 (1856).
- [2] L. Boltzmann, Wien. Ber. 76, 373 (1877).
- [3] R. Clausius Ann. Physik, **125**, 353 (1865).
- [4] D. ter Haar, Rev. Mod. Phys. 27, 289 (1955).
- [5] J.L. Lebowitz, Rev. Mod. Phys. 71, S346 (1999).
- [6] In classical physics, it is taken for granted as an axiom that a precise measurement and preparation can always be made, at least in principle, so that the state can be prepared in a unique microstate. In quantum mechanics, the same is also true of an isolated system. A quantum measurement, according to one of the axioms, will prepare the system in one of the eigenstates; the latter are identified as the microstates.
- [7] H. Poincaré Acta Math. 13,1-270 (1890); see also S. Chandrasekhar, Rev. Mod. Phys. 15, 1 (1943).
- [8] E. Zermelo, Ann. Physik, 57, 485-494 (1896); ibid 59, 793 (1896); Wied. Ann., 60, 392-398 (1897).
- [9] L. Boltzmann, Ann. Physik, 57, 773 (1896); ibid 60, 392 (1897).
- [10] M. von Smoluchowski, *Physik. Z.*, **13**, 1069 (1912); **14**, 261 (1913).
- [11] P.D. Gujrati, arXiv:0803.1099.
- [12] K. Huang, *Statistical Mechanics*, Second Edition, John Wiley and Sons, New York (1987).
- [13] The division in cells is to ensure that the number of microstates does not become infinite even for a finite system (finite N, E and V).
- [14] Even the collisions are deterministic in such a system.
- [15] An isolated quantum system, which is prepared in a particular microstate [6], always remains in the same microstate (if we do not consider the changes in the phase of the eigenstate relevant), so that we may set  $t_{\rm R} \equiv 0$ . Here, we are assuming that the eigenstates are mutually orthogonal. If the phase change  $Et/\hbar$  is considered relevant, the recurrence time will be  $t_{\rm R} \equiv h/E$  when the phase changes by  $2\pi$ .
- [16] As the system is no longer isolated because of its interaction with the environment, E, N, V need not remain constant and may fluctuate. However, as long as we are deal-

ing with very weak environmental noise, we can safely treat the system as quasi-isolated in that the widths of

their spread can be neglected.