CENTRE DE PHYSIQUE THÉORIQUE ¹

CNRS-Luminy, Case 907 13288 Marseille Cedex 9 FRANCE

Spectral action on $SU_q(2)$

B. $Iochum^{1,2}$, C. $Levy^{1,2}$ and A. $Sitarz^{3,4}$

Abstract

The spectral action on the equivariant real spectral triple over $\mathcal{A}(SU_q(2))$ is computed explicitly. Properties of the differential calculus arising from the Dirac operator are studied and the results are compared to the commutative case of the sphere \mathbb{S}^3 .

PACS numbers: 11.10.Nx, 02.30.Sa, 11.15.Kc MSC-2000 classes: 46H35, 46L52, 58B34

CPT-P06-2007

 $^{^{1}}$ UMR 6207

⁻ Unité Mixte de Recherche du CNRS et des Universités Aix-Marseille I, Aix-Marseille II et de l'Université du Sud Toulon-Var

[–] Laboratoire affilié à la FRUMAM – FR 2291

² Also at Université de Provence, iochum@cpt.univ-mrs.fr, levy@cpt.univ-mrs.fr

³ Institute of Physics, Jagiellonian University, Reymonta 4, 30-059 Kraków, Poland, sitarz@if.uj.edu.pl

 $^{^4}$ Partially supported by Polish Government grants 189/6. PRUE/2007/7; 115/E-343/SPB/6. PR UE/DIE and N 201 1770 33

Contents

| 1 Introduction | | | | | | | | |
|----------------|--|--|--|--|--|--|--|--|
| 2 | Spectral action in 3-dimension 2.1 Tadpole and cocycles | 3 3 6 | | | | | | |
| 3 | $SU_q(2)$ triple The spectral triple | | | | | | | |
| 4 | Reality operator and spectral action on $SU_q(2)$ 4.1 Spectral action in dimension 3 with $[F, A] \in OP^{-\infty}$ | 14 14 15 16 17 22 23 30 | | | | | | |
| 5 | $\begin{array}{llllllllllllllllllllllllllllllllllll$ | 31 33 36 37 40 | | | | | | |
| 6 | Examples of spectral action | | | | | | | |
| 7 | The commutative sphere \mathbb{S}^3 7.1 Translation of Dirac operator | | | | | | | |
| 8 | Conclusion | 45 | | | | | | |

1 Introduction

The quantum group $SU_q(2)$ has already a rather long history of studies [33] being one of the finest examples of quantum deformation. This includes an approach via the noncommutative notion of spectral triple introduced by Connes [10, 15] and various notions of Dirac operators were introduced in [2, 4, 6, 13, 28]. Finally, a real spectral triple, which was exhibited in [21], is invariant by left and right action of $\mathcal{U}_q(su(2))$ and satisfies almost all postulated axioms of triples except the commutant and first-order properties. These, however, remain valid only up to infinitesimal of arbitrary high order. The last presentation generalizes in a straightforward way all geometric construction details of the spinorial spectral triple for the classical three-sphere. In

particular, both the equivariant representation and the symmetries have a $q \to 1$ proper classical limit.

The goal of this article is to obtain the spectral action defined in [7] by

$$S(\mathcal{D}_A, \Phi, \Lambda) := \text{Tr}\left(\Phi(\mathcal{D}_A/\Lambda)\right) \tag{1}$$

where \mathcal{D} is the Dirac operator, A is a selfadjoint one-form, $\mathcal{D}_A = \mathcal{D} + A + JAJ^{-1}$ and J is the reality operator. Here, Φ is any even positive cut-off function which could be replaced by a step function up to some mathematical difficulties investigated in [23]. This means that \mathcal{S} counts the spectral values of $|\mathcal{D}_A|$ less than the mass scale Λ . Actually, as shown in [8]

$$\mathcal{S}(\mathcal{D}_A, \Phi, \Lambda) = \sum_{0 \le k \in Sd^+} \Phi_k \Lambda^k \int |D_A|^{-k} + \Phi(0) \zeta_{D_A}(0) + \mathcal{O}(\Lambda^{-1}), \tag{2}$$

$$\zeta_{D_A}(0) = \zeta_D(0) + \sum_{k=1}^d \frac{(-1)^k}{k} \int (A\mathcal{D}^{-1})^k.$$
(3)

where $D_A = \mathcal{D}_A + P_A$, P_A the projection on $\operatorname{Ker} \mathcal{D}_A$, $\Phi_k = \frac{1}{2} \int_0^\infty \Phi(t) \, t^{k/2-1} \, dt$, d is the spectral dimension of the triple and Sd^+ is the strictly positive part of the dimension spectrum Sd of $(\mathcal{A}, \mathcal{H}, \mathcal{D})$. Here, $Sd^+ = Sd = \{1, 2, 3\}$, so

$$\mathcal{S}(\mathcal{D}_A, \Phi, \Lambda) = \sum_{1 \le k \le 3} \Phi_k \Lambda^k \int |D_A|^{-k} + \Phi(0) \zeta_{D_A}(0). \tag{4}$$

Recall that the tadpole of order Λ^k is the linear term in $A \in \Omega^1_D(A)$ in the Λ^k part of (4). Note that there are no terms in Λ^{-k} , k > 0 because the dimension spectrum is bounded below by 1. This spectral action has been computed on few examples: [3,8,9,15,22,24–26,30,34].

Here, we compute (4) with the main difficulty which is to control the differential calculus generated by the Dirac operator. To proceed, we introduce two presentations of one-forms. The main ingredient is $F = \text{sign }(\mathcal{D})$ which appears to be a one-form up to $OP^{-\infty}$.

In section 2, we discuss the spectral action of an arbitrary 3-dimensional spectral triple using cocycles.

In sections 3 and 4 we recall the main results on $SU_q(2)$ of [21] and show that the full spectral action with reality operator given by (4) is completely determined by the terms

$$\oint A^q |\mathcal{D}|^{-p}, \quad 1 \le q \le p \le 3.$$

This question of computation of spectral action was addressed in the epilogue of [37]. In section 5, we establish a differential calculus up to some ideal in pseudodifferential operators and apply these results to the precise computation of previous noncommutative integrals. Section 6 is devoted to explicit examples, while in next section are given different comparisons with the commutative case of the 3-sphere corresponding to SU(2).

2 Spectral action in 3-dimension

2.1 Tadpole and cocycles

Let $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ be a spectral triple of dimension 3. For $n \in \mathbb{N}^*$ and $a_i \in \mathcal{A}$, define

$$\phi_n(a_0,\cdots,a_n) := \int a_0[\mathcal{D},a_1]\mathcal{D}^{-1}\cdots[\mathcal{D},a_n]\mathcal{D}^{-1}.$$

We also use notational integrals on the universal n-forms $\Omega_n^n(\mathcal{A})$ defined by

$$\int_{\phi_n} a_0 da_1 \cdots da_n := \phi_n(a_0, a_1, \cdots, a_n).$$

and the reordering fact that $(da_0)a_1 = d(a_0a_1) - a_0da_1$.

We use the b-B bicomplex defined in [10]: b is the Hochschild coboundary map (and b' is truncated one) defined on n-cochains ϕ by

$$b\phi(a_0,\ldots,a_{n+1}) := b'\phi(a_0,\ldots,a_{n+1}) + (-1)^{n+1}\phi(a_{n+1}a_0,a_1,\ldots,a_n),$$

$$b'\phi(a_0,\ldots,a_{n+1}) := \sum_{j=0}^{n} (-1)^j\phi(a_0,\ldots,a_ja_{j+1},\ldots,a_{n+1}).$$

Recall that B_0 is defined on the normalized cochains ϕ_n by

$$B_0\phi_n(a_0, a_1, \dots, a_{n-1}) := \phi_n(1, a_0, \dots, a_{n-1}), \text{ thus } \int_{\phi_n} d\omega = \int_{B_0\phi_n} \omega \text{ for } \omega \in \Omega_u^{n-1}(\mathcal{A}).$$

Then $B := NB_0$, where $N := 1 + \lambda + \dots \lambda^n$ is the cyclic skewsymmetrizer on the *n*-cochains and λ is the cyclic permutation $\lambda \phi(a_0, \dots, a_n) := (-1)^n \phi(a_n, a_0, \dots, a_{n-1})$. We will also encounter the cyclic 1-cochain $N\phi_1$:

$$N\phi_1(a_0, a_1) := \phi_1(a_0, a_1) - \phi_1(a_1, a_0)$$
 and $\int_{N\phi_1} a_0 da_1 := N\phi_1(a_0, a_1)$.

Remark 2.1. Assume the integrand of f is in OP^{-3} . Since $[\mathcal{D}^{-1}, a] = -\mathcal{D}^{-1}[\mathcal{D}, a]\mathcal{D}^{-1} \in OP^{-2}$, this commutator introduces a integrand in OP^{-4} so has a vanishing integral: under the integral, we can commute \mathcal{D}^{-1} with all $a \in \mathcal{A}$ and all one-forms.

Lemma 2.2. We have

- $(i) b\phi_1 = -\phi_2.$
- (ii) $b\phi_2 = 0.$
- (iii) $b\phi_3 = 0.$
- (iv) $B\phi_1 = 0.$
- (v) $B_0\phi_2 = -(1-\lambda)\phi_1.$
- (vi) $bB_0\phi_2 = 2\phi_2 + B_0\phi_3$.
- (vii) $B\phi_2 = 0.$
- (viii) $B_0\phi_3 = Nb'\phi_1$.
- $(ix) \quad B\phi_3 = 3B_0\phi_3.$

Proof. (i)

$$b\phi_1(a_0, a_1, a_2) = \int a_0 a_1[\mathcal{D}, a_2] \mathcal{D}^{-1} - \int a_0 (a_1[\mathcal{D}, a_2] + [\mathcal{D}, a_1] a_2) \mathcal{D}^{-1} + \int a_2 a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}$$

$$= \int a_0[\mathcal{D}, a_1] (\mathcal{D}^{-1} a_2 - a_2 \mathcal{D}^{-1}) = -\int a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}[\mathcal{D}, a_2] \mathcal{D}^{-1}$$

$$= -\phi_2(a_0, a_1, a_2)$$

where we have used the trace property of the noncommutative integral.

 $(ii) b\phi_2(a_0, a_1, a_2, a_3)$

$$\begin{split} &= \int a_0 a_1[\mathcal{D}, a_2] \mathcal{D}^{-1}[\mathcal{D}, a_3] \mathcal{D}^{-1} - \int a_0 \left(a_1[\mathcal{D}, a_2] + [\mathcal{D}, a_1] a_2 \right) \mathcal{D}^{-1}[\mathcal{D}, a_3] \mathcal{D}^{-1} \\ &+ \int a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}(a_2[\mathcal{D}, a_3] + [\mathcal{D}, a_2] a_3) \mathcal{D}^{-1} - \int a_3 a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}[\mathcal{D}, a_2] \mathcal{D}^{-1} \\ &= \int a_0[\mathcal{D}, a_1] \left(\mathcal{D}^{-1} a_2 - a_2 \mathcal{D}^{-1} \right) [\mathcal{D}, a_3] \mathcal{D}^{-1} + \int a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}[\mathcal{D}, a_2] \left(a_3 \mathcal{D}^{-1} - \mathcal{D}^{-1} a_3 \right) \\ &= - \int a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}[\mathcal{D}, a_2] \mathcal{D}^{-1}[\mathcal{D}, a_3] \mathcal{D}^{-1} + \int a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}[\mathcal{D}, a_2] \mathcal{D}^{-1}[\mathcal{D}, a_3] \mathcal{D}^{-1} \\ &= 0. \end{split}$$

(iii) Using Remark 2.1, we get $\phi_3(a_0, a_1, a_2, a_3) = \int a_0[\mathcal{D}, a_1][\mathcal{D}, a_2][\mathcal{D}, a_3]|\mathcal{D}|^{-3}$, so similar computations as for ϕ_2 gives $b\phi_3 = 0$.

(iv)
$$B_0\phi_1(a_0) = f[\mathcal{D}, a_0]\mathcal{D}^{-1} = f(\mathcal{D}a_0\mathcal{D}^{-1} - a_0) = 0.$$

$$(v) \quad B_0\phi_2(a_0, a_1) = \int [\mathcal{D}, a_0] \mathcal{D}^{-1}[\mathcal{D}, a_1] \mathcal{D}^{-1} = \int a_0 \mathcal{D}^{-1}[\mathcal{D}, a_1] - \int a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}$$
$$= \int a_0 a_1 - \int a_0 \mathcal{D}^{-1} a_1 \mathcal{D} - \int a_0[\mathcal{D}, a_1] \mathcal{D}^{-1}$$
$$= -\int a_1[\mathcal{D}, a_0] \mathcal{D}^{-1} - \int a_0[\mathcal{D}, a_1] \mathcal{D}^{-1} = -\phi_1(a_1, a_0) - \phi_1(a_0, a_1).$$

(vi) Since
$$-b\lambda\phi_1(a_0, a_1, a_2) = \phi_1(a_2, a_0a_1) - \phi_1(a_1a_2, a_0) + \phi_1(a_1, a_2a_0)$$
, one obtains that $-b\lambda\phi_1(a_0, a_1, a_2) = \int a_0a_1\mathcal{D}^{-1}a_2\mathcal{D} + a_0\mathcal{D}^{-1}a_1\mathcal{D}a_2 - a_0\mathcal{D}^{-1}a_1a_2\mathcal{D} - a_0a_1a_2$.

So by direct expansion, this is equal to $-\int a_0 \mathcal{D}^{-1}[\mathcal{D}, a_1] \mathcal{D}^{-1}[\mathcal{D}, a_2]$ which means that

$$-b\lambda\phi_1(a_0, a_1, a_2) = \int [\mathcal{D}^{-1}, a_0][\mathcal{D}, a_1]\mathcal{D}^{-1}[\mathcal{D}, a_2] - a_0[\mathcal{D}, a_1]\mathcal{D}^{-1}[\mathcal{D}, a_2]\mathcal{D}^{-1}$$
$$= -B_0\phi_3(a_0, a_1, a_2) - \phi_2(a_0, a_1, a_2).$$

Now the result follows from (i), (v).

(vii)
$$B\phi_2 = NB_0\phi_2 = -N(1-\lambda)\phi_1 = 0$$
 since $N(1-\lambda) = 0$.

$$(viii) \quad B_0\phi_3(a_0,a_1,a_2) = \int [\mathcal{D},a_0]\mathcal{D}^{-1}[\mathcal{D},a_1]\mathcal{D}^{-1}[\mathcal{D},a_2]\mathcal{D}^{-1}$$

$$= \int a_0\mathcal{D}^{-1}[\mathcal{D},a_1]\mathcal{D}^{-1}[\mathcal{D},a_2] - \int a_0[\mathcal{D},a_1]\mathcal{D}^{-1}[\mathcal{D},a_2]\mathcal{D}^{-1}$$

$$= \int a_0a_1\mathcal{D}^{-1}[\mathcal{D},a_2] - \int a_0\mathcal{D}^{-1}a_1[\mathcal{D},a_2] - \int a_0[\mathcal{D},a_1]\mathcal{D}^{-1}[\mathcal{D},a_2]\mathcal{D}^{-1}$$

$$= \int a_0a_1a_2 - \int a_0a_1\mathcal{D}^{-1}a_2\mathcal{D} - \int a_0\mathcal{D}^{-1}a_1\mathcal{D}a_2 + \int a_0\mathcal{D}^{-1}a_1a_2\mathcal{D}$$

$$- \int a_0[\mathcal{D},a_1]\mathcal{D}^{-1}[\mathcal{D},a_2]\mathcal{D}^{-1}$$

$$= \int a_0a_1a_2 - a_2\mathcal{D}a_1a_0\mathcal{D}^{-1} + a_1a_2\mathcal{D}a_0\mathcal{D}^{-1} + a_2\mathcal{D}a_0a_1\mathcal{D}^{-1}$$

$$- \left(a_0\mathcal{D}a_1a_2\mathcal{D}^{-1} - a_0\mathcal{D}a_1\mathcal{D}^{-1} - a_0a_1\mathcal{D}a_2\mathcal{D}^{-1} + a_0a_1a_2\right).$$

Expanding $(id + \lambda + \lambda^2)b'\phi_1(a_0, a_1, a_2)$, we recover previous expression. (ix) Consequence of (viii).

Scale-invariant term of the spectral action

We know from [8] that the scale-invariant term of the action can be written as

$$\zeta_{D_A}(0) - \zeta_D(0) = -\int A\mathcal{D}^{-1} + \frac{1}{2} \int A\mathcal{D}^{-1} A\mathcal{D}^{-1} - \frac{1}{3} \int A\mathcal{D}^{-1} A\mathcal{D}^{-1} A\mathcal{D}^{-1}.$$
 (5)

In fact, this action can be expressed in dimension 3 as contributions corresponding to tadpole and the Yang-Mills and Chern-Simons actions in dimension 4:

Proposition 2.3. For any one-form A,

$$\zeta_{D_A}(0) - \zeta_D(0) = -\frac{1}{2} \int_{N\phi_1} A + \frac{1}{2} \int_{\phi_2} (dA + A^2) - \frac{1}{2} \int_{\phi_3} (AdA + \frac{2}{3}A^3).$$
 (6)

To prove this, we calculate now each terms of the action.

Lemma 2.4. For any one-form A, we have

(i)
$$\int_{\phi_2} dA = \int_{B_0\phi_2} A = -\int_{\phi_1} A - \int_{\lambda\phi_1} A$$
.

$$(ii) \int A \mathcal{D}^{-1} = \int_{\phi_1} A = \frac{1}{2} \int_{N\phi_1} A - \frac{1}{2} \int_{\phi_2} dA.$$

$$(iii) \int A \mathcal{D}^{-1} A \mathcal{D}^{-1} = -\int_{\phi_3} A dA + \int_{\phi_2} A^2.$$

$$(iv) \int A \mathcal{D}^{-1} A \mathcal{D}^{-1} A \mathcal{D}^{-1} = \int_{\phi_3} A^3.$$

$$(iii) \int A \mathcal{D}^{-1} A \mathcal{D}^{-1} = - \int_{\partial \Omega} A dA + \int_{\partial \Omega} A^2.$$

$$(iv) \int A \mathcal{D}^{-1} A \mathcal{D}^{-1} A \mathcal{D}^{-1} = \int_{\phi_3} A^3$$

Proof. (i) and (ii) follow directly from Lemma 2.2 (v).

(iii) With the shorthand $A = a_i db_i$ (summation on i)

$$\begin{split} \int A \mathcal{D}^{-1} A \mathcal{D}^{-1} &= \int a_0 [\mathcal{D}, b_0] \mathcal{D}^{-1} a_1 [\mathcal{D}, b_1] \mathcal{D}^{-1} \\ &= - \int_{\phi_3} A dA + \int a_0 [\mathcal{D}, b_0] a_1 b_1 \mathcal{D}^{-1} - \int a_0 [\mathcal{D}, b_0] a_1 \mathcal{D}^{-1} b_1. \end{split}$$

We calculate further the remaining terms

$$\int a_0[\mathcal{D}, b_0] a_1 b_1 \mathcal{D}^{-1} - \int a_0[\mathcal{D}, b_0] a_1 \mathcal{D}^{-1} b_1 = \int a_0 \mathcal{D} b_0 a_1 b_1 \mathcal{D}^{-1} - \int a_0 b_0 \mathcal{D} a_1 b_1 \mathcal{D}^{-1} - \int a_0 b_0 \mathcal{D} a_1 \mathcal{D}^{-1} b_1 + \int a_0 b_0 \mathcal{D} a_1 \mathcal{D}^{-1} b_1,$$

which are compared with $\int_{\phi_2} A^2 = \int_{\phi_2} a_0(db_0)a_1db_1 = \int_{\phi_2} a_0d(b_0a_1)db_1 - a_0b_0da_1db_1$:

$$\int_{\phi_2} A^2 = \int a_0 [\mathcal{D}, b_0 a_1] \mathcal{D}^{-1} [\mathcal{D}, b_1] \mathcal{D}^{-1} - \int a_0 b_0 [\mathcal{D}, a_1] \mathcal{D}^{-1} [\mathcal{D}, b_1] \mathcal{D}^{-1}
= \int a_0 \mathcal{D} b_0 a_1 b_1 \mathcal{D}^{-1} - \int a_0 \mathcal{D} b_0 a_1 \mathcal{D}^{-1} b_1 - \int a_0 b_0 a_1 \mathcal{D} b_1 \mathcal{D}^{-1} + \int a_0 b_0 a_1 b_1
- \int a_0 b_0 \mathcal{D} a_1 b_1 \mathcal{D}^{-1} + \int a_0 b_0 \mathcal{D} a_1 \mathcal{D}^{-1} b_1 + \int a_0 b_0 a_1 \mathcal{D} b_1 \mathcal{D}^{-1} - \int a_0 b_0 a_1 b_1
= \int a_0 \mathcal{D} b_0 a_1 b_1 \mathcal{D}^{-1} - \int b_1 a_0 \mathcal{D} b_0 a_1 \mathcal{D}^{-1} - \int a_0 b_0 \mathcal{D} a_1 b_1 \mathcal{D}^{-1} + \int b_1 a_0 b_0 \mathcal{D} a_1 \mathcal{D}^{-1}.$$

(iv) Note that

$$\begin{split} \int_{\phi_3} A^3 &= \int_{\phi_3} a_0 (db_0) a_1 (db_1) a_2 db_2 = \int_{\phi_3} a_0 d(b_0 a_1) d(b_1 a_2) db_2 - a_0 b_0 da_1 d(b_1 a_2) db_2 \\ &- a_0 d(b_0 a_1 b_1) d(a_2 db_2 + a_0 b_0 d(a_1 b_1) da_2 db_2 \\ &= \int a_0 [\mathcal{D}, b_0 a_1] \mathcal{D}^{-1} [\mathcal{D}, b_1 a_2] \mathcal{D}^{-1} [\mathcal{D}, b_2] \mathcal{D}^{-1} - a_0 b_0 [\mathcal{D}, a_1] \mathcal{D}^{-1} [\mathcal{D}, b_1 a_2] \mathcal{D}^{-1} [\mathcal{D}, b_2] \mathcal{D}^{-1} \\ &- a_0 [\mathcal{D}, b_0 a_1 b_1] \mathcal{D}^{-1} [\mathcal{D}, a_2] \mathcal{D}^{-1} [\mathcal{D}, b_2] \mathcal{D}^{-1} + a_0 b_0 [\mathcal{D}, a_1 b_1] \mathcal{D}^{-1} [\mathcal{D}, a_2] \mathcal{D}^{-1} [\mathcal{D}, b_2] \mathcal{D}^{-1}. \end{split}$$

Summing up the first two terms and the last two ones gives

$$\int_{\phi_3} A^3 = \int a_0[\mathcal{D}, b_0] a_1 \mathcal{D}^{-1}[\mathcal{D}, b_1 a_2] \mathcal{D}^{-1}[\mathcal{D}, b_2] \mathcal{D}^{-1} - a_0[\mathcal{D}, b_0] a_1 b_1 \mathcal{D}^{-1}[\mathcal{D}, a_2] \mathcal{D}^{-1}[\mathcal{D}, b_2] \mathcal{D}^{-1}.$$

Using Remark 2.1, we can commute under the integral \mathcal{D}^{-1} with all $a \in \mathcal{A}$ and similarly

$$\int A \mathcal{D}^{-1} A \mathcal{D}^{-1} A \mathcal{D}^{-1} = \int a_0 [\mathcal{D}, b_0] a_1 \mathcal{D}^{-1} [\mathcal{D}, b_1] a_2 \mathcal{D}^{-1} [\mathcal{D}, b_2] \mathcal{D}^{-1}$$

which proves (iv).

We deduce Proposition 2.3 from (5) using the previous lemma.

3 The $SU_q(2)$ triple

3.1 The spectral triple

We briefly recall the main facts of the real spectral triple $(\mathcal{A}(SU_q(2)), \mathcal{H}, \mathcal{D})$ introduced in [21], see also [4, 5, 13].

The algebra:

Let $\mathcal{A} := \mathcal{A}(SU_q(2))$ be the *-algebra generated polynomially by a and b, subject to the following commutation rules with 0 < q < 1:

$$ba = q ab,$$
 $b^*a = q ab^*,$ $bb^* = b^*b,$ $a^*a + q^2 b^*b = 1,$ $aa^* + bb^* = 1.$ (7)

Lemma 3.1. For any representation π of A,

$$Spect(\pi(bb^*)) = \{0, q^{2k} \mid | k \in \mathbb{N} \} \text{ or } \pi(b) = 0,$$

 $Spect(\pi(aa^*)) = \{1, 1 - q^{2k} \mid k \in \mathbb{N} \} \text{ or } \pi(b) = 0 \text{ and } \pi(a) \text{ is a unitary.}$

Proof. [31] Since $\{0\} \cup \sigma(\pi(aa^*)) = \{0\} \cup \sigma(\pi(a^*a))$, we get

$$\{1\} \cup B = \{1\} \cup q^2 B \tag{8}$$

if $B:=\sigma\bigl(\pi(bb^*)\bigr)$. Since $0\leq\pi(bb^*)\leq 1$, so B is a closed subset of [0,1]. Assume $b\neq 0$. Let $s:=\sup(B)$ and suppose $s\neq 1$. Then $s=q^2x$ where $x\in B$. Thus $s=q^2x< x\leq s$ gives s=0 and the contradiction b=0, thus $1\in B$. Similar argument for $\inf(B)$ implies $0\in B$. Let $C:=\{0,q^{2k}\,|\,k\in\mathbb{N}\}\subset B$ and assume $B\setminus C\neq\emptyset$. Then $B\setminus C=(q^2B)\setminus C$ by (8) and this is equal to $q^2(B\setminus C)$ since $q^{-2}>1$. If $s:=\sup(B\setminus C)$, then $s=\lim_n(q^2x_n)$ where $x_n\in B\setminus C$ and $s=q^2\lim_n x_n\leq q^2s$ implying s=0. But $B\setminus C\subset\{0\}$ yields a contradiction, so $B\setminus C=\emptyset$. \square

This lemma is interesting since it shows the appearance of discreteness for $0 \le q < 1$ while for $q=1, SU_q(2)=SU(2)\simeq \mathbb{S}^3$ and the spectrum of the commuting operator $\pi(aa^*)$ and $\pi(bb^*)$ are equal to [0,1]. Moreover, all foregoing results on noncommutative integrals will involve q^2 and not q.

Any element of A can be uniquely decomposed as a linear combination of terms of the form $a^{\alpha}b^{\beta}b^{*\gamma}$ where $\alpha \in \mathbb{Z}, \beta, \gamma \in \mathbb{N}$, with the convention

$$a^{-|\alpha|} := a^{*|\alpha|}$$

The spinorial Hilbert space:

 $\mathcal{H} = \mathcal{H}^{\uparrow} \oplus \mathcal{H}^{\downarrow}$ has an orthonormal basis consisting of vectors $|j\mu n\uparrow\rangle$ with $j=0,\frac{1}{2},1,\ldots,$ $\mu=-j,\ldots,j$ and $n=-j^+,\ldots,j^+$, together with $|j\mu n\downarrow\rangle$ for $j=\frac{1}{2},1,\ldots,$ $\mu=-j,\ldots,j$ and $n=-j^-,\ldots,j^-$ (here $x^{\pm}:=x\pm\frac{1}{2}$).

It is convenient to use a vector notation, setting:

$$|j\mu n\rangle\rangle := \binom{|j\mu n\uparrow\rangle}{|j\mu n\downarrow\rangle} \tag{9}$$

and with the convention that the lower component is zero when $n = \pm (j + \frac{1}{2})$ or j = 0.

The representation π and its approximate π :

It is known that representation theory of $SU_q(2)$ is similar to that of SU(2) [39]. The representation π given in [21] is:

$$\pi(a) |j\mu n\rangle\rangle := \alpha_{j\mu n}^{+} |j^{+}\mu^{+}n^{+}\rangle\rangle + \alpha_{j\mu n}^{-} |j^{-}\mu^{+}n^{+}\rangle\rangle,$$

$$\pi(b) |j\mu n\rangle\rangle := \beta_{j\mu n}^{+} |j^{+}\mu^{+}n^{-}\rangle\rangle + \beta_{j\mu n}^{-} |j^{-}\mu^{+}n^{-}\rangle\rangle,$$

$$\pi(a^{*}) |j\mu n\rangle\rangle := \tilde{\alpha}_{j\mu n}^{+} |j^{+}\mu^{-}n^{-}\rangle\rangle + \tilde{\alpha}_{j\mu n}^{-} |j^{-}\mu^{-}n^{-}\rangle\rangle,$$

$$\pi(b^{*}) |j\mu n\rangle\rangle := \tilde{\beta}_{j\mu n}^{+} |j^{+}\mu^{-}n^{+}\rangle\rangle + \tilde{\beta}_{j\mu n}^{-} |j^{-}\mu^{-}n^{+}\rangle\rangle$$

$$(10)$$

where

$$\begin{split} \alpha_{j\mu n}^{+} &:= \sqrt{q^{\mu+n-1/2}[j+\mu+1]} \begin{pmatrix} q^{-j-1/2} \frac{\sqrt{[j+n+3/2]}}{[2j+2]} & 0 \\ q^{1/2} \frac{\sqrt{[j-n+1/2]}}{[2j+1][2j+2]} & q^{-j} \frac{\sqrt{[j+n+1/2]}}{[2j+1]} \end{pmatrix}, \\ \alpha_{j\mu n}^{-} &:= \sqrt{q^{\mu+n+1/2}[j-\mu]} \begin{pmatrix} q^{j+1} \frac{\sqrt{[j-n+1/2]}}{[2j+1]} & -q^{1/2} \frac{\sqrt{[j+n+1/2]}}{[2j][2j+1]} \\ 0 & q^{j+1/2} \frac{\sqrt{[j-n-1/2]}}{[2j]} \end{pmatrix}, \\ \beta_{j\mu n}^{+} &:= \sqrt{q^{\mu+n-1/2}[j+\mu+1]} \begin{pmatrix} \frac{\sqrt{[j-n+3/2]}}{[2j+2]} & 0 \\ -q^{-j-1} \frac{\sqrt{[j+n+1/2]}}{[2j+1][2j+2]} & q^{-1/2} \frac{\sqrt{[j-n+1/2]}}{[2j+1]} \\ -q^{-j} \frac{\sqrt{[j+n+1/2]}}{[2j+1]} & -q^{j} \frac{\sqrt{[j-n+1/2]}}{[2j][2j+1]} \\ 0 & -\frac{\sqrt{[j+n-1/2]}}{[2j]} \end{pmatrix}, \end{split}$$

with $\tilde{\alpha}_{j\mu n}^{\pm} := (\alpha_{j^{\pm}\mu^{-}n^{-}}^{\mp})^{*}$, $\tilde{\beta}_{j\mu n}^{\pm} := (\beta_{j^{\pm}\mu^{-}n^{+}}^{\mp})^{*}$ and with the q-number of $\alpha \in \mathbb{R}$ be defined as

$$[\alpha] := \frac{q^{\alpha} - q^{-\alpha}}{q - q^{-1}}.$$

For the purpose of this paper it is sufficient to use the approximate spinorial *-representation $\underline{\pi}$ of $SU_q(2)$ presented in [21,38] instead of the full spinorial one π . This approximate representation is

$$\underline{\pi}(a) := a_+ + a_-, \quad \underline{\pi}(b) := b_+ + b_-$$

with the following definitions:

$$a_{+} |j\mu n\rangle\rangle := q_{j^{+}+\mu^{+}} \begin{pmatrix} q_{j^{+}+n^{+}+1} & 0 \\ 0 & q_{j^{+}+n} \end{pmatrix} |j^{+}\mu^{+}n^{+}\rangle\rangle,$$

$$a_{-} |j\mu n\rangle\rangle := q^{2j^{+}\mu^{+}n^{+}\frac{1}{2}} \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix} |j^{-}\mu^{+}n^{+}\rangle\rangle,$$

$$b_{+} |j\mu n\rangle\rangle := q^{j^{+}n^{-}\frac{1}{2}} q_{j^{+}+\mu^{+}} \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix} |j^{+}\mu^{+}n^{-}\rangle\rangle,$$

$$b_{-} |j\mu n\rangle\rangle := -q^{j^{+}\mu} \begin{pmatrix} q_{j^{+}+n} & 0 \\ 0 & q_{j^{-}+n} \end{pmatrix} |j^{-}\mu^{+}n^{-}\rangle\rangle.$$
(11)

All disregarded terms are trace-class and do not influence residue calculations. More precisely, $\pi(x) - \underline{\pi}(x) \in \mathcal{K}_q$ where \mathcal{K}_q is the principal ideal generated by the operators

$$J_q |j\mu n\rangle\rangle := q^j |j\mu n\rangle\rangle. \tag{12}$$

Actually, \mathcal{K}_q is independent of q and is contained in all ideals of operators such that $\mu_n = o(n^{-\alpha})$ (infinitesimal of order α) for any $\alpha > 0$, and $\mathcal{K}_q \subset OP^{-\infty}$.

We define the alternative orthonormal basis $v_{m,l}^{j\uparrow}$ and $v_{m,l}^{j\downarrow}$ and the vector notation

$$v_{m,l}^{j} := \begin{pmatrix} v_{m,l}^{j\uparrow} \\ v^{j\downarrow} \end{pmatrix} \text{ where } v_{m,l}^{j\uparrow} := |j,m-j,l-j^{+},\uparrow\rangle, \quad v_{m,l}^{j\downarrow} := |j,m-j,l-j^{-},\downarrow\rangle.$$

Here $j \in \frac{1}{2}\mathbb{N}$, $0 \le m \le 2j$, $0 \le l \le 2j+1$ and $v_{m,l}^{\downarrow,j}$ is zero whenever j=0 or l=2j or 2j+1. The interest is that now, the operators a_{\pm} and b_{\pm} satisfy the simpler relations

$$a_{+} v_{m,l}^{j} = q_{m+1} q_{l+1} v_{m+1,l+1}^{j^{+}}, \quad a_{-} v_{m,l}^{j} = q^{m+l+1} v_{m,l}^{j^{-}}, b_{+} v_{m,l}^{j} = q^{l} q_{m+1} v_{m+1,l}^{j^{+}}, \quad b_{-} v_{m,l}^{j} = -q^{m} q_{l} v_{m,l-1}^{j^{-}}.$$

$$(13)$$

Thus

$$a_{+}^{*} v_{m,l}^{j} = q_{m} q_{l} v_{m-1,l-1}^{j-}, \qquad a_{-}^{*} v_{m,l}^{j} = q^{m+l+1} v_{m,l}^{j+}, b_{+}^{*} v_{m,l}^{j} = q^{l} q_{m} v_{m-1,l}^{j-}, \qquad b_{-}^{*} v_{m,l}^{j} = -q^{m} q_{l+1} v_{m,l+1}^{j+}.$$

$$(14)$$

Moreover, we have

$$a_{-}a_{+} = q^{2} a_{+} a_{-}, \quad b_{-}b_{+} = q^{2} b_{+}b_{-}, \qquad b_{+}a_{+} = q a_{+}b_{+}, \quad b_{-}a_{-} = q a_{-}b_{-},$$

$$a_{-}^{*}a_{+} = q^{2} a_{+}a_{-}^{*}, \quad a_{-}^{*}a_{-} = a_{-}a_{-}^{*}, \qquad a_{-}^{*}b_{+} = q b_{+}a_{-}^{*}, \quad a_{-}^{*}b_{-} = q b_{-}a_{-}^{*},$$

$$a_{+}^{*}a_{-} = q^{2} a_{-}a_{+}^{*}, \quad b_{-}^{*}b_{+} = b_{+}b_{-}^{*}, \qquad b_{-}^{*}a_{+} = q a_{+}b_{-}^{*}, \quad a_{-}b_{+} = q b_{+}a_{-}.$$

$$(15)$$

Note for instance that

$$a_{+}a_{+}^{*}v_{m,l}^{j} = q_{m}^{2}q_{l}^{2}v_{m,l}^{j}, \quad a_{+}^{*}a_{+}v_{m,l}^{j} = q_{m+1}^{2}q_{l+1}^{2}v_{m,l}^{j},$$

$$b_{+}b_{+}^{*}v_{m,l}^{j} = q^{2l}q_{m}^{2}v_{m,l}^{j}, \quad b_{+}^{*}b_{+}v_{m,l}^{j} = q^{2l}q_{m+1}^{2}v_{m,l}^{j},$$

so applied to $v_{m,l}^{j}$, we get the first relation (and similarly for the others)

$$a_{+}^{*}a_{+} - q^{2} a_{+}a_{+}^{*} + q^{2} (b_{+}^{*}b_{+} - b_{+}b_{+}^{*}) = 1 - q^{2},$$
 (16)

$$a_{+}a_{+}^{*} + a_{-}a_{-}^{*} + b_{+}b_{+}^{*} + b_{-}b_{-}^{*} = 1, (17)$$

$$a_{+}^{*}a_{+} + a_{-}^{*}a_{-} + q^{2}(b_{+}^{*}b_{+} + b_{-}^{*}b_{-}) = 1,$$

$$(18)$$

$$a_{-}^{*}a_{-} - q^{2} a_{-} a_{-}^{*} + q^{2} b_{-}^{*} b_{-} - q^{2} b_{-} b_{-}^{*} = 0,$$

$$(19)$$

$$a_{+}a_{-}^{*} + b_{-}^{*}b_{+} = 0,$$
 $a_{-}^{*}a_{+} + q^{2}b_{-}^{*}b_{+} = 0,$ (20)

$$a_{-}a_{+}^{*} + b_{+}^{*}b_{-} = 0,$$
 $a_{+}^{*}a_{-} + q^{2}b_{+}^{*}b_{-} = 0,$ (21)

$$b_{+}b_{+}^{*} - b_{+}^{*}b_{+} + b_{-}b_{-}^{*} - b_{-}^{*}b_{-} = 0, (22)$$

$$q a_{+}b_{-} - b_{-}a_{+} + q a_{-}b_{+} - b_{+}a_{-} = 0. (23)$$

And two others:

Note that we also use two other infinite dimensional *-representations π_{\pm} of \mathcal{A} on $\ell^2(\mathbb{N})$ defined as follows on the orthonormal basis $\{\varepsilon_n : n \in \mathbb{N}\}$ of $\ell^2(\mathbb{N})$ by

$$\pi_{\pm}(a)\,\varepsilon_n := q_{n+1}\,\varepsilon_{n+1}, \qquad \pi_{\pm}(b)\,\varepsilon_n := \pm q^n\,\varepsilon_n,$$

$$q_n := \sqrt{1 - q^{2n}}.$$
(24)

These representations are irreducible but not faithful since for instance $\pi_{\pm}(b-b^*)=0$.

The Dirac operator:

It is chosen the same as in the classical case of a 3-sphere with the round metric:

$$\mathcal{D}|j\mu n\rangle\rangle := \begin{pmatrix} 2j + \frac{3}{2} & 0\\ 0 & -2j - \frac{1}{2} \end{pmatrix}|j\mu n\rangle\rangle, \tag{25}$$

which means, with our convention, that $\mathcal{D} v_{ml}^j = \begin{pmatrix} 2j + \frac{3}{2} & 0 \\ 0 & -2j - \frac{1}{2} \end{pmatrix} v_{ml}^j$. Note that this operator is asymptotically diagonal with linear spectrum and

the eigenvalues $2j + \frac{1}{2}$ for $j \in \frac{1}{2}\mathbb{N}$, have multiplicities (2j+1)(2j+2), the eigenvalues $-(2j+\frac{1}{2})$ for $j \in \frac{1}{2}\mathbb{N}^*$, have multiplicities 2j(2j+1).

So this Dirac operator coincide exactly with the classical one on the 3-sphere (see [1,32]) with a gap around 0.

Let $\mathcal{D} = F|\mathcal{D}|$ be the polar decomposition of \mathcal{D} , thus

$$|\mathcal{D}||j\mu n\rangle\rangle = \begin{pmatrix} d_{j+} & 0\\ 0 & d_{j} \end{pmatrix}|j\mu n\rangle\rangle, \quad d_{j} := 2j + \frac{1}{2},$$
 (26)

$$F|j\mu n\rangle\rangle = \begin{pmatrix} 1 & 0\\ 0 & -1 \end{pmatrix}|j\mu n\rangle\rangle, \tag{27}$$

and it follows from (11) and (27) that

$$F \text{ commutes with } a_{\pm}, b_{\pm}.$$
 (28)

The reality operator:

This antilinear operator J is defined on the basis of \mathcal{H} by

$$J|j,\mu,n,\uparrow\rangle := i^{2(2j+\mu+n)}|j,-\mu,-n,\uparrow\rangle, \qquad J|j,\mu,n,\downarrow\rangle := i^{2(2j-\mu-n)}|j,-\mu,-n,\downarrow\rangle \tag{29}$$

thus it satisfies

$$J^{-1} = -J = J^* \text{ and } \mathcal{D}J = J\mathcal{D},$$

 $J v_{m,l}^{j\uparrow} = i^{2(m+l)-1} v_{2j-m,2j+1-l}^{j\uparrow}, \qquad J v_{m,l}^{j\downarrow} = i^{-2(m+l)+1} v_{2j-m,2j-1-l}^{j\downarrow}.$

The Hopf map r

For the explicit calculations of residues, we need a *-homomorphism $r: X \to \pi_+(A) \otimes \pi_-(A)$ defined by the tensor product in the sense of Hopf algebras of representations π_+ and π_- :

$$r(a_{+}) := \pi_{+}(a) \otimes \pi_{-}(a), \qquad r(a_{-}) := -q \,\pi_{+}(b) \otimes \pi_{-}(b^{*}),$$

$$r(b_{+}) := -\pi_{+}(a) \otimes \pi_{-}(b), \qquad r(b_{-}) := -\pi_{+}(b) \otimes \pi_{-}(a^{*}). \tag{30}$$

In fact, \mathcal{A} is a Hopf *-algebra under the coproduct $\Delta(a) := a \otimes a - q b \otimes b^*$, $\Delta(b) := a \otimes b + b \otimes a^*$. These homomorphisms appeared in [39] with the translation $\alpha \leftrightarrow a^*$, $\gamma \leftrightarrow -b$. In particular, if $U := \begin{pmatrix} a & b \\ -ab^* & a^* \end{pmatrix}$ is the canonical generator of the $K_1(\mathcal{A})$ -group $(\Delta a, \Delta b) = (a, b) \otimes U$ where the last $\dot{\otimes}$ means the matrix product of tensors of components.

The grading:

According to the shift $j \to j^{\pm}$ appearing in formulae (13), (14), we get a \mathbb{Z} -grading on X defined by the degree +1 on a_+, b_+, a_-^*, b_-^* and -1 on a_-, b_-, a_+^*, b_+^* .

Any operator $T \in X$ can be (uniquely) decomposed as $T = \sum_{i \in J \subset \mathbb{Z}} T_i$ where T_i is homogeneous of degree j.

For $T \in X$, T° will denote the 0-degree part of T for this grading and by a slight abuse of notations, we write $r(T)^{\circ}$ instead of $r(T^{\circ})$.

The symbol map:

We also use the *-homomorphism $\sigma: \pi_{\pm}(\mathcal{A}) \to C^{\infty}(S^1)$ defined for $z \in S^1$ on the generators by

$$\sigma\big(\pi_{\pm}(a)\big)(z):=z,\quad \sigma\big(\pi_{\pm}(a^*)\big)(z):=\bar{z},\quad \sigma\big(\pi_{\pm}(b)\big)(z)=\sigma\big(\pi_{\pm}(b^*)\big)(z):=0.$$

The application $(\sigma \otimes \sigma) \circ r$ is defined on X (and so on \mathcal{B}) with values in $C^{\infty}(S^1) \otimes C^{\infty}(S^1)$.

We define

$$dT := [\mathcal{D}, T]$$
 and $\delta(T) := [|\mathcal{D}|, T]$.

Lemma 3.2. a_{\pm} , b_{\pm} are bounded operators on \mathcal{H} such that for all $p \in \mathbb{N}$,

(i)
$$\delta(a_{\pm}) = \pm a_{\pm}, \quad \delta(b_{\pm}) = \pm b_{\pm},$$

(i)
$$\delta(a_{\pm}) = \pm a_{\pm}$$
, $\delta(b_{\pm}) = \pm b_{\pm}$,
(ii) $\delta^{p}(\underline{\pi}(a)) = a_{+} + (-1)^{p}a_{-}$, $\delta^{p}(\underline{\pi}(b)) = b_{+} + (-1)^{p}b_{-}$,
(iii) $\delta(a_{\pm}^{p}) = \pm p \, a_{\pm}^{p}$, $\delta(b_{\pm}^{p}) = \pm p \, b_{\pm}^{p}$.

(iii)
$$\delta(a_{\perp}^{p}) = \pm p a_{\perp}^{p}$$
, $\delta(b_{\perp}^{p}) = \pm p b_{\perp}^{p}$

Proof. (i) By definition, $a_{\pm} |j\mu n\rangle\rangle = \begin{pmatrix} \alpha_{\pm} & 0 \\ 0 & \beta_{\pm} \end{pmatrix} |j^{\pm}\mu^{+}n^{+}\rangle\rangle$ where the numbers α_{\pm} and β_{\pm} depend on j, μ , n and q, so we get by (26)

$$\delta(a_{\pm})|j\mu n\rangle\rangle = \begin{pmatrix} (d_{j^{+\pm}})\alpha_{\pm} & 0\\ 0 & d_{j^{\pm}}\beta_{\pm} \end{pmatrix}|j^{\pm}\mu^{+}n^{+}\rangle\rangle - \begin{pmatrix} (d_{j^{+}})\alpha_{\pm} & 0\\ 0 & d_{j}\beta_{\pm} \end{pmatrix}|j^{\pm}\mu^{+}n^{+}\rangle\rangle$$
$$= \begin{pmatrix} \pm \alpha_{\pm} & 0\\ 0 & +\beta_{\pm} \end{pmatrix}|j^{\pm}\mu^{+}n^{+}\rangle\rangle = \pm a_{\pm}|j\mu n\rangle\rangle$$

and similar proofs for b_+ .

(ii) and (iii) are straightforward consequences of (i) and definition of $\underline{\pi}$.

We note

 \mathcal{B} the *-subalgebra of $\mathcal{B}(\mathcal{H})$ generated by the operators in $\delta^k(\pi(\mathcal{A}))$ for all $k \in \mathbb{N}$,

 $\Psi_0^0(\mathcal{A})$ the algebra generated by $\delta^k(\pi(\mathcal{A}))$ and $\delta^k([\mathcal{D},\pi(\mathcal{A})])$ for all $k\in\mathbb{N}$,

X the *-subalgebra of $\mathcal{B}(\mathcal{H})$ algebraically generated by the set $\{a_{\pm}, b_{\pm}\}.$

Remark 3.3. By Lemma 3.2, we see that, modulo $OP^{-\infty}$, X is equal to \mathcal{B} and in particular contains $\pi(\mathcal{A})$.

Using (28), we get that $\mathcal{B} \subset \Psi_0^0(\mathcal{A}) \subset algebra generated by \mathcal{B}$ and $\mathcal{B}F$.

Note that, despite the last inclusion, F is not a priori in $\Psi_0^0(A)$.

3.2 The noncommutative integrals

Recall that for any pseudodifferential operator $T, f T := \underset{s=0}{\operatorname{Res}} \zeta_{\mathcal{D}}^{T}(s)$ where $\zeta_{\mathcal{D}}^{T}(s) := \operatorname{Tr}(T|\mathcal{D}|^{-s})$.

Theorem 3.4. The dimension spectrum (without reality structure given by J) of the spectral triple $(A(SU_q(2)), \mathcal{H}, D)$ is simple and equal to $\{1, 2, 3\}$. Moreover, the corresponding residues for $T \in \mathcal{B}$ are

$$\int T|\mathcal{D}|^{-3} = 2(\tau_1 \otimes \tau_1) (r(T)^{\circ}),$$

$$\int T|\mathcal{D}|^{-2} = 2(\tau_1 \otimes \tau_0 + \tau_0 \otimes \tau_1) (r(T)^{\circ}),$$

$$\int T|\mathcal{D}|^{-1} = (2 \tau_0 \otimes \tau_0 - \frac{1}{2} \tau_1 \otimes \tau_1) (r(T)^{\circ}),$$

$$\int FT|\mathcal{D}|^{-3} = 0,$$

$$\int FT|\mathcal{D}|^{-2} = 0,$$

$$\int FT|\mathcal{D}|^{-1} = (\tau_0 \otimes \tau_1 - \tau_1 \otimes \tau_0) (r(T)^{\circ}),$$

where the functionals τ_0 , τ_1 are defined for $x \in \pi_{\pm}(A)$ by

$$\tau_0(x) := \lim_{N \to \infty} \left(\operatorname{Tr}_N x - (N+1) \tau_1(x) \right), \qquad \tau_1(x) := \frac{1}{2\pi} \int_0^{2\pi} \sigma(x) (e^{i\theta}) d\theta,$$

with $\operatorname{Tr}_N x = \sum_{n=0}^N \langle \varepsilon_n, x \, \varepsilon_n \rangle$.

Proof. Consequence of [38, Theorem 4.1 and (4.3)].

Remark 3.5. Since F is not in \mathcal{B} , the equation of Theorem 3.4 are not valid for all $T \in \Psi_0^0(\mathcal{A})$. But when $T \in \Psi_0^0(\mathcal{A})$, $f T |\mathcal{D}|^{-k} = 0$ for $k \notin \{1, 2, 3\}$ since the dimension spectrum is $\{1, 2, 3\}$ [38].

Compared to [38] where we had

$$\tau_0^{\uparrow}(x) := \lim_{N \to \infty} \operatorname{Tr}_N x - (N + \frac{3}{2}) \, \tau_1(x), \quad \tau_0^{\downarrow}(x) := \lim_{N \to \infty} \operatorname{Tr}_N x - (N + \frac{1}{2}) \, \tau_1(x),$$

we replaced them with τ_0 :

$$\tau_0^{\uparrow} = \tau_0 - \frac{1}{2}\tau_1, \quad \tau_0^{\downarrow} = \tau_0 + \frac{1}{2}\tau_1.$$

Note that τ_1 is a trace on $\pi_{\pm}(\mathcal{A})$ such that $\tau_1(1) = 1$, while τ_0 is not since $\tau_0(1) = 0$ and

$$\tau_0(\pi_{\pm}(aa^*)) = \lim_{N \to \infty} \sum_{n=0}^{\infty} (1 - q^{2n}) - (N+1) = -\frac{1}{1-q^2},\tag{31}$$

so, because of the shift, the replacement $a \leftrightarrow a^*$ gives

$$\tau_0(\pi_{\pm}(a^*a)) = q^2 \,\tau_0(\pi_{\pm}(aa^*)). \tag{32}$$

3.3 The tadpole

Lemma 3.6. For $SU_q(2)$, the condition of the vanishing tadpole (see [15]) is not satisfied.

Proof. For example, an explicit calculation gives $\int \pi(b)[\mathcal{D}, \pi(b^*)]\mathcal{D}^{-1} = \frac{2}{1-q^2}$: Let $x, y \in \underline{\pi}(\mathcal{A})$. Since [F, x] = 0, we have

$$\int x[\mathcal{D}, y] D^{-1} = \int x \delta(y) |\mathcal{D}|^{-1} = \tau' \left(r(x \delta(y))^{0} \right)$$

where $\tau' := 2 \tau_0 \otimes \tau_0 - \frac{1}{2} \tau_1 \otimes \tau_1$.

By Lemma 3.2, $\underline{\pi}(b)\delta(\underline{\pi}(b^*)) = (b_+ + b_-)((b_-)^* - (b_+)^*) = -b_+b_+^* + b_-b_-^* + b_+b_-^* - b_-b_+^*$. Since only the first two terms have degree 0, we get, using the formulae from Theorem 3.4

$$\tau'(r(-b_{+}b_{+}^{*})) = -\tau'(\pi_{+}(aa^{*}) \otimes \pi_{-}(bb^{*}))$$
$$= -2\tau_{0}(\pi_{+}(aa^{*}))\tau_{0}(\pi_{-}(bb^{*})) + \frac{1}{2}\tau_{1}(\pi_{+}(aa^{*}))\tau_{1}(\pi_{-}(bb^{*}))$$

and $\tau_1(\pi_+(aa^*)) = \frac{1}{2\pi} \int_0^{2\pi} 1 d\theta = 1$, $\tau_1(\pi_-(bb^*)) = 0$. Similarly, using (32)

$$\tau'(r(-b_-b_-^*)) = 2\tau_0(\pi_+(bb^*))\tau_0(\pi_-(a^*a)) = 2q^2\tau_0(\pi_-(aa^*))\tau_0(\pi_+(bb^*).$$

Since $\tau_0(\pi_{\pm}(bb^*)) = \text{Tr}(\pi_{\pm}(bb^*)) = \sum_{n=0}^{\infty} q^{2n} = \frac{1}{1-q^2}$ and (31),

$$\oint \pi(b)[\mathcal{D}, \pi(b^*)] \mathcal{D}^{-1} = 2\frac{1}{1-q^2} \frac{1}{1-q^2} + 2q^2 \frac{1}{1-q^2} \frac{1}{1-q^2} = \frac{2}{1-q^2}.$$

In particular the pairing of the tadpole cyclic cocycle ϕ_1 with the generator of K_1 -group is nontrivial:

Remark 3.7. Other examples: with the shortcut x instead of $\underline{\pi}(x)$,

$$(\tau_{1} \otimes \tau_{1}) r (a\delta(a^{*})^{\circ}) = -1,$$

$$(\tau_{1} \otimes \tau_{1}) r (a^{*}\delta(a)^{\circ}) = 1,$$

$$(\tau_{0} \otimes \tau_{0}) r (a\delta(a^{*})^{\circ}) = \frac{1}{q^{2}-1},$$

$$(\tau_{0} \otimes \tau_{0}) r (a^{*}\delta(a)^{\circ}) = \frac{q^{2}}{q^{2}-1},$$

$$(\tau_{0} \otimes \tau_{0}) r (a^{*}\delta(a)^{\circ}) = \frac{q^{2}}{q^{2}-1},$$

$$\int a\delta(a^{*})|\mathcal{D}|^{-1} = \frac{q^{2}+3}{2(q^{2}-1)},$$

$$\int b^{*}\delta(a)|\mathcal{D}|^{-1} = \frac{3q^{2}+1}{2(q^{2}-1)},$$

$$\int b^{*}\delta(b^{*})|\mathcal{D}|^{-1} = 0,$$

$$\int b^{*}\delta(b^{*})|\mathcal{D}|^{-1} = \frac{-2}{q^{2}-1}.$$

In particular, $N\Phi_1$ does not vanish on 1-forms since $\int_{N\Phi_1} ada^* = N\Phi_1(a, a^*) = -1$.

Let U be the canonical generator of the $K_1(\mathcal{A})$ -group, $U = \begin{pmatrix} a \\ -qb^* \end{pmatrix}$ acting on $\mathcal{H} \otimes \mathbb{C}^2$. Then for $A_U := \sum_{k,l=1}^2 \pi(U_{kl}) d\pi(U^*_{kl})$, using above remark, $\int_{\phi_1} A_U = -2$ as obtained in [38, page 391]: in fact, with $P := \frac{1}{2}(1+F)$,

$$\psi_1(U, U^*) := 2\sum_{k,l} \int U_{kl} \delta(U_{kl}^*) P|\mathcal{D}|^{-1} - \int U_{kl} \delta^2(U_{kl}^*) P|\mathcal{D}|^{-2} + \frac{2}{3} \int U_{kl} \delta^3(U_{kl}^*) P|\mathcal{D}|^{-3}$$

satisfies $\psi_1(U, U^*) = 2 \sum_{k,l} \int U_{kl} \delta(U_{kl}^*) P |\mathcal{D}|^{-1} = \int_{\phi_1} A_U.$

4 Reality operator and spectral action on $SU_q(2)$

4.1 Spectral action in dimension 3 with $[F, A] \in OP^{-\infty}$

Let $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ a be real spectral triple of dimension 3. Assume that $[F, \mathcal{A}] \in OP^{-\infty}$, where $F := \mathcal{D}|\mathcal{D}|^{-1}$ (we suppose \mathcal{D} invertible). Let \mathbb{A} be a selfadjoint one form, so \mathbb{A} is of the form $\mathbb{A} = \sum_i a_i db_i$ where $a_i, b_i \in \mathcal{A}$.

Thus, $\mathbb{A} \simeq AF \mod OP^{-\infty}$ where $A := \sum_i a_i \delta(b_i)$ is the δ -one-form associated to \mathbb{A} . Note that A and F commute modulo $OP^{-\infty}$.

We define

$$D_{\mathbb{A}} := \mathcal{D}_{\mathbb{A}} + P_{\mathbb{A}}, \quad P_{\mathbb{A}} \text{ the projection on } \operatorname{Ker} \mathcal{D}_{\mathbb{A}},$$

 $\mathcal{D}_{\mathbb{A}} := \mathcal{D} + \widetilde{\mathbb{A}}, \quad \widetilde{\mathbb{A}} := \mathbb{A} + J \mathbb{A} J^{-1}.$

Theorem 4.1. The coefficients of the full spectral action (with reality operator) on any real spectral triple $(A, \mathcal{H}, \mathcal{D})$ of dimension 3 such that $[F, A] \in OP^{-\infty}$ are

(i)
$$\int |D_{\mathbb{A}}|^{-3} = \int |\mathcal{D}|^{-3}.$$

(ii)
$$\int |D_{\mathbb{A}}|^{-2} = \int |\mathcal{D}|^{-2} - 4 \int A|\mathcal{D}|^{-3}.$$

$$(iii) \qquad \int |D_{\mathbb{A}}|^{-1} = \int |\mathcal{D}|^{-1} - 2 \int A|\mathcal{D}|^{-2} + 2 \int A^2 |\mathcal{D}|^{-3} + 2 \int AJAJ^{-1} |\mathcal{D}|^{-3}.$$

$$(iv) \qquad \zeta_{D_{\mathbb{A}}}(0) = \zeta_{D}(0) - 2 \int A|\mathcal{D}|^{-1} + \int A(A + JAJ^{-1})|\mathcal{D}|^{-2} + \int \delta(A)(A + JAJ^{-1})|\mathcal{D}|^{-3} - \frac{2}{3} \int A^{3}|\mathcal{D}|^{-3} - 2 \int A^{2}JAJ^{-1}|\mathcal{D}|^{-3}.$$

Proof. (i) We apply [22, Proposition 4.9].

(ii) By [22, Lemma 4.10 (i)], we have $\int |D_{\mathbb{A}}|^{-2} = \int |\mathcal{D}|^{-2} - \int (\widetilde{\mathbb{A}}\mathcal{D} + \mathcal{D}\widetilde{\mathbb{A}} + \widetilde{\mathbb{A}}^2)|\mathcal{D}|^{-4}$. By the trace property of the noncommutative integral and the fact that $\widetilde{\mathbb{A}}^2|\mathcal{D}|^{-4}$ is trace-class, we get $\int |D_{\mathbb{A}}|^{-2} = \int |\mathcal{D}|^{-2} - 2\int \widetilde{\mathbb{A}}\mathcal{D}|\mathcal{D}|^{-4} = \int |\mathcal{D}|^{-2} - 4\int \mathcal{A}\mathcal{D}|\mathcal{D}|^{-4}$. Since $\mathcal{A}\mathcal{D} \sim A|\mathcal{D}| \mod OP^{-\infty}$, we get the result.

(iii) By [22, Lemma 4.10 (ii)], we have

$$\int |D_{\mathbb{A}}|^{-1} = \int |\mathcal{D}|^{-1} - \frac{1}{2} \int (\widetilde{\mathbb{A}}\mathcal{D} + \mathcal{D}\widetilde{\mathbb{A}} + \widetilde{\mathbb{A}}^2) |\mathcal{D}|^{-3} + \frac{3}{8} \int (\widetilde{\mathbb{A}}\mathcal{D} + \mathcal{D}\widetilde{\mathbb{A}} + \widetilde{\mathbb{A}}^2)^2 |\mathcal{D}|^{-5}.$$

Following arguments of (ii), we get

$$\int (\widetilde{\mathbb{A}}\mathcal{D} + \mathcal{D}\widetilde{\mathbb{A}} + \widetilde{\mathbb{A}}^2)|\mathcal{D}|^{-3} = 4 \int A|\mathcal{D}|^{-2} + 2 \int A^2|\mathcal{D}|^{-3} + 2 \int AJAJ^{-1}|\mathcal{D}|^{-3},$$

$$\int (\widetilde{\mathbb{A}}\mathcal{D} + \mathcal{D}\widetilde{\mathbb{A}} + \widetilde{\mathbb{A}}^2)^2|\mathcal{D}|^{-5} = 8 \int A^2|\mathcal{D}|^{-3} + 8 \int AJAJ^{-1}|\mathcal{D}|^{-3},$$

and the result follows.

(iv) By [22, Lemma 4.5] gives $\zeta_{D_{\mathbb{A}}}(0) = \sum_{j=1}^{3} \frac{(-1)^{j}}{j} f(\widetilde{\mathbb{A}}\mathcal{D}^{-1})^{j}$. Moreover, we have $f\widetilde{\mathbb{A}}\mathcal{D}^{-1} = 2fA|\mathcal{D}|^{-1}$ and $f(\widetilde{\mathbb{A}}\mathcal{D}^{-1})^{2} = 2f(A|\mathcal{D}|^{-1})^{2} + 2fA|\mathcal{D}|^{-1}JAJ^{-1}|\mathcal{D}|^{-1}$. Since $\delta(A) \in OP^{0}$, we can check that $f(A|\mathcal{D}|^{-1})^{2} = fA^{2}|\mathcal{D}|^{-2} + f\delta(A)A|\mathcal{D}|^{-3}$ and, with the same argument, that $fA|\mathcal{D}|^{-1}JAJ^{-1}|\mathcal{D}|^{-1} = fAJAJ^{-1}|\mathcal{D}|^{-2} + f\delta(A)JAJ^{-1}|\mathcal{D}|^{-3}$. Thus, we get

$$f(\widetilde{\mathbb{A}}\mathcal{D}^{-1})^2 = 2f A(A + JAJ^{-1})|\mathcal{D}|^{-2} + 2f \delta(A)(A + JAJ^{-1})|\mathcal{D}|^{-3}.$$
 (33)

The third term to be computed is

$$f(\widetilde{\mathbb{A}}\mathcal{D}^{-1})^3 = 2f(A|\mathcal{D}|^{-1})^3 + 4f(A|\mathcal{D}|^{-1})^2JAJ^{-1}|\mathcal{D}|^{-1} + 2fA|\mathcal{D}|^{-1}JAJ^{-1}|\mathcal{D}|^{-1}A|\mathcal{D}|^{-1}.$$

Any operator in OP^{-4} being trace-class here, we get

$$f(\widetilde{\mathbb{A}}\mathcal{D}^{-1})^3 = 2f A^3 |\mathcal{D}|^{-3} + 4f A^2 J A J^{-1} |\mathcal{D}|^{-3} + 2f A J A J^{-1} A |\mathcal{D}|^{-3}.$$
 (34)

Since $\int AJAJ^{-1}A|\mathcal{D}|^{-3} = \int A^2JAJ^{-1}|\mathcal{D}|^{-3}$ by trace property and the fact that $\delta(A) \in OP^0$, the result follows then from (33) and (34).

Corollary 4.2. For the spectral action of \mathbb{A} without the reality operator (i.e. $\mathcal{D}_{\mathbb{A}} = \mathcal{D} + \mathbb{A}$), we get

$$\begin{split}
& \int |D_{\mathbb{A}}|^{-2} = \int |\mathcal{D}|^{-2} - 2 \int A|\mathcal{D}|^{-3}, \\
& \int |D_{\mathbb{A}}|^{-1} = \int |\mathcal{D}|^{-1} - \int A|\mathcal{D}|^{-2} + \int A^{2}|\mathcal{D}|^{-3}, \\
& \zeta_{D_{\mathbb{A}}}(0) = \zeta_{\mathcal{D}}(0) - \int A|\mathcal{D}|^{-1} + \frac{1}{2} \int A^{2}|\mathcal{D}|^{-2} + \frac{1}{2} \int \delta(A)A|\mathcal{D}|^{-3} - \frac{1}{3} \int A^{3}|\mathcal{D}|^{-3}.
\end{split}$$

Spectral action on $SU_q(2)$: main result

On $SU_q(2)$, since F commutes with a_{\pm} and b_{\pm} , the previous lemma can be used for the spectral action computation.

Here is the main result of this section

Theorem 4.3. In the full spectral action (4) (with the reality operator) of $SU_q(2)$ for a one-form

A and A its associated δ -one-form, the coefficients are:

$$\begin{split} & \int |D_{\mathbb{A}}|^{-3} = 2 \,, \\ & \int |D_{\mathbb{A}}|^{-2} = -4 \int A|\mathcal{D}|^{-3}, \\ & \int |D_{\mathbb{A}}|^{-1} = -\frac{1}{2} + 2 \left(\int A^{2}|\mathcal{D}|^{-3} - \int A|\mathcal{D}|^{-2} \right) + \left| \int A|\mathcal{D}|^{-3} \right|^{2}, \\ & \zeta_{D_{\mathbb{A}}}(0) = -2 \int A|\mathcal{D}|^{-1} + \int A^{2}|\mathcal{D}|^{-2} - \frac{2}{3} \int A^{3}|\mathcal{D}|^{-3} \\ & + \overline{\int A|\mathcal{D}|^{-3}} \left(\frac{1}{2} \int A|\mathcal{D}|^{-2} - \int A^{2}|\mathcal{D}|^{-3} \right) + \frac{1}{2} \int A|\mathcal{D}|^{-3} \overline{\int A|\mathcal{D}|^{-2}} \,. \end{split}$$

In order to prove this theorem, we will use a decomposition of one-forms in the Poincaré-Birkhoff–Witt basis of \mathcal{A} with an extension of previous representations to operators like $TJT'J^{-1}$ where T and T' are in X.

4.3 Balanced components and Poincaré–Birkhoff–Witt basis of A

Our objective is to compute all integrals in term of A and the computation will lead to functions of A which capture certain symmetries on A.

For convenience, let us introduce now these functions:

Let $\mathbb{A} = \sum_i \pi(x^i) d\pi(y^i)$ on $SU_q(2)$ be one-form and A the associated δ -one-form. The x^i and y^i are in \mathcal{A} and as such they can be uniquely written as finite sums $x^i = \sum_{\alpha} x^i_{\alpha} m^{\alpha}$ and $y^i = \sum_{\beta} y^i_{\beta} m^{\beta}$ where $m^{\alpha} := a^{\alpha_1} b^{\alpha_2} b^{*\alpha_3}$ is the canonical monomial of \mathcal{A} with $\alpha, \beta \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N}$ based on a fixed Poincaré–Birkhoff–Witt type basis of \mathcal{A} .

Remark 4.4. Any one-form $\mathbb{A} = \sum_i \pi(x^i) d\pi(y^i)$ on $SU_q(2)$ is characterized by a complex valued matrix $A^{\beta}_{\alpha} = \sum_i x^i_{\alpha} y^i_{\beta}$ where $\alpha, \beta \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N}$. This matrix is such that

$$A = A_{\alpha}^{\beta} M_{\beta}^{\alpha}$$

where $M^{\alpha}_{\beta} := \pi(m^{\alpha})\delta(\pi(m^{\beta}))$. In the following, we note

$$\bar{A} := \bar{A}^{\beta}_{\alpha} M^{\alpha}_{\beta}$$

so for any $p \in \mathbb{N}$, $\int \overline{A}|\mathcal{D}|^{-p} = \overline{\int A|\mathcal{D}|^{-p}}$.

This presentation of one-forms is not unique modulo $OP^{-\infty}$ since, as we will see in section 5, $F = \sum_i x_i dy_i$ where $x_i, y_i \in \mathcal{A}$, thus for any generator z, $[F, z] = \sum_i x_i d(y_i z) - x_i y_i dz - z x_i dy_i = 0$ mod $OP^{-\infty}$. We do not know however if this presentation is unique when the $OP^{-\infty}$ part is taken into account.

The δ -one-forms M^{α}_{β} are said to be *canonical*. Any product of n canonical δ -one forms, where $n \in \mathbb{N}^*$, is called a *canonical* δ^n -one-form. Thus, if A is a δ -one-form, $A^n = (A^n)^{\bar{\beta}}_{\bar{\alpha}} M^{\bar{\alpha}}_{\bar{\beta}}$ where $\bar{\alpha} = (\alpha, \alpha', \cdots, \alpha^{(n-1)}), \ \bar{\beta} = (\beta, \beta', \cdots, \beta^{(n-1)})$ are in $\mathbb{Z}^n \times \mathbb{N}^n \times \mathbb{N}^n$, $(A^n)^{\bar{\beta}}_{\bar{\alpha}} := A^{\beta}_{\alpha} \cdots A^{\beta^{(n-1)}}_{\alpha^{(n-1)}}$ and $M^{\bar{\alpha}}_{\bar{\beta}}$ is the canonical δ^n -one form equals to $M^{\alpha}_{\beta} \cdots M^{\alpha^{(n-1)}}_{\beta^{(n-1)}}$.

Definition 4.5. A canonical δ^n -one-form is a-balanced if it is of the form

$$a^{\alpha_1}\delta(a^{\beta_1})\cdots a^{\alpha_1^{(n-1)}}\delta(a^{\beta_1^{(n-1)}})$$

where $\sum_{i=0}^{n-1} \alpha_1^{(i)} + \beta_1^{(i)} = 0$.

For any δ -one-form A, the a-balanced components of A^n are noted $B_a(A^n)_{\bar{\alpha}}^{\bar{\beta}}$.

Note that

$$B_a(A)_{\bar{\alpha}}^{\bar{\beta}} = A_{-\beta_100}^{\beta_100} \, \delta_{\alpha_1 + \beta_1, 0} \, \delta_{\alpha_2 + \alpha_3 + \beta_2 + \beta_3, 0}.$$

Definition 4.6. A canonical δ^n -one-form is balanced if it is of the form

$$m^{\alpha}\delta(m^{\beta})\cdots m^{\alpha^{(n-1)}}\delta(m^{\beta^{(n-1)}})$$

where $\sum_{i=0}^{n-1} \alpha_1^{(i)} + \beta_1^{(i)} = 0$ and $\sum_{i=0}^{n-1} \alpha_2^{(i)} + \beta_2^{(i)} = \sum_{i=0}^{n-1} \alpha_3^{(i)} + \beta_3^{(i)}$. For any δ -one-form A, the balanced components of A^n are noted $B(A^n)_{\bar{\alpha}}^{\bar{\beta}}$.

Note that

$$B(A)_{\bar{\alpha}}^{\bar{\beta}} = A_{-\beta_1\alpha_2\alpha_3}^{\beta_1\beta_2\beta_3} \, \delta_{\alpha_1+\beta_1,0} \, \delta_{\alpha_2+\beta_2,\alpha_3+\beta_3}.$$

As we will show, a contribution to the k^{th} -coefficient in the spectral action, is only brought by one-forms \mathbb{A} such that A^k is balanced (and even a-balanced in the case k=1). Note also that if A is balanced, then A^k for $k \geq 1$ is also balanced, whereas the converse is false.

4.4 The reality operator J on $SU_q(2)$

For any $n, p \in \mathbb{N}$,

$$q_n := \sqrt{1 - q^{2n}},$$
 $q_{-n} := 0 \text{ if } n > 0,$ $q_{n,p}^{\uparrow} := q_{n+1} \cdots q_{n+p},$ $q_{n,p}^{\downarrow} := q_n \cdots q_{n-(p-1)},$

with the convention $q_{n,0}^{\uparrow} = q_{n,0}^{\downarrow} := 1$. Thus, we have the relations

$$\pi_{\pm}(a^p) \ \varepsilon_n = q_{n,p}^{\uparrow} \ \varepsilon_{n+p} , \qquad \qquad \pi_{\pm}(a^{*p}) \ \varepsilon_n = q_{n,p}^{\downarrow} \ \varepsilon_{n-p} ,$$

$$\pi_{\pm}(b^p) \ \varepsilon_n = (\pm q^n)^p \ \varepsilon_n , \qquad \qquad \pi_{\pm}(b^{*p}) \ \varepsilon_n = (\pm q^n)^p \ \varepsilon_n ,$$

where $\varepsilon_k := 0$ if k < 0.

The sign of $x \in \mathbb{R}$ is noted η_x . By convention, $a_j := a$, $a_{\pm,j} := a_{\pm}$ if $j \geq 0$ and $a_j := a^*$, $a_{\pm,j} := a^*_{\pm}$ if j < 0. Note that, with convention

$$q_{n,p}^{\uparrow_{\alpha_1}} := q_{n,p}^{\uparrow} \text{ if } \alpha_1 > 0, \quad q_{n,p}^{\uparrow_{\alpha_1}} := q_{n,p}^{\downarrow} \text{ if } \alpha_1 < 0, \text{ and } q_{n,p}^{\uparrow_0} := 1,$$

we have for any $\alpha_1 \in \mathbb{Z}$ and $p \leq \alpha_1$, $\pi_{\pm}(a_{\alpha_1}^p) \varepsilon_n = q_{n,p}^{\uparrow_{\alpha_1}} \varepsilon_{n+\eta_{\alpha_1}p}$. Recall that the reality operator J is defined by

$$J\,v_{m,l}^{j\uparrow} = i^{2(m+l)-1}v_{2j-m,2j+1-l}^{j\uparrow}\,, \qquad J\,v_{m,l}^{j\downarrow} = i^{-2(m+l)+1}v_{2j-m,2j-1-l}^{j\downarrow}\,,$$

thus the real conjugate operators

$$\widehat{a}_{\pm} := J a_{\pm} J^{-1}, \quad \widehat{b}_{\pm} := J b_{\pm} J^{-1}$$

satisfy

$$\begin{split} \widehat{a}_{+} \, v_{m,l}^{j} &:= -q_{2j+1-m} \left(\begin{smallmatrix} q_{2j+2-l} & 0 \\ 0 & q_{2j-l} \end{smallmatrix} \right) v_{m,l}^{j^{+}}, \qquad \widehat{a}_{-} \, v_{m,l}^{j} &:= -q^{2j-m} \left(\begin{smallmatrix} q^{2j+2-l} & 0 \\ 0 & q^{2j-l} \end{smallmatrix} \right) v_{m-1,l-1}^{j^{-}}, \\ \widehat{b}_{+} \, v_{m,l}^{j} &:= q_{2j+1-m} \left(\begin{smallmatrix} q^{2j+1-l} & 0 \\ 0 & q^{2j-1-l} \end{smallmatrix} \right) v_{m,l+1}^{j^{+}}, \qquad \widehat{b}_{-} \, v_{m,l}^{j} &:= -q^{2j-m} \left(\begin{smallmatrix} q_{2j+1-l} & 0 \\ 0 & q_{2j-1-l} \end{smallmatrix} \right) v_{m-1,l}^{j^{-}}. \end{split}$$

So the real conjugate operator behave differently on the up and down part of the Hilbert space. The difference comes from the fact that the index l is not treated uniformly by J on up and down parts.

We note \widehat{X} the algebra generated by $\{\widehat{a}_{\pm}, \widehat{b}_{\pm}\}$, \widetilde{X} the algebra generated by $\{a_{\pm}, b_{\pm}, \widehat{a}_{\pm}, \widehat{b}_{\pm}\}$ and $\mathcal{H}' := \ell^2(\mathbb{N}) \otimes \ell^2(\mathbb{Z})$ and we construct two *-representations $\widehat{\pi}_{\pm}$ of \mathcal{A} :

The representation $\widehat{\pi}_+$ gives bounded operators on \mathcal{H}' while $\widehat{\pi}_-$ represents \mathcal{A} into $\mathcal{B}(\mathcal{H}'\otimes\mathbb{C}^2)$. The representation $\hat{\pi}_{+}$ is defined on the generators by:

$$\widehat{\pi}_{+}(a)\,\varepsilon_{m}\otimes\varepsilon_{2j}:=q_{2j+1-m}\,\varepsilon_{m}\otimes\varepsilon_{2j+1},\qquad \widehat{\pi}_{+}(b)\,\varepsilon_{m}\otimes\varepsilon_{2j}:=-q^{2j-m}\,\varepsilon_{m+1}\otimes\varepsilon_{2j+1}$$

while $\widehat{\pi}_{-}$ is defined by:

$$\widehat{\pi}_{-}(a)\,\varepsilon_{l}\otimes\varepsilon_{2j}\otimes\varepsilon_{\uparrow\downarrow}:=-q_{2j+1\pm1-l}\,\varepsilon_{l}\otimes\varepsilon_{2j+1}\otimes\varepsilon_{\uparrow\downarrow}\,,$$

$$\widehat{\pi}_{-}(b)\,\varepsilon_{l}\otimes\varepsilon_{2j}\otimes\varepsilon_{\uparrow\downarrow}:=-q^{2j\pm1-l}\,\varepsilon_{l+1}\otimes\varepsilon_{2j+1}\otimes\varepsilon_{\uparrow\downarrow}\,,$$

where $\varepsilon_{\uparrow\downarrow}$ is the canonical basis of \mathbb{C}^2 and the + in \pm corresponds to \uparrow in $\uparrow\downarrow$.

The link between $\hat{\pi}_{\pm}$ and π_{\pm} which explains the notations about these intermediate objects and the fact that $\hat{\pi}_{\pm}$ are representations on different Hilbert spaces, is in the parallel between equations (30), (35) and (36).

Let us give immediately a few properties $(x_{\beta} \text{ equals } x \text{ if the sign } \beta \text{ is positive and equals } x^*$ otherwise)

$$\widehat{\pi}_{+}(a_{\beta})^{p} \, \varepsilon_{m} \otimes \varepsilon_{2j} = q_{2j-m,p}^{\uparrow_{\beta}} \, \varepsilon_{m} \otimes \varepsilon_{2j+\eta_{\beta}p} \,,$$

$$\widehat{\pi}_{-}(a_{\beta})^{p} \, \varepsilon_{l} \otimes \varepsilon_{2j} \otimes \varepsilon_{\uparrow\downarrow} = (-1)^{p} \, q_{2j\pm1-l,p}^{\uparrow_{\beta}} \, \varepsilon_{l} \otimes \varepsilon_{2j+\eta_{\beta}p} \otimes \varepsilon_{\uparrow\downarrow} \,,$$

$$\widehat{\pi}_{+}(b_{\beta})^{p} \, \varepsilon_{m} \otimes \varepsilon_{2j} = (-1)^{p} \, q^{(2j-m)p} \, \varepsilon_{m+\eta_{\beta}p} \otimes \varepsilon_{2j+\eta_{\beta}p} \,,$$

$$\widehat{\pi}_{-}(b_{\beta})^{p} \, \varepsilon_{l} \otimes \varepsilon_{2j} \otimes \varepsilon_{\uparrow\downarrow} = (-1)^{p} \, q^{(2j\pm1-l)p} \, \varepsilon_{l+\eta_{\beta}p} \otimes \varepsilon_{2j+\eta_{\beta}p} \otimes \varepsilon_{\uparrow\downarrow} \,.$$

Note that the $\hat{\pi}_+$ representations still contain the shift information, contrary to representations π_{\pm} . Moreover, $\widehat{\pi}_{\pm}(b) \neq \widehat{\pi}_{\pm}(b^*)$ while $\pi_{\pm}(b) = \pi_{\pm}(b^*)$. The operators $\widehat{a}_{\pm}, \widehat{b}_{\pm}$ are coded on $\mathcal{H}' \otimes \mathcal{H}' \otimes \mathbb{C}^2$ as the correspondence

$$\widehat{a}_{+} \longleftrightarrow \widehat{\pi}_{+}(a) \otimes \widehat{\pi}_{-}(a), \qquad \widehat{a}_{-} \longleftrightarrow -q \,\widehat{\pi}_{+}(b^{*}) \otimes \widehat{\pi}_{-}(b^{*}),
\widehat{b}_{+} \longleftrightarrow -\widehat{\pi}_{+}(a) \otimes \widehat{\pi}_{-}(b), \qquad \widehat{b}_{-} \longleftrightarrow -\widehat{\pi}_{+}(b^{*}) \otimes \widehat{\pi}_{-}(a^{*}). \tag{35}$$

We now set the following extension to $\mathcal{B}(\mathcal{H}')$ of π_+ and to $\mathcal{B}(\mathcal{H}' \otimes \mathbb{C}^2)$ of π_- by

$$\pi'_{+}(a) := \pi_{+}(a) \otimes V, \quad \pi'_{+}(b) := \pi_{+}(b) \otimes V \quad \text{(V is the shift of } \ell^{2}(\mathbb{Z})),$$

 $\pi'_{-}(a) := \pi_{-}(a) \otimes V \otimes 1_{2}, \quad \pi'_{-}(b) := \pi_{-}(b) \otimes V \otimes 1_{2}.$

So, we can define a canonical algebra morphism $\tilde{\rho}$ from \tilde{X} into the bounded operators on $\mathcal{H}' \otimes \mathcal{H}' \otimes \mathbb{C}^2$. This morphism is defined on the generators part $\{\widehat{a}_{\pm}, \widehat{b}_{\pm}\}$ of X by preceding correspondence and on the generators part $\{a_{\pm}, b_{\pm}\}$ by -see (30):

$$a_{+} \longleftrightarrow \pi'_{+}(a) \otimes \pi'_{-}(a), \qquad a_{-} \longleftrightarrow -q \, \pi'_{+}(b^{*}) \otimes \pi'_{-}(b^{*}), b_{+} \longleftrightarrow -\pi'_{+}(a) \otimes \pi'_{-}(b), \qquad b_{-} \longleftrightarrow -\pi'_{+}(b^{*}) \otimes \pi'_{-}(a^{*}).$$
(36)

We note S the canonical surjection from $\mathcal{H}' \otimes \mathcal{H}' \otimes \mathbb{C}^2$ onto \mathcal{H} . This surjection is associated to the parameters restrictions on m, j, l, j'. In particular, the index j' associated to the second $\ell^2(\mathbb{N})$ in $\mathcal{H}' \otimes \mathcal{H}' \otimes \mathbb{C}^2$ is set to be equal to j. Any vector in $\mathcal{H}' \otimes \mathcal{H}' \otimes \mathbb{C}^2$ not satisfying these restrictions is sent to 0 in \mathcal{H} .

Denote by I the canonical injection of \mathcal{H} into $\mathcal{H}' \otimes \mathcal{H}' \otimes \mathbb{C}^2$ (the index j is doubled). Thus, $S\widetilde{\rho}(\cdot)I$ is the identity on X.

In the computation of residues of $\zeta_{\mathcal{D}}^T$ functions, we can therefore replace the operator T by

We now extend τ_0 on $\pi'_{\pm}(\mathcal{A})\widehat{\pi}_{\pm}(\mathcal{A})$: For $x, y \in \mathcal{A}$, we set

$$\operatorname{Tr}_{N}\left(\pi'_{+}(x)\widehat{\pi}_{+}(y)\right) := \sum_{m=0}^{N} \left\langle \varepsilon_{m} \otimes \varepsilon_{N}, \, \pi'_{+}(x)\widehat{\pi}_{+}(y) \, \varepsilon_{m} \otimes \varepsilon_{N} \right\rangle,$$

$$\operatorname{Tr}_{N}^{\uparrow}\left(\pi'_{-}(x)\widehat{\pi}_{-}(y)\right) := \sum_{l=0}^{N} \left\langle \varepsilon_{l} \otimes \varepsilon_{N-1} \otimes \varepsilon_{\uparrow}, \, \pi'_{-}(x)\widehat{\pi}_{-}(y) \, \varepsilon_{l} \otimes \varepsilon_{N-1} \otimes \varepsilon_{\uparrow} \right\rangle,$$

$$\operatorname{Tr}_{N}^{\downarrow}\left(\pi'_{-}(x)\widehat{\pi}_{-}(y)\right) := \sum_{l=0}^{N} \left\langle \varepsilon_{l} \otimes \varepsilon_{N+1} \otimes \varepsilon_{\downarrow}, \, \pi'_{-}(x)\widehat{\pi}_{-}(y) \, \varepsilon_{l} \otimes \varepsilon_{N+1} \otimes \varepsilon_{\downarrow} \right\rangle.$$

Actually, a computation on monomials of \mathcal{A} shows that $\operatorname{Tr}_{N}^{\downarrow}\left(\pi'_{-}(x)\widehat{\pi}_{-}(y)\right) = \operatorname{Tr}_{N}^{\uparrow}\left(\pi'_{-}(x)\widehat{\pi}_{-}(y)\right)$. For convenience, we shall note $\operatorname{Tr}_N\left(\pi'_-(x)\widehat{\pi}_-(y)\right)$ this functional.

Lemma 4.7. Let $x, y \in A$. Then,

(i) $\tau_0(\pi'_+(x)\widehat{\pi}_{\pm}(y)) := \lim_{N\to\infty} U_N$ exists where

$$U_N := \operatorname{Tr}_N \left(\pi'_{\pm}(x) \widehat{\pi}_{\pm}(y) \right) - (N+1) \tau_1 \left(\pi_{\pm}(x) \right) \tau_1 \left(\pi_{\pm}(y) \right).$$

(ii)
$$U_N = \tau_0 \left(\pi'_{\pm}(x) \widehat{\pi}_{\pm}(y) \right) + \mathcal{O}(N^{-k})$$
 for all $k > 0$.

Proof. (i) We can suppose that x and y are monomials, since the result will follow by linearity. We will give a proof for the case of the π_+ representations, the case π_- being similar, with minor changes.

We have $\widehat{\pi}_+(y) = (\widehat{\pi}_+ a_{\beta_1})^{|\beta_1|} (\widehat{\pi}_+ b)^{\beta_2} (\widehat{\pi}_+ b^*)^{\beta_3}$. A computation gives

$$\widehat{\pi}_{+}(y)\,\varepsilon_{m}\otimes\varepsilon_{2j}=(-1)^{\beta_{2}+\beta_{3}}\,q^{(2j-m)(\beta_{2}+\beta_{3})}\,q_{2j-m,|\beta_{1}|}^{\uparrow_{\beta_{1}}}\,\varepsilon_{m-\beta_{3}+\beta_{2}}\otimes\varepsilon_{2j-\beta_{3}+\beta_{2}+\beta_{1}}$$

and with the notation $t_{2j,m} := \langle \varepsilon_m \otimes \varepsilon_{2j}, \pi'_{\pm}(x) \widehat{\pi}_{\pm}(y) \varepsilon_m \otimes \varepsilon_{2j} \rangle$ and $T_{2j} := \sum_{m=0}^{2j} t_{2j,m}$, we get

$$\begin{split} t_{2j,m} &= (-1)^{\beta_2 + \beta_3} \, q^{(2j-m)(\beta_2 + \beta_3)} \, q_{2j-m,|\beta_1|}^{\uparrow_{\beta_1}} \, q_{m-\beta_3 + \beta_2,|\alpha_1|}^{\uparrow_{\alpha_1}} \, q^{(m+\beta_2 - \beta_3)(\alpha_2 + \alpha_3)} \, \delta_{\alpha_1 + \beta_2 - \beta_3,0} \\ & \times \delta_{-\alpha_3 + \alpha_2 + \beta_1,0} \\ &= (-1)^{\alpha_1} \, q^{(2j-m)(\beta_2 + \beta_3) + (m-\alpha_1)(\alpha_2 + \alpha_3)} \, q_{2j-m,|\beta_1|}^{\uparrow_{\beta_1}} \, q_{m-\alpha_1,|\alpha_1|}^{\uparrow_{\alpha_1}} \, \delta_{\alpha_1 + \beta_2 - \beta_3,0} \, \delta_{\alpha_2 - \alpha_3 + \beta_1,0} \\ &=: f_{\alpha,\beta} \, q^{2j\lambda} \, t_{2j,m}' =: f_{\alpha,\beta} \, q^{2j\kappa} \, t_{2j,2j-m}'' \end{split}$$

where

$$t'_{2j,m} := q^{m(\kappa - \lambda)} q_{2j-m,|\beta_1|}^{\uparrow_{\beta_1}} q_{m-\alpha_1,|\alpha_1|}^{\uparrow_{\alpha_1}},$$

$$t''_{2j,m} := q^{m(\lambda - \kappa)} q_{m,|\beta_1|}^{\uparrow_{\beta_1}} q_{2j-m-\alpha_1,|\alpha_1|}^{\uparrow_{\alpha_1}},$$
(38)

$$t_{2j,m}'' := q^{m(\lambda - \kappa)} q_{m,|\beta_1|}^{\uparrow_{\beta_1}} q_{2j-m-\alpha_1,|\alpha_1|}^{\uparrow_{\alpha_1}}, \tag{38}$$

with $\lambda := \beta_2 + \beta_3 \ge 0$ and $\kappa := \alpha_2 + \alpha_3 \ge 0$. We will now prove that if $\lambda \ne \kappa$, then (T_{2j}) is a convergent sequence. Suppose $\kappa > \lambda$. Let us note $U'_{2j} := \sum_{m=0}^{2j} t'_{2j,m}$. Since the $t'_{2j,m}$ are positive and $t'_{2j+1,m} \ge t'_{2j,m}$ for all j, m, U'_{2j} is an increasing real sequence. The estimate

$$U'_{2j} \le \sum_{m=0}^{2j} q^{m(\kappa - \lambda)} \le \frac{1}{1 - q^{\kappa - \lambda}} < \infty$$

proves then that U'_{2j} is a convergent sequence. With $T_{2j} = f_{\alpha,\beta} \, q^{2j\lambda} \, U'_{2j}$, we obtain our result. Suppose now that $\lambda > \kappa$. Let us note $U''_{2j} := \sum_{m=0}^{2j} t''_{2j,m}$. Since the $t''_{2j,m}$ are positive and $t''_{2j+1,m} \ge t''_{2j,m}$ for all j,m,U''_{2j} is an increasing real sequence. The estimate

$$U_{2j}'' \le \sum_{m=0}^{2j} q^{m(\lambda - \kappa)} \le \frac{1}{1 - q^{\lambda - \kappa}} < \infty$$

proves then that U_{2j}'' is a convergent sequence. With $T_{2j} = f_{\alpha,\beta} q^{2j\kappa} U_{2j}''$, we have again our result. Moreover, note that if λ and κ are both different from zero, the limit of (T_{2j}) is zero and more precisely,

$$T_{2j} = \mathcal{O}(q^{2j\lambda}) \text{ if } \kappa > \lambda > 0,$$
 (39)

$$T_{2j} = \mathcal{O}(q^{2j\kappa}) \text{ if } \lambda > \kappa > 0.$$
 (40)

Suppose now that $\lambda = \kappa \neq 0$. In that case, (T_{2j}) also converges rapidly to zero. Indeed, let us fix $q < \varepsilon < 1$. we have $\varepsilon^{-2j\lambda} T_{2j} = \sum_{m=0}^{2j} c_m d_{2j-m} = c * d(2j)$ where $c_m := f_{\alpha,\beta} (q/\varepsilon)^{\lambda m} q_{m-\alpha_1,|\alpha_1|}^{\uparrow_{\alpha_1}}$ and $d_m := (q/\varepsilon)^{\lambda m} q_{m,|\beta_1|}^{\uparrow_{\beta_1}}$. Since both $\sum_m c_m$ and $\sum_m d_m$ are absolutely convergent series, their Cauchy product $\sum_{2j} \varepsilon^{-2j\lambda} T_{2j}$ is convergent. In particular, $\lim_{j\to\infty} \varepsilon^{-2j\lambda} T_{2j} = 0$, and

$$T_{2j} = \mathcal{O}(\varepsilon^{2j\lambda}). \tag{41}$$

Finally, T_{2j} has a finite limit in all cases except possibly when $\lambda = \kappa = 0$, which is the case when $\alpha_1 = \alpha_2 = \alpha_3 = \beta_1 = \beta_2 = \beta_3 = 0$. In that case, $t_{2j,m} = 1$.

A straightforward computation gives $\tau_1(\pi_{\pm}(x))$ $\tau_1(\pi_{\pm}(y)) = \delta_{\alpha_1,0} \, \delta_{\beta_1,0} \, \delta_{\alpha_2,0} \, \delta_{\beta_2,0} \, \delta_{\alpha_3,0} \, \delta_{\beta_3,0}$. Thus,

$$U_{2j} = T_{2j} - (2j+1)\delta_{\alpha_1,0}\,\delta_{\beta_1,0}\,\delta_{\alpha_2,0}\,\delta_{\beta_2,0}\,\delta_{\alpha_3,0}\,\delta_{\beta_3,0}$$

has always a finite limit when $j \to \infty$.

(ii) The result is clear if $\lambda = \kappa = 0$ (in that case $U_N = \tau_0 = 0$). Suppose λ or κ is not zero. In that case $U_{2j} = T_{2j}$. By (40), (39) and (41), we see that if $\lambda > \kappa > 0$ or $\kappa > \lambda > 0$ or $\kappa = \lambda$, (T_{2j}) converges to 0 with a rate in $\mathcal{O}(\varepsilon^{2j\alpha})$ where $\alpha > 0$ and $q \leq \varepsilon < 1$. Thus, it only remains to check the cases $(\kappa > 0, \lambda = 0)$ and $(\kappa = 0, \lambda > 0)$. In the first one, we get from (37), $U_{2j} = f_{\alpha,\beta} \sum_{m=0}^{2j} q^{m\kappa} q_{2j-m,|\beta_1|}^{\beta_1}$. If $\beta_1 = 0$, we are done.

Suppose $\beta_1 > 0$. We have $q_{2j-m,|\beta_1|}^{\uparrow_{\beta_1}} = \sum_{p=0}^{\infty} l_p \, q^{r_p} \, q^{2|p|_1(2j-m)}$ where $p = (p_1, \dots, p_{\beta_1})$ and $l_p = (-1)^{|p|_1} \left(\frac{1}{p}\right), \ r_p := 2p_1 + \dots + 2\beta_1 p_{\beta_1}$. Thus, cutting the sum in two, we get, noting $L_{2j} := f_{\alpha,\beta} \sum_{m=0}^{2j} q^{m\kappa}$,

$$U_{2j} - L_{2j} = f_{\alpha,\beta} \sum_{|p|_1 > \kappa/2} l_p \, q^{r_p} \, \frac{q^{4|p|_1 j} - q^{(2j+1)\kappa - 2|p|_1}}{1 - q^{\kappa - 2|p|_1}} + f_{\alpha,\beta} \sum_{0 \neq |p|_1 \leq \kappa/2} l_p \, q^{r_p} \, q^{4|p|_1 j} \sum_{m=0}^{2j} q^{m(\kappa - 2|p|_1)}.$$

Since $\sum_{0\neq |p|_1\leq \kappa/2} l_p q^{r_p} q^{4|p|_1 j} \sum_{m=0}^{2j} q^{m(\kappa-2|p|_1)}$ is in $\mathcal{O}_{j\to\infty}(jq^{4j})$, we have, modulo a rapidly decreasing sequence,

$$U_{2j} - L_{2j} \sim f_{\alpha,\beta} \sum_{|p|_1 > \kappa/2} l_p \, q^{r_p} \, \frac{q^{4|p|_1 j} - q^{(2j+1)\kappa - 2|p|_1}}{1 - q^{\kappa - 2|p|_1}} =: f_{\alpha,\beta} q^{2\kappa j} V_{2j}$$

with

$$V_{2j} = \sum_{|p|_1 > \kappa/2} l_p \, q^{r_p} \, \frac{1 - q^{(2|p|_1 - \kappa)(2j+1)}}{1 - q^{2|p|_1 - \kappa}} = \sum_{|p|_1 > \kappa/2} \sum_{m=0}^{2j} l_p \, q^{r_p} \, q^{(2|p|_1 - \kappa)m}.$$

The family $v_{m,p}:=(l_p\,q^{r_p}\,q^{(2|p|_1-\kappa)m})_{(p,m)\in I}$, where $I=\{(p,m)\in\mathbb{N}^{\beta_1}\times\mathbb{N}:|p|_1>\kappa/2\}$ is (absolutely) summable. Indeed $|v_{m,p}|\leq |l_p|q^{r_p}\,q^m$ so $|v_{m,p}|$ is summable as the product of two summable families. As a consequence, $\lim_{j\to\infty}V_{2j}$ exists and is finite, which proves that $(q^{2\kappa j}V_{2j})$, and thus $(U_{2j}-L_{2j})$ converge rapidly to 0.

Suppose now that $\beta_1 < 0$. In that case, $q_{2j-m,|\beta_1|}^{\uparrow_{\beta_1}} = q_{2j-m,|\beta_1|}^{\downarrow} = q_{2j-m,|\beta_1|}^{\uparrow} = q_{2j-(m+|\beta_1|),|\beta_1|}^{\uparrow}$ and by (37), we get $U_{2j} = f_{\alpha,\beta} \sum_{m=0}^{2j} q^{m\kappa} q_{2j-(m+|\beta_1|),|\beta_1|}^{\uparrow} = f_{\alpha,\beta} q^{-|\beta_1|\kappa} \sum_{m=|\beta_1|}^{2j+|\beta_1|} q^{m\kappa} q_{2j-m,|\beta_1|}^{\uparrow}$, so the same arguments as in case $\beta_1 > 0$ apply here, the summation on m simply shifted of $|\beta_1|$.

The same proof can be applied for the other case ($\kappa = 0, \lambda > 0$). This time, we only need to use (38) instead of (37) and the preceding arguments follow by replacing κ by λ and β_1 by α_1 .

Remark 4.8. Contrary to the preceding τ_0 , the new functional contains the shift information. In particular, it filters the parts of nonzero degree.

If $T \in X\widehat{X}$, $\widetilde{\rho}(T) \in \pi_{+}(\mathcal{A})\widehat{\pi}_{+}(\mathcal{A}) \otimes \pi_{-}(\mathcal{A})\widehat{\pi}_{-}(\mathcal{A})$. For notational convenience, we define τ_{1} on $\pi'_{+}(\mathcal{A})\widehat{\pi}_{\pm}(\mathcal{A})$ as

$$\tau_1(\pi'_{\pm}(x)\widehat{\pi}_{\pm}(y)) := \tau_1(\pi_{\pm}(x)) \tau_1(\pi_{\pm}(y)).$$

In the following, the symbol \sim_e means equals modulo a entire function.

Theorem 4.9. Let $T \in X\widehat{X}$. Then

(i)
$$\zeta_{\mathcal{D}}^{T}(s) \sim_{e} 2(\tau_{1} \otimes \tau_{1}) \left(\widetilde{\rho}(T)\right) \zeta(s-2) + 2(\tau_{0} \otimes \tau_{1} + \tau_{1} \otimes \tau_{0}) \left(\widetilde{\rho}(T)\right) \zeta(s-1) + 2(\tau_{0} \otimes \tau_{0} - \frac{1}{2}\tau_{1} \otimes \tau_{1}) \left(\widetilde{\rho}(T)\right) \zeta(s),$$

(ii)
$$\int T|\mathcal{D}|^{-3} = 2(\tau_1 \otimes \tau_1)(\widetilde{\rho}(T)),$$

(iii)
$$\int T|\mathcal{D}|^{-2} = 2(\tau_0 \otimes \tau_1 + \tau_1 \otimes \tau_0) \left(\widetilde{\rho}(T)\right),$$

$$(iv) \quad \int T|\mathcal{D}|^{-1} = 2(\tau_0 \otimes \tau_0 - \frac{1}{2}\tau_1 \otimes \tau_1) \left(\widetilde{\rho}(T)\right).$$

Proof. (i) Since $T \in X\widehat{X}$, $\widetilde{\rho}(T)$ is a linear combination of terms like $\pi'_{+}(x)\widehat{\pi}_{+}(y) \otimes \pi'_{-}(z)\widehat{\pi}_{-}(t)$, where $x, y, z, t \in \mathcal{A}$. Such a term is noted in the following $T_{+} \otimes T_{-}$. Linear combination of these term is implicit. With the shortcut $T_{c_{1}, \dots, c_{p}} := \langle \varepsilon_{c_{1}} \otimes \dots \otimes \varepsilon_{c_{p}}, T \varepsilon_{c_{1}} \otimes \dots \otimes \varepsilon_{c_{p}} \rangle$, recalling that

 $v_{m,l}^{j,\downarrow}$ is 0 when j=0, or $l\geq 2j$, we get

$$\zeta_{\mathcal{D}}^{T}(s) = \sum_{2j=0}^{\infty} \sum_{m=0}^{2j} \sum_{l=0}^{2j+1} \left\langle {v_{m,l}^{i,\uparrow} \choose 0}, S\widetilde{\rho}(T) I {v_{m,l}^{i,\uparrow} \choose 0} \right\rangle d_{j+}^{-s} + \left\langle {0 \choose v_{m,l}^{j,\downarrow}}, S\widetilde{\rho}(T) I {0 \choose v_{m,l}^{j,\downarrow}} \right\rangle d_{j}^{-s}
= \sum_{2j=0}^{\infty} \sum_{m=0}^{2j} \sum_{l=0}^{2j+1} \widetilde{\rho}(T)_{m,2j,l,2j,\uparrow} d_{j+}^{-s} + \sum_{2j=1}^{\infty} \sum_{m=0}^{2j} \sum_{l=0}^{2j-1} \widetilde{\rho}(T)_{m,2j,l,2j,\downarrow} d_{j}^{-s}
= \sum_{2j=0}^{\infty} \left(\operatorname{Tr}_{2j}(T_{+}) \operatorname{Tr}_{2j+1}^{\uparrow}(T_{-}) + \operatorname{Tr}_{2j+1}(T_{+}) \operatorname{Tr}_{2j}^{\downarrow}(T_{-}) \right) d_{j+}^{-s}.$$

By Lemma 4.7 (ii), for all k > 0,

$$\operatorname{Tr}_{2j}(T_{\pm}) = (2j + \frac{3}{2})\tau_1(T_{\pm}) + \tau_0(T_{\pm}) - \frac{1}{2}\tau_1(T_{\pm}) + \mathcal{O}((2j)^{-k}),$$

$$\operatorname{Tr}_{2j+1}(T_{\pm}) = (2j + \frac{3}{2})\tau_1(T_{\pm}) + \tau_0(T_{\pm}) + \frac{1}{2}\tau_1(T_{\pm}) + \mathcal{O}((2j)^{-k}).$$

The result follows by noting that the difference of the Hurwitz zeta function $\zeta(s, \frac{3}{2})$ and Riemann zeta function $\zeta(s)$ is an entire function.

$$(ii, iii, iv)$$
 are direct consequences of (i) .

4.5 The smooth algebra $C^{\infty}(SU_a(2))$

In [13, 38], the smooth algebra $C^{\infty}(SU_q(2))$ is defined by pulling back the smooth structure $C^{\infty}(D_{q^{\pm}}^2)$ into the C^* -algebra generated by \mathcal{A} , through the morphism ρ and the application λ (the compression which gives an operator on \mathcal{H} from an operator on $l^2(\mathbb{N}) \otimes l^2(\mathbb{N}) \otimes l^2(\mathbb{Z}) \otimes \mathbb{C}^2$). The important point is that with [13, Lemma 2, p. 69], this algebra is stable by holomorphic calculus. By defining $\rho := \tilde{\rho} \circ c$ and $\lambda(\cdot) := S(\cdot)I$, the same lemma (with same notation) can be applied to our setting, with $c := \pi(x) \mapsto \underline{\pi}(x)$ and

$$\mathcal{C}:=C^{\infty}(D^2_{q^+})\otimes C^{\infty}(S^1)\otimes C^{\infty}(D^2_{q^+})\otimes C^{\infty}(S^1)\otimes \mathcal{M}_2(\mathbb{C})$$

as algebra stable by holomorphic calculus containing the image of $\widetilde{\rho}$. Here, we use Schwartz sequences to define the smooth structures. We finally obtain $C^{\infty}(SU_q(2))$ with real structure as a subalgebra stable by holomorphic calculus of the C^* -algebra generated by $\pi(\mathcal{A}) \cup J\pi(\mathcal{A})J^{-1}$ and containing $\pi(\mathcal{A}) \cup J\pi(\mathcal{A})J^{-1}$.

Corollary 4.10. The dimension spectrum of the real spectral triple $(C^{\infty}(SU_q(2)), \mathcal{H}, D)$ is simple and given by $\{1, 2, 3\}$. Its KO-dimension is 3.

Proof. Since F commutes with $\underline{\pi}(\mathcal{A})$, the pseudodifferential operators of order 0 (without the real structure and in the sense of [22]) are exactly (modulo $OP^{-\infty}$) the operators in $\mathcal{B} + \mathcal{B}F$. From Theorem 3.4 we see that the dimension spectrum of $SU_q(2)$ without taking into account the reality operator J is $\{1,2,3\}$, in other words, the possible poles of $\zeta_D^b: s \mapsto \operatorname{Tr}(bF^\varepsilon |\mathcal{D}|^{-s})$ (with $\varepsilon \in \{0,1\}, b \in \mathcal{B}$) are in $\{1,2,3\}$. Theorem 4.9 (i) shows that the possible poles are still $\{1,2,3\}$ when we take into account the real structure of $SU_q(2)$, that is to say, when \mathcal{B} is enlarged to $\mathcal{B}J\mathcal{B}J^{-1}$. Indeed, any element of $\mathcal{B}J\mathcal{B}J^{-1}$ is in $X\widehat{X}$ and it is clear from the preceding proof that adding F in the previous zeta function do not add any pole to $\{1,2,3\}$. All arguments goes true from the polynomial algebra $\mathcal{A}(SU_q(2))$ to the smooth pre-C*-algebra $C^\infty(SU_q(2))$.

KO-dimension refers just to $J^2 = -1$ and $\mathcal{D}J = J\mathcal{D}$ since there is no chirality because spectral dimension is 3.

4.6 Noncommutative integrals with reality operator and one-forms on $SU_q(2)$

The goal of this section is to obtain the following suppression of J:

Theorem 4.11. Let A and B be δ -one-forms. Then

(i)
$$\int AJBJ^{-1}|\mathcal{D}|^{-3} = \frac{1}{2}\int A|\mathcal{D}|^{-3}\overline{\int B|\mathcal{D}|^{-3}},$$

(ii) $\int AJBJ^{-1}|\mathcal{D}|^{-2} = \frac{1}{2}\int A|\mathcal{D}|^{-2}\overline{\int B|\mathcal{D}|^{-3}} + \frac{1}{2}\int A|\mathcal{D}|^{-3}\overline{\int B|\mathcal{D}|^{-2}},$
(iii) $\int A^2JBJ^{-1}|\mathcal{D}|^{-3} = \frac{1}{2}\int A^2|\mathcal{D}|^{-3}\overline{\int B|\mathcal{D}|^{-3}},$
(iv) $\int \delta(A)A|\mathcal{D}|^{-3} = \int \delta(A)JAJ^{-1}|\mathcal{D}|^{-3} = 0.$

We gather at the beginning of this section the main notations for technical lemmas which will follow.

For any pair $(k,p) \in \mathbb{N}^3 \times \mathbb{N}^3$ such that $k_i \leq |\alpha_i|$, $p_i \leq |\beta_i|$, where $\alpha, \beta \in \mathbb{Z} \times \mathbb{N} \times \mathbb{N}$, we define

$$\begin{split} v_{k,p} &:= g(p) \, \binom{|\alpha_1|}{k_1}_{q^{2\eta_{\alpha_1}}} \binom{\alpha_2}{k_2} \binom{\alpha_3}{k_3} \binom{|\beta_1|}{p_1}_{q^{2\eta_{\beta_1}}} \binom{\beta_2}{p_2} \binom{\beta_3}{p_3} (-1)^{k_1 + p_1 + \alpha_2 + \alpha_3 + \beta_2 + \beta_3} \, q^{\sigma_{k,p}}, \\ h_{k,p} &:= \alpha_1 + \alpha_2 - \alpha_3 - 2(\eta_{\alpha_1} k_1 + k_2 - k_3) + g(p) \,, \\ g(p) &:= \beta_1 + \beta_2 - \beta_3 - 2(\eta_{\beta_1} p_1 + p_2 - p_3) \,, \\ \sigma_{k,p} &:= k_1 + p_1 + \sigma_{k,p}^t + \sigma_{k,p}^u \,, \\ \sigma_{k,p}^t &:= k_1 \widehat{k}_2 - \widehat{k}_3 (k_1 + k_2) + \eta_{\beta_1} \widehat{p}_1 |k|_1 + \widehat{p}_2 (|k|_1 + p_1) - \widehat{p}_3 (|k|_1 + p_1 + p_2) \,, \\ \sigma_{k,p}^u &:= (k_3 + \eta_{\beta_1} \widehat{p}_1 - p_2 + p_3) (k_1 + \widehat{k}_2 + \widehat{k}_3) - k_2 (k_1 + \widehat{k}_2) + (p_1 + \widehat{p}_2) (-p_2 + p_3) + \widehat{p}_3 p_3 \,, \\ t_{k,p} &= a_{\alpha_1}^{\widehat{k}_1} \, a^{\widehat{k}_2} \, a^{*\widehat{k}_3} \, a_{\beta_1}^{\widehat{p}_1} \, a^{*p_2} \, a^{*\widehat{p}_3} \, b^{|k|_1 + |p|_1} \,, \\ u_{k,p} &= a_{\alpha_1}^{\widehat{k}_1} \, a^{*k_2} \, a^{k_3} \, a_{\beta_1}^{\widehat{p}_1} \, a^{*p_2} \, a^{p_3} \, b^{|\widehat{k}|_1 + |\widehat{p}|_1} \,. \end{split}$$

where we used the notation

$$\widehat{k}_i := |\alpha_i| - k_i, \, \widehat{p}_i := |\beta_i| - p_i,$$

so $0 \le \hat{k}_i \le |\alpha_i|$, $0 \le \hat{p}_i \le |\beta_i|$. We will also use the shortcut $\tilde{k} := (k_1, \hat{k}_2, \hat{k}_3)$. For $\beta_1 \in \mathbb{Z}$ and $j \in \mathbb{N}$, we define

$$w_1(\beta_1, j) := \sum_{n=0}^{\infty} \left(q^{2jn} (q_{n, |\beta_1|}^{\uparrow_{\beta_1}})^2 - \delta_{j, 0} \right),$$

$$w_{\beta}^{\alpha} := 2\beta_1 q^{\beta_1 (2\alpha_3 + \beta_3 - \beta_2)} w_1(\beta_1, \alpha_3 + \beta_3).$$

We introduce the following notations:

$$\begin{split} q_{k,p,n}^+ &:= q^{n(|k|_1 + |p|_1)} q_{n+r_{k,p}^+ - \eta_{\alpha_1} \widehat{k}_1, \widehat{k}_1}^{\uparrow_{\alpha_1}} q_{n-\widehat{k}_3 + \eta_{\beta_1} \widehat{p}_1 + \widehat{p}_2 - \widehat{p}_3, \widehat{k}_2}^{\uparrow} q_{n+\eta_{\beta_1} \widehat{p}_1 + \widehat{p}_2 - \widehat{p}_3, \widehat{k}_3}^{\uparrow_{\beta_1}} q_{n+\widehat{p}_2 - \widehat{p}_3, \widehat{k}_3}^{\uparrow_{\beta_1}} q_{n-\widehat{p}_3, \widehat{p}_2}^{\uparrow} q_{n-\widehat{p}_3, \widehat{p}_2}^{\downarrow} q_{n-\widehat{p}_3, \widehat{p}_2}^{\uparrow_{\beta_1}} q_{n+p_3, p_2}^{\downarrow} q_{n+p_3, p_2}^{\uparrow_{\beta_1}} q_{n+p_3, p_3}^{\uparrow_{\beta_1}} q_{n+p_3, p_3}$$

Thus, $\pi_+(t_{k,p})\varepsilon_n = q_{k,p,n}^+\varepsilon_{n+r_{k,p}^+}$ and $\pi_-(u_{k,p})\varepsilon_n = q_{k,p,n}^-\varepsilon_{n+r_{k,p}^-}$.

Lemma 4.12. We have

$$r\left((M_{\beta}^{\alpha})^{\circ}\right) = \sum_{k,p} \delta_{h_{k,p},0} v_{k,p} \pi_{+}(t_{k,p}) \otimes \pi_{-}(u_{k,p})$$

where the summation is done on k_i, p_i in \mathbb{N} such that $k_i \leq |\alpha_i|, p_i \leq |\beta_i|$ for $i \in \{1, 2, 3\}$.

 $\textit{Proof. Since } \underline{\pi}(m^{\alpha}) = (a_{+} + a_{-})^{\alpha_{1}}(b_{+} + b_{-})^{\alpha_{2}}(b_{+}^{*} + b_{-}^{*})^{\alpha_{3}}, \text{ with } v_{k} := \binom{|\alpha_{1}|}{k_{1}}_{q^{2\eta_{\alpha_{1}}}}\binom{\alpha_{2}}{k_{2}}\binom{\alpha_{3}}{k_{3}},$

$$\underline{\pi}(m^{\alpha}) = \sum_{k} v_{k} c_{k} \text{ where } c_{k} := a_{+,\alpha_{1}}^{|\alpha_{1}|-k_{1}} a_{-,\alpha_{1}}^{k_{1}} b_{+}^{\alpha_{2}-k_{2}} b_{-}^{k_{2}} b_{+}^{k_{3}-k_{3}} b_{-}^{* k_{3}}.$$

By Lemma 3.2 (iii) we see that $\delta(\underline{\pi}(m^{\beta})) = \sum_{p} w_{p} d_{p}$ where we introduce $w_{p} := \binom{|\beta_{1}|}{p_{1}}_{q^{2\eta_{\beta_{1}}}} \binom{\beta_{2}}{p_{2}} \binom{\beta_{3}}{p_{3}}$ and $d_{p} := g(p) a_{+,\beta_{1}}^{|\beta_{1}|-p_{1}} a_{-,\beta_{1}}^{p_{1}} b_{+}^{\beta_{2}-p_{2}} b_{+}^{p_{2}} b_{+}^{*\beta_{3}-p_{3}} b_{-}^{*p_{3}}$. As a consequence, $(M_{\beta}^{\alpha})^{\circ} = \sum_{k,p} \delta_{h(k,p),0} g(p) v_{k} w_{p} c_{k,p}$ where

$$c_{k,p} = a_{+,\alpha_1}^{\hat{k}_1} a_{-,\alpha_1}^{k_1} b_{+}^{\hat{k}_2} b_{+}^{k_2} b_{+}^{*\hat{k}_3} b_{-}^{*\hat{k}_3} a_{+,\beta_1}^{\widehat{p}_1} a_{-,\beta_1}^{p_1} b_{+}^{\widehat{p}_2} b_{-}^{p_2} b_{+}^{*\hat{p}_3} b_{-}^{*p_3}$$

$$(42)$$

With (42), we get $r(c_{k,p}) = (-1)^{k_1+p_1+\alpha_2+\alpha_3+\beta_2+\beta_3} q^{k_1+p_1} \pi_+(t'_{k,p}) \otimes \pi_-(u'_{k,p})$ where

$$\begin{split} t'_{k,p} &= a_{\alpha_1}^{\widehat{k}_1} \, b^{k_1} \, a^{\widehat{k}_2} \, b^{k_2} \, a^{*\widehat{k}_3} \, b^{k_3} a_{\beta_1}^{\widehat{p}_1} \, b^{p_1} \, a^{\widehat{p}_2} \, b^{p_2} \, a^{*\widehat{p}_3} \, b^{p_3} \,, \\ u'_{k,p} &= a_{\alpha_1}^{\widehat{k}_1} \, b^{k_1} \, b^{\widehat{k}_2} \, a^{*k_2} \, b^{\widehat{k}_3} a^{k_3} \, a_{\beta_1}^{\widehat{p}_1} \, b^{p_1} \, b^{\widehat{p}_2} \, a^{*p_2} \, b^{\widehat{p}_3} \, a^{p_3} \,. \end{split}$$

A recursive use of relation $ba_j = q^{\eta_j} a_j b$ yields the result.

Lemma 4.13. We have

(i) $(\tau_1 \otimes \tau_1) (r(M_{\beta}^{\alpha})^{\circ}) = \beta_1 \, \delta_{\alpha_1,-\beta_1} \, \delta_{\alpha_2,0} \, \delta_{\alpha_3,0} \, \delta_{\beta_2,0} \, \delta_{\beta_3,0}$. (ii) $(\tau_1 \otimes \tau_0 + \tau_0 \otimes \tau_1) (r(M_{\beta}^{\alpha})^{\circ}) = \delta_{\alpha_1,-\beta_1} \, \delta_{\alpha_2+\beta_2,\alpha_3+\beta_3} \, w_{\beta}^{\alpha}$. In particular, if A is a δ -one-form, we have

$$\int A|\mathcal{D}|^{-3} = 2\beta_1 A_{-\beta_1 00}^{\beta_1 00},
\int A|\mathcal{D}|^{-2} = 2w_{\beta}^{\alpha} B(A)_{\alpha}^{\beta}.$$

where we implicitly summed on all α, β indices.

Proof. (i) Using same notations of Lemma 4.12, we obtain by definition of τ_1 ,

$$\tau_1(\pi_+(t_{k,p})) = \delta_{k,0} \,\delta_{p,0} \,\delta_{\alpha_1 + \alpha_2 - \alpha_3 + \beta_1 + \beta_2 - \beta_3,0},$$
(43)

$$\tau_1(\pi_-(u_{k,p})) = \delta_{\widetilde{k},0} \, \delta_{\widetilde{p},0} \, \delta_{\alpha_1 - \alpha_2 + \alpha_3 + \beta_1 - \beta_2 + \beta_3,0} \,. \tag{44}$$

We get $\tau_1(\pi_+(t_{k,p}))$ $\tau_1(\pi_-(u_{k,p})) = \delta_{k,0} \, \delta_{p,0} \, \delta_{\alpha_2,0} \, \delta_{\alpha_3,0} \, \delta_{\beta_2,0} \, \delta_{\beta_3,0} \, \delta_{\alpha_1,-\beta_1}$, so Lemma 4.12 gives the result.

(ii) Since $\pi_+(t_{k,p})\varepsilon_n=q_{k,p,n}^+\varepsilon_{n+r_{k,p}^+}$ and $\pi_-(u_{k,p})\varepsilon_n=q_{k,p,n}^-\varepsilon_{n+r_{k,p}^-}$, we get,

$$\tau_0(\pi_+(t_{k,p})) = \delta_{r_{k,p}^+,0} \sum_{n=0}^{\infty} (q_{k,p,n}^+ - \delta_{k,0} \, \delta_{p,0} \, \delta_{\alpha_1 + \alpha_2 - \alpha_3 + \beta_1 + \beta_2 - \beta_3,0}), \qquad (45)$$

$$\tau_0(\pi_-(u_{k,p})) = \delta_{r_{k,p}^-,0} \sum_{n=0}^{\infty} (q_{k,p,n}^- - \delta_{\tilde{k},0} \delta_{\tilde{p},0} \delta_{\alpha_1 - \alpha_2 + \alpha_3 + \beta_1 - \beta_2 + \beta_3,0}). \tag{46}$$

With (43) and (46) we get

$$\tau_{1}(\pi_{+}(t_{k,p})) \tau_{0}(\pi_{-}(u_{k,p})) = \delta_{k,0} \, \delta_{p,0} \, \delta_{\alpha_{2}+\beta_{2},\alpha_{3}+\beta_{3}} \, \delta_{\alpha_{1},-\beta_{1}} \sum_{n=0}^{\infty} \left(\delta_{k,0} \, \delta_{p,0} \, q_{k,p,n}^{-} - \delta_{\alpha_{3}+\beta_{3},0}\right)$$
$$= \delta_{k,0} \, \delta_{p,0} \, \delta_{\alpha_{2}+\beta_{2},\alpha_{3}+\beta_{3}} \, \delta_{\alpha_{1},-\beta_{1}} w_{1}(\beta_{1},\alpha_{3}+\beta_{3}).$$

Using (44) and (45),

$$\tau_{0}(\pi_{+}(t_{k,p})) \tau_{1}(\pi_{-}(u_{k,p})) = \delta_{\widetilde{k},0} \, \delta_{\widetilde{p},0} \, \delta_{\alpha_{2}+\beta_{2},\alpha_{3}+\beta_{3}} \, \delta_{\alpha_{1},-\beta_{1}} \sum_{n=0}^{\infty} \left(\delta_{\widetilde{k},0} \, \delta_{\widetilde{p},0} \, q_{k,p,n}^{+} - \delta_{\alpha_{3}+\beta_{3},0} \right) \\ = \delta_{\widetilde{k},0} \, \delta_{\widetilde{p},0} \, \delta_{\alpha_{2}+\beta_{2},\alpha_{3}+\beta_{3}} \, \delta_{\alpha_{1},-\beta_{1}} w_{1}(\beta_{1},\alpha_{3}+\beta_{3}).$$

Lemma 4.12 yields the result.

With notations of Lemma 4.12, it is direct to check that for given $\bar{\alpha} = (\alpha, \alpha', \dots, \alpha^{(n-1)})$ and $\bar{\beta} = (\beta, \beta', \dots, \beta^{(n-1)})$,

$$r\left(\left(M_{\bar{\beta}}^{\bar{\alpha}}\right)^{\circ}\right) = \sum_{K,P} \delta_{h_{K,P},0} \, v_{K,P} \, \pi_{+}(t_{K,P}) \otimes \pi_{-}(u_{K,P}) \tag{47}$$

where
$$K = (k, k', \dots k^{(n-1)}), P = (p, p', \dots, p^{(n-1)})$$
 with $0 \le k_i^{(j)} \le |\alpha_i^{(j)}|, 0 \le p_i^{(j)} \le |\beta_i^{(j)}|,$

$$t_{K,P} := t_{k,p} t_{k',p'} \dots t_{k^{(n-1)},p^{(n-1)}}, \qquad u_{K,P} := u_{k,p} u_{k',p'} \dots u_{k^{(n-1)},p^{(n-1)}},$$

$$v_{K,P} := v_{k,p} v_{k',p'} \dots v_{k^{(n-1)},p^{(n-1)}}, \qquad h_{K,P} := h_{k,p} + h_{k',p'} + \dots h_{k^{(n-1)},p^{(n-1)}}.$$

In the following, we will use the shortcuts $A_i := \alpha_i + \alpha'_i + \dots + \alpha_i^{(n-1)}$, $B_i := \beta_i + \beta'_i + \dots + \beta_i^{(n-1)}$. In the case n = 2, we also note $r_{K,P}^{\pm} := r_{k,p}^{\pm} + r_{k',p'}^{\pm}$ and $q_{K,P,n}^{\pm} := q_{k',p',n}^{\pm} q_{k,p,n+r_{k',p'}}^{\pm}$.

Thus, we have $\pi_+(t_{K,P}) \, \varepsilon_m = q_{K,P,m}^+ \, \varepsilon_{m+r_{K,P}^+}$ and $\pi_-(u_{K,P}) \, \varepsilon_m = q_{K,P,n}^- \, \varepsilon_{m+r_{K,P}^-}$. We also introduce, still for n=2,

$$\begin{split} v_{\beta_1,\alpha_1',\beta_1'}(l,j) := \sum_{n=0}^{\infty} \left(q^{l+2nj} \, q_{n+\beta_1'+\alpha_1'+\beta_1,|\beta_1'+\alpha_1'+\beta_1|}^{\uparrow_{\alpha_1'-\alpha_1'-\beta_1}} \, q_{n+\beta_1'+\alpha_1',|\beta_1|}^{\uparrow_{\beta_1}} q_{n+\beta_1',|\alpha_1'|}^{\uparrow_{\alpha_1'}} q_{n,|\beta_1'|}^{\uparrow_{\beta_1'}} - \delta_{j,0} \right), \\ V_{\bar{\beta}}^{\bar{\alpha}} := 2 [\beta_1 \beta_1' + (\beta_2 - \beta_3)(\beta_2' - \beta_3')] \, q^{2\beta_1(\alpha_2 + \alpha_3) + 2\beta_1'(\alpha_2' + \alpha_3')} \\ & \times v_{\beta_1,\alpha_1',\beta_1'} ((\alpha_2 + \beta_2 + \alpha_3 + \beta_3)(\alpha_1' + \beta_1'), A_3 + B_3). \end{split}$$

Lemma 4.14. We have

(i) $(\tau_1 \otimes \tau_1) \left(r(M_{\beta}^{\alpha} M_{\beta'}^{\alpha'})^{\circ} \right) = \beta_1 \beta_1' \delta_{A_1, -B_1} \delta_{A_2, 0} \delta_{A_3, 0} \delta_{B_2, 0} \delta_{B_3, 0}$.

(ii)
$$(\tau_1 \otimes \tau_0 + \tau_0 \otimes \tau_1) (r(M^{\alpha}_{\beta} M^{\alpha'}_{\beta'})^{\circ}) = \delta_{A_2 + B_2, A_3 + B_3} \delta_{A_1, -B_1} V^{\bar{\alpha}}_{\bar{\beta}}$$
.

$$(iii) \quad (\tau_1 \otimes \tau_1) \left(r(M_{\beta}^{\alpha} M_{\beta'}^{\alpha'} M_{\beta''}^{\alpha''})^0 \right) = \beta_1 \beta_1' \beta_1'' \, \delta_{A_1, -B_1} \, \delta_{A_2, 0} \, \delta_{A_3, 0} \, \delta_{B_2, 0} \delta_{B_3, 0}.$$

$$(iv) \quad (\tau_1 \otimes \tau_1) \, \left(r(\delta(M_\beta^\alpha) M_{\beta'}^{\alpha'})^0 \right) = -(\alpha_1' + \beta_1') \beta_1 \beta_1' \, \delta_{A_1, -B_1} \, \delta_{A_2, 0} \, \delta_{A_3, 0} \, \delta_{B_2, 0} \delta_{B_3, 0}.$$

(v) In particular, if A is a δ -one-form,

$$\oint A^{2} |\mathcal{D}|^{-3} = 2\beta_{1} \beta_{1}' B_{a} (A^{2})_{\bar{\alpha}}^{\bar{\beta}},$$

$$\oint A^{2} |\mathcal{D}|^{-2} = 2V_{\bar{\beta}}^{\bar{\alpha}} B(A^{2})_{\bar{\alpha}}^{\bar{\beta}},$$

$$\oint A^{3} |\mathcal{D}|^{-3} = 2\beta_{1} \beta_{1}' \beta_{1}'' B_{a} (A^{3})_{\bar{\alpha}}^{\bar{\beta}},$$

$$\oint \delta(A) A |\mathcal{D}|^{-3} = \oint A \delta(A) |\mathcal{D}|^{-3} = 0.$$

Proof. We have

$$\tau_1(\pi_+(t_{K,P})) = \delta_{K,0} \,\delta_{P,0} \,\delta_{A_1 + A_2 - A_3 + B_1 + B_2 - B_3,0}, \tag{48}$$

$$\tau_1(\pi_-(u_{K,P})) = \delta_{\widetilde{K} \ 0} \, \delta_{\widetilde{P} \ 0} \, \delta_{A_1 - A_2 + A_3 + B_1 - B_2 + B_3, 0} \,. \tag{49}$$

and

$$\tau_0(\pi_+(t_{K,P})) = \delta_{r_{K,P}^+,0} \sum_{n=0}^{\infty} \left(q_{K,P,n}^+ - \delta_{K,0} \, \delta_{P,0} \, \delta_{A_1 + A_2 - A_3 + B_1 + B_2 - B_3,0} \right),\tag{50}$$

$$\tau_0(\pi_-(u_{K,P})) = \delta_{r_{K,P}^-,0} \sum_{n=0}^{\infty} \left(q_{K,P,n}^- - \delta_{\widetilde{K},0} \, \delta_{\widetilde{P},0} \, \delta_{A_1 - A_2 + A_3 + B_1 - B_2 + B_3,0} \right). \tag{51}$$

(i) Equations (48) and (49) give $(\tau_1 \otimes \tau_1) r(\underline{A} \underline{A}')^0 = \delta_{A_1,-B_1} \delta_{A_2,0} \delta_{A_3,0} \delta_{B_2,0} \delta_{B_3,0} \lambda_{0,0}$. A computation of $v_{0,0}$ with $\delta_{A_1,-B_1} \delta_{A_2,0} \delta_{A_3,0} \delta_{B_2,0} \delta_{B_3,0} = 1$ gives the result.

(ii) Equations (48) and (51) yield

$$\tau_{1}(\pi_{+}(t_{K,P})) \tau_{0}(\pi_{-}(u_{K,P})) = \delta_{K,0} \delta_{P,0} \delta_{A_{2}+B_{2},A_{3}+B_{3}} \delta_{A_{1},-B_{1}} \times v_{\beta_{1},\alpha'_{1},\beta'_{1}}((\alpha_{2}+\beta_{2}+\alpha_{3}+\beta_{3})(\alpha'_{1}+\beta'_{1}), A_{3}+B_{3}).$$

Equations (50) and (49) yield

$$\tau_0(\pi_+(t_{K,P})) \tau_1(\pi_-(u_{K,P})) = \delta_{\widetilde{K},0} \delta_{\widetilde{P},0} \delta_{A_2+B_2,A_3+B_3} \delta_{A_1,-B_1} \times v_{\beta_1,\alpha_1',\beta_1'}((\alpha_2+\beta_2+\alpha_3+\beta_3)(\alpha_1'+\beta_1'), A_3+B_3)$$

and the result follows.

(iii) With (47) a direct computation gives

$$\tau_1(\pi_+(t_{K,P})) = \delta_{K,0} \, \delta_{P,0} \, \delta_{A_1 + A_2 - A_3 + B_1 + B_2 - B_3,0} \,, \tag{52}$$

$$\tau_1(\pi_-(u_{K,P})) = \delta_{\widetilde{K},0} \, \delta_{\widetilde{P},0} \, \delta_{A_1 - A_2 + A_3 + B_1 - B_2 + B_3,0} \,. \tag{53}$$

Using (52) and (53), $(\tau_1 \otimes \tau_1) (r(\underline{A}\underline{A'}\underline{A''})^{\circ}) = \delta_{A_1,-B_1} \delta_{A_2,0} \delta_{A_3,0} \delta_{B_2,0} \delta_{B_3,0} v_{0,0}$. A computation of $v_{0,0}$ with $\delta_{A_1,-B_1} \delta_{A_2,0} \delta_{A_3,0} \delta_{B_2,0} \delta_{B_3,0} = 1$ gives the result.

(iv) We have $\delta(M^{\alpha}_{\beta})M^{\alpha'}_{\beta'} = \delta(x)\delta(y)x'\delta(y') + x\delta^2(y)x'\delta(y')$ where x, x', y, y' are monomials ($\underline{\pi}$ omitted). Since

$$\underline{\pi}(x) = \sum_{k} {\alpha \choose k} a_{+,\alpha_1}^{\hat{k}_1} a_{-,\alpha_1}^{k_1} b_{+}^{\hat{k}_2} b_{-}^{k_2} b_{+}^{*\hat{k}_3} b_{-}^{*k_3} =: \sum_{k} {\alpha \choose k} c_k,$$

we get
$$\delta(\underline{\pi}(x)) = \sum_{k} g(k) \binom{\alpha}{k} c_k$$
.
Similarly, $\delta(\underline{\pi}(y)) = \sum_{p} g(p) \binom{\beta}{p} c_p$ and $\delta^2(\underline{\pi}(y)) = \sum_{p} g(p)^2 \binom{\beta}{p} c_p$.
Thus, with $c_{K,P} := c_k c_p c_{k'} c_{p'}$,

$$\begin{split} &\delta(x)\delta(y)x'\delta(y') = \sum_{K,P} g(k)\,g(p)\,g(p')\binom{\alpha}{K}\binom{\beta}{P}\,c_{K,P}\,,\\ &x\delta^2(y)x'\delta(y') = \sum_{K,P} g(p)^2g(p')\binom{\alpha}{K}\binom{\beta}{P}\,c_{K,P}\,,\\ &r(\delta(M^\alpha_\beta)M^{\alpha'}_{\beta'})^0 = \sum_{K,P} \delta_{h_{K,P},0}\big(g(k)+g(p)\big)\,g(p)\,g(p')\binom{\alpha}{K}\binom{\beta}{P}\,r(c_{K,P}) =: \sum_{K,P} \lambda_{K,P}\,\,r(c_{K,P})\,. \end{split}$$

Since $r(c_k) = (-q)^{k_1}(-1)^{\alpha_2+\alpha_3}\pi_+(t_k) \otimes \pi_-(u_k)$ with t_k, u_k defined by

$$t_k := a_{\alpha_1}^{\widehat{k}_1} \, b^{k_1} \, a^{\widehat{k}_2} \, b^{k_2} \, a^{*\widehat{k}_3} \, b^{k_3} \text{ and } u_k := a_{\alpha_1}^{\widehat{k}_1} \, b^{k_1} \, b^{\widehat{k}_2} \, a^{*k_2} \, b^{\widehat{k}_3} \, a^{k_3},$$

we get

$$r(\delta(M_{\beta}^{\alpha})M_{\beta'}^{\alpha'})^{0} = \sum_{K,P} \lambda_{K,P} (-q)^{k_{1} + k_{1}' + p_{1} + p_{1}'} (-1)^{A_{2} + A_{3} + B_{2} + B_{3}} \pi_{+}(t_{K,P}) \otimes \pi_{-}(u_{K,P})$$

where $t_{K,P} = t_k t_p t_{k'} t_{p'}$ and $u_{K,P} = u_k u_p u_{k'} u_{p'}$. Direct computations yield

$$\tau_1(\pi_+(t_{K,P})) = \delta_{K,0} \, \delta_{P,0} \, \delta_{A_1 + A_2 - A_3 + B_1 + B_2 - B_3,0}, \tau_1(\pi_-(u_{K,P})) = \delta_{\widetilde{K},0} \, \delta_{\widetilde{P},0} \, \delta_{A_1 - A_2 + A_3 + B_1 - B_2 + B_3,0}.$$

The result follows.

(v) For the last equality, note that by (iv)

$$\int \delta(A)A|\mathcal{D}|^{-3} = -2\sum_{\alpha_1,\alpha_1',\beta_1,\beta_1'} (\alpha_1' + \beta_1')\beta_1\beta_1' A_{\alpha_1000}^{\beta_100} A_{\alpha_1'00}^{\beta_1'00} \delta_{\alpha_1+\alpha_1'+\beta_1+\beta_1',0}.$$

The following change of variables $\alpha_1 \leftrightarrow \alpha'_1$, $\beta_1 \leftrightarrow \beta'_1$, implies by symmetry that this is equal to zero.

For a given δ -1-form A, we say that A is homogeneous of degree in a equal to $n \in \mathbb{Z}$ if it is a linear combination of M_{β}^{α} such that $\alpha_1 + \beta_1 = n$. From Lemma 4.14 (iv) we get,

Corollary 4.15. Let A, A' be two δ -1-forms, then

$$\begin{split} & \int (A|\mathcal{D}|^{-1})^2 = \int A^2|\mathcal{D}|^{-2} \,, \\ & \int A|\mathcal{D}|^{-1}A'|\mathcal{D}|^{-1} = \int AA'|\mathcal{D}|^{-2} - n \int AA'|\mathcal{D}|^{-3} \,, \text{ when } A' \text{ homogenous of degree } n. \end{split}$$

Lemma 4.16. We have

(i)
$$(\tau_{1} \otimes \tau_{1}) \widetilde{\rho} (M_{\beta}^{\alpha} J M_{\beta'}^{\alpha'} J^{-1}) = \beta_{1} \beta'_{1} \delta_{\alpha_{1}, -\beta_{1}} \delta_{\alpha'_{1}, -\beta'_{1}} \delta_{A_{2}, 0} \delta_{A_{3}, 0} \delta_{B_{2}, 0} \delta_{B_{3}, 0}.$$

(ii) $(\tau_{0} \otimes \tau_{1} + \tau_{1} \otimes \tau_{0}) \widetilde{\rho} (M_{\beta}^{\alpha} J M_{\beta'}^{\alpha'} J^{-1}) = \delta_{\alpha_{1}, -\beta_{1}} \delta_{\alpha'_{1}, -\beta'_{1}} (\beta'_{1} w_{\beta}^{\alpha} \delta_{\alpha'_{2} + \beta'_{2} + \alpha'_{3} + \beta'_{3}, 0} \delta_{\alpha_{2} + \beta_{2}, \alpha_{3} + \beta_{3}} + \beta_{1} w_{\beta'}^{\alpha'} \delta_{\alpha_{2} + \beta_{2} + \alpha_{3} + \beta_{3}, 0} \delta_{\alpha'_{2} + \beta'_{2}, \alpha'_{3} + \beta'_{3}}).$

$$(iii) \ (\tau_1 \otimes \tau_1) \ \widetilde{\rho} \big(M_\beta^\alpha M_{\beta'}^{\alpha'} J M_{\beta''}^{\alpha''} J^{-1} \big) = \beta_1 \beta_1' \beta_1'' \ \delta_{\alpha_1 + \alpha_1', -\beta_1 - \beta_1'} \ \delta_{\alpha_1'', -\beta_1''} \ \delta_{A_2, 0} \ \delta_{A_3, 0} \ \delta_{B_2, 0} \delta_{B_3, 0} \ .$$

$$(iv) (\tau_1 \otimes \tau_1) \widetilde{\rho} (\delta(M_{\beta}^{\alpha}) J M_{\beta'}^{\alpha'} J^{-1}) = -(\alpha_1' + \beta_1') \beta_1 \beta_1' \delta_{\alpha_1, -\beta_1} \delta_{\alpha_1', -\beta_1'} \delta_{A_2, 0} \delta_{A_3, 0} \delta_{B_2, 0} \delta_{B_3, 0}.$$

(v) In particular, if A and A' are δ -one forms,

$$\int AJA'J^{-1}|\mathcal{D}|^{-3} = 2(\beta_1 A_{-\beta_100}^{\beta_100})(\beta_1 \bar{A'}_{-\beta_100}^{\beta_100}),$$

$$\int AJA'J^{-1}|\mathcal{D}|^{-2} = 2(\beta_1 \bar{A'}_{-\beta_100}^{\beta_100})(w_{\beta}^{\alpha}B(A)_{\alpha}^{\beta}) + 2(\beta_1 A_{-\beta_100}^{\beta_100})(w_{\beta}^{\alpha}B(\bar{A'})_{\alpha}^{\beta}),$$

$$\int A^2 JA'J^{-1}|\mathcal{D}|^{-3} = 2(\beta_1 \bar{A'}_{-\beta_100}^{\beta_100})(\beta_1 \beta'_1 B_a(A^2)_{\bar{\alpha}}^{\bar{\beta}}),$$

$$\int \delta(A)JAJ^{-1} = 0.$$

Proof. (i) Following notations of Lemma 4.12, we have

$$M_{\beta}^{\alpha} J M_{\beta'}^{\alpha'} J^{-1} = \sum_{K,P} v_{K,P} c_{k,p} J c_{k',p'} J^{-1}$$

where $K=(k,k'),\,P=(p,p'),\,\lambda_{K,P}=g(p)g(p')v_kv_{k'}w_pw_{p'}.$ Thus,

$$\widetilde{\rho}(M_{\beta}^{\alpha}JM_{\beta'}^{\alpha'}J^{-1}) = (-1)^{A_2 + A_3 + B_2 + B_3} \sum_{K,P} (-q)^{k_1 + k_1' + p_1 + p_1'} \lambda_{K,P} \, T_{K,P}^+ \otimes T_{K,P}^-$$

where $T_{K,P}^+ := \pi'_+(t_k t_p) \widehat{\pi}_+(t_{k'} t_{p'})$ and $T_{K,P}^- := \pi'_-(u_k u_p) \widehat{\pi}_-(u_{k'} u_{p'})$ with

$$t_k := a_{\alpha_1}^{\hat{k}_1} b_{\alpha_1}^{*k_1} a^{\hat{k}_2} b^{*k_2} a^{*\hat{k}_3} b^{k_3},$$

$$u_k := a_{\alpha_1}^{\hat{k}_1} b_{\alpha_1}^{*k_1} b^{\hat{k}_2} a^{*k_2} b^{*\hat{k}_3} a^{k_3}.$$

A direct computation leads to

$$\tau_1(T_{K,P}^+) = \delta_{K,0} \, \delta_{P,0} \, \delta_{\alpha_1 + \alpha_2 - \alpha_3 + \beta_1 + \beta_2 - \beta_3,0} \, \delta_{\alpha_1' + \alpha_2' - \alpha_3' + \beta_1' + \beta_2' - \beta_3',0}$$

$$\tau_1(T_{K,P}^-) = \delta_{\widetilde{K},0} \, \delta_{\widetilde{P},0} \, \delta_{\alpha_1 - \alpha_2 + \alpha_3 + \beta_1 - \beta_2 + \beta_3,0} \, \delta_{\alpha_1' - \alpha_2' + \alpha_2' + \beta_1' - \beta_2' + \beta_2',0}$$

which gives the result.

(ii) Using the commutation relations on \mathcal{A} , we see that there are real functions of (K, P), noted $\sigma_{K,P}^t$ and $\sigma_{K,P}^u$ such that

$$\begin{split} T_{K,P}^{+} &= q^{\sigma_{K,P}^{t}} \, \pi'_{+}(t_{k,p}) \widehat{\pi}_{+}(t_{k',p'}), \\ T_{K,P}^{-} &= q^{\sigma_{K,P}^{u}} \, \pi'_{-}(u_{k,p}) \widehat{\pi}_{-}(u_{k',p'}), \\ t_{k,p} &:= a_{\alpha_{1}}^{\widehat{k}_{1}} \, a^{\widehat{k}_{2}} \, a^{*\widehat{k}_{3}} \, a_{\beta_{1}}^{\widehat{p}_{1}} \, a^{\widehat{p}_{2}} \, a^{*\widehat{p}_{3}} \, b^{*k_{1}}_{\alpha_{1}} \, b^{*p_{1}}_{\beta_{1}} \, b^{*k_{2}+p_{2}} b^{k_{3}+p_{3}}, \\ u_{k,p} &:= a_{\alpha_{1}}^{\widehat{k}_{1}} \, a^{*k_{2}} \, a^{k_{3}} \, a_{\beta_{1}}^{\widehat{p}_{1}} \, a^{*p_{2}} \, a^{p_{3}} \, b^{*k_{1}}_{\alpha_{1}} \, b^{*p_{1}}_{\beta_{1}} \, b^{\widehat{k}_{2}+\widehat{p}_{2}} b^{*\widehat{k}_{3}+\widehat{p}_{3}}. \end{split}$$

We have, under the hypothesis $\tau_1(T_{K,P}^-) = 1$,

$$\begin{split} \widehat{\pi}_{+}(t_{k',p'})\varepsilon_{m,2j} &= (-1)^{\lambda'}q^{(2j-m)\lambda'}\,q_{2j-m-s+\beta'_{1},|\alpha'_{1}|}^{\uparrow_{\alpha'_{1}}}\,q_{2j-m-s,|\beta'_{1}|}^{\uparrow_{\beta'_{1}}}\,\varepsilon_{m+s,2j}\,,\\ s &:= -\alpha'_{2} + \alpha'_{3} - \beta'_{2} + \beta'_{3} = \alpha'_{1} + \beta'_{1}\,,\\ \lambda' &:= \alpha'_{2} + \alpha'_{3} + \beta'_{2} + \beta'_{3}\,,\\ \lambda &:= \alpha_{2} + \alpha_{3} + \beta_{2} + \beta_{3}\\ \tau_{1}(T_{K,P}^{+}) &= \delta_{\lambda,0}\,\delta_{\lambda',0}\,. \end{split}$$

and then,

$$\begin{split} (T_{K,P}^{+})_{m,2j} &= q^{\sigma_{K,P}^{t}+s\lambda}(-1)^{\lambda'}q^{(2j-m)\lambda'+m\lambda} F_{m} F_{2j-m}' \delta_{A_{1}+B_{1},0} , \\ F_{2j-m}' &:= q_{2j-m-\alpha'_{1},|\alpha'_{1}|}^{\uparrow_{\alpha'_{1}}} q_{2j-m-\alpha'_{1}-\beta'_{1},|\beta'_{1}|}^{\uparrow_{\beta'_{1}}} , \\ F_{m} &:= q_{m-\alpha_{1},|\alpha_{1}|}^{\uparrow_{\alpha_{1}}} q_{m-\beta_{1}-\alpha_{1},|\beta_{1}|}^{\uparrow_{\beta_{1}}} . \end{split}$$

Following the proof of Lemma 4.7, we see that $\tau_0(T_{K,P}^+)$ is possibly nonzero only in the two cases $\lambda' = 0$ or $\lambda = 0$.

Suppose first $\lambda = \lambda' = 0$. In that case, we have

$$\tau_0(T_{K,P}^+) = \lim_{2j \to \infty} \sum_{m=0}^{2j} \left((q_{m,|\beta_1|}^{\uparrow_{\beta_1}} q_{2j-m,|\beta_1'|}^{\uparrow_{\beta_1'}})^2 - 1 \right) = \sum_{m=0}^{\infty} \left((q_{m,|\beta_1|}^{\uparrow_{\beta_1}})^2 - 1 \right) + \sum_{m=0}^{\infty} \left((q_{m,|\beta_1|}^{\uparrow_{\beta_1'}})^2 - 1 \right)$$

where the second equality comes from Lemma 4.17.

In the case $(\lambda = 0, \lambda' > 0)$, we get $\alpha'_1 = -\beta'_1$ and thus,

$$(T_{K,P}^+)_{m,2j} = q^{\sigma_{K,P}^t} q^{m\lambda} \left(q_{m,|\beta_1|}^{\uparrow_{\beta_1}} q_{2j-m,|\beta_1'|}^{\uparrow_{\beta_1'}} \right)^2 \delta_{\alpha_1+\beta_1,0}$$

Let us note $U_{2j} = \sum_{m=0}^{2j} q^{m\lambda} (q_{m,|\beta_1|}^{\uparrow_{\beta_1}} q_{2j-m,|\beta_1'|}^{\uparrow_{\beta_1'}})^2$ and $L_{2j} = \sum_{m=0}^{2j} q^{m\lambda} (q_{m,|\beta_1|}^{\uparrow_{\beta_1}})^2$.

Suppose $\beta_1' > 0$. Since $(q_{2j-m,|\beta_1'|}^{\uparrow_{\beta_1'}})^2 - 1 = \sum_{|p|_1 \neq 0, p_i \in \{0,1\}} (-1)^{|p|_1} q^{r_p} q^{2(2j-m)|p|_1}$ where we have $r_p = 2 + \dots + 2\beta_1'$. As in the proof of Lemma 4.7 (ii), we can conclude that $U_{2j} - L_{2j}$ converges to 0. The case $\beta_1' \leq 0$ is similar.

In the other case $(\lambda > 0, \lambda' = 0)$, the arguments are the same, replacing λ by λ' and α_1 , β_1 by α'_1 , β'_1 . Finally,

$$\tau_{0}(T_{K,P}^{+})\tau_{1}(T_{K,P}^{-}) = \delta_{\widetilde{K},0} \, \delta_{\widetilde{P},0} \, \delta_{\alpha_{1},-\beta_{1}} \, \delta_{\alpha'_{1},-\beta'_{1}}(\delta_{\lambda',0} \, \delta_{\alpha_{2}+\beta_{2},\alpha_{3}+\beta_{3}} \, s_{\alpha,\beta} + \delta_{\lambda,0} \, \delta_{\alpha'_{2}+\beta'_{2},\alpha'_{3}+\beta'_{3}} \, s_{\alpha',\beta'}),$$

$$s_{\alpha\beta} := q^{\beta_{1}(\alpha_{3}-\alpha_{2})} \, \sum_{m=0}^{\infty} \left(q^{m\lambda} \, (q_{m,|\beta_{1}|}^{\uparrow_{\beta_{1}}})^{2} - \delta_{\lambda,0} \right).$$

A similar computation of $\tau_0(T_{K,P}^-)$ can be done following the same arguments. We find eventually

$$\tau_1(T_{K,P}^+)\tau_0(T_{K,P}^-) = \delta_{K,0}\,\delta_{P,0}\,\delta_{\alpha_1,-\beta_1}\,\delta_{\alpha_1',-\beta_1'}(\delta_{\lambda',0}\,\delta_{\alpha_2+\beta_2,\alpha_3+\beta_3}\,s_{\alpha,\beta} + \delta_{\lambda,0}\,\delta_{\alpha_2'+\beta_2',\alpha_3'+\beta_3'}\,s_{\alpha',\beta'})$$

and the result follows.

- (iii) The same arguments of (i) apply here with minor changes.
- (iv) follows from a slight modification of the proof of Lemma 4.14 (iv).
- (v) is a straightforward consequence of (i, ii, iii, iv).

Lemma 4.17. Let $\beta, \beta' \in \mathbb{Z}$. Then,

$$\lim_{2j \to \infty} \sum_{m=0}^{2j} \left((q_{m,|\beta|}^{\uparrow_{\beta}} q_{2j-m,|\beta'|}^{\uparrow_{\beta'}})^2 - 1 \right) = \sum_{m=0}^{\infty} \left((q_{m,|\beta|}^{\uparrow_{\beta}})^2 - 1 \right) + \sum_{m=0}^{\infty} \left((q_{m,|\beta'|}^{\uparrow_{\beta'}})^2 - 1 \right).$$

Proof. We give a proof for β and $\beta' > 0$, the other cases being similar. Since $(q_{m,|\beta|}^{\uparrow_{\beta}})^2 = \sum_{p_i \in \{0,1\}} (-1)^{|p|_1} q^{r_p} q^{2|p|_1 m}$ where $p = (p_1, \dots, p_{\beta})$ and $r_p := 2(p_1 + \dots + \beta p_{\beta})$, we get, with the notations $\lambda_{p,p'} := (-1)^{|p+p'|_1} q^{r_p+r_{p'}}$ and $U_{2j} := \sum_{m=0}^{2j} (q_{m,|\beta|}^{\uparrow_{\beta}} q_{2j-m,|\beta'|}^{\uparrow_{\beta'}})^2 - 1$,

$$U_{2j} = \sum_{m=0}^{2j} \sum_{|p+p'|_1>0} \lambda_{p,p'} q^{2|p|_1 m + 2|p'|_1(2j-m)}$$

$$= \sum_{|p|_1 \ge |p'|_1, |p|_1>0} \lambda_{p,p'} V_{2j,p,p'} + \sum_{|p|_1 < |p'|_1, |p'|_1>0} \lambda_{p,p'} V'_{2j,p,p'}$$

where

$$V_{2j,p,p'} = q^{4j|p'|_1} \sum_{m=0}^{2j} q^{2(|p|_1 - |p'|_1)m}, \qquad V'_{2j,p,p'} = q^{4j|p|_1} \sum_{m=0}^{2j} q^{2(|p'|_1 - |p|_1)m}.$$

It is clear that $V_{2j,p,p'}$ has 0 for limit when $j \to \infty$ when $|p'|_1 > 0$, and $V'_{2j,p,p'}$ has 0 for limit when $j \to \infty$ when $|p|_1 > 0$. As a consequence,

$$U_{2j} = \sum_{|p|_1 > 0} \lambda_{p,0} V_{2j,p,0} + \sum_{|p'|_1 > 0} \lambda_{0,p'} V'_{2j,0,p'} + o(1).$$

The result follows as
$$\sum_{m=0}^{2j} \left((q_{m,|\beta|}^{\uparrow_{\beta}})^2 - 1 \right) = \sum_{|p|_1 > 0} \lambda_{p,0} V_{2j,p,0} \text{ and } \sum_{m=0}^{2j} \left((q_{m,|\beta'|}^{\uparrow_{\beta'}})^2 - 1 \right) = \sum_{|p'|_1 > 0} \lambda_{0,p'} V'_{2j,0,p'}.$$

Proof of Theorem 4.11. The result follows from Lemmas 4.13, 4.14 (v) and 4.16 (v).

Proof of Theorem 4.3 and corollaries

Lemma 4.18. We have on $SU_q(2)$,

- $\int |\mathcal{D}|^{-3} = 2.$
- (ii) $\int |\mathcal{D}|^{-2} = 0.$ (iii) $\int |\mathcal{D}|^{-1} = -\frac{1}{2}.$

Proof. (iv) We have by definition

$$\zeta_{\mathcal{D}}(s) := \text{Tr}(|\mathcal{D}|^{-s}) = \sum_{2j=0}^{\infty} \sum_{m=0}^{2j} \sum_{l=0}^{2j+1} \langle v_{m,l}^j, |\mathcal{D}|^{-s} v_{m,l}^j \rangle.$$

Since $|\mathcal{D}|^{-s} v_{m,l}^j = \begin{pmatrix} d_{j+}^{-s} & 0 \\ 0 & d_j^{-s} \end{pmatrix} v_{m,l}^j$ where $d_j := 2j + \frac{1}{2}$, we get

$$\zeta_{\mathcal{D}}(s) = \sum_{2j=0}^{\infty} (2j+1)(2j+2) d_{j+}^{-s} + \sum_{2j=1}^{\infty} (2j+1)(2j) d_{j}^{-s} = 2\sum_{2j=0}^{\infty} (2j+1)(2j) d_{j}^{-s}.$$

With the equalities $(2j+1)(2j) = d_j^2 - \frac{1}{4}$ and $\zeta(s,\frac{1}{2}) = (2^s-1)\zeta(s)$ (here $\zeta(s,x) := \sum_{n \in \mathbb{N}} \frac{1}{(n+x)^s}$ is the Hurwitz zeta function and $\zeta(s) := \zeta(s,1)$ is the Riemann zeta function) we get

$$\zeta_{\mathcal{D}}(s) = 2(2^{s-2} - 1)\zeta(s-2) - \frac{1}{2}(2^s - 1)\zeta(s)$$
(54)

which entails that $\zeta_{\mathcal{D}}(0) = 0$.

(i, ii, iii) are direct consequences of equation (54).

Proof of Theorem 4.3. It is a consequence of Lemma 4.18 and Theorems 4.1, 4.11.

As we have seen, the computation of noncommutative integral on $SU_q(2)$ leads to certain function of A which filter some symmetry on the degree in a, a^* , b, b^* of the canonical decomposition. Precisely, it is the balanced features that appear and the following functions of A^n , $n \in \{1, 2, 3\}$:

$$\oint A^n |\mathcal{D}|^{-p} \tag{55}$$

where $1 \le n \le p \le 3$. We will see in the next section a method for the computation of these integrals.

Corollary 4.19. Let u be a unitary in $C^{\infty}(SU_q(2))$ and $\gamma_u(\mathbb{A}) := \pi(u)\mathbb{A}\pi(u^*) + \pi(u)d\pi(u^*)$ be a gauge-variant of \mathbb{A} . Then the following term of Theorem 4.3 are gauge invariant

$$\int A |\mathcal{D}|^{-3} \,, \qquad \int A^2 |\mathcal{D}|^{-3} \,- \int A |\mathcal{D}|^{-2} \,, \qquad \qquad -2 \int A |\mathcal{D}|^{-1} \,+ \int A^2 |\mathcal{D}|^{-2} \,- \, \tfrac{2}{3} \int A^3 |\mathcal{D}|^{-3} \,.$$

Proof. It is sufficient to remark that all terms $\int |D_{\mathbb{A}}|^{-k}$ and $\zeta_{\mathcal{D}_{\mathbb{A}}}(0)$ in spectral action (4) are gauge invariant. This can also be seen via the computation $D_{\gamma_u(\mathbb{A})} = V_u \mathcal{D} V_u^* + V_u P_0 V_u^*$ where P_0 is the projection on Ker \mathcal{D} and $V_u = \pi(u)J\pi(u)J^{-1}$ and $\int |D_{\mathbb{A}}|^{n-k} = \operatorname{Res}_{s=n-k} \operatorname{Tr} \left(|D_{\mathbb{A}}|^{n-k}\right)$ (see [22, Prop. 5.1 (iii) and Prop. 4.8].)

Corollary 4.20. In the case of the spectral action without the reality operator (i.e. $D_{\mathbb{A}} = \mathcal{D} + \mathbb{A}$), we get

$$\begin{split} & \int |D_{\mathbb{A}}|^{-3} = 2, \qquad \int |D_{\mathbb{A}}|^{-2} = -2 \int A|\mathcal{D}|^{-3} \,, \qquad \int |D_{\mathbb{A}}|^{-1} = -\tfrac{1}{2} - \int A|\mathcal{D}|^{-2} + \int A^2|\mathcal{D}|^{-3}, \\ & \zeta_{D_{\mathbb{A}}}(0) = -\int A|\mathcal{D}|^{-1} + \tfrac{1}{2} \int A^2|\mathcal{D}|^{-2} - \tfrac{1}{3} \int A^3|\mathcal{D}|^{-3} \,. \end{split}$$

As a consequence, if \mathbb{A} is a one-form such that $\int A|\mathcal{D}|^{-3} = 0$, then the scale invariant term of the spectral action with or without J is exactly the same modulo a global factor of 2.

5 Differential calculus on $SU_q(2)$ and applications

5.1 The sign of \mathcal{D}

There are multiple differential calculi on $SU_q(2)$, see [33,39]. Thanks to [36, Theorem 3], the 3D and $4D_{\pm}$ differential calculi do not coincide with the one considers here: the right multiplication of one-forms by an element in the algebra A is a consequence of the chosen Dirac operator which was introduced according to some equivariance properties with respect to the duality between the two Hopf algebras $SU_q(2)$ and $U_q(su(2))$.

It is known that the Fredholm module associated to $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is one-summable since $[F, \pi(x)]$ is trace-class for all $x \in \mathcal{A}$. In fact, more can be said about F^{-1} :

Proposition 5.1. Since

$$F = \frac{1}{1-q^2} \left(\underline{\pi}(a^*) \, d\underline{\pi}(a) + q^2 \, \underline{\pi}(b) \, d\underline{\pi}(b^*) + q^2 \, \underline{\pi}(a) \, d\underline{\pi}(a^*) + q^2 \, \underline{\pi}(b^*) \, d\underline{\pi}(b) \right), \tag{56}$$

F is a central one-form modulo $OP^{-\infty}$.

¹Note that a similar result for a different spectral triple over $SU_q(2)$ when q=0 was obtained in [13, eq. (48)]

Proof. Forgetting $\underline{\pi}$, this follows from

$$a^* \delta a + q^2 b \delta b^* + q^2 a \delta a^* + q^2 b^* \delta b$$

$$= (a_+^* + a_-^*)(a_+ - a_-) + q^2 (b_+ + b_-)(b_-^* - b_+^*) + q^2 (a_+ + a_-)(a_-^* - a_+^*)$$

$$+ q^2 (b_+^* + b_-^*)(b_+ - b_-)$$

$$= [a_+^* a_+ - q^2 a_+ a_+^* + q^2 b_+^* b_+ - q^2 b_+ b_+^*] + R = (1 - q^2) + R$$
(57)

by (16) where we check that the remainder R is zero:

$$R = -\left[a_{+}^{*}a_{-} + q^{2}b_{+}^{*}b_{-}\right] + \left[a_{-}^{*}a_{+} + q^{2}b_{-}^{*}b_{+}\right] - \left[a_{-}^{*}a_{-} - q^{2}a_{-}a_{-}^{*} + q^{2}b_{-}^{*}b_{-} - q^{2}b_{-}b_{-}^{*}\right] + \left(q^{2}a_{+}a_{-}^{*} + q^{2}q_{-}^{*}b_{+}\right) - \left(a_{+}^{*}a_{-} + q^{2}b_{+}^{*}b_{-}\right),$$

thus, applying (19), (20), (21), $R = +(q^2 a_+ a_-^* + q^2 q_-^* b_+) - (a_+^* a_- + q^2 b_+^* b_-) = 0$ using commutation relations (15).

Now, replacing δ by d in (57) gives (56) since F commute with a_{\pm} , b_{\pm} and F is central by (28).

Proposition 5.2. The one-form in (56) is in fact exactly a function of the Dirac operator D:

$$\pi(a^*) d\pi(a) + q^2 \pi(b) d\pi(b^*) + q^2 \pi(a) d\pi(a^*) + q^2 \pi(b^*) d\pi(b) = \xi_q(\mathcal{D}) = F \xi_q(|\mathcal{D}|), \tag{58}$$

where $\xi_q(s) := q \frac{[2s]-2s}{[s+1/2][s-1/2]}$. Moreover, $F = \lim_{g \to 0} \xi_g(D)$.

Proof. First, let us observe that the one-form ω in (58) is invariant under the action of the $\mathcal{U}_q(su(2)) \times \mathcal{U}_q(su(2))$: $h \triangleright \omega = \epsilon(h) \omega$ for any $h \in \mathcal{U}_q(su(2)) \times \mathcal{U}_q(su(2))$. For instance, using notations of [21]

$$e \triangleright \omega = q^{\frac{1}{2}}a^*db + q^2\left(-q^{\frac{1}{2}-1}bda^* + q^{-\frac{1}{2}}bda^* - q^{-1-\frac{1}{2}}a^*db\right) = 0 = \epsilon(e)\,\omega.$$

Therefore, since both the representation π as well as the operator D are equivariant, the image of ω must be diagonal in the spinorial base. A tedious computation with the full spinorial representation π given in (10) yields

$$\begin{split} \langle v_{ml}^{j\uparrow},\,\omega\,v_{ml}^{j\uparrow}\rangle &= \frac{q^{8j+8}-q^{8j+6}-(4j+3)\,q^{4j+6}+(8j+6)\,q^{4j+4}-(4j+3)\,q^{4j+2}-q^2+1}{(q^{4j+4}-1)(q^{4j+2}-1)} = \xi_q(2j+\frac{3}{2}),\\ \langle v_{ml}^{j\downarrow},\,\omega\,v_{ml}^{j\downarrow}\rangle &= \frac{-q^{8j+4}+q^{8j+2}+(4j+1)\,q^{4j+4}-(8j+2)\,q^{4j+2}+(4j+1)\,q^{4j}+q^2-1}{(q^{4j+2}-1)(q^{4j}-1)} = -\xi_q(2j+\frac{1}{2}). \end{split}$$

These expressions have a clear q=0 limit equal respectively to 1 and -1, so $\omega \to F$ as $q\to 0$.

In the q = 1 limit, these expressions yields identically 0, which is confirmed by the fact that all one-forms are central, it could be expressed as $d(aa^* + bb^*) = d1$.

Note that since the invariant one-form we constructed differs by $OP^{-\infty}$ from F, hence any commutator with it will be itself in $OP^{-\infty}$.

We do not know if a central form ω is automatically invariant by the action of both $U_q(su(2))$, that is: $h \triangleright \omega = \epsilon(h)\omega$.

Proposition 5.3. The order one calculus up to $OP^{-\infty}$ is not universal.

Proof. Let us take the one-form ω_F from (56), which gives F. Then, for any $x \in \mathcal{A}(SU_q(2))$ we have $\underline{\pi}(x\omega_F - \omega_F x) = 0$.

Corollary 5.4. Still modulo $OP^{-\infty}$, $1 \in \pi(\Omega_u^2(\mathcal{A}))$.

Proof.
$$1 = F^2$$
 is by definition in $\pi(\Omega_n^2(\mathcal{A}))$.

In fact, one checks, using (16), (19), (22) that

$$q^2 da da^* - da^* da = 1 - q^2 (59)$$

showing again that $1 \in \pi(\Omega_n^2(\mathcal{A}))$.

Similarly, using (15) and (17), (22), (23), we get still up to $OP^{-\infty}$

$$q \, da \, db = db \, da,$$
 $q \, da \, db^* = db^* \, da,$ $da^* \, db = q \, db \, da^*,$ $da^* \, db^* = q \, db^* \, da^*$ $da \, da^* + db \, db^* = -1.$ (60)

The use of the last equality of (60) and (59) gives

Proposition 5.5. Up to $OP^{-\infty}$, F is not a (universal) closed one-form, as

$$da^* da + q^2 da da^* + q^2 db^* db + q^2 db db^* = -1 - q^2.$$
(61)

5.2 The ideal \mathcal{R}

In order to perform explicit calculations of all terms of the spectral action, we observe that each δ -one-form could be expressed in terms of $x\delta(z)y$, where z is one of the generators a, a^*, b, b^* and x, y are some elements of the algebra $\mathcal{A}(SU_q(2))$.

Then, for the computation of $\int xdzy|D|^{-1}$ we can use the trace property of the noncommutative integral to get:

$$\int x \delta(z) y \, |D|^{-1} = \int y x \delta(z) \, |D|^{-1} + \int x \delta(z) \, |D|^{-1} \delta(y) \, |D|^{-1}.$$

Therefore, the problem of calculating the tadpole-like integral could be in effect reduced to the calculation of much simpler integrals: $\int x \delta(z)$ for all generators z and the integrals of higher order in $|D|^{-1}$.

However, it appears that the calculations of higher-order terms simplify a lot, when we further restrict the algebra by introducing an ideal, which is *invisible* to the parts of integral at dimension 2 and 3. For instance, consider the space of pseudodifferential operators $T \in \Psi^0(\mathcal{A})$ of order less or equal to zero (see [16]), which satisfy

$$\oint T t |D|^{-2} = \oint t T |D|^{-2} = \oint T t |D|^{-3} = \oint t T |D|^{-3} = 0, \ \forall t \in \Psi_0^0(\mathcal{A}). \tag{62}$$

The elements a_- , b_-b_+ , $b_-b_+^*$ and their adjoints are in this space up to $OP^{-\infty}$: this is due to the fact that in Theorem 3.4, $\tau_1 \otimes \tau_1(r(x)) = 0$ when $r(x) \in \pi_{\pm}(\mathcal{A}) \otimes \pi_{\pm}(\mathcal{A}) \mod OP^{-\infty}$ contains tensor products of $\pi_{\pm}(b)$ or $\pi_{\pm}(b^*)$ since these elements are in the kernel of the grading σ .

Definition 5.6. Let R be the kernel in X of $(\sigma \otimes \sigma) \circ r$ where r is the Hopf-map defined in (30) and σ is the symbol map and let R be the vector space generated by R and RF.

Note that R is a *-ideal in X and

$$a_{-}, b_{-}b_{+}(=q^{2}b_{+}b_{-}), b_{-}b_{+}^{*} \text{ are in } \mathcal{R}.$$

By construction and Theorem 3.4, any $T \in \mathcal{R}$ satisfies (62) and \mathcal{R} is invariant by F.

Moreover, by (19), $[b_-, b_-^*] \in R$, so by (16) and (22), $a_+^* a_+ - q^2 a_+ a_+^* - (1 - q^2) \in R$ and by (23), $q a_+ b_- - b_- a_+ \in R$.

It is interesting to quote, thanks to Theorem 3.4 that if $x \in R$, then $\int F x |\mathcal{D}|^{-1} = 0$ while a priori, $\int x |\mathcal{D}|^{-1} \neq 0$.

Note that $F \in \Psi^0(\mathcal{A})$ also satisfies (62) by Theorem 3.4 while $F \notin \mathcal{R}$ since $F^2 = 1$. Moreover other elements are in \mathcal{R} like for instance $d(b^*b) = d(bb^*)$:

$$\delta(bb^*) = -\delta(aa^*) = -\delta a \, a^* - a \, \delta a^* = -(a_+ - a_-)(a_+^* + a_-^*) - (a_+ + a_-)(a_-^* - a_+^*)$$
$$= 2(a_+ a^* - a_- a_+^*)$$

is in R since $a_- \in R$ yielding $d(bb^*) \in RF$.

We do not know if \mathcal{R} is equal to the subset of the algebra generated by \mathcal{B} and $\mathcal{B}F$ satisfying (62).

Lemma 5.7. \mathcal{R} is a *-ideal in $\Psi^0(\mathcal{A})$ which is invariant by F, d, δ .

Proof. Since R is an ideal in $X = \mathcal{B} \mod OP^{-\infty}$ (see Remark 3.3), \mathcal{R} appears to be an ideal in $\Psi^0(\mathcal{A}) \subset \text{algebra generated by } \mathcal{B} \text{ and } \mathcal{B} F$. Since \mathcal{R} is invariant by F, its invariance by d follows from its invariance by δ which is true on the generators of R.

Note that, according to Theorem 4.13, $\int da |D|^{-2} = \int da |D|^{-3} = 0$ while $\int a^* da |D|^{-3} = 2$ which emphasize the role of "for all t" in (62).

Lemma 5.8. For any $t \in \Psi_0^0(A)$ and $T \in \mathcal{R}$, we have $\int t T |D|^{-1} = \int T t |D|^{-1}$.

Proof. For any $t \in \mathcal{B}$, we have $\int T t |D|^{-1} = \int t T |D|^{-1} + \int T |D|^{-1} \delta(t) |D|^{-1}$ and moreover $\int T |D|^{-1} \delta(t) |D|^{-1} = \int T \delta(t) |D|^{-2} - \int T \delta^2(t) |D|^{-3}$ which comes from

$$|D|^{-1}\delta(t)|D|^{-1} = \delta(t)|D|^{-2} + [|D|^{-1}, \delta(t)]|D|^{-1} = \delta(t)|D|^{-2} - |D|^{-1}\delta^{2}(t)|D|^{-2}$$
$$= \delta(t)|D|^{-2} - \delta^{2}(t)|D|^{-3} + |D|^{-1}\delta^{3}(t)|D|^{-3}.$$

So we get the result because T satisfies (62).

Lemma 5.9. If \simeq means equality up to the ideal \mathcal{R} , the following rules with $d(.) = [\mathcal{D}, .]$ of the first-order differential calculus hold (forgetting $\underline{\pi}$)

Moreover

$$a^* da - q^2 da \, a^* \simeq (1 - q^2) \, F,$$
 $q^2 a \, da^* - da^* \, a \simeq (1 - q^2) \, F.$ (63)

Proof. The table follows from relations (7) and Lemma 3.2 with (28) (one can also use (15).) For instance, since $a_{-} \in \mathcal{R}$, using the fact that \mathcal{R} is invariant by F,

$$b da = (b_+ + b_-)(a_+ - a_-) F \simeq (b_+ + b_-)(a_+ + a_-) F = ba F = q ab F \simeq q (a_+ - a_-) F b$$

= $q da b$

or similarly, $a^* da = (a_+^* + a_-^*)(a_+ - a_-)F \simeq (a_+^* - a_-^*)(a_+ + a_-)F = -da^* a$. The second of equivalence of (63) is just the adjoint of the first one that we prove now:

$$a^* da - q^2 da \, a^* = (a_+^* + a_-^*)(a_+ - a_-)F - q^2 (a_+ - a_-)F(a_+^* + a_-^*)$$

$$\simeq (a_+^* + a_-^*)(a_+ + a_-)F - q^2 (a_+ + a_-)(a_+^* + a_-^*)F = (a^* a - q^2 a a^*) F$$

$$= (1 - q^2) F.$$

Remark 5.10. The above written rules remain valid if dx is replaced by $\delta(x)$ and F by 1.

Working modulo \mathcal{R} simplifies the writing of a one-form:

Lemma 5.11. (i) Every one-form A can be, up to elements from \mathcal{R} , presented as

$$A \simeq x_a da + da^* x_{a^*} + x_b db + db^* x_{b^*},$$

where all x_* are the elements of A.

(ii) When A is selfadjoint, A can be written up to R (not in a unique way, though) as

$$A \simeq x_a da - da^* (x_a)^* + x_b db - db^* (x_b)^*$$
,

where x_a, x_b are arbitrary elements of A.

Proof. (i) A basis for one-forms consists of the following forms: $a^{\alpha}b^{\beta}(b^*)^{\gamma}d(a^{\alpha'}b^{\beta'}(b^*)^{\gamma'})$, where $\alpha, \alpha' \in \mathbb{Z}$ and $\beta, \gamma, \beta', \gamma' \in \mathbb{N}$.

Using the Leibniz rule and the commutation rules within the algebra (up to the \mathcal{R} according to Lemma 5.9), we reduce the problem to the case of the forms: $(a^{\alpha}b^{\beta}(b^*)^{\gamma})dx(a^{\alpha'}b^{\beta'}(b^*)^{\gamma'})$, where x can be either of the generators a, a^*, b, b^* . If x = b or $x = b^*$, the straightforward application of the rules of the differential calculus leads to the answer that the one-form could be expressed as: $a^{\alpha}b^{\beta}(b^*)^{\gamma}db$ and $db^*a^{\alpha}b^{\beta}(b^*)^{\gamma}$.

Similar considerations for the case $x = a, a^*$ lead to the remaining terms.

Note that the presentation is not unique, since there still might remain terms, which are in \mathcal{R} , for example: $b^*db + db^*b = d(bb^*) \in \mathcal{R}$.

(ii) is direct.
$$\Box$$

Next we can start explicit calculation of the integrals, beginning with the tadpole terms. Application of the Leibniz rule yields to a presentation of one-forms which is different from the one of previous lemma. Each δ -one-form could be expressed, as a finite sum of the terms $x\delta(z)y$, where z is one of the generators a, a^*, b, b^* and x, y are some elements of the algebra $\mathcal{A}(SU_q(2))$.

Proposition 5.12. For all $x, y \in \mathcal{A}(SU_q(2))$ and $z \in \{a, a^*, b, b^*\}$ we have

$$\oint x\delta(z)y\,|D|^{-1} = \oint yx\delta(z)\,|D|^{-1} + \oint x\delta(z)\delta(y)\,|D|^{-2} - \oint x\delta(z)\delta^2(y)\,|\mathcal{D}|^{-3}.$$

Proof. This is just the application of the trace property of the noncommutative integral, together with the identity: $|D|^{-1}\delta(z)|D|^{-1} = -[|D|^{-1},z]$.

Remark 5.13. The computation of tadpole-like integrals is reduced to the integrals $\int x\delta(z)|\mathcal{D}|^{-1}$ for all generators z and the integrals of higher order in $|\mathcal{D}|^{-2}$. However, the calculations of higher-order terms simplify a lot after we use the relations which hold up to the ideal \mathcal{R} : this erases parts of integral depending on $|\mathcal{D}|^{-2}$ and $|\mathcal{D}|^{-3}$. Thus, beside $\int x\delta(z)|\mathcal{D}|^{-1}$, we only need to compute $\int x\delta(z)\delta(z')|\mathcal{D}|^{-2}$ where z and z' are generators, since all the $|\mathcal{D}|^{-3}$ integrals have already been explicitly computed in section 4.6 (these integrals do not depend on q.)

Besides the tadpole, the only integrals that need to be computed are $\int A|\mathcal{D}|^{-2}$ and $\int A^2|\mathcal{D}|^{-2}$ where A is a δ -1-form. Working modulo \mathcal{R} and using again Leibniz rule, we only need to compute $\int x\delta(z)|\mathcal{D}|^{-2}$ and the previous integrals $\int x\delta(z)\delta(z')|\mathcal{D}|^{-2}$.

5.2.1 Operators L_q and M_q

In the notation $v_{l,m}^j$ of \mathcal{H} , we have already use the j dependence in (12) with $J_q v_{m,l}^j := q^j v_{m,l}^j$. Let L_q and M_q be the similar diagonal operators

$$\begin{split} L_q \, v^j_{m,l} &:= q^{2l} \, v^j_{m,l} \,, \\ M_q \, v^j_{m,l} &:= q^{2m} \, v^j_{m,l} \,. \end{split}$$

We immediately get

Lemma 5.14. For $n \in \mathbb{N}^*$, $f(L_q)^n |\mathcal{D}^{-2}| = f(M_q)^n |\mathcal{D}^{-2}| = \frac{2}{1-q^{2n}}$.

Proof. We have

$$\operatorname{Tr}\left(L_{q}^{n}|\mathcal{D}|^{-2-s}\right) = \sum_{2j=0}^{\infty} \sum_{m=0}^{2j} \sum_{l=0}^{2j+1} \langle v_{m,l}^{j}, L_{q}^{n}|\mathcal{D}|^{-2-s} v_{m,l}^{j} \rangle$$

$$= \sum_{2j=0}^{\infty} (2j+1) \frac{1-q^{2n(2j+2)}}{1-q^{2n}} d_{j+}^{-2-s} + \sum_{2j=0}^{\infty} (2j+1) \frac{1-q^{2n(2j+2)}}{1-q^{2n}} d_{j}^{-2-s}$$

$$\sim_{0} \frac{1}{1-q^{2n}} \left(\zeta(s+1, \frac{3}{2}) + \zeta(s+1, \frac{1}{2}) \sim_{e} \frac{2}{1-q^{2n}} \zeta(s+1) \right).$$

where \sim_0 means modulo a function holomorphic at 0. This gives the result for L_q^n and a similar computation can be done for M_q^n .

The interest of these operators stems in

Lemma 5.15. We have $L_qM_q \in \mathcal{R}$. Moreover,

$$\begin{split} b\,\delta b^* &\simeq M_q - L_q, \quad b^*\delta b \simeq L_q - M_q, \quad bb^* \simeq L_q + M_q, \\ a\,\delta (a^*) &\simeq -aa^* \simeq L_q + M_q - 1, \quad a^*\,\delta a \simeq a^*a \simeq 1 - q^2(L_q + M_q), \\ da\,da^* &\simeq L_q + M_q - 1, \quad da^*\,da \simeq q^2(L_q + M_q) - 1, \\ b^{n-2}(b^*)^n\,db\,db \simeq (L_q)^n + (M_q)^n, \\ b^{n-1}\,(b^*)^{n-1}\,db\,db^* \simeq -(L_q)^n - (M_q)^n, \\ b^n(b^*)^{n-2}\,db^*\,db^* \simeq (L_q)^n + (M_q)^n. \end{split}$$

Proof. Since $L_q M_q = q^2 a_- a_-^* \in \mathcal{R}$, we compute up to the ideal \mathcal{R}

$$b \, \delta b^* = (b_+ + b_-)(b_-^* - b_+^*) \simeq -b_+ b_+^* + b_- b_-^* = M_q - L_q + L_q M_q (1 - q^2) \simeq M_q - L_q$$

and similarly for the other relations.

5.2.2 Automorphisms of the algebra and symmetries of integrals

Proposition 5.16. For any $n \in \mathbb{N}^*$,

$$\int (bb^*)^n |\mathcal{D}|^{-1} = \frac{-2(1+q^{2n})}{(1-q^{2n})^2},$$

$$\int (bb^*)^n b^* \, \delta b \, |\mathcal{D}|^{-1} = \int (bb^*)^n b \, \delta b^* \, |\mathcal{D}|^{-1} = \frac{2}{1-q^{2n+2}},$$

$$\int (bb^*)^n a \, da^* \, \mathcal{D}^{-1} = \frac{-2q^{4n+2}-2q^{4n}-2q^{2n+2}+6q^{2n}}{(1-q^{2n})^2(1-q^{2n+2})},$$

$$\int (bb^*)^n a^* \, da \, \mathcal{D}^{-1} = \frac{6q^{2n+2}-2q^{2n}-2q^2-2}{(1-q^{2n})^2(1-q^{2n+2})}.$$

Note that the knowledge of these integral is enough for the computation of any term of the form $\int x \delta(z) |\mathcal{D}|^{-1}$, where z is a generator, since any other δ -one-form will be unbalanced.

To show this proposition, we will use few symmetries, properties of the ideal \mathcal{R} and replacement of δ -one-forms in terms of L_q , M_q as above.

Let U be the following unitary operator on the Hilbert space:

$$U\,v_{m,l}^{j\uparrow} = (-1)^{m+l}\,v_{l,m}^{j+\downarrow}, \quad U\,v_{m,l}^{j\downarrow} = (-1)^{m+l}\,v_{l,m}^{j-\uparrow}.$$

Then, by explicit computations we have

$$U^*aU = a$$
, $U^*a^*U = a^*$, $U^*bU = b^*$, $U^*b^*U = b$, and $U^*\mathcal{D}U = -\mathcal{D}$.

Lemma 5.17. Each noncommutative integral (55) of an element of the algebra or differential forms is (up to sign) invariant under the algebra automorphism ρ defined by

$$\rho(a) := a, \quad \rho(a^*) := a^*, \quad \rho(b) := b^*, \quad \rho(b^*) := b.$$
(64)

Proof. For any homogeneous polynomial p and any $k \in \mathbb{N}$,

$$\int p(a, a^*, b, b^*, \mathcal{D}) \, \mathcal{D}^{-k} = \int U^* p(a, a^*, b, b^*, \mathcal{D}) \, \mathcal{D}^{-n} U
= (-1)^k \int p(U^* a U, U^* a^* U, U^* b U, U^* b^* U, U^* \mathcal{D} U) \, \mathcal{D}^{-k}
= (-1)^{k+d} \int p(\rho(a), \rho(a^*), \rho(b), \rho(b^*), \mathcal{D}) \, \mathcal{D}^{-k},$$

where d is the degree of p with respect to D.

Corollary 5.18. For any $n \in \mathbb{N}$, $f(bb^*)^n b^* db \mathcal{D}^{-1} = f(bb^*)^n b db^* \mathcal{D}^{-1}$.

Lemma 5.19. For any $x, y \in \Psi^0(\mathcal{A})$,

(i)
$$\int xy |\mathcal{D}|^{-1} = \int yx |\mathcal{D}|^{-1} + \int x\delta(y) |\mathcal{D}|^{-2} - \int x\delta^2(y) |\mathcal{D}|^{-2}$$
.

(ii)
$$\int z \, x \, \mathcal{D}^{-1} y \, \mathcal{D}^{-1} = \int z \, xy \, \mathcal{D}^{-2}$$
, where $z \in \mathcal{A}$ contains b or b^* .

Proof. (i) is direct consequence of the trace property of f and the fact that OP^{-4} operators are trace-class.

ii) We calculate:

$$\int z \, x \, \mathcal{D}^{-1} y \, \mathcal{D}^{-1} = \int z \, x \, (y \, \mathcal{D}^{-1} - \mathcal{D}^{-1} [\mathcal{D}, y] \, \mathcal{D}^{-1}) \, \mathcal{D}^{-1} = \int z \, x y \, \mathcal{D}^{-2} - \int z \, x \mathcal{D}^{-1} [\mathcal{D}, y] \, \mathcal{D}^{-2} \\
= \int z \, x y \, \mathcal{D}^{-2}.$$

The last step is based on the observation that any integral with \mathcal{D}^{-3} vanishes if the expression integrated contains b or b^* .

Lemma 5.20. For any $n \in \mathbb{N}$,

(i)
$$\int (bb^*)^n b^* db \mathcal{D}^{-1} = \frac{2}{1-q^{2n+2}}$$
.

(ii)
$$\int (bb^*)^n d(bb^*) \mathcal{D}^{-1} = 0.$$

(iii)
$$f(bb^*)^n |D|^{-1} = \frac{-2(1+q^{2n})}{(1-q^{2n})^2}$$
.

Proof. (i) With n > 1, we begin with $\int d((bb^*)^n) \mathcal{D}^{-1} = 0$, which follows directly from the trace property of the noncommutative integral. Expanding the expression using Leibniz rule and the commutation

$$x\mathcal{D}^{-1} = \mathcal{D}^{-1}x + \mathcal{D}^{-1}[D, x]\mathcal{D}^{-1},$$
 (65)

we obtain

$$0 = \sum_{k=0}^{n-1} \int b^k \, db \, b^{n-k-1} (b^*)^n \, \mathcal{D}^{-1} + \sum_{k=0}^{n-1} \int b^n (b^*)^k \, db^* \, (b^*)^{n-k-1} \, \mathcal{D}^{-1}$$

$$= n \left(\int b^{n-1} (b^*)^n \, db \, \mathcal{D}^{-1} + \int b^n (b^*)^{n-1} \, db^* \, \mathcal{D}^{-1} \right)$$

$$+ \sum_{k=0}^{n-1} \int \left(b^k \, db \, \mathcal{D}^{-1} d (b^{n-k-1} (b^*)^n) \mathcal{D}^{-1} + b^n (b^*)^k \, db^* \, \mathcal{D}^{-1} d ((b^*)^{n-k-1}) \mathcal{D}^{-1} \right).$$

Using Lemma 5.19,

$$0 = n \int (bb^*)^{n-1} (b^* db + b db^*) \mathcal{D}^{-1}$$

+
$$\int \left(\frac{1}{2}n(n-1)b^{n-2}(b^*)^n db db + n^2b^{n-1} (b^*)^{n-1} db db^* + \frac{1}{2}n(n-1)b^n(b^*)^{n-2} db^* db^*\right) \mathcal{D}^{-2}.$$

The integrals with \mathcal{D}^{-2} could be easily calculated when we take restrict ourselves to calculations modulo ideal \mathcal{R} :

$$n \int (bb^*)^{n-1} (b^* db + b db^*) \mathcal{D}^{-1} = -2 \left(n(n-1) - 2n^2 + n(n-1) \right) \frac{1}{1 - q^{2n}} = 4n \frac{1}{1 - q^{2n}}.$$

Hence $f(bb^*)^{n-1}(b^*db+bdb^*)\mathcal{D}^{-1}=\frac{4}{1-q^{2n}}$, which together with Corollary 5.18 proves i). (ii) In a similar way, $f(bb^*)^{n-1}d(bb^*)\mathcal{D}^{-1}=0=f(bb^*)^{n-1}d(aa^*)\mathcal{D}^{-1}$ implies:

$$0 = \sum_{k=0}^{n-1} (bb^*)^{n-k-1} d(bb^*) (bb^*)^k \mathcal{D}^{-1}$$

$$= n \int (bb^*)^{n-1} d(bb^*) \mathcal{D}^{-1} + \frac{1}{2} n(n-1) \int (bb^*)^{n-2} d(bb^*) d(bb^*) \mathcal{D}^{-2}$$

$$= n \int (bb^*)^{n-1} d(bb^*) \mathcal{D}^{-1},$$

where in the last step we used that $d(bb^*) \in \mathcal{R}$. The identity (ii) now follows from the equality $aa^* = 1 - bb^*$.

(iii) Using Lemma 5.19, we get

$$A_n := \oint (bb^*)^n |D|^{-1} = \oint (bb^*)^n (aa^* + bb^*) |D|^{-1}$$

and we push now a^* through $|D|^{-1}$ and from cyclicity of the trace through $(bb^*)^n$,

$$= A_{n+1} + \int (bb^*)^n q^{2n} a^* a |D|^{-1} + \int (bb^*)^n q^{2n} a \delta(a^*) |D|^{-2}$$

the last term being calculated explicitly, since up to ideal \mathcal{R} , $a\delta(a^*) \simeq L_q + M_q - 1$,

$$= A_{n+1}(1 - q^{2n+2}) + q^{2n}A_n + 4\left(\frac{1}{1 - q^{2n+2}} - \frac{1}{1 - q^{2n}}\right),$$

which leads to

$$A_n(1-q^{2n}) + \frac{4}{1-q^{2n}} = A_{n+1}(1-q^{2n+2}) + \frac{4}{1-q^{2n+2}}$$
.

Assuming $A_n = \frac{f_n}{(1-q^{2n})^2}$ we have $\frac{f_n+4}{1-q^{2n}} = \frac{f_{n+1}+4}{1-q^{2n+2}}$, and taking into account that $A_0 = -2\frac{1+q^2}{(1-q^2)^2}$, we obtain $A_n = -2\frac{1+q^{2n}}{(1-q^{2n})^2}$.

Finally, to get Proposition 5.16, it remains to prove

Lemma 5.21. *For* $n \ge 1$,

$$\oint (bb^*)^n a \, da^* \, \mathcal{D}^{-1} = \frac{-2q^{4n+2} - 2q^{4n} - 2q^{2n+2} + 6q^{2n}}{(1-q^{2n})^2 (1-q^{2n+2})},$$

$$\oint (bb^*)^n a^* \, da \, \mathcal{D}^{-1} = \frac{6q^{2n+2} - 2q^{2n} - 2q^{2} - 2}{(1-q^{2n})^2 (1-q^{2n+2})}.$$

Proof. First, using Leibniz rule, (65) and Lemma 5.19 we have (for $n \ge 1$)

$$\int (bb^*)^n a \, da^* \, \mathcal{D}^{-1} = -q^{2n} \int (bb^*)^n a^* \, da - \int (bb^*)^n da \, da^* \, \mathcal{D}^{-2}.$$

Further, we use the identity (56):

$$\int (bb^*)^n \left(a^* da + q^2 a da^* + q^2 b db^* + q^2 b^* db\right) \mathcal{D}^{-1} = (1 - q^2) \int (bb^*)^n |D|^{-1}.$$

taking into account that $F \mathcal{D} = |\mathcal{D}|$.

These equations give together a system of linear equations

$$\int (bb^*)^n a \, da^* \, \mathcal{D}^{-1} + q^{2n} \int (bb^*)^n a^* \, da \, \mathcal{D}^{-1} = -4 \left(\frac{1}{1 - q^{2n+2}} - \frac{1}{1 - q^{2n}} \right),$$

$$q^2 \int (bb^*)^n a \, da^* \, \mathcal{D}^{-1} + \int (bb^*)^n a^* \, da \, \mathcal{D}^{-1} = -2(1 - q^2) \frac{1 + q^{2n}}{(1 - q^{2n})^2} - \frac{4q^2}{1 - q^{2n+2}}$$

which is solved by the expressions stated in the lemma.

5.2.3 The noncommutative integrals at $|D|^{-2}$

We need to separate this task into two problems. First, we shall to calculate all integrals $\int x \, \delta(z) |D|^{-2}$, with $x \in \mathcal{A}(SU_q(2))$ and z being one of the generators. The second problem is to calculate $\int x \, \delta(y) \, \delta(z) \, |D|^{-2}$, with both y and z being the generators $\{a, a^*, b, b^*\}$.

Lemma 5.22. The only a priori non-vanishing integrals of the type $\int x \, \delta(z) |D|^{-2}$ are for $n \in \mathbb{N}$:

$$\int (bb^*)^n b^* \delta(b) |\mathcal{D}|^{-2} = \int (bb^*)^n b \delta(b^*) |\mathcal{D}|^{-2} = 0,$$

$$\int (bb^*)^n a \delta(a^*) |\mathcal{D}|^{-2} = \frac{4q^{2n}(1-q^2)}{(q^{2n+2}-1)(1-q^{2n})}, \quad n > 0$$

$$\int (bb^*)^n a^* \delta(a) |\mathcal{D}|^{-2} = \frac{4(1-q^2)}{(1-q^{2n+2})(1-q^{2n})}.$$

Proof. Since $a\delta(a^*) \simeq L_q + M_q - 1$ and $(bb^*)^n \simeq L_q^n + M_q^n$, we get

$$(bb^*)^n a\delta(a^*) \simeq L_q^{n+1} + M_q^{n+1} - L_q^n - M_q^n$$

and the second result is obtained from Lemma 5.14. The other integrals are computed in a similar way. \Box

Lemma 5.23. The only a priori non-vanishing integrals of the type $\int x \, dy \, dz \, |D|^{-2}$ are for $n \in \mathbb{N}$:

$$\int (bb^*)^n (b^*)^2 db db |\mathcal{D}|^{-2} = \frac{4}{1 - q^{2n+4}},$$

$$\int (bb^*)^n db db^* |\mathcal{D}|^{-2} = \frac{4}{1 - q^{2n+2}},$$

$$\int (bb^*)^n (a^*b^*) (da db) |\mathcal{D}|^{-2} = 0,$$

$$\int (bb^*)^n (ab^*) (da^* db) |\mathcal{D}|^{-2} = 0,$$

$$\int (bb^*)^n (a^*b) (da db^*) |\mathcal{D}|^{-2} = 0,$$

$$\int (bb^*)^n (ab) (da^* db^*) |\mathcal{D}|^{-2} = 0,$$

$$\int (bb^*)^n (da da^*) |\mathcal{D}|^{-2} = \frac{4(q^{2n+2} - q^{2n})}{(1 - q^{2n+2})(1 - q^{2n})}, \quad n > 0$$

$$\int (bb^*)^n (da^* da) |\mathcal{D}|^{-2} = \frac{4(q^2 - 1)}{(1 - q^{2n+2})(1 - q^{2n})}.$$

Proof. This follows from Lemma 5.14 with the equivalences up to \mathcal{R} gathered in Lemma 5.15. \square

6 Examples of spectral action

It is clear from Theorem 4.3 that any one-form of the form ada, bdb, adb, a^*db , etc... do not contribute to the spectral action. Indeed, only the balanced parts of one-forms give a possibly nonzero term in the coefficients. Let us now give the values of the terms $\int A^n |\mathcal{D}|^{-p}$ and the full $\zeta_{\mathcal{D}_{\mathbb{A}}}(0)$ for few examples

Table 1: Values of noncommutative integrals

| A | $\int A \mathcal{D} ^{-3}$ | $\int A^2 \mathcal{D} ^{-3}$ | $\oint A^3 \mathcal{D} ^{-3}$ | $\int A \mathcal{D} ^{-2}$ | $\int A^2 \mathcal{D} ^{-2}$ | $\int A \mathcal{D} ^{-1}$ | $\zeta_{\mathcal{D}_{\mathbb{A}}}(0)$ |
|---------------------------------|-----------------------------|-------------------------------|--|---|---|--|---|
| a^*da b^*db ada^* bdb^* | 2 0 -2 0 | 2 0 2 0 | $ \begin{array}{c} 2 \\ 0 \\ -2 \\ 0 \end{array} $ | $ \frac{4q^2}{q^2 - 1} $ $ 0 $ $ \frac{-4}{q^2 - 1} $ $ 0 $ | $\begin{array}{c} \frac{4q^2(q^2+2)}{q^4-1} \\ \frac{-4}{q^4-1} \\ \frac{4(2q^2+1)}{q^4-1} \\ \frac{-4}{q^4-1} \end{array}$ | $\begin{array}{c} \frac{3q^2+1}{2(q^2-1)} \\ \frac{-2}{q^2-1} \\ \frac{q^2+3}{2(q^2-1)} \\ \frac{-2}{q^2-1} \end{array}$ | $\frac{11q^4 + 36q^2 + 13}{3(q^4 - 1)}$ $\frac{4q^2}{q^4 - 1}$ $\frac{13q^4 + 36q^2 + 11}{3(q^4 - 1)}$ $\frac{4q^2}{q^4 - 1}$ |

1) Clearly the spectral action depends on q: for instance,

$$\mathcal{S}(\mathcal{D}_{a^*da}, \Phi, \Lambda) = 2 \Phi_3 \Lambda^3 - 8 \Phi_2 \Lambda^2 + \frac{q^2 + 15}{2(1 - q^2)} \Phi_1 \Lambda^1 + \frac{11q^4 + 36q^2 + 13}{3(q^4 - 1)} \Phi(0).$$

2) Moreover, for $B := a \, \delta a^*$ and $A := B + B^*$, we get since $B \simeq B^* \mod \mathcal{R}$,

$$\int A^p |\mathcal{D}|^{-k} = 2^p \int B^p |\mathcal{D}|^{-k}, \quad 1 \le p \le k \le 3.$$
(66)

Thus the spectral action of the selfadjoint one-form $A := ada^* + (ada^*)^*$ is

$$\mathcal{S}(\mathcal{D}_{\mathbb{A}}, \Phi, \Lambda) = 2 \Phi_3 \Lambda^3 + 16 \Phi_2 \Lambda^2 + \frac{q^2 - 33}{2(1 - q^2)} \Phi_1 \Lambda^1 + \frac{122q^4 + 168q^2 - 2}{3(q^4 - 1)} \Phi(0).$$

3) When $B_n := (bb^*)^n b \, \delta b^*$, then by Lemma (5.15), $B_n \simeq B_n^*$, so for $A_n := B_n + B_n^*$, the equation (66) is still valid and $\int B_n^p |\mathcal{D}|^{-k}$ are all zero but $\int B_n |\mathcal{D}|^{-1} = \frac{2}{1-q^{2n+2}}$ and $\int B_n^p |\mathcal{D}|^{-2} = \frac{4}{1-q^{4n+4}}$, so

$$S(\mathcal{D}_{\mathbb{A}_n}, \Phi, \Lambda) = 2 \Phi_3 \Lambda^3 - \frac{1}{2} \Phi_1 \Lambda^1 + \frac{8}{1 + \sigma^{2n+2}} \Phi(0).$$
 (67)

Remark that this spectral action still exists as $q \to 1$!

Note however that the symmetrization process (66) is not true in general, for instance if $B:=a\,\delta b$ and $A:=B+B^*$, then $\int A^2|\mathcal{D}|^{-1}=\frac{8(q^4-q^2-1)}{(1-q^4)^2}$ while $\int B^2|\mathcal{D}|^{-1}=0$ or $\int [B,\,B^*]|\mathcal{D}|^{-1}=\frac{4}{1-q^4}$.

4) The spectral action can be also independent of q: for instance, if $\mathbb{A} = \frac{1}{1-q^2} \xi(\mathcal{D})$ is the q-dependent selfadjoint one-form given in (58), then,

$$\mathcal{S}(\mathcal{D}_{\mathbb{A}}, \Phi, \Lambda) \, = \, 2 \, \Phi_3 \, \Lambda^3 - 8 \, \Phi_2 \, \Lambda^2 + \tfrac{15}{2} \, \Phi_1 \, \Lambda^1 - \tfrac{13}{3}.$$

7 The commutative sphere \mathbb{S}^3

Since $SU(2) \simeq \mathbb{S}^3$, we get a concrete spinorial representation of the algebra $\mathcal{A} := C^{\infty}(\mathbb{S}^3)$ on the same Hilbert space \mathcal{H} and same Dirac operator \mathcal{D} with (10) where q = 1 which means that q-numbers are trivial: $[\alpha] = \alpha$. So

$$\pi(a) |j\mu n\rangle\rangle := \alpha_{j\mu n}^{+} |j^{+}\mu^{+}n^{+}\rangle\rangle + \alpha_{j\mu n}^{-} |j^{-}\mu^{+}n^{+}\rangle\rangle,$$

$$\pi(b) |j\mu n\rangle\rangle := \beta_{j\mu n}^{+} |j^{+}\mu^{+}n^{-}\rangle\rangle + \beta_{j\mu n}^{-} |j^{-}\mu^{+}n^{-}\rangle\rangle,$$

$$\pi(a^{*}) |j\mu n\rangle\rangle := \tilde{\alpha}_{j\mu n}^{+} |j^{+}\mu^{-}n^{-}\rangle\rangle + \tilde{\alpha}_{j\mu n}^{-} |j^{-}\mu^{-}n^{-}\rangle\rangle,$$

$$\pi(b^{*}) |j\mu n\rangle\rangle := \tilde{\beta}_{j\mu n}^{+} |j^{+}\mu^{-}n^{+}\rangle\rangle + \tilde{\beta}_{j\mu n}^{-} |j^{-}\mu^{-}n^{+}\rangle\rangle$$
(68)

where

$$\alpha_{j\mu n}^{+} := \sqrt{j + \mu + 1} \begin{pmatrix} \frac{\sqrt{j + n + 3/2}}{2j + 2} & 0\\ \frac{\sqrt{j - n + 1/2}}{(2j + 1)(2j + 2)} & \frac{\sqrt{j + n + 1/2}}{2j + 1} \end{pmatrix},$$

$$\alpha_{j\mu n}^{-} := \sqrt{j - \mu} \begin{pmatrix} \frac{\sqrt{j - n + 1/2}}{2j + 1} & -\frac{\sqrt{j + n + 1/2}}{2j(2j + 1)}\\ 0 & \frac{\sqrt{j - n - 1/2}}{2j} \end{pmatrix},$$

$$\beta_{j\mu n}^{+} := \sqrt{j + \mu + 1} \begin{pmatrix} \frac{\sqrt{j - n + 3/2}}{2j + 2} & 0\\ -\frac{\sqrt{j + n + 1/2}}{(2j + 1)(2j + 2)} & \frac{\sqrt{j - n + 1/2}}{2j + 1} \\ -\frac{\sqrt{j - n + 1/2}}{2j + 1} & -\frac{\sqrt{j - n + 1/2}}{2j(2j + 1)} \\ 0 & -\frac{\sqrt{j + n - 1/2}}{2j} \end{pmatrix},$$

with $\tilde{\alpha}^{\pm}_{j\mu n} := (\alpha^{\mp}_{j^{\pm}\mu^{-}n^{-}})^{*}, \, \tilde{\beta}^{\pm}_{j\mu n} := (\beta^{\mp}_{j^{\pm}\mu^{-}n^{+}})^{*}.$

Note that the representation on the vectors $v_{m,l}^j$ is not as convenient as in (11). One can check that the generators $\pi(a)$, $\pi(b)$ and their adjoint commute and that $[x, [\mathcal{D}', y]] = 0$ for any $x, y \in \mathcal{A}$.

7.1 Translation of Dirac operator

In general the Dirac operator is defined in a more symmetric way than that we did. So, although not absolutely necessary here, we define for the interested reader the unbounded self-adjoint translated operator \mathcal{D}' on \mathcal{H} by the constant λ as

$$\mathcal{D}' := \mathcal{D} + \lambda$$
.

For instance, this gives for $\lambda = -\frac{1}{2}$ in the case of \mathbb{S}^3 , see [32], $\mathcal{D}' v_{m,l}^j = (2j+1) \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} v_{m,l}^j$ so $v_{m,l}^j$ is an eigenvector of $|\mathcal{D}'|$.

As the following lemma shows, the computation of noncommutative integrals involving \mathcal{D} can be reduced to the computation of certain noncommutative integrals involving \mathcal{D}' :

Lemma 7.1. If $f'T := \operatorname{Res}_{s=0} \operatorname{Tr} (T|\mathcal{D}'|^{-s})$, then for any 1-form A on a spectral triple of dimension n,

$$\int A |\mathcal{D}|^{-(n-2)} = \int' A |\mathcal{D}'|^{-(n-2)} + \lambda (n-2) \int' A \mathcal{D}' |\mathcal{D}'|^{-n} + \lambda^2 \frac{(n-1)(n-2)}{2} \int' A |\mathcal{D}'|^{-n},
\int A \mathcal{D}^{-(n-2)} = \int' A \mathcal{D}'^{-(n-2)} + \lambda (n-2) \int' A \mathcal{D}'^{-(n-1)} + \lambda^2 \frac{(n-1)(n-2)}{2} \int' A \mathcal{D}'^{-n}.$$

Proof. Recall from [22, Proposition 4.8] that for any pseudodifferential operator P,

$$\int P|\mathcal{D}|^{-r} = \operatorname{Res}_{s=0} \operatorname{Tr} \left(P|\mathcal{D}|^{-r}|\mathcal{D}'|^{-s} \right).$$

Moreover, by [22, Lemma 4.3], for any $s \in \mathbb{C}$ and $N \in \mathbb{N}^*$

$$|\mathcal{D}|^{-s} = |\mathcal{D}'|^{-s} + \sum_{p=1}^{N} K_{p,s} Y^p |\mathcal{D}'|^{-s} \mod OP^{-N-1-\Re(s)}$$
(69)

where $Y = \sum_{k=1}^{N} \frac{(-1)^{k+1}}{k} (-2\lambda \mathcal{D}' + \lambda^2)^k \mathcal{D}'^{-2k} \mod OP^{-N-1}$ and $K_{p,s}$ are complex numbers that can be explicitly computed. Precisely, we find $K_{p,s} = (-\frac{s}{2})^p V(p)$ where V(p) is the volume of the p-simplex. Since the spectral dimension is n, we work modulo $OP^{-(n+1)}$, and for s = n-2, we get from (69): $|\mathcal{D}|^{-(n-2)} = |\mathcal{D}'|^{-(n-2)} + \lambda(n-2)\mathcal{D}'|\mathcal{D}'|^{-n} + \lambda^2 \frac{(n-1)(n-2)}{2} |\mathcal{D}'|^{-n} \mod OP^{-(n+1)}$. As a consequence, we have for $P \in OP^0$ (the OP^0 spaces are the same for \mathcal{D} or \mathcal{D}'),

$$\int P|\mathcal{D}|^{-(n-2)} = \int' P|\mathcal{D}'|^{-(n-2)} + \lambda(n-2)\int' P\mathcal{D}'|\mathcal{D}'|^{-n} + \lambda^2 \frac{(n-1)(n-2)}{2}\int' P|\mathcal{D}'|^{-n}.$$

Since A and AF are in OP^0 , we get both formulae.

7.2 Tadpole and spectral action on \mathbb{S}^3

We consider now the commutative spectral triple $(C^{\infty}(\mathbb{S}^3), \mathcal{H}, \mathcal{D})$. It is 1-summable since $\langle \langle j\mu n \, s \, | \, [F, \pi(x)] \, | j\mu n \, s \rangle \rangle = 0$ when $x = a, a^*, b, b^*$ for any $j, \mu, n, s = \uparrow, \downarrow$. All integrals of above lemma are zero for \mathbb{S}^3 :

Proposition 7.2. There is no tadpole of any order on the commutative real spectral triple $(C^{\infty}(\mathbb{S}^3), \mathcal{H}, \mathcal{D})$. In fact, for any one-form or δ -one-form A, $\oint A\mathcal{D}^{-p} = 0$ for $p \in \{1, 2, 3\}$.

Proof. We first want to prove $\int A\mathcal{D}^{-p} = 0$ for $p \in \{1, 2, 3\}$ and any one-form A. Since the representation is real, that is any matrix elements of the generators are real, so must be the trace of $A\mathcal{D}^{-p}$. Hence $\int A\mathcal{D}^{-p} = \int A^*\mathcal{D}^{-p}$.

The reality operator J introduced in (29) satisfies, when q=1, the commutative relation $JxJ^{-1}=x^*$ for $x\in\mathcal{A}$. Thus $JAJ^{-1}=-A^*$, so $\int A\mathcal{D}^{-p}=\int J\left(A^*\mathcal{D}^{-p}\right)J^{-1}=-\int A^*\mathcal{D}^{-p}$ and $\int A\mathcal{D}^{-p}=0$. Similar proof for the δ -form AF.

For any selfadjoint one-form A, $\mathcal{D}_A := \mathcal{D} + \widetilde{A} = \mathcal{D}$. Thus, the spectral action for the real spectral triple $(C^{\infty}(\mathbb{S}^3), \mathcal{H}, \mathcal{D})$ for \mathcal{D}_A is trivialized by

$$\mathcal{S}(\mathcal{D}_A, \Phi, \Lambda) = 2 \Phi_3 \Lambda^3 - \frac{1}{2} \Phi_1 \Lambda^1 + \mathcal{O}(\Lambda^{-1}). \tag{70}$$

But it is more natural to compare with the spectral action of $\mathcal{D}+A$. This is obtained respectively from Lemma 4.18 and general heat kernel approach [27]:

$$S(D+A,\Phi,\Lambda) \,=\, 2\,\Phi_3\,\Lambda^3 + \int |{\cal D}+A|^{-1}\,\Phi_1\,\Lambda^1 + {\cal O}(\Lambda^{-1})$$

since all terms of (2) in Λ^{n-k} are zero for k odd and $\zeta_{\mathcal{D}+A}(0)=0$ when n is odd: as a verification, $\int |\mathcal{D}+A|^{-2}$ is zero according to [22, Lemma 4.10], Lemmas 4.18 and Proposition 7.2. Similarly, $\zeta_{\mathcal{D}+A}(0)=0$ because in (3), all terms with k odd are zero (same proof as in Proposition 7.2) but for k even, it is not that easy to show that $\int A\mathcal{D}^{-1}A\mathcal{D}^{-1}=0$. Moreover, the curvature term does not depend on A:

Lemma 7.3. For any one-form A on a commutative spectral triple of dimension n based on a compact Riemannian spin^c manifold without boundary, we have

$$\oint |\mathcal{D} + A|^{-(n-2)} = \oint |\mathcal{D}|^{-(n-2)}.$$
(71)

Proof. Follows from [23, first formula page 511] with $\rho := A = A^*$, $N(\rho) = \rho$ (the constraint $J\rho J^{-1} = \pm \rho$ is not used.)

One can also use
$$[15, Proposition 1.149]$$
.

From [22, Lemma 4.10] $f | \mathcal{D} + A|^{-(n-2)} = f |\mathcal{D}|^{-(n-2)} + \frac{n(n-2)}{4} f(AF)^2 |\mathcal{D}|^{-3} + \frac{(n-2)^2}{4} f A^2 |\mathcal{D}|^{-3}$ using $X := A\mathcal{D} + \mathcal{D}A + A^2$ and $[|\mathcal{D}|, A] \in OP^0$, but again, it is not that easy to show that the last two terms cancelled: for instance here, for $B = b[\mathcal{D}, b^*]$, we obtain by direct computation (using the easiest translated Dirac operator \mathcal{D}')

$$\operatorname{Tr}(B^{2}|\mathcal{D}'|^{-3-s}) = \operatorname{Tr}((B^{*})^{2}|\mathcal{D}'|^{-3-s}) = \frac{1}{2}\operatorname{Tr}(BB^{*}|\mathcal{D}'|^{-3-s}) = \frac{1}{2}\operatorname{Tr}(B^{*}B|\mathcal{D}'|^{-3-s})$$
$$= \frac{4}{3}\sum_{2j\in\mathbb{N}}\frac{j+1}{(2j+1)^{2+s}},$$

so $\int B^2 |\mathcal{D}'|^{-3} = \frac{2}{3}$. Similarly, one checks that $\int (BF)^2 |\mathcal{D}|^{-3} = \frac{1}{2} \int BFB^*F |\mathcal{D}|^{-3} = -\frac{2}{9}$. Thus if $A := B + B^*, \int A^2 |\mathcal{D}|^{-3} = \int A^2 |\mathcal{D}'|^{-3} = 4$ and $\int (AF)^2 |\mathcal{D}|^{-3} = -\frac{4}{3}$ which yields to (71). Thus for any one-form A on the 3-sphere,

$$\mathcal{S}(\mathcal{D} + A, \Phi, \Lambda) = 2 \Phi_3 \Lambda^3 - \frac{1}{2} \Phi_1 \Lambda^1 + \mathcal{O}(\Lambda^{-1}, A)$$

which as (70) is not identical to (67) which contains a nonzero constant term Λ^0 for q=1.

8 Conclusion

We computed in this paper the full spectral action on the $SU_q(2)$ spectral triple of [21] with the reality operator J (notice the change of definition for pseudodifferential operators.) The dimension spectrum being a finite set, there is only a finite number of terms in the spectral action expansion. The tadpole hypothesis is not satisfied on $SU_q(2)$. We saw that that the action depends on q and the limit $q \to 1$ does not exist automatically. When it exists, such limit does not lead to the associated action on the commutative sphere \mathbb{S}^3 . The sign F of the Dirac operator has special properties: first, it commutes modulo $OP^{-\infty}$ with elements of the algebra, and second, it can be seen as a one-form, giving terms independent of q in the spectral action. Here, we were interested in the computation of the spectral action of a quantum group. Naturally, it would be interesting to investigate other related cases like the Podleś spheres [17, 19] or the Euclidean quantum spheres [20, 35], especially the 4-sphere [18].

References

- [1] C. Bär, "The Dirac operator on homogeneous spaces and its spectrum on 3-dimensional lens spaces", Arch. Math. **59** (1992), 65–79.
- [2] P.N. Bibikov and P.P. Kulish, "Dirac operators on the quantum group $SU_q(2)$ and the quantum sphere", Zap. Nauchn. Sem. St. Petersburg. Otdel. Mat. Inst. Steklov. **245** (1997), 49, Vopr. Kvant. Teor. Polya i Stat. fiz. **14** (1997), 49–65; translated in J. Math. Sci. **100** (2000), 2039–2050.
- [3] L. Carminati, B. Iochum and T. Schücker, "Noncommutative Yang-Mills and noncommutative relativity: a bridge over troubled water", Eur. Phys. J. C 8 (1999), 697–709.
- [4] P. S. Chakraborty and A. Pal, "Equivariant spectral triples on the quantum SU(2) group", K-Theory **28** (2003), 107–126.
- [5] P. S. Chakraborty and A. Pal, "On equivariant Dirac operator for $SU_q(2)$ ", Proc. Indian Acad. Sci. **116** (2003), 531–541.
- [6] P. S. Chakraborty and A. Pal, "Spectral triples and associated Connes-de Rham complex for the quantum SU(2) and the quantum sphere," Commun. Math. Phys. **240** (2003), 447–456.
- [7] A. Chamseddine and A. Connes, "The spectral action principle", Commun. Math. Phys. **186** (1997), 731–750.
- [8] A. Chamseddine and A. Connes, "Inner fluctuations of the spectral action", J. Geom. Phys. **57** (2006), 1–21.
- [9] A. Chamseddine, A. Connes and M. Marcolli, "Gravity and the standard model with neutrino mixing", [arXiv:hep-th/0610241].
- [10] A. Connes, Noncommutative Geometry, Academic Press, London and San Diego, 1994.
- [11] A. Connes, "Geometry from the spectral point of view", Lett. Math. Phys. **34** (1995), 203–238.

- [12] A. Connes, "Noncommutative geometry and reality", J. Math. Phys. 36 (1995), 6194–6231.
- [13] A. Connes, "Cyclic cohomology, quantum group symmetries and the local index formula for $SU_q(2)$ ", J. Inst. Math. Jussieu **3** (2004), 17–68.
- [14] A. Connes and G. Landi, "Noncommutative manifolds, the instanton algebra and isospectral deformations", Commun. Math. Phys. **221** (2001), 141–159.
- [15] A. Connes and M. Marcolli, *Noncommutative Geometry, Quantum Fields and Motives*, to appear.
- [16] A. Connes and H. Moscovici, "The local index formula in noncommutative geometry", Geom. Funct. Anal. 5 (1995), 174–243.
- [17] F. D'Andrea and L. Dąbrowski, "Local index formula on the equatorial Podleś sphere", Lett. Math. Phys. **75** (2006), 235–254.
- [18] F. D'Andrea, L. Dąbrowski and G. Landi, "The isospectral Dirac operator on the 4-dimensional quantum Euclidean sphere", arXiv:math/0611100.
- [19] F. D'andrea, L. Dabrowki, G. Landi and E. Wagner, "Dirac operators on all Podleś spheres", J. Noncommut. Geom. 1 (2007), 213–239.
- [20] L. Dabrowski, "Geometry of quantum spheres", J. Geom. Phys. 56 (2005), 86–107.
- [21] L. Dąbrowski, G. Landi, A. Sitarz, W. van Suijlekom and J. Várilly, "The Dirac operator on $SU_q(2)$ ", Commun. Math. Phys. **259** (2005), 729–759.
- [22] D. Essouabri, B. Iochum, C. Levy and A. Sitarz, "Spectral action on noncommutative torus", J. Noncommut. Geom. 2 (2008), 53–123.
- [23] R. Estrada, J. M. Gracia-Bondía and J. C. Várilly, "On summability of distributions and spectral geometry", Commun. Math. Phys. 191 (1998), 219–248.
- [24] V. Gayral and B. Iochum, "The spectral action for Moyal plane", J. Math. Phys. 46 (2005), no. 4, 043503, 17 pp.
- [25] V. Gayral, B. Iochum and J. C. Várilly, "Dixmier traces on noncompact isospectral deformations", J. Funct. Anal. **237** (2006), 507–539.
- [26] V. Gayral, B. Iochum and D. V. Vassilevich, "Heat kernel and number theory on NC-torus", Commun. Math. Phys. **273** (2007), 415–443.
- [27] P. B. Gilkey, Asymptotic Formulae in Spectral Geometry, Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [28] D. Goswami, "Some noncommutative geometric aspects of $SU_q(2)$ ", math-ph/018003.
- [29] J. M. Gracia-Bondía, J. C. Várilly and H. Figueroa, *Elements of Noncommutative Geometry*, Birkhäuser Advanced Texts, Birkhäuser, Boston, 2001.
- [30] H. Grosse and R. Wulkenhaar, "8D-spectral triple on 4D-Moyal space and the vacuum of noncommutative gauge theory", arXiv:0709.0095.

- [31] E. Hawkins and G. Landi, "Fredholm modules for quantum Euclidean spheres", J. Geom. Phys. **49** (2004), 272–293.
- [32] Y. Homma, "A representation of Spin(4) on the eigenspinors of the Dirac operator on S^3 ", Tokyo J. Math. **23** (2000), 453–472.
- [33] A. Klimyk and K. Schmüdgen, Quantum Groups and Their Representations, Text and Monographs in Physics, Springer-Verlag, Berlin, 1997.
- [34] M. Knecht and T. Schücker, "Spectral action and big desert", Phys. Lett. B **640** (2006), 272–277.
- [35] G. Landi, "Noncommutative spheres and instantons", in *Quantum field theory and non-commutative geometry*, U. Carow-Watamura, Y. Maeda, S. Watamura, Lecture Notes in Physics, Springer, 2005, 3–56.
- [36] K. Schmüdgen, "Commutator representations of differential calculi on the quantum group $SU_q(2)$ ", J. Geom. Phys. **31** (1999), 241–264.
- [37] W. van Suijlekom, *The Geometry of Noncommutative Spheres and their Symmetries*, PhD thesis, Trieste 2005.
- [38] W. van Suijlekom, L. Dąbrowski, G. Landi, A. Sitarz and J. C. Várilly, "The local index formula for $SU_q(2)$ ", K-Theory **35** (2005), 375–394.
- [39] S. Woronowicz, "Twisted SU(2) group. An example of a non-commutative differential calculus", Publ. RIMS, Kyoto Univ. **23** (1987), 117–181.