

Photoassociation of a Bose-Einstein Condensate near a Feshbach Resonance

M. Junker, D. Dries, C. Welford, J. Hitchcock, Y. P. Chen and R. G. Hulet

Department of Physics and Astronomy and Rice Quantum Institute, Rice University, Houston, TX 77251, USA

(Dated: February 25, 2019)

We measure the effect of a magnetically-tuned Feshbach resonance on the rate and light-induced frequency shift of a photoassociation resonance in ultracold ${}^7\text{Li}$. The photoassociation rate constant depends strongly on magnetic field, varying by more than a factor of 10^4 for fields near the Feshbach resonance. Intensity-dependent saturation and rollover in the rate constant are observed in a thermal gas, while saturation is observed for the first time in a Bose-Einstein condensate. The frequency shift is also strongly field-dependent and exhibits an anomalous blue-shift for fields just below the Feshbach resonance.

PACS numbers: 03.75.Nt, 34.50.Rk, 33.80.Gj, 34.50.-s

Photoassociation (PA) and magnetically-tuned Feshbach resonance are two ways in which gases of ultracold atoms may be connected to the molecular bound states of their underlying two-body interaction potentials. A Feshbach resonance (FR) is realized by tuning a molecular bound state near the scattering threshold [1]. The bound state perturbs the scattering state which can, in turn, strongly affect the rate of PA [2], as was seen experimentally [3, 4]. Heightened interest in creating ultracold molecules via the association of ultracold atoms motivates a fundamental understanding of association processes and especially, of their interplay when occurring together. The rate of association of an Bose-Einstein condensate (BEC) is of particular interest, as a universal regime is predicted at extremely high rates of association where the rate is limited by many-body effects, independent of the underlying microscopic mechanisms [5, 6, 7, 8].

Several groups have measured absolute PA rate constants that could be compared with theory [9, 10, 11, 12, 13]. Of these, however, only two were performed with a quantum degenerate gas. In ref. [9], the rate of PA of a sodium BEC was found to increase linearly with intensity, with no indication of saturation. We previously investigated PA of ${}^7\text{Li}$ at the critical temperature for condensation T_c , with a negligibly small condensate fraction, and observed saturation at a value consistent with the maximum two-body collision rate imposed by quantum mechanical unitarity [10]. Other experiments using thermal gases have made similar observations [12, 14, 15, 16]. A light-induced spectral red shift of the PA resonance position has been observed [9, 10, 17, 18], and ascribed to light-induced coupling to the continuum [19, 20]. In this paper, measurements of the PA rate constant, K_p , and the spectral shift are presented in the vicinity of a magnetic FR of ${}^7\text{Li}$ for both a thermal gas and a BEC. The effect of the FR on K_p is striking as the measured values of K_p vary by more than a factor of 10^4 for fields near the FR. By exploiting the huge FR-enhancement, the first evidence for saturation of the rate of association in a condensate is obtained. Furthermore, a light-induced

spectral *blue* shift is observed for the first time.

Atomic ${}^7\text{Li}$ in the $|F=1, m_F=1\rangle$ state is prepared in a single-beam optical dipole trap, as previously described [21, 22]. A uniform magnetic field can be set between 480–900 G to access both sides of the Feshbach resonance located near 736 G [23]. Efficient evaporative cooling is achieved by setting the field to 715 G where the scattering length $a \simeq +150 a_o$ (a_o is the Bohr radius), and reducing the optical trap power. Either Bose condensates, with no discernable thermal fraction, or thermal gases, with $T > T_c$, can be produced depending on the final optical trap depth.

The PA laser is tuned to resonance with a vibrational level v of the $1^3\Sigma_g^+$ excited molecular state. For most of the measurements we tune to the $v = 83$ level, located 60 GHz below the $2^2S_{1/2} + 2^2P_{1/2}$ dissociation limit [24]. The $v = 83$ level is a good compromise between strong free-bound overlap and relatively weak off-resonance scattering from the atomic transition. Excited molecules created by the PA laser decay into pairs of energetic atoms that generally escape the trap. The duration of the PA laser pulse τ is adjusted at each intensity and field to maintain the fractional loss of atoms between 25–35%. The number of remaining atoms following the laser pulse is measured using on-resonant absorption imaging.

The on-resonance rate constant, K_p , is defined by the time evolution of the density distribution: $\dot{n}(t, \mathbf{r}) = -K_p n^2(t, \mathbf{r})$. For the BEC data (Fig. 4), τ is much less than either trap period, and collisional redistribution during the PA pulse can be neglected. In this case, we solve for the time-dependent density distribution analytically and extract K_p by spatially integrating the density and matching the observed and calculated loss after a time τ [9]. For much of the thermal data (Figs. 1 and 3), however, τ is comparable to the radial trap period and redistribution is not negligible. Without accounting for the details of the time evolution, we make two estimates for K_p for this data: first, by assuming no redistribution, as for the BEC case, and secondly, by taking the opposite limit where the distribution remains Boltzmann throughout. Since the true value lies between these two extremes,




FIG. 1: Rate constants for resonant PA to $v=83$ for a thermal gas. The intensity I was fixed at 1.65 W/cm^2 , while τ was adjusted between 0.07 and 270 ms to keep the fractional loss approximately constant. The fitted temperatures range between 9–18 μK . The spread in initial peak density, 0.5×10^{12} to $1 \times 10^{13} \text{ cm}^{-3}$, reflects a significant decrease in density near the FR, which we ascribe to enhanced three-body loss. The axial and radial trapping frequencies were 20 Hz and 3 kHz, respectively. The systematic uncertainty in K_p of 25% reflects the uncertainty in the evolution of the density, as described in the text, as well as uncertainties in probe detuning and intensity. The measurement of the highest values of K_p (near 750 G) are saturated (Fig. 3) and are therefore lower limits, while the lowest values of K_p should be considered upper limits, since the statistical uncertainties in these points are large. The inset shows K_p (same scale as main figure) for three different excited state vibrational levels.

we take K_p to be the average of the two estimates. Due to the small fractional loss, we find the difference between the two extremes to also be small ($<25\%$), and account for it in the stated uncertainties.

Figure 1 shows the field dependence of K_p on both sides of the FR in the case of a thermal gas. The data show a pronounced dip in K_p at 710 G, and a maximum, which is more than 10^4 times larger, at 745 G. Measurements could not be made closer to the FR due to high inelastic collisional loss rates, presumably from a three-body process. A condensate could not be probed on the high-field side of the FR due to its instability for $a < 0$, but at a given field on the low-field side of the FR, K_p for a condensate was found to be approximately half that of a thermal gas, consistent with quantum statistics [25].

The variation in K_p can be understood in terms of the Franck-Condon principle and the dependence of the scattering state wavefunction $f(R)$ with field (R is the internuclear separation). PA predominately occurs at the Condon radius, R_C , defined as the outer classical turning point of the excited vibrational energy state.

Since $K_p \propto |f(R_C)|^2$, a node in $f(R)$ at R_C results in a minimum in K_p . Such minima have been previously observed [26], although with much lower contrast. For sufficiently low T that the collisional wave vector k satisfies $k(R-a) \ll 1$, the asymptotic scattering state $f(R) \sim k^{1/2}(R-a)$, and nodes in $f(R)$ appear at $R \simeq a$ [27]. The inset to Fig. 1 shows that the location of the K_p minima shift to higher fields (larger a) with increasing v , consistent with increasing R_C with v . The peak in K_p near 740 G, on the other hand, is a manifestation of $f(R)$ when $|a|$ is large. The ratio of the asymptotic values of K_p on the high and low field sides of the FR is ~ 3.5 , which can be understood from the ratio of $|f(R)|^2$ evaluated using the values of a at the two asymptotic fields. Using a coupled-channel calculation [28] with previously determined potentials for lithium [29], we find the scattering lengths at 600 G and 900 G to be $a_l \simeq 8 a_o$ and $a_h \simeq -43 a_o$, respectively, and $R_C \simeq 103 a_o$ for $v = 83$. With these values, the expected ratio is $(R_C - a_h)^2 / (R_C - a_l)^2 \simeq 2.4$, in reasonable agreement with measurement.

The effect of the FR on $f(R)$ is also evident in the light-induced spectral shifts. We measured spectral shifts for both a thermal gas and a condensate for fields around the FR. The location of a PA resonance at a particular field and intensity I is determined by fitting the resonance lineshape to a Lorentzian. By taking resonance curves for several values of I , we observe that the spectral shift is linear in I , and extract the slope, Σ . Figure 2 displays measurements of Σ for approximately the same range of fields as in Fig. 1. Far from the FR, the shift is red ($\Sigma < 0$), in agreement with all previous measurements [9, 10, 17, 18]. As the FR is approached from low field, however, Σ changes sign, becoming large and positive just below the FR, and large and negative just above it in a dispersive-like manner. The field where Σ vanishes (710 G), coincides with the location of the zero in K_p shown in Fig. 1. To within our uncertainties, we find no difference in Σ between a condensate and a thermal gas.

The spectral shift is a consequence of light-induced mixing of nearby molecular states into $f(R)$ [20]. In all previous experiments, the shift was dominated by coupling to the continuum, resulting in a red shift. In the case of a FR, however, the shift is affected by the underlying closed-channel molecular state responsible for the FR. For fields below the FR, the contribution to the shift from this bound state is positive, while above resonance it is negative. The bound state contribution dominates for fields close to the FR. It is shown in ref. [20] that like K_p , the light shift may be understood in terms of $f(R)$, but in this case, the dependence is linear, $\Sigma \propto -f(R_C)$.

The data of Fig. 1 demonstrate that K_p can be extraordinarily large near the FR, making it an ideal system for exploring saturation. Figure 3 shows measured values of K_p vs. I for a thermal gas for a field (755 G) just above the FR. The rate saturates at $I \simeq 5 \text{ W/cm}^2$, while at




FIG. 2: Slope of the intensity-dependent spectral light shift Σ . Data points are obtained from both thermal gases and condensates. The error bars are the standard error of the fitted slope at each field. For most of the data, this statistical uncertainty is smaller than the size of the plotted data points. The systematic uncertainty in Σ is 10% due to uncertainty in I .

higher intensities, K_p decreases.

The data of Fig. 3 can be analyzed in terms of a Fano model of a bound state coupled to a continuum, where the on-resonance intensity-dependent loss rate coefficient can be written as [20]

$$K_p(I) = 4K_{max} \frac{I \cdot I_{sat}}{(I + I_{sat})^2}, \quad (1)$$

where K_{max} is the maximum loss rate coefficient and I_{sat} is the corresponding saturation intensity that accounts for broadening of the PA resonance. Because of the low temperatures in our experiment we neglect any energy dependence of K_p . The solid line in Fig. 3 is a fit of the data to Eq. 1. The data clearly show the predicted rollover in K_p that occurs when $I > I_{sat}$. We also observe significant power broadening that accompanies saturation. By fitting the measured lineshape to a power-broadened Lorentzian at 35 W/cm², for example, we obtain $I_{sat} = 4.3$ W/cm², in good agreement with the fit to Eq. 1.

The theory of Bohn and Julienne provides an estimate of the unitarity-limited maximum rate constant $K_u = 2k_B T \Lambda^3 / h$, where h is Planck's constant and Λ is the usual thermal de Broglie wavelength, but using the reduced mass $m/2$ rather than the atomic mass m [20]. The factor of 2 accounts for losing both atoms per PA event. For $T = 16$ μ K, $K_u = 8.5 \times 10^{-9}$ cm³/s, in excellent agreement with the observed maximum.

Figure 4 shows the measured intensity dependence of K_p for a BEC with no discernable thermal fraction. The




FIG. 3: K_p vs. I in a thermal gas at 755 G. $T \simeq 16$ μ K and the peak density is 7.5×10^{11} cm⁻³. τ is between 210–250 μ s for all of the data except at the lowest intensity, for which $\tau = 2$ ms. The error bars are the statistical uncertainties from the fitted T . The systematic uncertainties are 30% for K_p and 10% for I . The solid line is a weighted fit to the data using Eq. 1, with the fitted parameters $K_{max} = 9.5 \times 10^{-9}$ cm³/s and $I_{sat} = 4.1$ W/cm². A linear fit to the slope for small I gives $8.2 \pm 1.1 \times 10^{-9}$ cm³ s⁻¹/(W cm⁻²), where the uncertainty refers only to statistical uncertainty.

field of 732 G was chosen as a compromise between large K_p and a sufficiently small inelastic loss rate [30]. At this field, $a \simeq 1000 a_o$. By achieving extremely large FR-enhanced PA rates, saturation was observed in a condensate for the first time. The maximum K_p value of 1.4×10^{-7} cm³/s is nearly a factor of 10 larger than any previous PA rate measurement [10].

Various theories of saturation may be most easily compared with the data by expressing K_p in terms of a characteristic length, L , as $K_p = (\hbar/m)L$. For the unitarity limit, L should correspond to the smallest length scale of the condensate. If we take $L \sim 2R_{TF}$, where $R_{TF} \simeq 10$ μ m is the radial Thomas-Fermi radius, we obtain $K_u \sim 1.8 \times 10^{-7}$ cm³/s, in reasonable agreement with the measured value of 1.4×10^{-7} cm³/s.

Another relevant length scale is the average interatomic separation, evaluated at the peak density n_o . Taking $L = n_o^{-1/3}$, we obtain the “rogue photodissociation” rate limit, $K_{pd} \sim (\hbar/m)n_o^{-1/3}$, a universal regime where saturation occurs when excited molecules dissociate into the hot pair continuum rather than back into the atomic condensate [5]. Taking $n_o = 1.6 \times 10^{12}$ cm⁻³ gives $K_{pd} \sim 8 \times 10^{-9}$ cm³/s. Although there is an unknown numerical prefactor, our measured K_{max} is nearly 20 times larger than K_{pd} . According to the theory of ref. [8], however, the high-intensity region of Fig. 4 is well into the dissociative regime where K_p is no longer




FIG. 4: K_p vs. I in a BEC at 732 G. τ was set to between 3–50 μ s over the range of I . The peak density is 1.6×10^{12} cm^{-3} . The axial and radial trapping frequencies were 4.5 Hz and 200 Hz, respectively. The solid line is a fit to Eq. 1 with fit parameters $K_{max} = 1.4 \times 10^{-7}$ cm^3/s and $I_{sat} = 26$ W/cm^2 . A linear fit to the low intensity data gives a slope of 1.6×10^{-8} $\text{cm}^3 \text{ s}^{-1}/(\text{W cm}^{-2})$, as indicated by the dashed line. The error bars are the statistical uncertainties, due mainly to the measured Thomas-Fermi radius of the initial density distribution. The systematic uncertainty of 12% reflects the uncertainty in the image probe detuning and intensity. Because of the relatively weak optical trap following evaporation to a BEC, I could not be increased further without producing loss from PA beam induced dipole forces.

constant, but rather increasing with time as $t^{1/2}$ during the PA pulse. We have not yet investigated a possible time dependence of K_p experimentally.

Recent unpublished work has interpreted our experiment in terms of a three-level scheme in which the atomic condensate is coupled to *two* molecular condensates; one by PA and the other by magnetoassociation [31]. Good agreement between theory and experiment is found for the maximum K_p for a condensate, and for the location of the minimum in K_p on the low-field side of the FR.

For the future, the dissociative regime could be further explored by dynamical measurements of the excitation process. Feshbach enhanced PA may also prove useful in realizing the long-sought regime of coherent oscillation between atomic and molecular condensates [32] or for efficient production of ground-state molecules [33]. We envision an off-resonant two-photon process, where a FR-enhanced free-bound PA transition is followed by bound-bound excitation back to a ground molecular level [34].

We thank T. Corcovilos, R. Côté, P. Julienne, T. Killian, M. Mackie, P. Naidon, G. Partridge, and H. Stoof for valuable discussions. Support for this work was provided by the NSF, ONR, the Keck Foundation, and the Welch Foundation (C-1133).

- [1] E. Tiesinga, B. J. Verhaar, and H. T. C. Stoof, *Phys. Rev. A* **47**, 4114 (1993).
- [2] F. A. van Abeelen, D. J. Heinzen, and B. J. Verhaar, *Phys. Rev. A* **57**, R4102 (1998).
- [3] Ph. Courteille, R. S. Freeland, D. J. Heinzen, F. A. van Abeelen, and B. J. Verhaar, *Phys. Rev. Lett.* **81**, 69 (1998).
- [4] B. Laburthe Tolra, N. Hoang, B. T'Jampens, N. Vanhaecke, C. Drag, A. Crubellier, D. Comparat, and P. Pillet, *Europhys. Lett.* **64**, 171 (2003).
- [5] J. Javanainen and M. Mackie, *Phys. Rev. Lett.* **88**, 090403 (2002).
- [6] K. Góral, M. Gajda, and K. Rzażewski, *Phys. Rev. Lett.* **86**, 1397 (2001).
- [7] M. Holland, J. Park, and R. Walser, *Phys. Rev. Lett.* **86**, 1915 (2001).
- [8] P. Naidon, E. Tiesinga, and P. S. Julienne, *Phys. Rev. Lett.* **100**, 093001 (2008).
- [9] C. McKenzie *et al.*, *Phys. Rev. Lett.* **88**, 120403 (2002).
- [10] I. D. Prodan, M. Pichler, M. Junker, R. G. Hulet, and J. L. Bohn, *Phys. Rev. Lett.* **91**, 080402 (2003).
- [11] R. Wester, S. D. Kraft, M. Mudrich, M. U. Staudt, J. Lange, N. Vanhaecke, O. Dulieu, and M. Weidemüller, *App. Phys. B* **79**, 993 (2004).
- [12] S. D. Kraft, M. Mudrich, M. U. Staudt, J. Lange, O. Dulieu, R. Wester, and M. Weidemüller, *Phys. Rev. A* **71**, 013417 (2005).
- [13] P. G. Mickelson, Y. N. Martinez, A. D. Saenz, S. B. Nagel, Y. C. Chen, T. C. Killian, P. Pellegrini, and R. Côté *et al.*, *Phys. Rev. Lett.* **95**, 223002 (2005).
- [14] U. Schlöder, C. Silber, T. Deuschle, and C. Zimmermann, *Phys. Rev. A* **66**, 061403(R) (2002).
- [15] C. Haimberger, J. Kleinert, O. Dulieu, and N. P. Bigelow, *J. Phys. B* **39**, S957 (2006).
- [16] M. Jie, W. Li-Rong, J. Wei-Bang, X. Lian-Tuan, and J. Suo-Tang, *Chinese Phys. Lett.* **24**, 1904 (2007).
- [17] J. M. Gerton, B. J. Frew, and R. G. Hulet, *Phys. Rev. A* **64**, 053410 (2001).
- [18] M. Portier, S. Moal, J. Kim, M. Leduc, C. Cohen-Tannoudji, and O. Dulieu, *J. Phys. B* **39**, S881 (2006).
- [19] P. O. Fedichev, M. W. Reynolds, and G. V. Shlyapnikov, *Phys. Rev. Lett.* **77**, 2921 (1996).
- [20] J. L. Bohn and P. S. Julienne, *Phys. Rev. A* **60**, 414 (1999).
- [21] K. E. Strecker, G. B. Partridge, A. G. Truscott, and R. G. Hulet, *Nature* **417**, 150 (2002).
- [22] Y. P. Chen, J. Hitchcock, D. Dries, M. Junker, C. Welford, and R. G. Hulet, arXiv:0710.5187 (2007).
- [23] The data of Fig. 2 place the center of the FR at 736 ± 1 G, which is a somewhat higher field than obtained in a lower resolution measurement reported previously [21].
- [24] E. R. I. Abraham, N. W. M. Ritchie, W. I. McAlexander, and R. G. Hulet, *J. Chem. Phys.* **103**, 7773 (1995).
- [25] H. T. C. Stoof, A. M. L. Janssen, J. M. V. A. Koelman, and B. J. Verhaar, *Phys. Rev. A* **39**, 3157 (1989).
- [26] V. Vuletić, C. Chin, A. J. Kerman, and S. Chu, *Phys. Rev. Lett.* **83**, 943 (1999).
- [27] R. Côté, A. Dalgarno, Y. Sun, and R. G. Hulet, *Phys. Rev. Lett.* **74**, 3581 (1995).
- [28] M. Houbiers, H. T. C. Stoof, W. I. McAlexander, and R. G. Hulet, *Phys. Rev. A* **57**, R1497 (1998).

- [29] E. R. I. Abraham, W. I. McAlexander, J. M. Gerton, R. G. Hulet, R. Côté, and A. Dalgarno, *Phys. Rev. A* **55**, R3299 (1997).
- [30] We were able to tune closer to the FR for a BEC than for a thermal gas, because of significantly smaller inelastic losses. We believe this to be a consequence of quantum statistics, since the rate for a three-body process in a BEC is 6 times smaller than for a thermal gas [25].
- [31] M. Mackie, private communication.
- [32] D. J. Heinzen, R. Wynar, P. D. Drummond, and K. V. Kheruntsyan, *Phys. Rev. Lett.* **84**, 5029 (2000).
- [33] S. J. J. M. F. Kokkelmans, H. M. J. Vissers, and B. J. Verhaar, *Phys. Rev. A* **63**, 031601(R) (2001).
- [34] E. R. I. Abraham, W. I. McAlexander, C. A. Sackett, and R. G. Hulet, *Phys. Rev. Lett.* **74**, 1315 (1995).