Lie Algebraic Analysis and Control of Quantum Dynamics

Domenico D'Alessandro*

October 29, 2018

Abstract

In this paper, we show how to use the analysis of the Lie algebra associated with a quantum mechanical system to study its dynamics and facilitate the design of controls. We give algorithms to decompose the dynamics and describe their application to the control of two coupled spin $\frac{1}{2}$'s.

1 Introduction

For several quantum mechanical systems subject to a control action, an appropriate model is the *Schrodinger operator (matrix) equation*

$$\dot{X} = -iH(u(t))X, \qquad X(0) = \mathbf{1}. \tag{1}$$

In this equation, H is called the Hamiltonian and it is an Hermitian matrix function of the control $u \equiv u(t)$ and X = X(t) is a unitary matrix for every time t, with 1 denoting the identity. Equation (1) is an appropriate model for many quantum phenomena under three main assumptions: 1) The quantum mechanical system under consideration can be adequately approximated as a system having a finite number of energy levels; 2) The interaction with the external environment (decoherence) is negligible; 3) The control, which usually represents an appropriately shaped electro-magnetic field, can be treated as a classical field (semiclassical approximation). Many physical systems share a model of the form (1). Examples are given by systems of particles with spin subject to a control magnetic field such as in NMR and EPR, molecular systems where the control is an electric field and several implementations of quantum information processing. Here the control can be seen as an action allowing us to switch among different Hamiltonians to

^{*}Department of Mathematics, Iowa State University, Ames, Iowa, U.S.A. Electronic address: daless@iastate.edu. This work was supported by NSF, under CAREER Grant ECS0237925 and Grant ECCS0824085.

implement given quantum evolutions (quantum gates). The books [1], [3], [14] present physical examples of systems sharing the model (1).

It is well known (see, e.g., [3], [7]) that the set of operators X reachable for (1) by varying the control u is the connected Lie subgroup of U(n) corresponding to the Lie algebra \mathcal{L} generated by the set $\mathcal{F} := \{iH(u)|u \in \mathcal{U}\}$, that is, the smallest Lie subalgebra of u(n) containing \mathcal{F} (see the Appendix). The Lie algebra \mathcal{L} is called the dynamical Lie algebra associated with the system and the associated connected Lie subgroup of U(n) will be denoted here by $e^{\mathcal{L}.1}$ There exists a very simple algorithm to calculate \mathcal{L} : One starts with a basis of \mathcal{F} , F_1, \ldots, F_r . If $r = n^2$ or $r = n^2 - 1$, one stops because $\mathcal{L} = u(n)$ or $\mathcal{L} = su(n)$, respectively. In this case, $e^{\mathcal{L}} = U(n)$ or $e^{\mathcal{L}} = SU(n)$ and the system is said to be controllable. If this is not the case, one performs the Lie brackets of depth 1 $[F_i, F_k]$, $j \neq k$ and select the ones that are linearly independent together with $\{F_1,\ldots,F_r\}$, say D_1,\ldots,D_s , if any. Then one performs Lie brackets of depth 2 which are Lie brackets of D_1, \ldots, D_s with the F_i 's, $j = 1, \ldots, r$ and select matrices that are linearly independent together with $F_1, \ldots, F_r, D_1, \ldots, D_s$. One goes on this way until one does not find any new linearly independent matrices. The set of matrices thus found is a basis of the dynamical Lie algebra \mathcal{L} . If the size of this set is n^2 or n^2-1 , we are in the controllable case. Otherwise the system is not controllable.² However the Lie algebra \mathcal{L} gives us information about the nature of the dynamics as we shall see next.

In section 2 we use the Levi's decomposition of Lie algebras to obtain a decomposition of the dynamics for (1). We highlight the simplifications that follow from the fact that \mathcal{L} is a subalgebra of u(n). Levi's decomposition is a classical result in Lie algebra theory but its impact in quantum control has not been considered before.³ In section 3, we give algorithms to calculate such a decomposition. Algorithms for general Lie algebras are known [5] but simplified algorithms can be given in the case of interest here. In some cases, our algorithms will be a simplified version of the ones in [5] in some cases different algorithms will be given. We shall point out this as we go on. In section 4 we use this decomposition for a control problem for two spin $\frac{1}{2}$ particles.

2 Decomposition of Quantum Dynamics

Every Lie algebra \mathcal{L} over the field of reals \mathbb{R} is the semidirect sum of a semisimple Lie algebra \mathcal{S} and the maximal solvable ideal in \mathcal{L} , \mathcal{R} , called the radical, ⁴ that is,

$$\mathcal{L} = \mathcal{S} \oplus \mathcal{R}. \tag{2}$$

¹Extending this notation, we shall use the notation $e^{\mathcal{K}}$ for the connected Lie group associated with a Lie algebra \mathcal{K} . See the Appendix for definitions.

²See subsection 3.2.1 of [3] for further discussion of this procedure.

³One exception is the book [3]. However we shall go beyond what is in this book here by pointing out the simplifications in this decomposition in the quantum case and giving explicit algorithms for calculation of the decomposition.

⁴See definitions in the Appendix.

Semidirect sum means that

$$[S, S] \subseteq S, \qquad [S, R] \subseteq R, \qquad [R, R] \subseteq R.$$
 (3)

This is a classical result known as the *Levi decomposition* (see, e.g., [5]). S is called the *Levi subalgebra*. It is the direct sum of $p \ge 1$ simple subalgebras S_j , $j = 1, \ldots, p$, i.e.,

$$S = \bigoplus_{j=1}^{p} S_j. \tag{4}$$

Direct sum means that $[S_l, S_b] = \{0\}$, when $l \neq b$. In our case, the fact that the dynamical Lie algebra \mathcal{L} is a subalgebra of u(n) implies several important simplifications.

Theorem 1 ⁵ If $\mathcal{L} \subseteq u(n)$ then the semidirect sum in (2) is a direct sum, i.e., $[\mathcal{S}, \mathcal{R}] = \{0\}$, and \mathcal{R} is Abelian, i.e., $[\mathcal{R}, \mathcal{R}] = \{0\}$.

This theorem is a consequence of Lie's theorem (cf., e.g., [6], Corollary A in Section 4.1).

Theorem 2 (Lie's Theorem) Let $\mathcal{R}_{\otimes \mathbb{C}}$ be a solvable Lie algebra of $n \times n$ matrices over the complex field. Then there exists a change of coordinates (i.e., a similarity transformation) to put all the elements in $\mathcal{R}_{\otimes \mathbb{C}}$ in upper triangular form.

We now give the proof of Theorem 1.

Proof. We first prove that \mathcal{R} is Abelian. We shall consider the field extension (see the Appendix and, e.g., [15] for a more in depth discussion) of \mathcal{R} to the complex field, $\mathcal{R}_{\otimes \mathbb{C}}$. We shall show that $\mathcal{R}_{\otimes \mathbb{C}}$ is Abelian and this implies \mathcal{R} Abelian. In fact \mathcal{R} is solvable (Abelian) if and only if $\mathcal{R}_{\otimes \mathbb{C}}$ is solvable (Abelian). Since $\mathcal{R}_{\otimes \mathbb{C}}$ is solvable, according to Lie's Theorem 2 it can be realized as upper triangular matrices. As a consequence, it can be written as the sum of two subalgebras: a nilpotent Lie algebra \mathcal{N} , corresponding to strictly upper triangular matrices, and an Abelian Lie algebra \mathcal{T} , corresponding to diagonal matrices in the coordinates indicated in Lie's theorem. Moreover, \mathcal{N} is an ideal in $\mathcal{R}_{\mathbb{C}}$. Now consider R and P in \mathcal{N} . Since \mathcal{N} is nilpotent, there exists a k>0 such that, $ad_R^k P := [R, [R, [\cdots, [R, P]]]] = 0$ where the Lie bracket is taken k times. Now, since R is skew-Hermitian, there is no loss of generality in assuming that R is diagonal, i.e., $R := \operatorname{diag}(i\lambda_1, \ldots, i\lambda_n)$. Moreover we calculate

$$(ad_R^k P)_{j,l} = p_{j,l} i^k (\lambda_j - \lambda_l)^k, \qquad \forall j, l = 1, ..., n,$$
(5)

where $i:=\sqrt{-1}$. From this expression it follows that if $ad_X^k P_{j,l}$ is zero for some k, it must be zero for every k and in particular for k=1. Therefore R and P commute and $\mathcal N$ is Abelian. Consider now $N\in\mathcal N$ and $T\in\mathcal T$. Since $\mathcal N$ is an ideal [N,T] is in $\mathcal N$ and therefore [N,[N,T]]=0 since $\mathcal N$ is Abelian. The calculation (5) shows that N and T commute and $\mathcal R_{\otimes \mathbb C}$ is the sum of two commuting Abelian subalgebras and it is therefore Abelian.

⁵This fact is mentioned in the paper [13] but without a proof. We provide a proof here.

The proof that $[S, \mathcal{R}] = 0$ use the same calculation (5). Since \mathcal{R} is an Abelian ideal (cf. (3)), $ad_R^2 P = 0$ for every $R \in \mathcal{R}$ and $P \in S$, which, from (5), implies $ad_R P = 0$.

This decomposition of the dynamical Lie algebra \mathcal{L} has immediate consequences for the Lie group of possible evolutions $e^{\mathcal{L}}$ and for the dynamics of the quantum system (1). For every control u, the solution of (1) X = X(t) factorizes as

$$X(t) = R \prod_{j=1}^{p} S_j. \tag{6}$$

Here $S_j \in e^{S_j}$ and $R \in e^{\mathcal{R}}$ and all the factors in (6) commute. Moreover R is itself the product of elements belonging to one dimensional subgroups. Write \mathcal{R} as the sum of one dimensional Lie algebras $\mathcal{R} = \bigoplus_{l=1}^q \mathcal{R}_l$, then $R = \prod_{l=1}^q R_l$, with $R_l \in e^{\mathcal{R}_l}$. Controlling the system (1) means controlling in parallel the systems $\dot{S}_j = -iH_{Sj}(u)S_j$, $S_j(0) = 1$, $j = 1, \ldots, p$, and $\dot{R}_l = -iH_{Rl}(u)R_l$, $R_l(0) = 1$, $l = 1, \ldots, q$, where $-iH_{Sj}$ and $-iH_{Rl}$ are the components of -iH(u) in S_j , $j = 1, \ldots, p$, and \mathcal{R}_l , $l = 1, \ldots, q$, respectively.

Every, finite dimensional, quantum system (1) has the structure of p + q subsystems in parallel of Figure 1. The first p subsystems vary on simple Lie groups for which a classification is known [9], [11]. The remaining q subsystems vary on one dimensional Lie groups. The total evolution is the commuting product of the evolutions on the various subgroups. To obtain the decomposition of the dynamics, we need to find bases for the subalgebras, S_j and R_l , $j = 1, \ldots, p$, $l = 1, \ldots, q$, of \mathcal{L} from a basis of \mathcal{L} . Next, we give algorithms for this task.

3 Algorithms

The book [5] contains several algorithms for general Lie algebras. In our case, we only need to consider are subalgebras of u(n). This allows us in some cases to give new and simple algorithms and in other cases to simplify the algorithms of [5].

Consider a basis $\{L_1, \ldots, L_s\}$ of \mathcal{L} . The calculation of bases of the two subspaces \mathcal{S} and \mathcal{R} in (2) is very simple in the case of interest here. It is easily seen that \mathcal{R} is the center of \mathcal{L} , and therefore it is the space of the solutions of the system of $s = \dim \mathcal{L}$ equations $[R, L_j] = 0$, $j = 1, \ldots, s$, in the variable $R \in \mathcal{L}$. Moreover, using the fact that for every semisimple Lie algebra \mathcal{S} , $\mathcal{S} = [\mathcal{S}, \mathcal{S}]$, along with $[\mathcal{S}, \mathcal{R}] = 0$ and that \mathcal{R} is Abelian, we have

$$[\mathcal{L}, \mathcal{L}] = [\mathcal{S} \oplus \mathcal{R}, \mathcal{S} \oplus \mathcal{R}] = [\mathcal{S}, \mathcal{S}] = \mathcal{S}. \tag{7}$$

Therefore the set of $\binom{s}{2}$ matrices, $[L_j, L_k], j \neq k$, spans \mathcal{S} .

The algorithm of [5] to find the solvable radical (cf. Section 2.6 in [5]) first finds the product space $[\mathcal{L}, \mathcal{L}]$ and then the adjoint representations of all elements in a basis of \mathcal{L} . Then the algorithm solves a linear system of dim $[\mathcal{L}, \mathcal{L}]$ equations in dim \mathcal{L} unknowns obtained by using the Killing form (cf. the Appendix). Then one goes on to calculate the




Figure 1: Structure of a quantum control system. The control u drives simultaneously p+q systems on Lie groups (which are simple Lie groups or one-dimensional Lie groups). The total evolution X is the commuting product of the evolutions on the various subgroups.

Levi subalgebra (cf. Section 4.13 in [5]). In our case, we have avoided the calculation and storage of the adjoint representation. Moreover, once one knows $[\mathcal{L}, \mathcal{L}]$ there is no need to apply any algorithm to find the Levi's subalgebra \mathcal{S} as we have (7).

The calculation of the simple ideals of \mathcal{S} , \mathcal{S}_j , $j = 1, \ldots, p$, is more complicated. We follow the path indicated in [5]. A preliminary step is the calculation of the so-called *primary decomposition* of \mathcal{S} which is also of interest to understand the structure of \mathcal{S} .

3.1 Calculation of the primary decomposition of S

The following definition is of interest for general Lie algebras.

Definition 3.1 A Cartan subalgebra of a Lie algebra \mathcal{S} is a nilpotent subalgebra \mathcal{A} which is equal to its *normalizer*, that is $\mathcal{A} = \{S \in \mathcal{S} | [S, \mathcal{A}] \subseteq \mathcal{A}\}.$

In the case of $S \subseteq u(n)$, we have the following

Proposition 3.2 Every nilpotent subalgebra of u(n) is Abelian. In particular the Cartan subalgebra of S is Abelian.

Proof. The proof uses the same calculation (5) and argument in Theorem 1 to show that if $ad_S^k X = 0$, for some k, and two elements S and X in the Lie algebra, then it must be $ad_S X = 0$, i.e., S and X commute.

The following algorithm calculates the Cartan subalgebra for S semisimple and $S \subseteq u(n)$.

Algorithm 1

- 1. Given the semisimple Lie algebra \mathcal{S} , set $\mathcal{A} = \{0\}$.
- 2. Select an element $X \neq 0$ in S.
- 3. Calculate the set of elements in \mathcal{S} which commute with X. Call this set \mathcal{D} . Notice \mathcal{D} is also a subalgebra of u(n).⁶ Therefore (just like \mathcal{L} above), it has a Levi decomposition in its semisimple part which is equal to $[\mathcal{D}, \mathcal{D}]$ and the center, $C(\mathcal{D})$. This justifies the next step.
- 4. Write $\mathcal{D} = [\mathcal{D}, \mathcal{D}] \oplus C(\mathcal{D})$ where $C(\mathcal{D})$ is the center of \mathcal{D} .
- 5. Set $\mathcal{A} = \mathcal{A} \oplus C(\mathcal{D})$.
- 6. If $[\mathcal{D}, \mathcal{D}] = 0$ Stop and return \mathcal{A} as Cartan subalgebra, otherwise set $\mathcal{S} = [\mathcal{D}, \mathcal{D}]$ and go to step 2.

The algorithm converges because at each step \mathcal{S} is semisimple and \mathcal{D} is a proper subspace of \mathcal{S} , otherwise \mathcal{S} would have an element which commutes with all of \mathcal{S} which contradicts semisimplicity. To show that the algorithm gives in fact a Cartan subalgebra of \mathcal{S} we have to show two facts

- 1) The resulting \mathcal{A} is nilpotent (it is in fact Abelian).
- 2) Every $S \in \mathcal{S}$ which is such that $[S, \mathcal{A}] \subseteq \mathcal{A}$ is an element of \mathcal{A} .

Proof. (Proof of 1) and 2) above) Let us denote by \mathcal{A}_k and \mathcal{D}_k the Lie algebras \mathcal{A} and \mathcal{D} obtained after the k-th step is complete. We show 1) by induction on the steps of the algorithm. \mathcal{A}_1 is definitely Abelian as it is the center of \mathcal{D}_1 . Moreover it commutes with \mathcal{D}_1 . The inductive step shows that these two properties are true at each step. Assume they are true at step k-1. At step k, $\mathcal{A}_k = \mathcal{A}_{k-1} \oplus C(\mathcal{D}_k)$. However both \mathcal{A}_{k-1} and $\mathcal{C}(\mathcal{D}_k)$ are Abelian and $[\mathcal{A}_{k-1}, \mathcal{C}(\mathcal{D}_k)] = \{0\}$, because $\mathcal{D}_k \subset \mathcal{D}_{k-1}$, and we know by the inductive assumption that \mathcal{A}_{k-1} commutes with \mathcal{D}_{k-1} . This proves that \mathcal{A}_k is Abelian. Using again $\mathcal{A}_k = \mathcal{A}_{k-1} \oplus C(\mathcal{D}_k)$, and the fact that $\mathcal{D}_k \subset \mathcal{D}_{k-1}$, $[\mathcal{A}_k, \mathcal{D}_k] = \{0\}$ and the inductive step is complete.

To show 2) assume $[S, A] \subseteq A$. Then repeating the argument following (5) $[S, A] = \{0\}$. Consider X and \mathcal{D}_1 at the first step. Write $S = S_1 + S_0$ with $S_0 \in \mathcal{D}_1$ and S_1 in a complement of \mathcal{D}_1 in S. Since $X \in A$ and S commutes with A, then S commutes

⁶The fact that it is a Lie algebra, i.e., $[D_1, D_2] \in \mathcal{D}$, for $D_1, D_2 \in \mathcal{D}$ follows immediately from an application of the Jacobi identity. We have $[[D_1, D_2], X] = -[[D_2, X], D_1] - [[X, D_1], D_2] = 0$.

with X. Since S_0 commutes with X so does S_1 . So, S_1 is in \mathcal{D}_1 and in a complement of \mathcal{D}_1 which implies $S_1 = 0$. Therefore $S \in \mathcal{D}_1$ and we can write it as $S = D_1 + A_1$ with $D_1 \in [\mathcal{D}_1, \mathcal{D}_1]$ and $A_1 \in C(\mathcal{D}_1)$ and therefore $\in \mathcal{A}$. At the second step we pick another X which turns out to be again in \mathcal{A} . Since S commutes with \mathcal{A} , D_1 is in \mathcal{D}_2 . We can then write $D_1 = D_2 + A_2$ with $D_2 \in [\mathcal{D}_2, \mathcal{D}_2]$, and $A_2 \in \mathcal{A}$. Therefore $S = D_2 + A_2 + A_1$. Proceeding this way, if the procedure ends after r steps, we have that $[\mathcal{D}_r, \mathcal{D}_r] = \{0\}$ and S has the form $S = A_r + A_{r-1} + \cdots + A_1$, with $A_j \in \mathcal{A}$, for every j = 1, ..., r. Therefore $S \in \mathcal{A}$.

Remark 3.3 We remark that the above algorithm and proof is not derived as a special case of the corresponding algorithm in [5] (cf. Section 3.2 in [5]) but it is, to the best of the author's knowledge, new. It gives the Cartan subalgebra for the specific situation of interest in quantum control. It is simpler than the general algorithm both because it involves fewer notions of Lie algebra theory and because it involves fewer operations. ⁷

The following definition refers to a general Lie algebra over a general field (cf. [5] (Definitions 3.1.1 and 3.1.9)).

Definition 3.4 A collected primary decomposition of a semisimple Lie algebra \mathcal{S} with respect to a Cartan subalgebra \mathcal{A} is a vector space decomposition of the form

$$S := A \oplus V_1 \oplus V_2 \oplus \cdots V_r, \tag{8}$$

where

1. The subspaces V_j 's, j = 1, ..., r, are invariant under ad_X , for every $X \in \mathcal{A}$, that is

$$[\mathcal{A}, \mathcal{V}_j] \subseteq \mathcal{V}_j, \qquad j = 1, \dots, r.$$

- 2. For every $X \in \mathcal{A}$, and every \mathcal{V}_j , the minimum polynomial of ad_X restricted to \mathcal{V}_j is the power of an irreducible polynomial.
- 3. For any two subspaces V_j and V_k , there exists an $X \in \mathcal{A}$ such that the minimum polynomials of ad_X restricted to V_j and V_k are powers of two different irreducible polynomials.

Given A such a decomposition exists and is unique (Theorem 3.1.10 of [5]).

In our case, for every X, the minimum polynomial of ad_X restricted to \mathcal{V}_j must be of the type $(\lambda^2 + a^2)$, otherwise ad_X would have eigenvalues with nonzero real parts and-or eigenvalues with geometric multiplicity greater than one. This is not possible because we have (in an appropriate basis) $ad_X^T = -ad_X$ (see Proposition 4.1 in Appendix).

⁷The algorithm of [5] involves, among the other things, the tuning a parameter so that a certain vector space has a given dimension.

An algorithm to calculate the collected primary decomposition is given below. This algorithm was derived by applying the general algorithm presented in [5] to the case considered here (cf. Section 4.11 in [5]).

Algorithm 2

1. Select an element $X \in \mathcal{A}$ such that ad_X has $\dim \mathcal{L} - \dim \mathcal{A} + 1$ different eigenvalues. That is, except for the 0 eigenvalue (which has eigenspace equal to \mathcal{A}), ad_X is non-degenerate.

Such elements are called *splitting elements* and they exist (Corollary 4.11.3 of [5]). To find such an X, notice that, if $\{A_1, \ldots, A_m\}$ is a basis of A then $ad_{A_1}, \ldots, ad_{A_m}$ all commute and they can be simultaneously diagonalized. It is easier then to select real coefficients c_j , such that $\sum_{j=1}^m c_j ad_{A_j}$ has the desired property, and $X = \sum_{j=1}^m c_j A_j$. For higher dimensional problems it may be more convenient to use randomized algorithms.

Let X be the selected element, the minimum polynomial is of the form $m_{ad_X}(\lambda) = \prod_{j=1}^f (\lambda^2 + a_j^2)$, with the a_j all $\in \mathbb{R}$, all different from each other and with one of them equal to zero (0 is always an eigenvalue of ad_X , X being an eigenvector). Moreover, from the choice of X being splitting $f = \frac{\dim \mathcal{L} - \dim \mathcal{A}}{2} + 1$ and $m_{ad_X}(\lambda)$ is equal to the characteristic polynomial except (possibly) for the power of the monomial associated to the eigenvalue 0.8

2. Take as V_j the (two-dimensional) eigenspaces associated with the pair of purely imaginary eigenvalues corresponding to a_j^2 . That is $V_j = \{V \in \mathcal{L} | (ad_X^2 + a_j^2 \mathbf{1})V = 0\}, \quad j = 1, \ldots, \frac{\dim \mathcal{S} - \dim \mathcal{A}}{2}$.

We now prove that the decomposition obtained with the above algorithm is the collected primary decomposition associated with the Cartan subalgebra \mathcal{A} .

Proof. (Proof of 1, 2 and 3 in definition 3.4). Condition 1 is verified since if $X_2 \in \mathcal{A}$, and $V \in \mathcal{V}_j$, we have (since X and X_2 commute and therefore so do ad_X and ad_{X_2}) $(ad_X^2 + a_j^2 \mathbf{1})ad_{X_2}V = ad_{X_2}(ad_X^2 + a_j^2 \mathbf{1})V = 0$. Therefore $ad_{X_2}V \in \mathcal{V}_j$ as well. Condition 2 is also verified. Since \mathcal{V}_j is two dimensional, the minimum polynomial of ad_{X_2} restricted to \mathcal{V}_j , for every $X_2 \in \mathcal{A}$, must be of the form $\lambda^2 + b^2$ for real b. Any other form would imply that ad_{X_2} has eigenvalues with nonzero real part, which has to be excluded because of Proposition 4.1 in the Appendix. Condition 3 is verified taking as element X in \mathcal{A} precisely the splitting element.

⁸Notice in particular that the difference between the dimension of \mathcal{L} and that of \mathcal{A} must be an even number.

3.2 Calculation of the decomposition in simple ideals

The primary decomposition is a fundamental tool to explore the structure of a semisimple Lie algebra S. Using it, one can directly obtain the decomposition into simple ideals (4). The algorithm is given in [5] (cf., Section 4.12) and we report it using our notations.

Algorithm 3

1. For every j = 1, ..., r, calculate the spaces

$$\mathcal{I}_j := \bigoplus_{k=0}^{\infty} ad_{\mathcal{S}}^k \mathcal{V}_j, \tag{9}$$

where V_j are defined in (8).

 \mathcal{I}_j is the smallest ideal containing \mathcal{V}_j . $ad_{\mathcal{S}}^k \mathcal{V}_j$, is defined inductively, where $ad_{\mathcal{S}}^0 \mathcal{V}_j = \mathcal{V}_j$, and $ad_{\mathcal{S}}^k \mathcal{V}_j = [\mathcal{S}, ad_{\mathcal{S}}^{k-1} \mathcal{V}_j]$.

2. The simple ideals S_l , l = 1, ..., p in (4) are given by the ideals \mathcal{I}_j . Notice that some ideals may be coinciding.

Proof. (Proof of Algorithm 3) The main fact used to justify the algorithm is that the primary decomposition (8) is compatible with the decomposition in simple ideals (4). This is proved in [5] (Theorem 4.12.1) and it means the following: The Cartan subalgebra of \mathcal{S} , \mathcal{A} , splits in p subalgebras, \mathcal{A}_j , $j = 1, \ldots, p$, i.e., $\mathcal{A} = \bigoplus_{j=1}^p \mathcal{A}_j$, where each \mathcal{A}_j is a Cartan subalgebra of the corresponding \mathcal{S}_j . Moreover, each \mathcal{V}_j , $j = 1, \ldots, r$, in (8) is a subspace of one of the \mathcal{S}_k , $k = 1, \ldots, p$. From the latter fact, it follows immediately that each \mathcal{S}_k , $k = 1, \ldots, p$, is the smallest ideal generated by one \mathcal{I}_j , that is, it is of the form (9). In fact, each \mathcal{I}_j is contained in some \mathcal{S}_k and \mathcal{S}_k cannot contain any nontrivial ideal other than itself, being simple.

4 Example: Control of two interacting spin $\frac{1}{2}$'s

Consider the control of two interacting spin $\frac{1}{2}$ particles subject to a magnetic field. The state of particle 1 (2) lives in a 2-dimensional Hilbert space \mathcal{H}_1 (\mathcal{H}_2), so that the state of the total system lives in a 4-dimensional Hilbert space $\mathcal{H}_1 \otimes \mathcal{H}_2$. This type of systems are of interest for example in quantum computation when one wants to perform operations with two quantum bits (cf., e.g., [12]). In the model we shall consider an externally applied magnetic field is constant, has nonzero component in the x direction only and it only affects the first spin. It is however possible to control the interaction between the two spins. There are several ways to experimentally achieve this; see, e.g., [8]. The Hamiltonian H(u) in the system's equation (1) has the form

$$H = u_1(t)\sigma_z \otimes \sigma_z + u_2(t)\sigma_y \otimes \sigma_y + \sigma_x \otimes \mathbf{1}$$
(10)

where $\sigma_{x,y,z}$ are the Pauli matrices $\sigma_x := \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma_y := \frac{1}{2} \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$, $\sigma_z := \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, which satisfy the commutation relations

$$[i\sigma_x, i\sigma_y] = i\sigma_z, \qquad [i\sigma_y, i\sigma_z] = i\sigma_x, \qquad [i\sigma_z, i\sigma_x] = i\sigma_y.$$
 (11)

The solution X of (1) represents the evolution on $\mathcal{H}_1 \otimes \mathcal{H}_2$ as the state $\psi(t)$ evolves as $\psi(t) = X(t)\psi(0)$ with X(t) the solution of (1). The dynamical Lie algebra is generated by $\{i\sigma_z \otimes \sigma_z, i\sigma_y \otimes \sigma_y, i\sigma_x \otimes \mathbf{1}\}$. It is given by

$$\mathcal{L} := span\{i\sigma_x \otimes \mathbf{1}, i\mathbf{1} \otimes \sigma_x, i\sigma_z \otimes \sigma_z, i\sigma_y \otimes \sigma_y, i\sigma_z \otimes \sigma_y, i\sigma_y \otimes \sigma_z\}, \tag{12}$$

which is 6-dimensional. As the dimension of the full Lie algebra su(4) is 15, this shows that the system is not controllable. The Lie group $e^{\mathcal{L}}$ gives the set of reachable evolutions. In order to understand the nature of this set and perform control to any possible value in it, we apply the analysis developed in this paper. An application of the algorithm for the calculation of the (Abelian) radical \mathcal{R} discussed at the beginning of Section 3 shows that $\mathcal{R} = \{0\}$ and that \mathcal{L} is semisimple. To calculate the simple subalgebras, we apply the algorithms developed in Section 3. We apply Algorithm 1 to find the Cartan subalgebra \mathcal{A} . Selecting $X = i\sigma_x \otimes 1$ at Step 2 of that algorithm, we find

$$\mathcal{A} = span\{i\sigma_x \otimes \mathbf{1}, i\mathbf{1} \otimes \sigma_x\}. \tag{13}$$

To find the primary decomposition according to Algorithm 2 we have to select a splitting element in \mathcal{A} . We write the adjoint representations of $i\sigma_x \otimes \mathbf{1}$ and $i\mathbf{1} \otimes \sigma_x$ in the ordered basis indicated in (12). We have

Neither $i\sigma_x \otimes \mathbf{1}$ nor $i\mathbf{1} \otimes \sigma_x$ is a splitting element but $i\sigma_x \otimes \mathbf{1} + 2i\mathbf{1} \otimes \sigma_x$ is because $ad_{i\sigma_x \otimes \mathbf{1} + 2i\mathbf{1} \otimes \sigma_x} = ad_{i\sigma_x \otimes \mathbf{1}} + 2ad_{i\mathbf{1} \otimes \sigma_x}$ has dim $\mathcal{L} - \dim \mathcal{A} + 1 = 6 - 2 + 1 = 5$ different eigenvalues, $0, \pm 3i, \pm i$. The eigenspaces corresponding to $\pm 3i$ and ± 1 are given by $\mathcal{V}_1 = span\{i(\sigma_z \otimes \sigma_y + \sigma_y \otimes \sigma_z), i(\sigma_z \otimes \sigma_z - \sigma_y \otimes \sigma_y)\}$, and $\mathcal{V}_2 = span\{i(\sigma_z \otimes \sigma_y - \sigma_y \otimes \sigma_z), i(\sigma_z \otimes \sigma_z + \sigma_y \otimes \sigma_y)\}$. With \mathcal{A} in (13) and \mathcal{V}_1 and \mathcal{V}_2 above, the primary decomposition is $\mathcal{L} = \mathcal{A} \oplus \mathcal{V}_1 \oplus \mathcal{V}_2$. Using Algorithm 3, we obtain the simple component Lie algebras \mathcal{S}_A , \mathcal{S}_B , as the ideals generated by \mathcal{V}_1 and \mathcal{V}_2 . We have $\mathcal{L} = \mathcal{S}_A \oplus \mathcal{S}_B$, $[\mathcal{S}_A, \mathcal{S}_B] = 0$, with the simple Lie algebras \mathcal{S}_A and \mathcal{S}_B given by (cf. (9)) $\mathcal{S}_A = \bigoplus_{k=0}^{\infty} ad_{\mathcal{L}}^k \mathcal{V}_1 := span\{A_1, A_2, A_3\}, \quad \mathcal{S}_B = \bigoplus_{k=0}^{\infty} ad_{\mathcal{L}}^k \mathcal{V}_2 := span\{B_1, B_2, B_3\}$, with $A_1 := \frac{i}{2}(\sigma_x \otimes \mathbf{1} + \mathbf{1} \otimes \sigma_x), A_2 := \frac{i}{2}(\sigma_z \otimes \sigma_y + \sigma_y \otimes \sigma_z), A_3 := \frac{i}{2}(\sigma_z \otimes \sigma_z - \sigma_y \otimes \sigma_y)$, and $B_1 := \frac{i}{2}(-\sigma_x \otimes \mathbf{1} + \mathbf{1} \otimes \sigma_x), B_2 := \frac{i}{2}(\sigma_z \otimes \sigma_y - \sigma_y \otimes \sigma_z), B_3 := \frac{i}{2}(\sigma_z \otimes \sigma_z + \sigma_y \otimes \sigma_y)$. The commutation relations $[A_1, A_2] = A_3, [A_2, A_3] = A_1$,

 $[A_3,A_1]=A_2$ ($[B_1,B_2]=B_3$, $[B_2,B_3]=B_1$, $[B_3,B_1]=B_2$), compared with (11), show that \mathcal{S}_A and \mathcal{S}_B are both isomorphic to su(2). Writing the Hamiltonian (10) as $iH=iH_A+iH_B$, $iH_A=A_3(u_1-u_2)+A_1$, $iH_B=B_3(u_1+u_2)-B_1$, we find that the solution X of (1), (10) is the commuting product of the solutions U_A and U_B of the decoupled systems $\dot{U}_A=-iH_A(u_1-u_2)U_A$, $U_A(0)=1$, and $\dot{U}_B=-iH_B(u_1+u_2)U_B$, $U_B(0)=1$,i.e., $X(t)=U_A(t)U_B(t)=U_B(t)U_A(t)$. We can use u_1-u_2 and u_1+u_2 as independent controls to drive U_A and U_B , respectively. Because of the isomorphism between \mathcal{S}_A and \mathcal{S}_B with su(2), both control problems are equivalent to control problems on SU(2) for which there exists a large literature. One can for example use the Riemannian symmetric space argument of [10] to obtain the minimum (in fact infimum) time control if there is no bound on the control. Alternatively one can use an optimal, minimum energy, control over a finite time horizon which turns out to be given by elliptic functions as described in [4] or a Lie group decomposition technique as in [2] which can be applied when there are bounds on the magnitude of the controls.

References

- [1] A. G. Butkovskiy and Y. I. Samoilenko, *Control of Quantum Mechanical Processes and Systems*, Mathematics and its Applications, Kluwer Academic Publishers, Dordrecht, 1990.
- [2] D. D'Alessandro, Optimal evaluation of generalized Euler angles with application to control, *Automatica*, 40, 1997-2002, (2004).
- [3] D. D'Alessandro, Introduction to Quantum Control and Dynamics, CRC Press, Boca Raton, FL, 2007.
- [4] D. D'Alessandro and M. Dahleh, Optimal control of two level quantum systems, *IEEE Transactions on Automatic Control*, 45, No. 1, (2001).
- [5] W. A. de Graaf, Lie Algebras; Theory and Algirithms, North-Holland, 2000.
- [6] J. E. Humphreys, Introduction to Lie algebras and Representation Theory, Springer Verlag, New York, 1972.
- [7] V. Jurdjević and H. Sussmann, Control Systems on Lie groups, *Journal of Differential Equations*, 12, 1972 313-329.
- [8] B. E. Kane, A silicon-based nuclear spin quantum computer, *Nature*, **393**, 33 (1998).
- [9] W. B. M. Kay and J. Patera, Table of Dimension, Indices and Branching Rules for Representations of Simple Lie algebras, Marcel Dekker, New York, 1981.
- [10] N. Khaneja, R. Brockett and S. J. Glaser, Time optimal control of spin systems, *Phys. Rev. A*, 63, 032308, (2001).

- [11] A. Knapp, Lie Groups: Beyond an Introduction, 2nd ed., Birkhauser Beston, 2002.
- [12] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information,, Cambridge University Press, Cambridge UK, 2000.
- [13] T. Polack, H. Suchowski, and D. Tannor, Uncontrollable quantum systems: A classification scheme based on Lie subalgebras, unpublished, presented at the APS March Meeting, 2007.
- [14] S. A. Rice and M. Zhao, Optical Control of Molecular Dynamics, New York, Wiley, 2000.
- [15] A. A. Sagle and R. E. Walde, *Introduction to Lie Groups and Lie Algebras*, Academic Press, New York, 1973.

Appendix: Some facts about Lie algebras and Lie groups

A Lie algebra (see e.g. the textbooks [5], [6], [15]) is a vector space closed under with a binary operation called the commutator $(x,y) \to [x,y]$ which is bilinear with respect to the vector space sum, skew-symmetric ([x,y] = -[y,x]) and satisfies the Jacobi identity: [x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0. For Lie algebras of matrices that are the ones that interest us in this paper, the commutator is taken to be the usual commutator of two matrices [A, B] := AB - BA. The Lie algebra u(n) (su(n)) is the Lie algebra over the reals of $n \times n$ skew-Hermitian matrices (with trace zero). It has dimension n^2 $(n^2 - 1)$. Two Lie algebras \mathcal{L} and \mathcal{L}' are **isomorphic**, if there exists a linear one to one and onto map $\phi: \mathcal{L} \to \mathcal{L}'$, such that $\phi([A, B]_{\mathcal{L}}) = [\phi(A), \phi(B)]_{\mathcal{L}'}$, for every A and B in \mathcal{L} . Here $[\cdot,\cdot]_{\mathcal{L}}$ and $[\cdot,\cdot]_{\mathcal{L}'}$ denote the commutators in \mathcal{L} and \mathcal{L}' respectively. Given a Lie algebra \mathcal{L} over the reals, it is possible to define a Lie algebra over the complex field \mathcal{L}_{∞} Which is called the **field extension** of \mathcal{L} . $\mathcal{L}_{\otimes \mathbb{C}}$ has the same basis as \mathcal{L} and it has the same dimension over the complex numbers as $\mathcal L$ over the reals. The Lie brackets between two basis elements give the same result as for the real Lie brackets. Associated with a Lie algebra \mathcal{L} of matrices is a **Lie group** $e^{\mathcal{L}}$ which is defined as the set of finite products of the form $e^{L_1}e^{L_2}\cdots e^{L_f}$, with $L_1,\ldots,L_f\in\mathcal{L}$, for $f\geq 0$. A Lie group is a group in the algebraic sense and it is a differentiable manifold. Naturally, the open sets in $e^{\mathcal{L}}$ are defined by requiring that the map $\pi: \mathbb{R}^n \to e^{\mathcal{L}}, \ \pi(t_1, \dots, t_n) = e^{A_1 t_1} \cdots e^{A_n t_n}, \text{ for any } \{A_1, \dots, A_n\}$ basis in \mathcal{L} is open, i.e., maps open sets in \mathbb{R}^n into open sets in $e^{\mathcal{L}}$. The Lie group associated with the Lie algebra u(n) (su(n)) is the Lie group of $n \times n$ unitary matrices U(n) $(n \times n \text{ unitary matrices with determinant equal to 1, <math>SU(n)$). A Lie subgroup **S** of $e^{\mathcal{L}}$ is a subgroup in the algebraic which is also a sub-manifold in the sense that the topology of S coincides with the one induced by the one of $e^{\mathcal{L}}$. If K is a subalgebra of \mathcal{L} , then the Lie group $e^{\mathcal{K}}$ is a Lie subgroup of $e^{\mathcal{L}}$ provided that the last condition on the topology is satisfied. Consider a Lie algebra \mathcal{L} , and define inductively the following

⁹This is tacitly assumed anytime we talk about a Lie subgroup in the paper. The result mentioned at the beginning of the introduction has to be slightly modified if this last assumption is not verified by

sequence of subalgebras $\mathcal{L}^0 := \mathcal{L}$, $\mathcal{L}^{(k+1)} = [\mathcal{L}^{(k)}, \mathcal{L}^{(k)}]$. \mathcal{L} is called **solvable** if there exists a k such that $\mathcal{L}^{(k)} = \{0\}$. Another sequence is given by $\mathcal{L}_0 = \mathcal{L}$, $\mathcal{L}_{k+1} = [\mathcal{L}, \mathcal{L}_k]$. A Lie algebra is **nilpotent** if, there exists a k such that $\mathcal{L}_k = \{0\}$. It is called **Abelian** if $\mathcal{L}_1 = \{0\}$. If a Lie algebra is Abelian it is nilpotent. If it is nilpotent it is solvable. An ideal of \mathcal{L} is a subspace \mathcal{I} , such that $[\mathcal{L}, \mathcal{I}] \subseteq \mathcal{I}$. A Lie algebra \mathcal{L} is called simple if it has dimension > 1 and it contains no ideals except the trivial ones, $\{0\}$ and \mathcal{L} . A Lie algebra is **semisimple** if it is the direct sum (i.e., the sum of vector spaces which commute with each other) of simple Lie algebras. Semisimple Lie algebras \mathcal{S} have the property that [S, S] = S. Consider a subspace N of the Lie algebra \mathcal{L} . The normalizer of \mathcal{N} , is the set $N_{\mathcal{L}}(\mathcal{N}) = \{L \in \mathcal{L} | [L, \mathcal{N}] \subseteq \mathcal{N} \}$. The **center** of a Lie algebra \mathcal{L} is the set $C(\mathcal{L}) := \{L \in \mathcal{L} | [L, \mathcal{L}] = 0\}$. It is clear that the center is a subspace of the normalizer and both of them (using Jacobi identity) are closed under commutation and therefore are subalgebras of \mathcal{L} . Let \mathcal{L} be a Lie algebra over the field of reals. The **adjoint representation** of the Lie algebra \mathcal{L} is a function $ad : \mathcal{L} \to \mathcal{M}_{\dim(\mathcal{L}),\dim(\mathcal{L})}^{10}$ which maps $X \in \mathcal{L}$ to ad_X , where ad_X is a linear map $\mathcal{L} \to \mathcal{L}$ defined by $ad_X L := [X, L]$. The adjoint representation is a **representation** of the Lie algebra \mathcal{L} in that it preserves the basic Lie algebra operations. We have $ad_{[X,Y]} = [ad_X, ad_Y]$ and ad_X is a matrix acting on $\mathbb{R}^{\dim \mathcal{L}}$. The following fact is used in the paper.

Proposition 4.1 If \mathcal{L} is semisimple, for every $X \in \mathcal{L}$, there exists a basis in \mathcal{L} such that $ad_X^T = -ad_X$.

Proof. Consider the Killing form on \mathcal{L} , $\langle Y, Z \rangle_K$ defined as $\langle Y, Z \rangle_K := Tr(ad_Yad_Z)$. Since \mathcal{L} is semisimple according to Cartan criterion (see, e.g., [5]) the Killing form is non-degenerate. Consider a basis of \mathcal{L} , which is orthogonal with respect to the Killing form. If we define the transposed of a linear operator O by $\langle Y, O^T(Z) \rangle_K = \langle OY, (Z) \rangle_K$, it is easily seen that the matrix form of the transposed in the orthonormal basis is the usual transposed of the matrix form of O. We have, given X, for every Y and Z, $\langle Y, ad_X^T Z \rangle_K := \langle ad_X Y, Z \rangle_K = \langle [Z, X], Y \rangle_K = \langle Y, [Z, X] \rangle_K = -\langle Y, ad_X Z \rangle_K$. The first equality follows from the definition of transposed. Then, we used the cyclic property of the Killing form in the third equality and its symmetry in the fourth equality. Summarizing, we have $\langle Y, ad_X^T Z \rangle_K = -\langle Y, ad_X Z \rangle_K$. Since the Killing form is non-degenerate, we must have $ad_X^T Z = -ad_X Z$ and since this is true for every Z the claim follows.

saying that the set of reachable values of X for (1) is dense (in the topology of U(n)) in the Lie group associated to the dynamical Lie algebra \mathcal{L} . Therefore from a practical point of view there is no difference whether $e^{\mathcal{L}}$ is or is not a Lie subgroup of U(n). The author wishes to thank Francesco Ticozzi for useful discussions on this point.

 $^{^{10}\}mathcal{M}_{n,n}$ denotes the Lie algebra of $n \times n$ real matrices with the commutator given by the standard matrix commutator.

¹¹The author thanks Richard Ng for working out this proof and for helpful discussions on this paper.