

# EXAMPLES OF AREA-MINIMIZING SURFACES IN THE SUBRIEMANNIAN HEISENBERG GROUP $\mathbb{H}^1$ WITH LOW REGULARITY

MANUEL RITORÉ

ABSTRACT. We give new examples of entire area-minimizing  $t$ -graphs in the subriemannian Heisenberg group  $\mathbb{H}^1$ . Most of the examples are locally lipschitz in Euclidean sense. Some regular examples have prescribed singular set consisting of either a horizontal line or a finite number of horizontal halflines extending from a given point. Amongst them, a large family of area-minimizing cones is obtained.

## 1. INTRODUCTION

Variational problems related to the subriemannian area in the Heisenberg group  $\mathbb{H}^1$  have received great attention recently. A major question in this theory is the regularity of minimizers. A related one is the construction of examples with low regularity properties. The study of minimal surfaces in subriemannian geometry was initiated in the paper by Garofalo and Nhieu [22]. Later Pauls [27] constructed minimal surfaces in  $\mathbb{H}^1$  as limits of minimal surfaces in Nil manifolds, the riemannian Heisenberg groups. Cheng, Hwang and Yang [9] have studied the weak solutions of the minimal surface equation for  $t$ -graphs and have proven existence and uniqueness results. Regularity of minimal surfaces, assuming that they are least  $C^1$ , has been treated in the papers by Pauls [28] and Cheng, Hwang and Yang [10]. We would like also to mention the recently distributed notes by Bigolin and Serra Cassano [5], where they obtain regularity properties of an  $\mathbb{H}$ -regular surface from regularity properties of its horizontal unit normal. Interesting examples of minimal surfaces which are not area-minimizing are obtained in [11]. See also [13]. Smoothness of lipschitz minimal intrinsic graphs in Heisenberg groups  $\mathbb{H}^n$ , for  $n > 1$ , has been recently obtained by Capogna, Citti and Manfredini [6].

Characterization in  $\mathbb{H}^1$  of solutions of the Bernstein problem for  $C^2$  surfaces has been obtained by Cheng, Hwang, Malchiodi and Yang [8], and Ritoré and Rosales [29] for  $t$ -graphs, and by Barone Adessi, Serra Cassano and Vittone [4] and Garofalo and Pauls [23] for vertical graphs.

Additional contributions concerning variational problems related to the subriemannian area in the Heisenberg groups include [26], [2], [8], [9], [10], [21], [21], [20], [19], [18], [17], [16], [25], [29]. The recent monograph by Capogna, Danielli, Pauls and Tyson [7] gives a recent overview of the subject with an exhaustive list of references. We would like

---

*Date:* October 26, 2018.

*2000 Mathematics Subject Classification.* 53C17, 49Q20.

*Key words and phrases.* Sub-Riemannian geometry, Heisenberg group, minimal surfaces, minimal cones.

Research supported by MEC-Feder grant MTM2007-61919.

to stress that, in  $\mathbb{H}^1$ , the condition  $H \equiv 0$  is not enough to guarantee that a given surface of class  $C^2$  is even a stationary point for the area functional, see Ritoré and Rosales [29], and Cheng, Hwang and Yang [9] for minimizing  $t$ -graphs.

The aim of this paper is to provide new examples in  $\mathbb{H}^1$  of Euclidean locally lipschitz area-minimizing entire graphs over the  $xy$ -plane.

In section 3 we construct the basic examples. We start from a given horizontal line  $L$ , and a monotone angle function  $\alpha : L \rightarrow (0, \pi)$  over this line. For each  $p \in L$ , we consider the two horizontal halflines extending from  $p$  making an angle  $\pm\alpha(p)$  with  $L$ . We prove that in this way we always obtain an entire graph over the  $xy$ -plane which is Euclidean locally lipschitz and area-minimizing. The angle function  $\alpha$  is only assumed to be continuous and monotone. Of course, further regularity on  $\alpha$  yields more regularity on the graph. In case  $\alpha$  is at least  $C^2$  we get that the associated surface is  $C^{1,1}$ .

The surfaces in section 3 are the building blocks for our next construction in section 4. We fix a point  $p \in \mathbb{H}^1$ , and a family of counter-clockwise oriented horizontal halflines  $R_1, \dots, R_n$  extending from  $p$ . We choose the bisector  $L_i$  of the wedge determined by  $R_{i-1}$  and  $R_i$ , and we consider angle functions  $\alpha_i : L_i \rightarrow (0, \pi)$  which are continuous, nonincreasing as a function of the distance to  $p$ , and such that  $\alpha(p)$  is equal to the angle between  $L_i$  and  $R_i$ . For every  $q \in L_i$ , we consider the halflines extending from  $q$  with angles  $\pm\alpha_i(q)$ . In this way we also a family of area-minimizing  $t$ -graphs which are Euclidean locally lipschitz. In case the obtained surface is regular enough we have that the singular set is precisely  $\bigcup_{i=1}^n L_i$ . If the angle functions  $\alpha_i$  are constant, then we obtain area-minimizing cones (the original motivation of this paper), which are Euclidean locally  $C^{1,1}$  minimizers, and  $C^\infty$  outside the singular set  $\bigcup_{i=1}^n L_i$ . For a single halfline  $L$  extending from the origin and an angle function  $\alpha : L \rightarrow (0, \pi)$ , continuous and nonincreasing as a function of the distance to 0, we patch the graph obtained over a wedge of the  $xy$ -plane with the plane  $t = 0$  along the halflines extending from 0 making an angle  $\alpha(0)$  with  $L$ . When  $\alpha$  is constant we get again an area-minimizing cone which is Euclidean locally lipschitz. These cones are a generalization of the one obtained by Cheng, Hwang and Yang [9, Ex. 7.2].

An interesting consequence of this construction is that we get a large number of Euclidean locally  $C^{1,1}$  area-minimizing cones with prescribed singular set consisting on either a horizontal line or a finite number of horizontal halflines extending from a given point. It is an open question to decide if these examples are the only area-minimizing cones, together with vertical halfspaces and the example by Cheng, Hwang and Yang [9, Ex. 7.2] with a singular halfline and its generalizations in the last section. The importance of tangent cones has been recently stressed in [1].

## 2. PRELIMINARIES

The *Heisenberg group*  $\mathbb{H}^1$  is the Lie group  $(\mathbb{R}^3, *)$ , where the product  $*$  is defined, for any pair of points  $[z, t], [z', t'] \in \mathbb{R}^3 \equiv \mathbb{C} \times \mathbb{R}$ , as

$$[z, t] * [z', t'] := [z + z', t + t' + \operatorname{Im}(z\bar{z}')], \quad (z = x + iy).$$

For  $p \in \mathbb{H}^1$ , the *left translation* by  $p$  is the diffeomorphism  $L_p(q) = p * q$ . A basis of left invariant vector fields (i.e., invariant by any left translation) is given by

$$X := \frac{\partial}{\partial x} + y \frac{\partial}{\partial t}, \quad Y := \frac{\partial}{\partial y} - x \frac{\partial}{\partial t}, \quad T := \frac{\partial}{\partial t}.$$

The *horizontal distribution*  $\mathcal{H}$  in  $\mathbb{H}^1$  is the smooth planar one generated by  $X$  and  $Y$ . The *horizontal projection* of a vector  $U$  onto  $\mathcal{H}$  will be denoted by  $U_H$ . A vector field  $U$  is called *horizontal* if  $U = U_H$ . A *horizontal curve* is a  $C^1$  curve whose tangent vector lies in the horizontal distribution.

We denote by  $[U, V]$  the Lie bracket of two  $C^1$  vector fields  $U, V$  on  $\mathbb{H}^1$ . Note that  $[X, T] = [Y, T] = 0$ , while  $[X, Y] = -2T$ . The last equality implies that  $\mathcal{H}$  is a bracket generating distribution. Moreover, by Frobenius Theorem we have that  $\mathcal{H}$  is nonintegrable. The vector fields  $X$  and  $Y$  generate the kernel of the (contact) 1-form  $\omega := -y dx + x dy + dt$ .

We shall consider on  $\mathbb{H}^1$  the (left invariant) Riemannian metric  $g = \langle \cdot, \cdot \rangle$  so that  $\{X, Y, T\}$  is an orthonormal basis at every point, and the associated Levi-Civita connection  $D$ . The modulus of a vector field  $U$  will be denoted by  $|U|$ .

Let  $\gamma : I \rightarrow \mathbb{H}^1$  be a piecewise  $C^1$  curve defined on a compact interval  $I \subset \mathbb{R}$ . The *length* of  $\gamma$  is the usual Riemannian length  $L(\gamma) := \int_I |\dot{\gamma}|$ , where  $\dot{\gamma}$  is the tangent vector of  $\gamma$ . For two given points in  $\mathbb{H}^1$  we can find, by Chow's connectivity Theorem [24, p. 95], a horizontal curve joining these points. The *Carnot-Carathéodory distance*  $d_{cc}$  between two points in  $\mathbb{H}^1$  is defined as the infimum of the length of horizontal curves joining the given points. A *geodesic*  $\gamma : \mathbb{H}^1 \rightarrow \mathbb{R}$  is a horizontal curve which is a critical point of length under variations by horizontal curves. They satisfy the equation

$$(2.1) \quad D_{\dot{\gamma}} \dot{\gamma} + 2\lambda J(\dot{\gamma}) = 0,$$

where  $\lambda \in \mathbb{R}$  is the *curvature* of the geodesic, and  $J$  is the  $\pi/2$ -degrees oriented rotation in the horizontal distribution. Geodesics in  $\mathbb{H}^1$  with  $\lambda = 0$  are horizontal straight lines. The reader is referred to the section on geodesics in [29] for further details.

The *volume*  $|\Omega|$  of a Borel set  $\Omega \subseteq \mathbb{H}^1$  is the Riemannian volume of the left invariant metric  $g$ , which coincides with the Lebesgue measure in  $\mathbb{R}^3$ . We shall denote this volume element by  $dv_g$ . The *perimeter* of  $E \subset \mathbb{H}^1$  in an open subset  $\Omega \subset \mathbb{H}^1$  is defined as

$$(2.2) \quad |\partial E|(\Omega) := \sup \left\{ \int_{\Omega} \operatorname{div} U dv_g : U \text{ horizontal and } C^1, |U| \leq 1, \operatorname{supp}(U) \subset \Omega \right\},$$

where  $\operatorname{supp}(U)$  is the support of  $U$ . A set  $E \subset \mathbb{H}^1$  is of *locally finite perimeter* if  $\mathcal{P}(E, \Omega) < +\infty$  for any bounded open set  $\Omega \subset \mathbb{H}^1$ . A set of locally finite perimeter has a measurable *horizontal unit normal*  $\nu_E$ , that satisfies the following divergence theorem [17, Corollary 7.6]: if  $U$  is a horizontal vector field with compact support, then

$$\int_E \operatorname{div} U dv_g = \int_{\mathbb{H}^1} \langle U, \nu_E \rangle d|\partial E|.$$

If  $E \subset \mathbb{H}^1$  has Euclidean Lipschitz boundary, then [17, Corollary 7.7]

$$(2.3) \quad |\partial E|(\Omega) = \int_{\partial E \cap \Omega} |N_H| d\mathcal{H}^2,$$

where  $N$  is the outer unit normal to  $\partial E$ , defined  $\mathcal{H}^2$ -almost everywhere. Here  $\mathcal{H}^2$  is the 2-dimensional riemannian Hausdorff measure.

Let  $\Omega \subset \mathbb{H}^1$  be an open set. We say that  $E \subset \mathbb{H}^1$  of locally finite perimeter is *area-minimizing* in  $\Omega$  if, for any set  $F$  such that  $E = F$  outside  $\Omega$  we have

$$|\partial E|(\Omega) \leq |\partial F|(\Omega).$$

The following extension of the divergence theorem will be needed to prove the area-minimizing property of sets of locally finite perimeter

**Theorem 2.1.** *Let  $E \subset \mathbb{H}^1$  be a set of locally finite perimeter,  $B \subset \mathbb{H}^1$  a set with piecewise smooth boundary, and  $U$  a  $C^1$  horizontal vector field in  $\text{int}(B)$  that extends continuously to the boundary of  $B$ . Then*

$$(2.4) \quad \int_{E \cap B} \text{div } U \, dv_g = \int_B \langle U, \nu_E \rangle d|\partial E| + \int_E \langle U, \nu_B \rangle d|\partial B|.$$

*Proof.* The proof is modelled on [15, § 5.7]. Let  $s$  denote the riemannian distance function to  $\mathbb{H}^1 - B$ . For  $\varepsilon > 0$ , define

$$h_\varepsilon(p) := \begin{cases} 1, & \varepsilon \leq s(p), \\ s(p)/\varepsilon, & 0 \leq s(p) \leq \varepsilon, \end{cases}$$

Then  $h_\varepsilon$  is a lipschitz function (in riemannian sense). For any smooth  $h$  with compact support in  $B$  we have  $\text{div}(hU) = h \text{div}(U) + \langle \nabla h, U \rangle$ . By applying the divergence theorem for sets of locally finite perimeter [17] we get

$$\int_{\mathbb{H}^1} h \langle U, \nu_E \rangle d|\partial E| = \int_E h \text{div}(U) + \int_E \langle \nabla h, U \rangle.$$

By approximation, this formula is also valid for  $h_\varepsilon$ . Taking limits when  $\varepsilon \rightarrow 0$  we have  $\mathcal{H}_\varepsilon \rightarrow \chi_B$ . By the coarea formula for lipschitz functions

$$\frac{1}{\varepsilon} \int_{\{0 \leq s \leq \varepsilon\}} \chi_E \langle \nabla s, U \rangle = \frac{1}{\varepsilon} \int_0^\varepsilon \left\{ \int_{\{s=r\}} \chi_E \langle \nabla s, U \rangle d\mathcal{H}^2 \right\} dr,$$

and, taking again limits when  $\varepsilon \rightarrow 0$  and calling  $N_B$  to the riemannian outer unit normal to  $\partial B$  (defined except on a small set), we have

$$\lim_{\varepsilon \rightarrow 0} \int_E \langle \nabla h_\varepsilon, U \rangle = \int_{\partial B} \chi_E \langle N_B, U \rangle d\mathcal{H}^2 = \int_E \langle \nu_B, U \rangle d|\partial B|.$$

Hence (2.4) is proved.  $\square$

For a  $C^1$  surface  $\Sigma \subset \mathbb{H}^1$  the *singular set*  $\Sigma_0$  consists of those points  $p \in \Sigma$  for which the tangent plane  $T_p \Sigma$  coincides with the horizontal distribution. As  $\Sigma_0$  is closed and has empty interior in  $\Sigma$ , the *regular set*  $\Sigma - \Sigma_0$  of  $\Sigma$  is open and dense in  $\Sigma$ . It was proved in [14, Lemme 1], see also [3, Theorem 1.2], that, for a  $C^2$  surface, the Hausdorff dimension with respect to the Riemannian distance on  $\mathbb{H}^1$  of  $\Sigma_0$  is less than two.

If  $\Sigma$  is a  $C^1$  oriented surface with unit normal vector  $N$ , then we can describe the singular set  $\Sigma_0 \subset \Sigma$ , in terms of  $N_H$ , as  $\Sigma_0 = \{p \in \Sigma : N_H(p) = 0\}$ . In the regular part  $\Sigma - \Sigma_0$ ,

we can define the *horizontal unit normal vector*  $v_H$ , as in [12], [30] and [23] by

$$(2.5) \quad v_H := \frac{N_H}{|N_H|}.$$

Consider the *characteristic vector field*  $Z$  on  $\Sigma - \Sigma_0$  given by

$$(2.6) \quad Z := J(v_H).$$

As  $Z$  is horizontal and orthogonal to  $v_H$ , we conclude that  $Z$  is tangent to  $\Sigma$ . Hence  $Z_p$  generates the intersection of  $T_p\Sigma$  with the horizontal distribution. The integral curves of  $Z$  in  $\Sigma - \Sigma_0$  will be called *characteristic curves* of  $\Sigma$ . They are both tangent to  $\Sigma$  and horizontal. Note that these curves depend on the unit normal  $N$  to  $\Sigma$ . If we define

$$(2.7) \quad S := \langle N, T \rangle v_H - |N_H| T,$$

then  $\{Z_p, S_p\}$  is an orthonormal basis of  $T_p\Sigma$  whenever  $p \in \Sigma - \Sigma_0$ .

In the Heisenberg group  $\mathbb{H}^1$  there is a one-parameter group of *dilations*  $\{\varphi_s\}_{s \in \mathbb{R}}$  generated by the vector field

$$(2.8) \quad W := xX + yY + 2tT.$$

We may compute  $\varphi_s$  in coordinates to obtain

$$(2.9) \quad \varphi_s(x_0, y_0, t_0) = (e^s x_0, e^s y_0, e^{2s} t_0).$$

Conjugating with left translations we get the one-parameter family of dilations  $\varphi_{p,s} := L_p \circ \varphi_s \circ L_p^{-1}$  with center at any point  $p \in \mathbb{H}^1$ . A set  $E \subset \mathbb{H}^1$  is a *cone of center*  $p$  if  $\varphi_{p,s}(E) \subset E$  for all  $s \in \mathbb{R}$ .

Any isometry of  $(\mathbb{H}^1, g)$  leaving invariant the horizontal distribution preserves the area of surfaces in  $\mathbb{H}^1$ . Examples of such isometries are left translations, which act transitively on  $\mathbb{H}^1$ . The Euclidean rotation of angle  $\theta$  about the  $t$ -axis given by

$$(x, y, t) \mapsto r_\theta(x, y, t) = (\cos \theta x - \sin \theta y, \sin \theta x + \cos \theta y, t),$$

is also an area-preserving isometry in  $(\mathbb{H}^1, g)$  since it transforms the orthonormal basis  $\{X, Y, T\}$  at the point  $p$  into the orthonormal basis  $\{\cos \theta X + \sin \theta Y, -\sin \theta X + \cos \theta Y, T\}$  at the point  $r_\theta(p)$ .

### 3. EXAMPLES WITH ONE SINGULAR LINE

Consider the  $x$ -axis in  $\mathbb{H}^1 = \mathbb{R}^3$  parametrized by  $\Gamma(v) := (v, 0, 0)$ . Take a non-increasing continuous function  $\alpha : \mathbb{R} \rightarrow (0, \pi)$ . For every  $v \in \mathbb{R}$ , consider two horizontal halflines  $L_v^+, L_v^-$  extending from  $\Gamma(v)$  with angles  $\alpha(v)$  and  $-\alpha(v)$ , respectively. The tangent vectors to these curves at  $\Gamma(v)$  are given by  $\cos \alpha(v) X_{\Gamma(v)} + \sin \alpha(v) Y_{\Gamma(v)}$  and  $\cos \alpha(v) X_{\Gamma(v)} - \sin \alpha(v) Y_{\Gamma(v)}$ , respectively.

The parametric equations of this surface are given by

$$(3.1) \quad (v, w) \mapsto \begin{cases} (v + w \cos \alpha(v), w \sin \alpha(v), -vw \sin \alpha(v)), & w \geq 0, \\ (v + |w| \cos \alpha(v), -|w| \sin \alpha(v), v|w| \sin \alpha(v)), & w \leq 0, \end{cases}$$

One can eliminate the parameters  $v, w$  to get the implicit equation

$$t + xy - y|y| \cot \alpha \left( -\frac{t}{y} \right) = 0.$$

Letting  $\beta := \cot(\alpha)$ , we get that  $\beta$  is a continuous non-decreasing function, and that the surface  $\Sigma_\beta$  defined by the parametric equations (3.1) is given by the implicit equation

$$(3.2) \quad 0 = f_\beta(x, y, t) := t + xy - y|y| \beta \left( -\frac{t}{y} \right).$$

Observe that, because of the monotonicity condition on  $\alpha$ , the projection of relative interiors of the open horizontal halflines to the  $xy$ -plane together with the planar  $x$ -axis  $L_x$  produce a partition of the plane. Since  $\Sigma_\beta$  is the union of the horizontal lifting of these planar halflines and the  $x$ -axis to  $\mathbb{H}^1$ , it is the graph of a continuous function  $u_\beta : \mathbb{R}^2 \rightarrow \mathbb{R}$ . For  $(x, y) \in \mathbb{R}^2$ , the only point in the intersection of  $\Sigma_\beta$  with the vertical line passing through  $(x, y)$  is precisely  $(x, y, u_\beta(x, y))$ . Obviously

$$(3.3) \quad f_\beta(x, y, u_\beta(x, y)) = 0.$$

For any  $(x, y) \in \mathbb{R}^2$ , denote by  $\xi_\beta(x, y)$  the only value  $v \in \mathbb{R}$  so that either  $\Gamma(v) = (x, y, 0)$ , or  $(x, y, u_\beta(x, y))$  is contained in one of the two above described halflines leaving  $\Gamma(v)$ . Trivially  $\xi_\beta(x, 0) = x$ . Using (3.1) one checks that

$$(3.4) \quad \xi_\beta(x, y) = -\frac{u_\beta(x, y)}{y}, \quad y \neq 0.$$

Recalling that  $\alpha = \cot^{-1}(\beta)$ , we see that the mapping

$$(v, w) \mapsto \begin{cases} (v + w \cos \alpha(v), w \sin \alpha(v)), & w \geq 0, \\ (v + |w| \cos \alpha(v), -|w| \sin \alpha(v)), & w \leq 0, \end{cases}$$

is an homeomorphism of  $\mathbb{R}^2$  whose inverse is given by

$$(x, y) \mapsto (\xi_\beta(x, y), \operatorname{sgn}(y) |(x - \xi_\beta(x, y), y)|),$$

where  $\operatorname{sgn}(y) := y/|y|$  for  $y \neq 0$ . Hence  $\xi_\beta : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a continuous function. By (3.4), the function  $u_\beta(x, y)/y$  admits a continuous extension to  $\mathbb{R}^2$ .

Let us analyze first the properties of  $u_\beta$  for regular  $\beta$

**Lemma 3.1.** *Let  $\beta \in C^k(\mathbb{R})$ ,  $k \geq 2$ , be a non-decreasing function. Then*

- (i)  $u_\beta$  is a  $C^k$  function in  $\mathbb{R}^2 - L_x$ ,
- (ii)  $u_\beta$  is merely  $C^{1,1}$  near the  $x$ -axis when  $\beta \neq 0$ ,
- (iii)  $u_\beta$  is  $C^\infty$  in  $\xi^{-1}(I)$  when  $\beta \equiv 0$  on any open set  $I \subset \mathbb{R}$ , and
- (iv)  $\Sigma_\beta$  is area-minimizing.
- (v) The projection of the singular set of  $\Sigma_\beta$  to the  $xy$ -plane is  $L_x$ .

*Proof.* Along the proof we shall often drop the subscript  $\beta$  for  $f_\beta, u_\beta, \xi_\beta$  and  $\Sigma_\beta$ .

The proof of 1 is just an application of the Implicit Function Theorem since  $f_\beta$  is a  $C^k$  function for  $y \neq 0$  when  $\beta$  is  $C^k$ .

To prove 2 we compute the partial derivatives of  $u_\beta$  for  $y \neq 0$ . They are given by

$$(3.5) \quad (u_\beta)_x(x, y) = \frac{-y}{1 + |y| \beta'(\xi_\beta(x, y))},$$

$$(3.6) \quad (u_\beta)_y(x, y) = \frac{-x + |y| (2\beta(\xi_\beta(x, y)) - \beta'(\xi_\beta(x, y)) \xi_\beta(x, y))}{1 + |y| \beta'(\xi_\beta(x, y))}.$$

Since  $u_\beta(x, 0) = 0$  for all  $x \in \mathbb{R}$  we get  $(u_\beta)_x(x, 0) = 0$ . On the other hand

$$(u_\beta)_y(x, 0) = \lim_{y \rightarrow 0} \frac{u_\beta(x, y)}{y} = - \lim_{y \rightarrow 0} \xi_\beta(x, y) = -\xi_\beta(x, 0) = -x.$$

The limits, when  $y \rightarrow 0$ , of (3.5) and (3.6) can be computed using (3.4). We conclude that the first derivatives of  $u_\beta$  are continuous functions and so  $u_\beta$  is a  $C^1$  function on  $\mathbb{R}^2$ . To see that  $u_\beta$  is merely lipschitz, we get from (3.6) and (3.4)

$$\begin{aligned} (u_\beta)_{yy}(x, 0) &= \lim_{y \rightarrow 0^\pm} \frac{(u_\beta)_y(x, y) + x}{y} \\ &= \lim_{y \rightarrow 0^\pm} \frac{|y| (2\beta(\xi_\beta(x, y)) - \beta'(\xi_\beta(x, y)) \xi_\beta(x, y) + x\beta'(\xi_\beta(x, y)))}{y (1 + |y| \beta'(\xi_\beta(x, y)))} \\ &= \pm 2\beta(x). \end{aligned}$$

Hence side derivatives exist, but they do not coincide unless  $\beta(x) = 0$ .

As  $u_\beta|_{\xi^{-1}(I)} = -xy$ , 3 follows easily .

To prove 4 we use a calibration argument. We shall drop the subscript  $\beta$  to simplify the notation. Let  $F \subset \mathbb{H}^1$  such that  $F = E$  outside a Euclidean ball  $B$  centered at the origin. Let  $H^1 := \{(x, y, t) : y \geq 0\}$ ,  $H^2 := \{(x, y, t) : y \leq 0\}$ ,  $\Pi := \{(x, y, t) : y = 0\}$ . Vertical translations of the horizontal unit normal  $\nu_E$ , defined outside  $\Pi$ , provide two vector fields  $U^1$  on  $H^1$ , and  $U^2$  on  $H^2$ . They are  $C^2$  in the interior of the halfspaces and extend continuously to the boundary plane  $\Pi$ . As in the proof of Theorem 5.3 in [29], we see that

$$\operatorname{div} U^i = 0, \quad i = 1, 2,$$

in the interior of the halfspaces. Here  $\operatorname{div} U$  is the riemannian divergence of the vector field  $U$ . Observe that the vector field  $Y$  is the riemannian unit normal, and also the horizontal unit normal, to the plane  $\Pi$ . We may apply the divergence theorem to get

$$\begin{aligned} 0 &= \int_{E \cap \operatorname{int}(H^i) \cap B} \operatorname{div} U^i = \int_E \langle U^i, \nu_{\operatorname{int}(H^i) \cap B} \rangle d|\partial(\operatorname{int}(H^i) \cap B)| \\ &\quad + \int_{\operatorname{int}(H^i) \cap B} \langle U^i, \nu_E \rangle d|\partial E|. \end{aligned}$$

Let  $D := \Pi \cap \bar{B}$ . Then, for every  $p \in D$ , we have  $\nu_{\operatorname{int}(H^1) \cap B} = -Y$ ,  $\nu_{\operatorname{int}(H^2) \cap B} = Y$ , and  $U^1 = J(v)$ ,  $U^2 = J(w)$ , where  $v - w$  is proportional to  $Y$ , by the construction of  $\Sigma_\beta$ . Hence

$$\langle U^1, \nu_{\operatorname{int}(H^1) \cap B} \rangle + \langle U^2, \nu_{\operatorname{int}(H^2) \cap B} \rangle = \langle v - w, J(Y) \rangle = 0, \quad p \in D.$$

Adding the above integrals we obtain

$$0 = \sum_{i=1,2} \int_E \langle U^i, \nu_B \rangle d|\partial B| + \sum_{i=1,2} \int_{B \cap \text{int}(H^i)} \langle U^i, \nu_E \rangle d|\partial E|.$$

We apply the same arguments to the set  $F$  and, since  $E = F$  on  $\partial B$  we conclude

$$(3.7) \quad \sum_{i=1,2} \int_{B \cap \text{int}(H^i)} \langle U^i, \nu_E \rangle d|\partial E| = \sum_{i=1,2} \int_{B \cap \text{int}(H^i)} \langle U^i, \nu_F \rangle d|\partial F|.$$

As  $E$  is a subgraph,  $|\partial E|(\Pi) = 0$  and so

$$|\partial E|(B) = \sum_{i=1,2} \int_{B \cap \text{int}(H^i)} \langle U^i, \nu_B \rangle d|\partial E|.$$

Cauchy-Schwarz inequality and the fact that  $|\partial F|$  is a positive measure imply

$$\sum_{i=1,2} \int_{B \cap \text{int}(H^i)} \langle U^i, \nu_F \rangle d|\partial F| \leq |\partial F|(B),$$

which implies 4.

To prove 5 simply take into account that the projection of the singular set of  $\Sigma_\beta$  to the  $xy$ -plane is composed of those points  $(x, y)$  such that  $(u_\beta)_x - x - y = (u_\beta)_y + x = 0$ . From (3.5) we get that  $(u_\beta)_x - y = 0$  if and only if

$$y(2 + |y| \beta'(\xi_\beta(x, y))) = 0,$$

i.e, when  $y = 0$ . In this case, from (3.6), we see that equation  $(u_\beta)_y + x = 0$  is trivially satisfied.  $\square$

We now prove the general properties of  $\Sigma_\beta$  from Lemma 3.1

**Proposition 3.2.** *Let  $\beta : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous non-decreasing function. Let  $u_\beta$  be the only solution of equation (3.3),  $\Sigma_\beta$  the graph of  $u_\beta$ , and  $E_\beta$  the subgraph of  $u_\beta$ . Then*

- (i)  $u_\beta$  is locally lipschitz in Euclidean sense,
- (ii)  $E_\beta$  is a set of locally finite perimeter in  $\mathbb{H}^1$ , and
- (iii)  $\Sigma_\beta$  is area-minimizing in  $\mathbb{H}^1$ .

*Proof.* Let

$$\beta_\varepsilon(x) := \int_{\mathbb{R}} \beta(y) \eta_\varepsilon(x - y) dy$$

the usual convolution, where  $\eta$  is a Dirac function and  $\eta_\varepsilon(x) := \eta(x/\varepsilon)$ , see [15]. Then  $\beta_\varepsilon$  is a  $C^\infty$  non-decreasing function, and  $\beta_\varepsilon$  converges uniformly, on compact subsets of  $\mathbb{R}$ , to  $\beta$ . Let  $u = u_\beta$ ,  $u_\varepsilon = u_{\beta_\varepsilon}$ ,  $f = f_\beta$ ,  $f_\varepsilon = f_{\beta_\varepsilon}$ .

Let  $D \subset \mathbb{R}^2$  be a bounded subset. To check that  $u$  is lipschitz on  $D$  it is enough to prove that the first derivatives of  $u_\varepsilon$  are uniformly bounded on  $D$ .

From (3.3) we get

$$\xi(x, y) + |y| \beta(\xi(x, y)) = x, \quad y \neq 0.$$

For  $y$  fixed, define the continuous strictly increasing function

$$\rho_y(x) := x + |y| \beta(x).$$



Hence we get

$$(3.8) \quad \tilde{\zeta}(x, y) = \rho_y^{-1}(x).$$

We can also define  $(\rho_\varepsilon)_y(x) := x + |y|\beta_\varepsilon(x)$ . Equation (3.8) holds replacing  $u, \beta$  by  $u_\varepsilon, \beta_\varepsilon$ .

Since  $\rho_y^{-1}(x) = \tilde{\zeta}(x, y)$ , we conclude that  $\rho_y^{-1}$  is a continuous function that depends continuously on  $y$ .

Let us estimate

$$|(\rho_\varepsilon)_y^{-1}(x) - \rho_y^{-1}(x)|.$$

Let  $z_\varepsilon := (\rho_\varepsilon)_y^{-1}(x)$ ,  $z = \rho_y^{-1}(x)$ . Then  $x = (\rho_\varepsilon)_y(z_\varepsilon) = \rho_y(z)$  and we have, assuming  $z_\varepsilon \geq z$ .

$$\begin{aligned} 0 &= (\rho_\varepsilon)_y(z_\varepsilon) - \rho_y(z) = z_\varepsilon + |y|\beta_\varepsilon(z_\varepsilon) - (z + |y|\beta(z)) \\ &= (z_\varepsilon - z) + |y|(\beta_\varepsilon(z_\varepsilon) - \beta_\varepsilon(z)) + |y|(\beta_\varepsilon(z) - \beta(z)) \\ &\geq (z_\varepsilon - z) + |y|(\beta_\varepsilon(z) - \beta(z)). \end{aligned}$$

A similar computation can be performed for  $z_\varepsilon \leq z$ . The consequence is that

$$|z_\varepsilon - z| \leq |y| |\beta_\varepsilon(z) - \beta(z)|,$$

or, equivalently,

$$|(\rho_\varepsilon)_y^{-1}(x) - \rho_y^{-1}(x)| \leq |y| |\beta_\varepsilon(\rho_y^{-1}(x)) - \beta(\rho_y^{-1}(x))|.$$

As  $\beta_\varepsilon \rightarrow \beta$  uniformly on compact subsets of  $\mathbb{R}$ , we have uniform convergence of  $(\rho_\varepsilon)_y^{-1}(x)$  to  $\rho_y^{-1}(x)$  on compact subsets of  $\mathbb{R}^2$ . This also implies the uniform convergence of  $\tilde{\zeta}_\varepsilon(x, y)$  to  $\tilde{\zeta}(x, y)$  on compact subsets. Hence also  $u_\varepsilon(x, y)$  converges uniformly to  $u(x, y)$  on compact subsets of  $\mathbb{R}^2$ .

From (3.5) and (3.6) we have

$$\begin{aligned} |(u_\varepsilon)_x(x, y)| &\leq |y|, \\ |(u_\varepsilon)_y(x, y)| &\leq |x| + 2|y| |\beta_\varepsilon(\tilde{\zeta}_\varepsilon(x, y))| + |\tilde{\zeta}_\varepsilon(x, y)|. \end{aligned}$$

As  $\beta_\varepsilon \rightarrow \beta$  and  $\tilde{\zeta}_\varepsilon(x, y) \rightarrow \tilde{\zeta}(x, y)$  uniformly on compact subsets, we have that the first derivatives of  $u_\varepsilon$  are uniformly bounded on compact subsets. Hence  $u$  is locally lipschitz.

The subgraph of  $u_\beta$  is a set of locally finite perimeter in  $\mathbb{H}^1$  since its boundary is locally lipschitz by 1. This follows from [17] and proves 2.

To prove 3 we use approximation and the calibration argument. Let  $F \subset \mathbb{H}^1$  so that  $F = \bar{E}$  outside a Euclidean ball  $B$  centered at the origin. For the functions  $\beta_\varepsilon$ , consider the vector fields  $U_\varepsilon^i$  obtained by translating vertically the horizontal unit normal to the surface  $\Sigma_\varepsilon$ . We repeat the arguments on the proof of 4 in Lemma 3.1 to conclude as in (3.7) that

$$\sum_{i=1,2} \int_{B \cap \text{int}(H^i)} \langle U_\varepsilon^i, \nu_E \rangle d|\partial E| = \sum_{i=1,2} \int_{B \cap \text{int}(H^i)} \langle U_\varepsilon^i, \nu_F \rangle d|\partial F|.$$

Trivially we have

$$\sum_{i=1,2} \int_{B \cap \text{int}(H^i)} \langle U_\varepsilon^i, \nu_F \rangle d|\partial F| \leq |\partial F|(B).$$

On the other hand,  $U_\varepsilon^i$  converges uniformly, on compact subsets, to  $U^i$  by Lemma 3.3. Passing to the limit when  $\varepsilon \rightarrow 0$  and taking into account that  $U^i = \nu_E$  we conclude

$$|\partial E|(B) \leq |\partial F|(B),$$

as desired.  $\square$

**Lemma 3.3.** *Let  $\beta$  be a continuous non-decreasing function. Then the horizontal unit normal of  $\Sigma_\beta$  is given, in  $\{X, Y\}$ -coordinates, by*

$$(3.9) \quad \nu_\beta(x, y) = \left( \frac{1}{(1 + \beta^2)^{1/2}}, \frac{-\operatorname{sgn}(y) \beta}{(1 + \beta^2)^{1/2}} \right) (\xi_\beta(x, y)), \quad y \neq 0.$$

Moreover,  $\nu_\beta$  admits continuous extensions to  $y = 0$  from both sides of this line.

*Proof.* Since  $u_\beta$  is lipschitz, it is differentiable almost everywhere on  $\mathbb{R}^2$ . On these points,

$$\nu_\beta(x, y) = ((u_\beta)_x - y, (u_\beta)_y + x).$$

The function  $-u_\beta(x, y)/y$  is constant along the lines  $(x_0, 0) + \lambda(1 + \beta^2)^{-1/2}(\beta, \pm 1)(x_0)$ , for  $\lambda \geq 0$ . Let  $y \geq 0$ . From (3.2) we have

$$0 = -x_0 + x - y\beta(x_0).$$

Let  $v := (1 + \beta^2)^{-1/2}(\beta, 1)(x_0)$ . Then  $v(-u_\beta(x, y)/y) = 0$ . Hence for almost every point on almost every line, we have

$$\beta(x_0)(u_\beta)_x + (u_\beta)_y = -x_0.$$

Hence we have

$$(u_\beta)_y + x = -x_0 - \beta(x_0)(u_\beta)_x + x_0 + y\beta(x_0) = \beta(x_0)(-(u_\beta)_x + y).$$

We conclude that the horizontal unit normal is proportional to  $(1, -\beta)$ , which implies (3.9). The case  $y \leq 0$  is handled similarly.  $\square$

**Example 3.4.** Taking  $\beta(x) := x$  we get

$$u_\beta(x, y) = -\frac{xy}{1 + |y|},$$

which is a Euclidean  $C^{1,1}$  graph.

Another family of interesting examples are the minimal cones obtained by taking the constant function  $\beta(x) := \beta_0$ . In this case we get

$$u_\beta(x, y) = -xy + \beta_0 y|y|.$$

In this case  $\Sigma_\beta$  is a  $C^{1,1}$  surface which is invariant by the dilations centered at any point of the singular line.

Take now

$$\beta(x) := \begin{cases} 0, & x \leq 0, \\ x, & x \geq 0. \end{cases}$$

In this case we obtain the graph

$$u_\beta(x, y) := \begin{cases} -xy, & x \leq 0, \\ -\frac{xy}{1+|y|}, & x \geq 0, \end{cases}$$

which is simply locally Lipschitz.

This example was mentioned to me by Scott Pauls. Consider now a continuous non-decreasing function  $\beta : \mathbb{R} \rightarrow \mathbb{R}$ , constant outside the Cantor set  $C \subset [0, 1]$  with  $\beta(0) = 0$ ,  $\beta(1) = 1$ . Then the associated surface  $\Sigma_\beta$  is an area-minimizing surface in  $\mathbb{H}^1$ .

#### 4. EXAMPLES WITH SEVERAL SINGULAR HALFLINES MEETING AT A POINT

Let  $\alpha_1^0, \dots, \alpha_k^0$  be a family of positive angles so that

$$\sum_{i=1}^k \alpha_i^0 = \pi.$$

Let  $r_\beta$  be the rotation of angle  $\beta$  around the origin in  $\mathbb{R}^2$ . Consider a family of closed halflines  $L_i \subset \mathbb{R}^2$ ,  $i \in \mathbb{Z}_k$ , extending from the origin, so that  $r_{\alpha_i^0 + \alpha_{i+1}^0}(L_i) = L_{i+1}$ . Finally, define  $R_i := r_{\alpha_i^0}(L_i)$ . (An alternative way of defining this configuration is to start from a family of counter-clockwise oriented halflines  $R_i \subset \mathbb{R}^2$ ,  $i \in \mathbb{Z}_k$ , choosing  $L_i$ ,  $i \in \mathbb{Z}_k$ , as the bisector of the angle determined by  $R_{i-1}$  and  $R_i$ , and defining  $\alpha_i^0$  as the angle between  $L_i$  and  $R_i$ ). Define  $W_i$  as the closed wedge, containing  $L_i$ , bordered by  $R_{i-1}$  and  $R_i$ .




FIGURE 1. The initial configuration with three halflines  $L_1, L_2, L_3$ .

For every  $i \in \mathbb{Z}_k$ , let  $\alpha_i : [0, \infty) \rightarrow (0, \pi)$  be a continuous nonincreasing function so that  $\alpha_i(0) = \alpha_i^0$ , and define, as in the previous section,  $\beta_i := \cot(\alpha_i)$ . Let  $v_i \in \mathbb{S}^1$ ,  $i \in \mathbb{Z}_k$ , be such that  $L_i = \{sv_i : s \geq 0\}$ . For every  $i \in \mathbb{Z}_k$  and  $s \geq 0$ , we take the two closed halflines  $L_{s,i}^\pm$  in  $\mathbb{R}^2$  extending from the point  $sv_i$  with tangent vectors  $(\cos \alpha_i(s), \pm \sin \alpha_i(s))$ . In this way we cover all of  $\mathbb{R}^2$ . We shall define  $\alpha := (\alpha_1, \dots, \alpha_k)$ .

Lift  $L_1, \dots, L_k$  to horizontal halflines  $L'_1, \dots, L'_k$  in  $\mathbb{H}^1$  from the origin, and  $L_{s,i}^\pm$  to horizontal halflines in  $\mathbb{H}^1$  extending from the unique point in  $L'_i$  projecting onto  $sv_i$ . In this way we obtain a continuous function  $u_\alpha : \mathbb{R}^2 \rightarrow \mathbb{R}$ . The graph  $\Sigma_\alpha$  of  $u_\alpha$  is a topological surface in  $\mathbb{H}^1$ .

Obviously the angle functions  $\alpha_i(s)$  can be extended continuously and preserving the monotonicity, to an angle function  $\tilde{\alpha}_i : \tilde{L}_i \rightarrow (0, \pi)$ , where  $\tilde{L}_i$  is the straight line containing the halfline  $L_i$ . The graph of  $u_\alpha$  restricted to  $W_i$  coincides with the Euclidean locally lipschitz area-minimizing surface  $u_{\tilde{\beta}_i}$ , for  $\tilde{\beta}_i := \cot \tilde{\alpha}_i$ , constructed in the previous section. So the examples in this section can be seen as pieces of the examples of the previous one patched together.

**Theorem 4.1.** *Under the above conditions*

- (i) *The function  $u_\alpha$  is locally lipschitz in the Euclidean sense.*
- (ii) *The surface  $\Sigma_\alpha$  is area-minimizing.*

*Proof.* It is immediate that  $u_\alpha$  is a graph which is locally lipschitz in Euclidean sense: choose a disk  $D \subset \mathbb{R}^2$ . Let  $p, q \in D$ . Assume first that  $(p, q)$  intersects the halflines  $R_1, \dots, R_k$  transversally at the points  $x_1, \dots, x_n$ . Then  $[p, x_1], [x_1, x_2], \dots, [x_n, p]$  are contained in wedges and hence

$$\begin{aligned} |u_\alpha(p) - u_\alpha(q)| &\leq |u_\alpha(p) - u_\alpha(x_1)| + \dots + |u_\alpha(x_n) - u_\alpha(q)| \\ &C (|p - x_1| + \dots + |x_n - q|) = C |p - q|, \end{aligned}$$

where  $C$  is the supremum of the Lipschitz constants of  $u_{\tilde{\beta}_i}$  restricted to  $D$ . The general case is then obtained by approximating  $p$  and  $q$  by points in the condition of the assumption.

To prove that  $u_\alpha$  is area minimizing we first approximate  $\alpha_i$  by smooth angle functions  $(\alpha_i)_\varepsilon$  with  $(\alpha_i)_\varepsilon(0) = \alpha_i(0)$ . In this way we obtain a calibrating vector field which is continuous along the vertical planes passing through  $R_i$  by Lemma 3.3. This allows us to apply the calibration argument to prove the area-minimizing property of  $\Sigma_\alpha$ .  $\square$

**Example 4.2** (Minimizing cones). Let  $\alpha_i(s) = \alpha_i^0$  be a constant for all  $i$ . Then the subgraph of  $\Sigma_\alpha$  is a minimizing cone with center at 0. Restricted to the interior of the wedges  $W_i$ , the surface  $\Sigma_\alpha$  is  $C^{1,1}$ . An easy computation shows that, taking  $\beta(s) := \beta_0$  in the construction of the first section, the Riemannian normal to  $\Sigma_\beta$  along the halflines  $\beta_0|y| = x, x \geq 0$  (that make angle  $\pm \cot^{-1}(\beta_0)$  with the positive  $x$ -axis) is given by

$$N = \frac{-2yX + 2\beta_0|y|Y - T}{\sqrt{1 + 4y^2 + 4\beta_0^2y^2}} = \frac{-2yX + 2xY - T}{\sqrt{1 + 4x^2 + 4y^2}}.$$

This vector field is invariant by rotations around the vertical axis. Hence in our construction, the normal vector field to  $\Sigma_\alpha$  is continuous. It is straightforward to show that it is locally lipschitz in Euclidean sense.

**Example 4.3** (Area-minimizing surfaces with a singular halfline). These examples are inspired by [9, Example 7.2]. We consider a halfline  $L$  extending from the origin, and an angle function  $\alpha : L \rightarrow (0, \pi)$  continuous and nonincreasing as a function of the distance to the origin. We consider the union of the halflines  $L_{\alpha(q)}^+, L_{\alpha(q)}^-$  extending from  $q \in L$  with angles  $\alpha(q), -\alpha(q)$ , respectively. We patch the area-minimizing surface defined by  $\alpha$  in the wedge delimited by the halflines  $L_0^+, L_0^-$ , with the plane  $t = 0$ . In this way we get an entire area-minimizing  $t$ -graph, with lipschitz regularity. In case the angle function  $\alpha$  is

constant, we get an area-minimizing cone with center 0, which is defined by the equation

$$u(x, y) := \begin{cases} -xy + \beta_0 y|y|, & -xy + \beta_0 y|y| \geq 0, \\ 0, & -xy + \beta_0 y|y| \leq 0. \end{cases}$$

This surface is composed of two smooth pieces patched together along the halflines  $x = \beta_0|y|$ .

#### REFERENCES

- [1] Luigi Ambrosio, Bruce Kleiner, and Enrico Le Donne, *Rectifiability of sets of finite perimeter in Carnot groups: existence of a tangent hyperplane*, arXiv math.DG/0801.3741v1, 2008.
- [2] Luigi Ambrosio, Francesco Serra Cassano, and Davide Vittone, *Intrinsic regular hypersurfaces in Heisenberg groups*, J. Geom. Anal. **16** (2006), no. 2, 187–232. MR MR2223801 (2007g:49072)
- [3] Zoltán M. Balogh, *Size of characteristic sets and functions with prescribed gradient*, J. Reine Angew. Math. **564** (2003), 63–83. MR MR2021034 (2005d:43007)
- [4] Vittorio Barone Adesi, Francesco Serra Cassano, and Davide Vittone, *The Bernstein problem for intrinsic graphs in Heisenberg groups and calibrations*, Calc. Var. Partial Differential Equations **30** (2007), no. 1, 17–49. MR MR2333095
- [5] Francesco Bigolin and Francesco Serra Cassano, *Intrinsic regular graphs in Heisenberg groups vs. weak solutions of non linear first-order PDEs*, work in progress, October 2007.
- [6] Luca Capogna, Giovanna Citti, and Maria Manfredini, *Smoothness of lipschitz minimal intrinsic graphs in Heisenberg groups  $\mathbb{H}^n$ ,  $n > 1$* , preprint, 2008.
- [7] Luca Capogna, Donatella Danielli, Scott D. Pauls, and Jeremy T. Tyson, *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*, Progress in Mathematics, vol. 259, Birkhäuser Verlag, Basel, 2007. MR MR2312336
- [8] Jih-Hsin Cheng, Jenn-Fang Hwang, Andrea Malchiodi, and Paul Yang, *Minimal surfaces in pseudohermitian geometry*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **4** (2005), no. 1, 129–177. MR MR2165405 (2006f:53008)
- [9] Jih-Hsin Cheng, Jenn-Fang Hwang, and Paul Yang, *Existence and uniqueness for  $p$ -area minimizers in the Heisenberg group*, Math. Ann. **337** (2007), no. 2, 253–293. MR MR2262784
- [10] Jih-Hsin Cheng, Jenn-Fang Hwang, and Paul Yang, *Regularity of  $C^1$  smooth surfaces with prescribed  $p$ -mean curvature in the Heisenberg group*, arXiv math.DG/0709.1776 v1, 2007.
- [11] D. Danielli, N. Garofalo, and D. M. Nhieu, *A notable family of entire intrinsic minimal graphs in the Heisenberg group which are not perimeter minimizing*, Amer. J. Math. (to appear).
- [12] D. Danielli, N. Garofalo, and D. M. Nhieu, *Sub-Riemannian calculus on hypersurfaces in Carnot groups*, Adv. Math. **215** (2007), no. 1, 292–378. MR MR2354992
- [13] D. Danielli, N. Garofalo, D. M. Nhieu, and S. D. Pauls, *Instability of graphical strips and a positive answer to the Bernstein problem in the Heisenberg group*, J. Differential Geom. (to appear).
- [14] Maklouf Derridj, *Sur un théorème de traces*, Ann. Inst. Fourier (Grenoble) **22** (1972), no. 2, 73–83. MR MR0343011 (49 #7755)
- [15] Lawrence C. Evans and Ronald F. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1992. MR MR1158660 (93f:28001)
- [16] Bruno Franchi, Raul Serapioni, and Francesco Serra Cassano, *Sur les ensembles de périmètre fini dans le groupe de Heisenberg*, C. R. Acad. Sci. Paris Sér. I Math. **329** (1999), no. 3, 183–188. MR MR1711057 (2000e:49008)
- [17] ———, *Rectifiability and perimeter in the Heisenberg group*, Math. Ann. **321** (2001), no. 3, 479–531. MR MR1871966 (2003g:49062)
- [18] ———, *On the structure of finite perimeter sets in step 2 Carnot groups*, J. Geom. Anal. **13** (2003), no. 3, 421–466. MR MR1984849 (2004i:49085)
- [19] ———, *Regular hypersurfaces, intrinsic perimeter and implicit function theorem in Carnot groups*, Comm. Anal. Geom. **11** (2003), no. 5, 909–944. MR MR2032504 (2004m:28008)
- [20] ———, *Intrinsic Lipschitz graphs in Heisenberg groups*, J. Nonlinear Convex Anal. **7** (2006), no. 3, 423–441. MR MR2287539
- [21] ———, *Regular submanifolds, graphs and area formula in Heisenberg groups*, Adv. Math. **211** (2007), no. 1, 152–203. MR MR2313532

- [22] Nicola Garofalo and Duy-Minh Nhieu, *Isoperimetric and Sobolev inequalities for Carnot-Carathéodory spaces and the existence of minimal surfaces*, *Comm. Pure Appl. Math.* **49** (1996), no. 10, 1081–1144. MR MR1404326 (97i:58032)
- [23] Nicola Garofalo and Scott D. Pauls, *The Bernstein Problem in the Heisenberg Group*, arXiv math.DG/0209065 v2, 2002.
- [24] Mikhael Gromov, *Carnot-Carathéodory spaces seen from within*, *Sub-Riemannian geometry*, *Progr. Math.*, vol. 144, Birkhäuser, Basel, 1996, pp. 79–323. MR MR1421823 (2000f:53034)
- [25] Roberto Monti and Francesco Serra Cassano, *Surface measures in Carnot-Carathéodory spaces*, *Calc. Var. Partial Differential Equations* **13** (2001), no. 3, 339–376. MR MR1865002 (2002j:49052)
- [26] Pierre Pansu, *Une inégalité isopérimétrique sur le groupe de Heisenberg*, *C. R. Acad. Sci. Paris Sér. I Math.* **295** (1982), no. 2, 127–130. MR MR676380 (85b:53044)
- [27] Scott D. Pauls, *Minimal surfaces in the Heisenberg group*, *Geom. Dedicata* **104** (2004), 201–231. MR MR2043961 (2005g:35038)
- [28] ———, *H-minimal graphs of low regularity in  $\mathbb{H}^1$* , *Comment. Math. Helv.* **81** (2006), no. 2, 337–381. MR MR2225631 (2007g:53032)
- [29] Manuel Ritoré and César Rosales, *Area-stationary surfaces in the Heisenberg group  $\mathbb{H}^1$* , arXiv.org:math/0512547 v2, 2005.
- [30] ———, *Rotationally invariant hypersurfaces with constant mean curvature in the Heisenberg group  $\mathbb{H}^n$* , *J. Geom. Anal.* **16** (2006), no. 4, 703–720. MR MR2271950

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, UNIVERSIDAD DE GRANADA, E-18071 GRANADA, ESPAÑA

E-mail address: ritore@ugr.es