## Self-consistent crystalline condensate in chiral Gross-Neveu and Bogoliubov-de Gennes systems

Gökçe Başar and Gerald V. Dunne

Physics Department, University of Connecticut, Storrs CT 06269

We derive a new exact self-consistent crystalline condensate in the 1+1 dimensional chiral Gross-Neveu model. This also yields a new exact crystalline solution for the one dimensional Bogoliubov-de Gennes equations and the Eilenberger equation of semiclassical superconductivity. We show that the functional gap equation can be reduced to a solvable nonlinear equation, and discuss implications for the temperature-chemical potential phase diagram.

Interacting fermion systems describe a wide range of physical phenomena, from particle physics, to solid state and atomic physics [1, 2, 3, 4]. Important paradigms include the Peierls-Frohlich model of conduction [5], the Gorkov-Bogoliubov-de Gennes approach to superconductivity [6], and the Nambu-Jona Lasinio (NJL) model of symmetry breaking in particle physics [7]. A 1 + 1dimensional version of the NJL model, the NJL<sub>2</sub> model [also known as the *chiral* Gross-Neveu model,  $\chi GN_2$ ] has been widely studied as it exhibits asymptotic freedom, dynamical mass generation, and chiral symmetry breaking [8, 9, 10, 11]. Surprisingly, the temperature-density phase diagram of this system is not yet fully understood. A gap equation analysis based on a homogeneous condensate suggests its phase diagram is the same as its discrete-chiral cousin, the original Gross-Neveu  $(GN_2)$ model [8], while recent work finds an inhomogeneous Larkin-Ovchinikov-Fulde-Ferrell (LOFF) helical complex condensate ("chiral spiral") below a critical temperature [12]. In fact, the phase diagram of the discrete-chiral version, the  $GN_2$  model, has only recently been solved in the particle physics literature, analytically [13], and on the lattice [14]. There is a crystalline phase at low temperature and high density, and this phase is characterized by a periodically inhomogeneous (real) condensate that solves exactly the gap equation. This phase is not seen in the old phase diagram which was based on a uniform condensate [15]. Interestingly, this discrete-chiral  $GN_2$ model (with vanishing bare fermion mass) is mathematically equivalent to several models in condensed matter physics: the real periodic condensate may be identified with a polaron crystal in conducting polymers [1, 16, 17], with a periodic pair potential in guasi 1D superconductors [18, 19], and with the real order parameter for superconductors in a ferromagnetic field [20]. This system also is a paradigm of the phenomenon of fermion number fractionalization [21, 22]. Variants of such models also apply to ultracold fermionic systems, for which there are interesting new theoretical and experimental developments [4, 23, 24].

In this paper we present an analogous *complex* crystalline condensate for the *chiral* GN system, the  $NJL_2$  model. This condensate is an exact inhomogeneous solution to the gap equation, and also provides a new self-

consistent solution to the Bogoliubov-de Gennes (BdG) [6] and Eilenberger [25] equations of superconductivity. Our solution may also be relevant for chiral superconductors and for incommensurate charge density waves in quasi 1D systems [26], which have chiral symmetry and an inherently complex order parameter.

Consider the massless  $NJL_2$  model with Lagrangian

$$\mathcal{L} = \bar{\psi} \, i \, \partial \!\!\!/ \psi + \frac{g^2}{2} \left[ \left( \bar{\psi} \psi \right)^2 + \left( \bar{\psi} i \gamma^5 \psi \right)^2 \right] \quad , \qquad (1)$$

which has a continuous chiral symmetry  $\psi \to e^{i\gamma^5 \alpha}\psi$ . We have suppressed summation over N flavors, which makes the semiclassical gap equation analysis exact in the  $N \to \infty$  limit, a limit in which we can consistently discuss chiral symmetry breaking in 2D. The original GN<sub>2</sub> model [8], without the pseudoscalar interaction term  $(\bar{\psi}i\gamma^5\psi)^2$ , has a discrete chiral symmetry  $\psi \to \gamma^5\psi$ . There are two equivalent ways to find self-consistent static condensates. First, introduce bosonic condensate fields,  $S \equiv \bar{\psi}\psi$  and  $P \equiv \bar{\psi}i\gamma^5\psi$ , which we combine into a complex condensate field:  $\Delta \equiv S - i P \equiv M e^{i\chi}$ . Integrating out the fermion fields we obtain an effective action for the condensate as

$$S_{\text{eff}} = -\frac{1}{2Ng^2} \int |\Delta|^2 - i \, \ln \det \left[ i \, \partial \!\!\!/ - M \, e^{-i\gamma^5 \chi} \right] \quad (2)$$

The corresponding (complex) gap equation is

$$\Delta(x) = -2iNg^2 \frac{\delta}{\delta\Delta(x)^*} \ln \det \left[ i \partial - M(x) e^{-i\gamma^5 \chi(x)} \right]$$
(3)

One of the main results of this paper is that this gap equation can be reduced in an elementary manner to an ordinary differential equation, which moreover is soluble. A second approach to finding a self-consistent condensate is to solve the relativistic Hartree-Fock problem  $H\psi = E\psi$ , subject to the consistency condition  $\langle \bar{\psi}\psi \rangle - i \langle \bar{\psi}i\gamma^5\psi \rangle = -\Delta/g^2$ , with single-particle Hamiltonian

$$H = -i\gamma^5 \frac{d}{dx} + \gamma^0 M(x) e^{-i\gamma^5 \chi(x)} = \begin{pmatrix} -i\frac{d}{dx} & \Delta(x) \\ \Delta^*(x) & i\frac{d}{dx} \end{pmatrix}$$
(4)

This is the Bogoliubov-de Gennes (BdG), or Andreev, hamiltonian, with  $\Delta(x)$  playing the role of the order parameter [6]. We have chosen Dirac matrices  $\gamma^0 = \sigma_1$ ,  $\gamma^1 = -i\sigma_2$ , and  $\gamma^5 = \sigma_3$ , to emphasize the natural complex combination  $\Delta = S - iP$ . The key object in our analysis is the Gork'ov Green's function, and in particular its coincident-point limit, the "diagonal resolvent":

$$R(x; E) \equiv \langle x | \frac{1}{H - E} | x \rangle \tag{5}$$

This is a  $2 \times 2$  matrix, and the spectral function is  $\rho(E) = \frac{1}{\pi} \text{Im } \text{Tr}_{D,x} R(x; E + i\epsilon)$ , where the trace is both a Dirac and spatial trace. Approximation methods, such as the gradient and semiclassical expansions, of R(x; E) have been widely studied [27]. In one spatial dimension, R(x; E) can be written in terms of two independent solutions  $\psi_{1,2}$  to the Dirac/BdG equation  $H\psi = E\psi$ . That is,  $R(x; E) = (\psi_1\psi_2^T + \psi_2\psi_1^T) \sigma_1/(2W)$ , with Wronskian  $W = -i\psi_1^T\sigma_2\psi_2$ . It follows immediately that R(x; E) satisfies the following first order equation:

$$R'\gamma^5 = i \left[\gamma^5 (E - \gamma^0 M e^{-i\gamma^5 \chi}), R\gamma^5\right]$$
(6)

In superconductivity, (6) is known as the Eilenberger equation [25], and in mathematical physics as the Dik'ii equation [28]. It is also straightforward to show that  $R = R^{\dagger}$ , det R = 1/4, and tr $(R \gamma^5) = 0$ .

Our main observation is that the gap equation (3) motivates a self-consistent ansatz form of the  $2 \times 2$  matrix R(x; E), and when this is combined with the identity (6), the exact self-consistent condensate and associated resolvent are completely determined. To see this, note that the gap equation (3) can be viewed in two ways. First, write the log det term as

$$\int_{-\infty}^{\infty} dE \,\rho(E) \frac{1}{\beta} \ln\left(1 + e^{-\beta(E-\mu)}\right) \tag{7}$$

All dependence on  $\Delta(x)$  resides in the spectral function  $\rho(E)$ , via Tr R, and so the simplest solution to the gap equation (3) is for *diagonal* entries of the 2 × 2 matrix R [recall they are equal since  $\operatorname{tr}(R\gamma^5) = 0$ ] to be linear in  $|\Delta|^2$ . On the other hand, we can also express the gap equation, by performing the functional derivatives, as

$$\Delta(x) = iNg^2 \operatorname{tr}_{D,E} \left[ \left( \gamma^0 + \gamma^1 \right) R(x; E) \right]$$
(8)

This suggests that the *off-diagonal* entries of R be proportional to  $\Delta$  and  $\Delta^*$ . In fact, consistency between (6) and (8) introduces derivative terms, leading to

$$R(x; E) = \mathcal{N}(E) \begin{pmatrix} a(E) + |\Delta|^2 & b(E)\Delta - i\Delta' \\ b(E)\Delta^* + i\Delta'^* & a(E) + |\Delta|^2 \end{pmatrix}$$
(9)

The gap equation is satisfied since we find  $\operatorname{tr}_E \mathcal{N}(E) = 0$ . With this ansatz, the diagonal entries of (6) are identically satisfied, while the off-diagonal entries imply that the condensate  $\Delta$  satisfy the equation

$$\Delta'' - 2|\Delta|^2 \,\Delta + i \,(b - 2E) \,\Delta' - 2 \,(a - Eb) \,\Delta = 0 \quad (10)$$

This nonlinear Schrödinger equation (NLSE) is analytically soluble, and all previously known examples of selfconsistent condensates in  $GN_2$  and  $NJL_2$  are special cases. Furthermore, the corresponding Dirac/BdG equation  $H\psi = E\psi$  is also exactly soluble, and has a spectrum consisting of a *single band* in the gap [30]. This provides an elementary explanation of the result from inverse scattering [9, 10, 11] that self-consistent ground state condensates are reflectionless (or finite-gap) with a single state (or band). For example, if we specialize to a real condensate, as is relevant for the [discrete-chiral] GN<sub>2</sub> model, the [rescaled] self-consistent solution is  $\Delta(x) = \sqrt{\nu} \operatorname{sn}(x; \nu)$ , which satisfies  $\Delta'' - 2\Delta^3 = -(1+\nu)\Delta$ , where sn is a Jacobi elliptic function, with elliptic parameter  $0 < \nu < 1$ [16, 17, 18, 19, 20]. The fermion spectrum has a single band in the gap, centered on E = 0, reflecting the chargeconjugation symmetry of the GN<sub>2</sub> system. As  $\nu \to 1$ , the period becomes infinite and we obtain the famous kink,  $\Delta(x) = \tanh(x)$ , with a single bound state at E = 0. This mid-gap zero mode has many interesting physical consequences in polymer systems, and is the paradigm of the fractional fermion number phenomenon [21]. It is also worth noting that for the [discrete-chiral] GN<sub>2</sub> model with a bare fermion mass, the NLSE (10) acquires an inhomogeneous term, in which case the general solution is written as  $\Delta(x) = \zeta(\gamma) + \zeta(x) - \zeta(x + \gamma)$ , where  $\zeta$  is the Weierstrass zeta function, and this represents a kink-antikink crystal (or bipolaron crystal in the polymer language [29]), which is a periodic generalization [13] of the Dashen-Hasslacher-Neveu (DHN) kink-antikink solution [9]:  $\Delta(x) = \operatorname{coth}(b) + \tanh(x) - \tanh(x+b)$ .

All these are *real* condensates, and are well-known. The only previously known complex condensates are (i) the simple "chiral spiral" or LOFF solution  $\Delta(x) = Ae^{iqx}$ ; and (ii) Shei's "twisted kink" solution [10], which we can express in complex form

$$\Delta(x) = e^{i\theta/2} \frac{\cosh\left(x\,\sin\frac{\theta}{2} - i\frac{\theta}{2}\right)}{\cosh\left(x\,\sin\frac{\theta}{2}\right)} \quad . \tag{11}$$

 $\theta$  is the angle through which the phase of the condensate rotates as x goes from  $-\infty$  to  $+\infty$ . The single-particle fermion spectrum has a single bound state within the gap, located at  $E = \cos\left(\frac{\theta}{2}\right)$ , as shown by the dashed line in Figure 1. Consistency with the gap equation requires vanishing of the coefficient of  $\Delta'$ , which places conditions on parameters of the solution. For example, for Shei's solution the condition is that  $\theta/(2\pi)$  is equal to the filling fraction of the bound level [10, 11]. We point out a simple physical interpretation of this condition in terms of conserved currents (see also [31]). Both NJL<sub>2</sub> and GN<sub>2</sub> models have a conserved current  $j^{\mu} = \bar{\psi} \gamma^{\mu} \psi$ . Since  $\langle j^{\mu}(x) \rangle = -iN \operatorname{tr}_{D,E}[\gamma^{0} \gamma^{\mu} R(x)]$ , we see that, for a static condensate, current conservation follows trivially since  $\langle i^1(x) \rangle = 0$ ; this is the physical origin of the identity  $\operatorname{tr}_{\mathrm{D}}(\gamma^5 R) = 0$ . The NJL<sub>2</sub> model also has a conserved axial current,  $j_5^{\mu} = \bar{\psi} \gamma^{\mu} \gamma^5 \psi$ , and the Eilenberger equation




FIG. 1: Band spectrum of the BdG Hamiltonian (4) for the complex crystal condensate (13), as a function of the twist parameter  $\theta$ . The dashed line denotes the infinite period limit [in this case  $\nu \to 0$ ], which is Shei's solution (11). At  $\theta = 2\mathbf{K}$  we recover the symmetric spectrum of the real kink crystal, relevant to the GN<sub>2</sub> system.

(6) implies

$$\partial_{\mu}\langle j_{5}^{\mu}\rangle \equiv -iN\mathrm{tr}(R') = 2S(x)\langle\bar{\psi}\,i\gamma^{5}\psi\rangle - 2P(x)\langle\bar{\psi}\psi\rangle\,(12)$$

Axial current conservation then follows from the gap equation, using precisely the same condition on the coefficient of  $\Delta'$ .

We have found a nontrivial crystalline solution to (10)

$$\Delta(x) = A \frac{\sigma(iAx + \theta/2 - \mathbf{K})}{\sigma(iAx - \mathbf{K})\sigma(\theta/2)}$$
(13)  
 
$$\times \exp\left[iAx\left(-\zeta(\theta/2) + \csc(\theta/2)\right) + \theta\eta/2 + i\operatorname{am}(\theta/2)\right]$$

Here, 
$$\sigma$$
 and  $\zeta$  are Weierstrass sigma and zeta functions,  
 $A(\theta) = 2 \operatorname{sd}(\theta/4) \operatorname{cn}(\theta/4)$ , and  $\eta = \zeta(\mathbf{K})$ , with  $\mathbf{K}(\nu)$   
the elliptic half-period. For this condensate, both the  
amplitude and the phase are x-dependent, as shown in  
Figs. 2, 3. The essential parameters of the solution  
(13) are: (i) the parameter  $\theta$  which [via (15) below]  
characterizes the chiral twist of  $\Delta(x)$  over one period;  
and (ii) the elliptic parameter  $\nu$  which, together with  $\theta$ ,  
determines the crystal period. The spinor solutions of  
the Dirac/BdG equation can also be expressed explicitly  
in terms of elliptic functions, and one can perform the  
Hartree-Fock analysis, as in the GN<sub>2</sub> system, to prove  
self-consistency of the crystalline condensate [30]. The  
spectrum is that of a Dirac particle with a *single* band  
in the gap, as shown in Fig. 1. However, unlike the  
real case where the band lies symmetrically in the cen-  
ter of the gap, here the band is offset. Indeed, the band  
edges are given by:  $E_1 = -1$ ,  $E_2 = -1 + 2 \operatorname{cn}^2(\theta/4; \nu)$ ,  
 $E_3 = -1 + 2 \operatorname{cd}^2(\theta/4; \nu)$ ,  $E_4 = +1$ , as shown in Fig. 1.  
This spectrum is a band version of the Shei spectrum,  
reducing to the Shei solution in the infinite period limit.  
It is also a deformation of the kink crystal spectrum of  
the [discrete-chiral] GN<sub>2</sub> system, reducing to that case  
when  $\theta = 2\mathbf{K}$ . We find the *exact* diagonal resolvent (9)  
with

$$\mathcal{N}(E) = \frac{1}{4} \frac{1}{\sqrt{(E^2 - 1)(E - E_2)(E - E_3)}} \quad , \quad (14)$$

 $b(E) = 2E - (E_2 + E_3)$ , and  $a(E) = 2(E+1)(E-1 - (E_2 + E_3)/2) + 1 + (E_2 + E_3) - (E_2 - E_3)^2/4$ .

Under a shift through one period  $L = 2\mathbf{K}'/A$  of the crystal, the BdG Hamiltonian is invariant up to a global chiral rotation through an angle  $\varphi$ :

$$H(x+L) = e^{i\gamma^5\varphi} H(x) e^{-i\gamma^5\varphi} , \qquad (15)$$
  
$$\varphi = \mathbf{K}' \left[ -\zeta(\theta/2) + \operatorname{cs}(\theta/2) - i\theta \,\zeta(i\mathbf{K}')/(2\mathbf{K}') \right]$$

The solutions to  $H\psi = E\psi$  acquire a chiral rotation and a Bloch phase under a period shift,  $\psi(x + L) = e^{ikL}e^{i\varphi\gamma^5}\psi(x)$ . The [relativistic] Bloch momentum k is related to the the spectral function by  $\rho(E) = dk/dE$ .




FIG. 2: The amplitude M(x) [solid line] and phase  $\chi(x)$  [dashed line] of  $\Delta = Me^{i\chi}$  in (13), over several periods. The amplitude is periodic while the phase rotates by an angle  $2\varphi$  each period.




FIG. 3: Crystalline complex kink (13) plotted as a function of x over one period. The cross-section denotes  $\Delta(x) = S(x) - iP(x)$ , and indicates a net rotation through the twist parameter  $2\varphi$  over one period.

To conclude, we discuss briefly the implications of this self-consistent solution to the gap equation for the  $(T, \mu)$ phase diagram of the NJL<sub>2</sub> model. Recall that the corresponding real condensate characterizes the inhomogeneous crystalline phase of the GN<sub>2</sub> system [13]. For the NJL<sub>2</sub> model, the Ginzburg-Landau (GL) approach shows that the "chiral spiral" phase identified in [12] has a richer structure, characterized by a crystalline complex condensate of the form (13). Near the tricritical point of the massless NJL<sub>2</sub> system the GL effective Lagrangian is

$$\mathcal{L}_{GL} = c_0 + c_2 |\Delta|^2 + c_3 \text{Im} [\Delta(\Delta')^*] + c_4 [|\Delta|^4 + |\Delta'|^2] + c_5 \text{Im} [(\Delta'' - 3|\Delta|^2 \Delta) (\Delta')^*]$$
(16)  
+  $c_6 [2|\Delta|^6 + 8|\Delta|^2 |\Delta'|^2 + 2\text{Re} ((\Delta')^2 (\Delta^*)^2) + |\Delta''|^2]$ 

Here the coefficients  $c_n$  are known functions of T and  $\mu$ [32, 33]. (This GL approach has been used previously in [33] to describe the phase diagram of the massive and massless NJL<sub>2</sub> models, in the vicinity of the tricritical point. For the massive NJL<sub>2</sub> model no (complex) exact solution to the gap equation is known, so [33] is the current state-of-the-art for the massive model.) In the GN<sub>2</sub> model, which has a real condensate, there is a tricritical point at T = 0.3183,  $\mu = 0.6082$ ; given by the point  $c_2 = c_4 = 0$ . In the [chiral] NJL<sub>2</sub> model, which has a complex condensate, there is a tricritical point at T = 0.5669,  $\mu = 0$ ; given by  $c_2 = c_3 = 0$ . To search for possible crystalline phases near the tricritical point we keep terms up to  $c_4$ , and study the effective equation of motion

$$c_4 \Delta'' - i c_3 \Delta' - (c_2 + 2c_4 |\Delta|^2) \Delta = 0$$
 . (17)

Note that this has precisely the same form as the NLSE (10) found from the gap equation and the Eilenberger equation. Thus, we can use our solution (13) as a variational ansatz and compute the free energy. We have found that the resulting free energy is lower than that of the LOFF-form "chiral spiral" variational ansatz  $\Delta(x) = Ae^{iqx}$ , for  $T < T_c$ . Noting that our ansatz reduces to the chiral spiral in the perturbative limit (just as the GN<sub>2</sub> self-consistent crystal condensate reduces to the LOFF form  $\Delta(x) = A\sin(qx)$  in this limit [13]), we are led to the phase diagram shown in Fig. 4. For a detailed discussion of the phase diagram, based on the *full* free energy (beyond the GL approximation), using our exact spectral data, see [30].




FIG. 4: Phase diagram of the chiral NJL<sub>2</sub> model, from a Ginzburg-Landau analysis based on a crystalline condensate of the form in (13). Along the T axis (below  $T_c$ ) the condensate is of the form of Shei's twisted kink (11), while on the  $T = T_c$  line the condensate reduces to the form of the LOFF chiral spiral [12]. The condensate is depicted as in Fig. 3.

We thank the DOE for support through grant DE-FG02-92ER40716.

- D. K. Campbell and A. R. Bishop, "Soliton Excitations In Polyacetylene And Relativistic Field Theory Models," Nucl. Phys. B 200, 297 (1982).
- [2] K. Rajagopal and F. Wilczek, "The condensed matter physics of QCD," in At the Frontier of Particle Physics / Handbook of QCD, M. Shifman, ed., (World Scientific, 2001); arXiv:hep-ph/0011333.
- [3] R. Casalbuoni and G. Nardulli, "Inhomogeneous superconductivity in condensed matter and QCD," Rev. Mod. Phys. 76, 263 (2004). [arXiv:hep-ph/0305069].
- [4] S. Giorgini, L. P. Pitaevskii and S. Stringari, "Theory of ultracold Fermi gases", [arXiv:0706.3360], to appear in Rev. Mod. Phys.
- [5] R. Peierls, The Quantum Theory of Solids (Oxford, 1955).
- [6] P. G. de Gennes, Superconductivity of Metals and Alloys (Addison-Wesley, Redwood City, CA, 1989).
- [7] Y. Nambu and G. Jona-Lasinio, "Dynamical model of elementary particles based on an analogy with superconductivity," Phys. Rev. **122**, 345 (1961).
- [8] D. J. Gross and A. Neveu, "Dynamical Symmetry Breaking In Asymptotically Free Field Theories," Phys. Rev. D 10, 3235 (1974).
- [9] R. F. Dashen, B. Hasslacher and A. Neveu, "Semiclassical Bound States In An Asymptotically Free Theory," Phys. Rev. D 12, 2443 (1975).
- [10] S. S. Shei, "Semiclassical Bound States In A Model With Chiral Symmetry," Phys. Rev. D 14, 535 (1976).
- [11] J. Feinberg and A. Zee, "Dynamical Generation of Extended Objects in a 1 + 1 Dimensional Chiral Field Theory: Non-Perturbative Dirac Operator Resolvent Analysis," Phys. Rev. D 56, 5050 (1997) [arXiv:cond-mat/9603173]; "Generalized supersymmetric quantum mechanics and reflectionless fermion bags in 1+1 dimensions," in *Multiple facets of quantization and supersymmetry*, M. Olshanetsky *et al* (ed.) (World Scientific, 2002) arXiv:hep-th/0109045.
- [12] V. Schon and M. Thies, "Emergence of Skyrme crystal in Gross-Neveu and 't Hooft models at finite density," Phys. Rev. D 62, 096002 (2000). [arXiv:hep-th/0003195].
- [13] M. Thies, "Analytical solution of the Gross-Neveu model at finite density," Phys. Rev. D 69, 067703 (2004), [arXiv:hep-th/0308164]; "From relativistic quantum fields to condensed matter and back again: Updating the Gross-Neveu phase diagram," J. Phys. A 39, 12707 (2006). [arXiv:hep-th/0601049].
- [14] P. de Forcrand and U. Wenger, "New baryon matter in the lattice Gross-Neveu model," PoS LAT2006, 152 (2006). [arXiv:hep-lat/0610117].
- [15] U. Wolff, "The phase diagram of the infinite-N Gross-Neveu Model at finite temperature and chemical potential", Phys. Lett. **157B**, 303 (1985).
- [16] B. Horovitz, "Soliton Lattice in Polyacetylene, Spin-Peierls Systems, and Two-Dimensional Sine-Gordon Systems", Phys. Rev. Lett. 46, 742 (1981).
- [17] S. A. Brazovskii, S. A. Gordynin, and N. N. Kirova, "Exact solution of the Peierls model with an arbitrary number of electrons in the unit cell", Pis. Zh. Eksp. Teor. Fiz. **31**, 486 (1980) [JETP Lett. **31**, 456 (1980)]; S. A. Brazovskii and N. N. Kirova, "Excitons, polarons and bipolarons in conducting polymers", Pis. Zh. Eksp. Teor. Fiz.

**33**, 6 (1981) [JETP Lett. **33**, 4 (1981)].

- [18] J. Mertsching and H. J. Fischbeck, "The incommensurate Peierls phase of the quasi-dimensional Frohlich model with a nearly half-filled band", Phys. Stat. Sol. B 103, 783 (1981).
- [19] A. I. Buzdin and V. V. Tugushev, "Phase diagrams of electronic and superconducting transitions to soliton lattice states", Zh. Eksp. Teor. Fiz. 85, 735 (1983), [Sov. Phys. JETP 58, 428 (1983).
- [20] K. Machida and H. Nakanishi, "Superconductivity under a ferromagnetic molecular field", Phys. Rev. B 30, 122 (1984).
- [21] R. Jackiw and C. Rebbi, "Solitons With Fermion Number 1/2," Phys. Rev. D 13, 3398 (1976).
- [22] J. Goldstone and F. Wilczek, "Fractional Quantum Numbers On Solitons," Phys. Rev. Lett. 47, 986 (1981).
- [23] M. W. Zwierlein, A. Schirotzek, C. H. Schunck and W. Ketterle, "Fermionic Superfluidity with Imbalanced Spin Populations and the Quantum Phase Transition to the Normal State", Science **311**, 492 (2006).
- [24] G. B. Partridge *et al*, "Pairing and Phase Separation in a Polarized Fermi Gas", Science **311**, 503 (2006).
- [25] G. Eilenberger, "Transformation of Gorkov's Equation for Type II Superconductors into Transport Like Equations", Z. Phys. 214, 195 (1968).

- [26] Z. b. Su and B. Sakita, "Chiral symmetry and chiral anomaly in an incommensurate charge-density-wave system," Phys. Rev. Lett. 56, 780 (1986).
- [27] I. Kosztin, S. Kos, M. Stone and A. J. Leggett, "Free energy of an inhomogeneous superconductor: A wavefunction approach", Phys. Rev. B 58, 9365 (1998); S. Kos and M. Stone, "Gradient expansion for the free energy of a clean superconductor", Phys. Rev. B 59, 9545 (1999).
- [28] L. A. Dickey, Soliton Equations and Hamiltonian Systems, (World Scientific, Singapore, 1991).
- [29] S. A. Brazovskii, N. N. Kirova and S. I. Matveenko, "Peierls effect in conducting polymers", Zh. Eksp. Teor. Fiz. 86, 743 (1984) Sov. Phys. JETP 59, 434 (1984).
- [30] G. Basar and G. V. Dunne, in preparation.
- [31] F. Karbstein and M. Thies, "Divergence of the axial current and fermion density in Gross-Neveu models," Phys. Rev. D 76, 085009 (2007). [arXiv:0706.0424 [hep-th]].
- [32] A. I. Buzdin and H. Kachkachi, "Generalized Ginzburg-Landau theory for nonuniform FFLO superconductors", Phys. Lett. A 225, 341 (1997).
- [33] C. Boehmer, M. Thies and K. Urlichs, "Tricritical behavior of the massive chiral Gross-Neveu model," Phys. Rev. D 75, 105017 (2007).