

# Modular Classes of Loday Algebroids

Mathieu Stiénon \*

E.T.H. Zürich

stienon@math.ethz.ch

Ping Xu †

Penn State University

ping@math.psu.edu

## Abstract

We introduce the concept of Loday algebroids, a generalization of Courant algebroids. We define the *naive cohomology* and modular class of a Loday algebroid, and we show that the modular class of the double of a Lie bialgebroid vanishes. For Courant algebroids, we describe the relation between the naive and standard cohomologies and we conjecture that they are isomorphic when the Courant algebroid is transitive.

## 1 Naive Cohomology

Given a Courant algebroid  $(E, \rho, \llbracket \cdot, \cdot \rrbracket, \langle \cdot, \cdot \rangle)$ , let  $\Gamma(\wedge^k \ker \rho)$  denote the space of smooth sections of the (possibly singular) vector bundle  $\wedge^k \ker \rho$  (i.e. smooth sections  $\alpha$  of  $\wedge^k E$  such that  $\alpha|_m \in \wedge^k \ker \rho$  for each  $m \in M$ ). The extension of the pseudo-metric  $\langle \cdot, \cdot \rangle$  to  $\wedge^k E$  naturally induces an isomorphism  $\Xi : \wedge^k E \rightarrow \wedge^k E^*$ . Since, by definition,  $\langle \mathcal{D}f, e \rangle = \frac{1}{2}\rho(e)f$ , the sections of  $\wedge^k \ker \rho$  are characterized as the elements  $\varepsilon \in \Gamma(\wedge^k E)$  such that  $\check{y}_{\mathcal{D}f}\varepsilon = 0$ ,  $\forall f \in C^\infty(M)$ . Here  $\check{y}_{\mathcal{D}f} = \Xi^{-1} \circ i_{\mathcal{D}f} \circ \Xi$ , where  $i_{\mathcal{D}f} : \Gamma(\wedge^{k+1} E^*) \rightarrow \Gamma(\wedge^k E^*)$  is the usual contraction of exterior forms with the section  $\mathcal{D}f \in \Gamma(E)$ . Define an operator  $\check{d} : \Gamma(\wedge^k \ker \rho) \rightarrow \Gamma(\wedge^{k+1} E)$  by

$$\begin{aligned} (\check{d}\alpha)(e_0, \dots, e_k) &= \sum_{i=0}^k (-1)^i \rho(e_i) \alpha(e_0, \dots, \widehat{e}_i, \dots, e_k) \\ &\quad + \sum_{i < j} (-1)^{i+j} \alpha(\llbracket e_i, e_j \rrbracket, e_0, \dots, \widehat{e}_i, \dots, \widehat{e}_j, \dots, e_k), \end{aligned}$$

for all  $\alpha \in \Gamma(\wedge^k \ker \rho)$  and  $e_0, \dots, e_k \in \Gamma(E)$ . Here the pairing between  $\Gamma(\wedge^k \ker \rho)$  and  $\Gamma(\wedge^k E)$  is via the identification  $\Xi : \wedge^k E \rightarrow \wedge^k E^*$ . The following Lemma follows from the Courant algebroid properties, in particular the relations  $\rho(\mathcal{D}f) = 0$  and  $\llbracket \mathcal{D}f, e \rrbracket + \mathcal{D}\langle \mathcal{D}f, e \rangle = 0$ .

\*Research supported by the European Union through the FP6 Marie Curie R.T.N. ENIGMA (Contract number MRTN-CT-2004-5652).

†Research partially supported by NSF grant DMS-0605725 & NSA grant H98230-06-1-0047.

**Lemma 1.1** *We have  $\check{d}\Gamma(\wedge^k \ker \rho) \subset \Gamma(\wedge^{k+1} \ker \rho)$ . Moreover,  $(\Gamma(\wedge^\bullet \ker \rho), \check{d})$  is a cochain complex.*

The cohomology of this cochain complex is called the naive cohomology of  $E$  and is denoted  $H_{\text{naive}}^\bullet(E)$ .

**Remark 1.2** *It is easy to see that a 1-cochain  $\theta \in \Gamma(\ker \rho)$  is a 1-cocycle if, and only if,  $\langle \theta, \llbracket a, b \rrbracket \rangle = \rho(a)\langle \theta, b \rangle - \rho(b)\langle \theta, a \rangle$  for all  $a, b \in \Gamma(E)$ , and a 1-coboundary if, and only if,  $\theta = \mathcal{D}f$  for some  $f \in C^\infty(M)$ .*

**Remark 1.3** *Let  $V$  be the  $C^\infty(M)$ -module generated by  $\mathcal{D}(C^\infty(M))$ . Since  $\langle \mathcal{D}f, a \rangle = \frac{1}{2}\rho(a)f$ , we have  $V = \Gamma(\ker \rho^\perp)$  and  $\Xi(V) = \Gamma(\ker \rho^0) = \rho^*(\Gamma(T^*M))$ . Moreover  $V \subset \Gamma(\ker \rho)$ , for  $\rho \circ \mathcal{D} = 0$ . Therefore, when  $(E, \rho, \llbracket \cdot, \cdot \rrbracket, \langle \cdot, \cdot \rangle)$  is a regular Courant algebroid (i.e.  $\rho$  has constant rank),  $E/\Xi^{-1}(\rho^*T^*M)$  is a Lie algebroid and  $H_{\text{naive}}^\bullet(E)$  is the cohomology of this Lie algebroid. However, in general,  $\Gamma(E)/V$  is only a Lie-Rinehart algebra over  $C^\infty(M)$ . One can consider  $H_{\text{naive}}^\bullet(E)$  as its cohomology [4].*

**Example 1.4** *When  $E = TM \oplus T^*M$  is an exact Courant algebroid,  $H_{\text{naive}}^\bullet(E)$  is isomorphic to the de Rham cohomology of  $M$ .*

**Example 1.5** *If  $E$  is a Courant algebroid over a point, i.e. a Lie algebra equipped with a non-degenerate ad-invariant bilinear form,  $H_{\text{naive}}^\bullet(E)$  is simply the Lie algebra cohomology.*

## 2 Relation with standard cohomology

Courant algebroids can also be obtained as derived brackets [6][9] using degree two super-symplectic manifolds. More precisely, given a Courant algebroid  $(E, \rho, \llbracket \cdot, \cdot \rrbracket, \langle \cdot, \cdot \rangle)$ ,  $E[1]$  is a super-Poisson manifold, where the Poisson structure is induced by the pseudo-metric. There is a minimal symplectic realization  $X \xrightarrow{\pi} E[1]$  and a cubic function  $\Theta$  on  $X$  such that  $\{\Theta, \Theta\} = 0$  and, for all  $f \in C^\infty(M)$  and  $e_1, e_2 \in \Gamma(E)$ ,

$$\mathcal{D}f = \{\Theta, f\} \quad \text{and} \quad e_1 \circ e_2 = \{\{\Theta, e_1\}, e_2\},$$

where the symbol  $\circ$  denotes the *asymmetric Dorfman bracket* defined by the relation  $a \circ b = \llbracket a, b \rrbracket + \mathcal{D}\langle a, b \rangle$ . Here elements in  $\Gamma(\wedge^k E)$  are viewed as functions of degree  $k$  on  $X$  by considering them as functions on  $E[1]$  via the pseudo-metric  $\langle \cdot, \cdot \rangle$  and identifying them with their pull back by  $\pi$ . Similarly functions on  $M$  are also identified with their pull back in  $X$ . By  $\mathcal{A}^k$  we denote the space of functions on  $X$  of degree  $k$ . Then  $(\mathcal{A}^\bullet, \{\Theta, \cdot\})$  is a cochain complex. Its cohomology is called the *standard cohomology* by Roytenberg [9] and we shall denote it by  $H_{\text{std}}^\bullet(E)$ .

**Lemma 2.1** *1. If  $c \in \Gamma(\wedge^k \ker \rho)$ , then  $\{\Theta, c\} = \check{d}c$ ;*

*2. If  $c \in \Gamma(\wedge^k E)$  satisfies  $\{\Theta, c\} = 0$ , then  $c \in \Gamma(\wedge^k \ker \rho)$  and  $\check{d}c = 0$ .*

PROOF. (i) It suffices to prove the case when  $k = 1$ . The general situation follows from the Leibniz rule. Now since  $\rho(c) = 0$ , we have  $\forall e_1, e_2 \in \Gamma(E)$ ,

$$\begin{aligned}
& \langle c \circ e_2, e_1 \rangle - (\check{d}c)(e_1, e_2) \\
&= (-\langle e_2 \circ c, e_1 \rangle + 2\langle \mathcal{D}\langle c, e_2 \rangle, e_1 \rangle) - (\rho(e_1)\langle c, e_2 \rangle - \rho(e_2)\langle c, e_1 \rangle - \langle c, \llbracket e_1, e_2 \rrbracket \rangle) \\
&= \rho(e_2)\langle c, e_1 \rangle - \langle e_2 \circ c, e_1 \rangle - \langle c, \llbracket e_2, e_1 \rrbracket \rangle \\
&= \rho(e_2)\langle c, e_1 \rangle - \langle e_2 \circ c, e_1 \rangle - \langle c, e_2 \circ e_1 \rangle \\
&= 0.
\end{aligned}$$

It thus follows that  $\{\{\{\Theta, c\}, e_2\}, e_1\} - \{\{\check{d}c, e_2\}, e_1\} = 0$ , which implies that  $\{\Theta, c\} = \check{d}c$ .

(ii) Since  $\check{y}_{\mathcal{D}f}c = \{\mathcal{D}f, c\} = \{\{\Theta, f\}, c\} = \{f, \{\Theta, c\}\} = 0$  for all  $f \in C^\infty(M)$ , we have  $c \in \Gamma(\wedge^k \ker \rho)$ .  $\square$

As a consequence, we have a homomorphism  $\phi : H_{\text{naive}}^\bullet(E) \rightarrow H_{\text{std}}^\bullet(E)$ . Lemma 2.1 also implies that  $\phi$  is an isomorphism in degrees 0 and 1. It is natural to ask when  $\phi$  is an isomorphism in all degrees. When  $E$  is a Courant algebroid over a point,  $\phi$  is clearly an isomorphism. On the other hand, when  $E$  is the standard Courant algebroid  $TM \oplus T^*M$ , both  $H_{\text{naive}}^\bullet(E)$  and  $H_{\text{std}}^\bullet(E)$  are isomorphic to the de Rham cohomology of  $M$ . Hence  $\phi$  is also an isomorphism. This leads to the following

**Conjecture** When  $E$  is a transitive Courant algebroid,  $\phi$  is an isomorphism.

### 3 Lie derivatives and Loday algebroids

The Lie derivative of Courant algebroids was introduced in [10]. Let us recall its definition briefly. An infinitesimal automorphism of the vector bundle  $E \xrightarrow{\pi} M$  is a vector field on  $E$  — a derivation of the algebra  $C^\infty(E)$  — which preserves the subspaces  $\pi^*C^\infty(M)$  and  $\Gamma(E)$  (whose elements are identified with functions linear on the fibers of  $\pi$  through the pairing  $\langle \cdot, \cdot \rangle$ ). In other words, it is a covariant differential operator on  $E$ , i.e. a pair of differential operators  $\delta^0 : C^\infty(M) \rightarrow C^\infty(M)$  and  $\delta^1 : \Gamma(E) \rightarrow \Gamma(E)$  satisfying

$$\delta^0(fg) = f\delta^0(g) + \delta^0(f)g \quad \text{and} \quad \delta^1(fe) = f\delta^1(e) + \delta^0(f)e,$$

for any  $f, g \in C^\infty(M)$  and  $e \in \Gamma(E)$ . It is known [9] that the Lie algebra  $\mathfrak{aut}(E)$  of infinitesimal automorphisms of the Courant algebroid  $E$  consists of those covariant differential operators  $\delta = (\delta^0, \delta^1)$  on  $E$  which satisfy the additional properties:

$$\delta^0\langle e_1, e_2 \rangle = \langle \delta^1 e_1, e_2 \rangle + \langle e_1, \delta^1 e_2 \rangle \quad \text{and} \quad \delta^1 \llbracket e_1, e_2 \rrbracket = \llbracket \delta^1 e_1, e_2 \rrbracket + \llbracket e_1, \delta^1 e_2 \rrbracket,$$

for all  $e_1, e_2 \in \Gamma(E)$ .

For any  $e \in \Gamma(E)$ , the pair  $\delta_e = (\delta_e^0, \delta_e^1)$  defined by the relations  $\delta_e^0(f) = \rho(e)f$  and  $\delta_e^1(x) = e \circ x$  is an infinitesimal automorphism of the Courant algebroid  $E$ , i.e.  $\delta_e \in \mathbf{aut}(E)$ . Let us denote the (local) flow generated by the vector field on  $E$  corresponding to  $\delta_e$  by  $\phi_t$ . By abuse of notations, we use the same symbol  $\phi_t$  (resp.  $\phi_t^*$ ) to denote its induced flow on the tensor bundles  $E_j^i = (\otimes^i E) \otimes (\otimes^j E^*)$  ( $i, j \in \{0, 1, 2, \dots\}$ ) (resp. the induced action on the spaces of sections of the  $E_j^i$ 's). For any section  $\sigma \in \Gamma(E_j^i)$ , define the Lie derivative  $\mathcal{L}_z \sigma \in \Gamma(E_j^i)$  by  $\mathcal{L}_z \sigma = \frac{d}{d\tau} \phi_\tau^* \sigma \Big|_{\tau=0}$ . Thus we have the usual identity:  $\frac{d}{d\tau} \phi_\tau^* \sigma \Big|_{\tau=t} = \phi_t^* (\mathcal{L}_z \sigma)$ . In the following proposition, we give a list of important properties of this Lie derivative.

**Proposition 3.1** *For all  $f, g \in C^\infty(M)$  and  $x, y, z \in \Gamma(E)$ , we have:*

$$\mathcal{L}_z f = \rho(z)f \quad \mathcal{L}_z x = z \circ x \quad \mathcal{L}_{\mathcal{D}f} x = 0 \quad \mathcal{L}_x \mathcal{D}f = \mathcal{D}\mathcal{L}_x f, \quad (1)$$

$$[\delta, \mathcal{L}_z] = \mathcal{L}_{\delta^1 z} \quad \forall \delta \in \mathbf{aut}(E), \quad (2)$$

$$\mathcal{L}_z(\sigma \otimes \tau) = \mathcal{L}_z \sigma \otimes \tau + \sigma \otimes \mathcal{L}_z \tau \quad \forall \sigma, \tau \in \oplus_{i,j} E_j^i, \quad (3)$$

$$\mathcal{L}_{[[x,y]]} = [\mathcal{L}_x, \mathcal{L}_y], \quad (4)$$

$$\mathcal{L}_z [[x, y]] = [[\mathcal{L}_z x, y]] + [[x, \mathcal{L}_z y]], \quad (5)$$

$$\mathcal{L}_{f \cdot x} y = f \mathcal{L}_x y - (\rho(y)f) x + 2\langle x, y \rangle \mathcal{D}f, \quad (6)$$

$$\mathcal{L}_z \langle x, y \rangle = \langle \mathcal{L}_z x, y \rangle + \langle x, \mathcal{L}_z y \rangle. \quad (7)$$

**Definition 3.2** *A Loday algebroid consists of a vector bundle  $\pi : E \rightarrow M$ , a pseudo-metric  $\langle \cdot, \cdot \rangle$  on the fibers of  $\pi$ , a bundle map  $\rho : E \rightarrow TM$  and an  $\mathbb{R}$ -bilinear operation  $\circ$  on  $\Gamma(E)$  satisfying*

$$e_1 \circ (e_2 \circ e_3) = (e_1 \circ e_2) \circ e_3 + e_2 \circ (e_1 \circ e_3), \quad (8)$$

$$\rho(e_1 \circ e_2) = [\rho(e_1), \rho(e_2)], \quad (9)$$

$$e_1 \circ (f e_2) = (\rho(e_1)f) e_2 + f(e_1 \circ e_2), \quad (10)$$

$$e_1 \circ e_2 + e_2 \circ e_1 = 2\mathcal{D}\langle e_1, e_2 \rangle, \quad (11)$$

$$\mathcal{D}f \circ e = 0, \quad (12)$$

where  $\mathcal{D} : C^\infty(M) \rightarrow \Gamma(E)$  is the  $\mathbb{R}$ -linear map defined by  $\langle \mathcal{D}f, e \rangle = \frac{1}{2} \rho(e)f$ .

**Remark 3.3** *1. According to [11], for Courant algebroids, axioms (9) and (10) are redundant. It would be interesting to investigate if it is also the case for Loday algebroids.*

*2. The Leibniz algebroids studied by several authors [3, 2, 12] are a more general notion.*

*3. A Courant algebroid is a Loday algebroid satisfying the additional axiom  $\rho(e)\langle e_1, e_2 \rangle = \langle e \circ e_1, e_2 \rangle + \langle e_1, e \circ e_2 \rangle$ .*

**Lemma 3.4** *If  $(E, \rho, \circ, \langle \cdot, \cdot \rangle)$  is a Loday algebroid, then  $\rho(\mathcal{D}f) = 0$  and  $[[\mathcal{D}f, e]] + \mathcal{D}\langle \mathcal{D}f, e \rangle = 0$ , for all  $f \in C^\infty(M)$  and  $e \in \Gamma(E)$ . Here  $[[x, y]] = \frac{1}{2}(x \circ y - y \circ x)$  as in a Courant algebroid.*

PROOF. Applying  $\rho$  to both sides of (11) and making use of (9), we get  $\rho(\mathcal{D}\langle e_1, e_2 \rangle) = 0$  for any  $e_1, e_2 \in \Gamma(E)$  and thus also  $\rho(\mathcal{D}\langle f e_1, e_2 \rangle) = 0$  for any  $f \in C^\infty(M)$ . The Leibniz rule  $\mathcal{D}(fg) = g\mathcal{D}f + f\mathcal{D}g$  implies that  $\rho(\mathcal{D}f) = 0$ . The other relation follows immediately from (12) and (11).  $\square$

As a consequence, the definition of the naive cohomology extends from Courant algebroids to Loday algebroids.

Let  $(E, \rho, \circ, \langle \cdot, \cdot \rangle)$  be a Loday algebroid. Given a section  $z \in \Gamma(E)$ , set  $\mathcal{L}_z f = \rho(z)f$  for  $f \in C^\infty(M)$  and  $\mathcal{L}_z x = z \circ x$  for  $x \in \Gamma(E)$  and extend  $\mathcal{L}_z$  to  $\Gamma(\wedge^k E)$  by the Leibniz rule.

**Proposition 3.5** *Identities (4), (5) and (6) still hold for any Loday algebroid.*

**Remark 3.6** *It is unknown if the standard cohomology can be defined for Loday algebroids. Indeed, it would be interesting to see if there exists a derived bracket in the sense of Kosmann-Schwarzbach [6] for a Loday algebroid.*

## 4 Modular classes

A Loday algebroid module is a vector bundle  $S \rightarrow M$  endowed with an  $\mathbb{R}$ -linear map  $\Gamma(E) \otimes \Gamma(S) \rightarrow \Gamma(S) : e \otimes s \mapsto \nabla_e s$  satisfying

$$\nabla_{\mathcal{D}f} s = 0 \quad \nabla_e (fs) = f\nabla_e s + (\rho(e)f)s \quad (13)$$

$$\nabla_{f e} s = f\nabla_e s \quad \nabla_{e_1}(\nabla_{e_2} s) - \nabla_{e_2}(\nabla_{e_1} s) = \nabla_{[[e_1, e_2]]} s \quad (14)$$

for any  $f \in C^\infty(M)$ ,  $e, e_1, e_2 \in \Gamma(E)$  and  $s \in \Gamma(S)$ .

Now let  $S$  be a real line bundle which is a module of the Loday algebroid  $E$ . Assume that there exists a nowhere zero section  $s \in \Gamma(S)$ . The relation  $D_e s = \langle \theta_s, e \rangle s$  defines a section  $\theta_s \in \Gamma(E)$ . From  $\nabla_{\mathcal{D}f} s = 0$ , it follows that  $\rho(\theta_s) = 0$ . And from (14), it follows that  $\theta_s$  is a naive 1-cocycle. Finally, (13) implies that, for any nowhere vanishing function  $f \in C^\infty(M)$ ,  $\theta_{fs} = f\theta_s + 2\mathcal{D}(\ln|f|)$ . Thus the class  $[\theta_s] \in H_{\text{naive}}^1(E)$  is independent of the chosen section  $s$  and only depends on the module  $S$ . We will denote this class by  $\theta_S$ . As in [1, 7], when the line bundle  $S$  is not trivial, we set  $\theta_S = \frac{1}{2}\theta_{S \otimes S}$ , where  $S \otimes S$  is necessarily a trivial real line bundle. We call  $\theta_S$  the modular class of the module  $S$ .

**Theorem 4.1** *Given a Loday algebroid  $(E, \rho, \circ, \langle \cdot, \cdot \rangle)$ ,  $\wedge^{\text{top}} E$  is an  $E$ -module with  $\nabla = \mathcal{L}$ .*

PROOF. It remains to prove that  $\mathcal{L}_{f e} s = f\mathcal{L}_e s$  for any  $f \in C^\infty(M)$  and  $s \in \Gamma(\wedge^{\text{top}} E)$ . According to Proposition 3.1, for any  $f \in C^\infty(M)$  and  $e, a \in \Gamma(E)$ , we have

$$\begin{aligned} \mathcal{L}_{f e} a &= f\mathcal{L}_e a - (\rho(a)f)e + 2\langle e, a \rangle \mathcal{D}f = f\mathcal{L}_e a - 2\langle \mathcal{D}f, a \rangle e + 2\langle e, a \rangle \mathcal{D}f \\ &= (f\mathcal{L}_e - 2(e\wedge)\circ\check{i}_{\mathcal{D}f} + 2(\mathcal{D}f\wedge)\circ\check{i}_e)(a) \end{aligned}$$

Note that, as differential operators on  $\Gamma(\wedge^\bullet E)$ ,  $\mathcal{L}_{fe}$ ,  $f\mathcal{L}_e$ ,  $2(e\wedge)\circ\check{\mathcal{D}}_f$  and  $2(Df\wedge)\circ\check{\mathcal{L}}_e$  all are derivations of degree 0 with respect to the wedge product on  $\Gamma(\wedge^\bullet E)$ . Since  $\mathcal{L}_{fe}$  and  $f\mathcal{L}_e - 2(e\wedge)\circ\check{\mathcal{D}}_f + 2(Df\wedge)\circ\check{\mathcal{L}}_e$  are equal when acting both on sections of  $E$  and on functions on  $M$ , they are also equal when extended to  $\Gamma(\wedge^* E)$ . In particular, if  $s \in \Gamma(\wedge^{\text{top}} E)$ ,

$$\mathcal{L}_{fe}s = f\mathcal{L}_e s - 2(e\wedge)\circ\check{\mathcal{D}}_f s + 2(Df\wedge)\circ\check{\mathcal{L}}_e s = f\mathcal{L}_e s - 2\langle e, \mathcal{D}f \rangle s + 2\langle \mathcal{D}f, e \rangle s = f\mathcal{L}_e s.$$

□

The modular class  $[\theta_{\wedge^{\text{top}} E}] \in H_{\text{naive}}^1(E)$  of the  $E$ -module  $\wedge^{\text{top}} E$  is called the *modular class of the Loday algebroid*  $E$ .

## 5 Examples

Let  $E = A \oplus A^*$  be the double of a Lie bialgebroid  $(A, A^*)$  [5]. In this case,  $\mathcal{D} = \frac{1}{2}(d + d_*)$  and, for all  $X, Y \in \Gamma(A)$  and  $\xi, \eta \in \Gamma(A^*)$ , the bracket on  $\Gamma(E)$  is defined by

$$\begin{aligned} \llbracket X, \xi \rrbracket &= (-L_\xi X + \frac{1}{2}d_*(\xi, X)) + (L_X \xi - \frac{1}{2}d(\xi, X)), \\ \llbracket X, Y \rrbracket &= [X, Y], \quad \llbracket \xi, \eta \rrbracket = [\xi, \eta]. \end{aligned}$$

Now  $\wedge^{\text{top}} E \cong (\wedge^{\text{top}} A) \otimes (\wedge^{\text{top}} A^*)$  is a trival line bundle. For the sake of simplicity, we assume that there exists a nowhere vanishing section  $V \in \Gamma(\wedge^{\text{top}} A)$ . Let  $\Omega \in \Gamma(\wedge^{\text{top}} A^*)$  be its dual section. For any  $X \in \Gamma(A)$  and  $\xi \in \Gamma(A^*)$ , one has  $\mathcal{L}_X \xi = \llbracket X, \xi \rrbracket + \mathcal{D}\langle X, \xi \rangle = -L_\xi X + d_*(\xi, X) + L_X \xi = -i_\xi d_* X + L_X \xi$ . It follows from the Leibniz rule (see Proposition 3.1) that, for any  $\sigma \in \Gamma(\wedge^k A^*)$ ,  $\mathcal{L}_X \sigma = L_X \sigma + \lambda$  where  $\lambda \in \Gamma(A \otimes (\wedge^{k-1} A^*))$ . On the other hand, since  $A$  is isotropic with respect to  $\langle \cdot, \cdot \rangle$ , we have that  $\mathcal{L}_X \tau = L_X \tau$  if  $\tau \in \Gamma(\wedge^k A)$ . It thus follows that

$$\mathcal{L}_X(V \wedge \Omega) = (\mathcal{L}_X V) \wedge \Omega + V \wedge (\mathcal{L}_X \Omega) = (L_X V) \wedge \Omega + V \wedge (L_X \Omega) = 0.$$

Similarly we have  $\mathcal{L}_\xi(V \wedge \Omega) = 0$ , for all  $\xi \in \Gamma(A^*)$ . Thus we have proved

**Theorem 5.1** *If a Courant algebroid is the double of a Lie bialgebroid, then its modular class vanishes.*

**Example 5.2** *If  $E = TM \oplus T^*M$  is an exact Courant algebroid, the Courant bracket is given by*

$$\llbracket X + \xi, Y + \eta \rrbracket = [X, Y] + i_{X \wedge Y} \phi + L_X \eta - L_Y \xi + \frac{1}{2}d((\xi, Y) - (\eta, X)),$$

for all  $X, Y \in \mathfrak{X}(M)$  and  $\xi, \eta \in \Omega^1(M)$ . Here  $\phi$  is a closed 3-form.

Take a nowhere zero  $V \in \Gamma(\wedge^{\text{top}} TM)$  and its dual  $\Omega \in \Omega^{\text{top}}(M)$ . One easily sees that  $\mathcal{L}_X V = L_X V + V'$ , where  $V' \in \Gamma(T^*M \otimes \wedge^{\text{top}-1} TM)$  and  $\mathcal{L}_X \Omega = L_X \Omega$ . Thus it follows that  $\mathcal{L}_X(V \wedge \Omega) = L_X(V \wedge \Omega) = 0$ ,  $\forall X \in \Gamma(TM)$ . One also sees that  $\mathcal{L}_\xi(V \wedge \Omega) = 0$ ,  $\forall \xi \in \Gamma(T^*M)$ . Therefore the modular class vanishes.

**Acknowledgments** We thank Zhang-Ju Liu, Alan Weinstein and the referee for useful comments.

## References

- [1] Evens, S., Lu, J.-H., and Weinstein, A., Transverse measures, the modular class, and a cohomology pairing for Lie algebroids, *Quart. J. Math. Oxford (2)* **50** (1999), 417-436.
- [2] Grabowski, J. and Urbański, P., Algebroids—general differential calculi on vector bundles. *J. Geom. Phys.* **31** (1999), no. 2-3, 111–141.
- [3] Ibanez, R., de Leon, M., Marrero, J. C., and Padrón, E., Leibniz algebroid associated with a Nambu-Poisson structure, *J. Phys. A* **32** (1999), 8129–8144.
- [4] Huebschmann, J., Duality for Lie-Rinehart algebras and the modular class, *J. Reine Angew. Math.* **510** (1999), 103–159.
- [5] Liu, Z.-J., Weinstein, A. and Xu, P., Manin triples for Lie bialgebroids, *J. Differential Geom.* **45** (1997), 547–574.
- [6] Kosmann-Schwarzbach, Y., From Poisson algebras to Gerstenhaber algebras, *Ann. Inst. Fourier* **46** (1996), 1243-1274.
- [7] Kosmann-Schwarzbach, Y. and Weinstein, A., Relative modular classes of Lie algebroids *C. R. Math. Acad. Sci. Paris* **341** (2005) 8, 509-514.
- [8] Loday, J.-L., Une version non commutative des algèbres de Lie: les algèbres de Leibniz, *Enseign. Math.* **39** (1993), 269–293.
- [9] Roytenberg, D., On the structure of graded symplectic supermanifolds and Courant algebroids, Quantization, Poisson brackets and beyond (Manchester, 2001), 169–185, *Contemp. Math.* **315** Amer. Math. Soc., Providence, RI, 2002.
- [10] Stiénon, M., Moser lemma in generalized complex geometry, arXiv:math/0702718.
- [11] Uchino, K., Remarks on the Definition of a Courant Algebroid, *Letters in Math. Phys.* **60** (2002), 171-175.
- [12] Wade, A., On some properties of Leibniz algebroids, Infinite dimensional Lie groups in geometry and representation theory (Washington, DC, 2000), 65–78, World Sci. Publ., River Edge, NJ, 2002.