Perfect transference of a *d*-level quantum state over pseudo-distance-regular networks

M. A. Jafarizadeh^{a,b,c} * R. Sufiani^{a,b} † S. F. Taghavi^a and E. Barati^a

^aDepartment of Theoretical Physics and Astrophysics, University of Tabriz, Tabriz 51664, Iran.

^bInstitute for Studies in Theoretical Physics and Mathematics, Tehran 19395-1795, Iran.

^cResearch Institute for Fundamental Sciences, Tabriz 51664, Iran.

September 9, 2021

 $^{^{*}}E\text{-mail:jafarizadeh@tabrizu.ac.ir}$

 $^{^{\}dagger}\text{E-mail:sofiani@tabrizu.ac.ir}$

Abstract

Following the prescription of Ref. [1] in which perfect state transference (PST) of a qubit over distance regular spin networks was discussed, in this paper PST of an arbitrary *d*-level quantum state (qudit) over antipodes of more general networks called pseudo distance-regular networks, is investigated. In fact, the spectral analysis techniques used in the previous work [1], and algebraic structures of pseudo distance-regular graphs are employed to give an explicit formula for suitable coupling constants in the Hamiltonians so that the state of a particular qudit initially encoded on one site will evolve freely to the opposite site without any dynamical control, i.e., we show that how to derive the parameters of the system so that PST can be achieved.

Keywords:Perfect state transfer, *d*-level quantum state, Stratification, Pseudodistance-regular network

PACs Index: 01.55.+b, 02.10.Yn

1 Introduction

The transference of quantum information, encoded in a quantum state, from one part of a physical unit, e.g., a qubit, to another part is a crucial ingredient for many quantum information processing protocols [2]. There are various physical systems that can serve as quantum channels, one of them being a quantum spin system. Quantum communication over short distances through a spin chain, in which adjacent qubits are coupled by equal strength has been studied in detail, and an expression for the fidelity of quantum state transfer has been obtained [3, 4]. Similarly, in Ref. [5], near perfect state transfer was achieved for uniform couplings provided a spatially varying magnetic field was introduced. After the work of Bose [3], in which the potentialities of the so-called spin chains have been shown, several strategies were proposed to increase the transmission fidelity [6] and even to achieve, under appropriate conditions, perfect state transfer [7, 8, 9, 10, 11, 12]. All of these proposals refer to ideal spin chains in which only nearest-neighbor couplings are present. In Refs. [7, 8], the d-dimensional hypercube with 2^d vertices has been projected to a linear chain with d+1 sites so that, by considering fixed but different couplings between the qubits assigned to the sites, the PST can be achieved over arbitrarily long distances in the chain. In Ref. [1], the so called distance-regular graphs have been considered as spin networks (in the sense that with each vertex of a distance-regular graph a qubit or a spin was associated) and perfect state transference (PST) over them has been investigated, where a procedure for finding suitable coupling constants in some particular spin Hamiltonians has been given so that perfect transference of a quantum state between antipodes of the networks can be achieved. The aim of this paper is to extend this proposal to systems of particles with arbitrary number of levels, the so-called qudits. These systems can be appeared in condensed matter and solid state physics such as the fermionic SU(N) Hubbard model [13, 14, 15]. In Ref. [16], state transference over spin chains of arbitrary spin has been discussed so that an arbitrary unknown qudit be transferred through a chain with rather good fidelity by the natural dynamics of the chain. In this work, we focus on the situation in which state transference is perfect, i.e., the fidelity is unity. Furthermore, we consider more general graphs called pseudo distance regular graphs or QD type graphs [17, 18] (distance regular graphs are special kinds of pseudo distance regular ones) as underlying networks and investigate PST over antipodes of these networks. In fact this work is an extension of the previous work [1] to PST of a qudit over a pseudo distance regular network. We use the techniques such as stratification of the graphs [17],[19]-[23] and spectral distribution associated with the graphs. Then, we consider particular hamiltonians with nonlinear terms and give a method for finding a suitable set of coupling constants so that PST over antipodes of the networks be possible. In fact, we give an explicit formula for suitable coupling constants so that PST between the first node of the networks and the opposite one can be achieved. As examples we will consider some important pseudo distance regular networks such as the G_n networks, the modified G_n networks, Hadamard network, etc.

The organization of the paper is as follows: In section 2, we review some preliminary facts about graphs and their stratification, pseudo distance-regular graphs and spectral distribution associated with graphs. Section 3 is devoted to PST of a qudit over antipodes of pseudo distance-regular networks, where a method for finding suitable coupling constants in particular spin Hamiltonians so that PST be possible, is given. The paper is ended with a brief conclusion and one appendix.

2 Preliminaries

In this section we recall some preliminaries related to graphs, their stratifications and the notion of pseudo-distance-regularity (as a generalization of distance regularity) of graphs.

2.1 Graphs and their stratifications

In this section, we review the stratification of the graphs and the notion of pseudo-distanceregularity.

A graph is a pair $\Gamma = (V, E)$, where V is a non-empty set called the vertex set and E is a subset of $\{(x, y) : x, y \in V, x \neq y\}$ called the edge set of the graph. Two vertices $x, y \in V$ are called adjacent if $(x, y) \in E$, and in that case we write $x \sim y$. For a graph $\Gamma = (V, E)$, the adjacency matrix A is defined as

$$(A)_{\alpha,\beta} = \begin{cases} 1 & \text{if } \alpha \sim \beta \\ 0 & \text{otherwise} \end{cases}.$$
 (2-1)

Conversely, for a non-empty set V, a graph structure is uniquely determined by such a matrix indexed by V.

The degree or valency of a vertex $x \in V$ is defined by

$$\kappa(x) = |\{y \in V : y \sim x\}| \tag{2-2}$$

where, $|\cdot|$ denotes the cardinality. The graph is called regular if the degree of all of the vertices be the same. In this paper, we will assume that graphs under discussion are regular. A finite sequence $x_0, x_1, ..., x_n \in V$ is called a walk of length n (or of n steps) if $x_{i-1} \sim x_i$ for all i = 1, 2, ..., n. Let $l^2(V)$ denote the Hilbert space of C-valued square-summable functions on V. With each $\beta \in V$ we associate a vector $|\beta\rangle$ such that the β -th entry of it is 1 and all of the other entries of it are zero. Then $\{|\beta\rangle : \beta \in V\}$ becomes a complete orthonormal basis of $l^2(V)$. The adjacency matrix is considered as an operator acting in $l^2(V)$ in such a way that

$$A|\beta\rangle = \sum_{\alpha \sim \beta} |\alpha\rangle.$$
(2-3)

Now, we recall the notion of stratification for a given graph Γ . To this end, let $\partial(x, y)$ be the length of the shortest walk connecting x and y for $x \neq y$. By definition $\partial(x, x) = 0$ for all $x \in V$. The graph becomes a metric space with the distance function ∂ . Note that

 $\partial(x, y) = 1$ if and only if $x \sim y$. We fix a vertex $o \in V$ as an origin of the graph, called the reference vertex. Then, the graph Γ is stratified into a disjoint union of strata (with respect to the reference vertex o) as

$$V = \bigcup_{i=0}^{\infty} \Gamma_i(o), \quad \Gamma_i(o) := \{ \alpha \in V : \partial(\alpha, o) = i \}$$
(2-4)

Note that $\Gamma_i(o) = \emptyset$ may occur for some $i \ge 1$. In that case we have $\Gamma_i(o) = \Gamma_{i+1}(o) = \dots = \emptyset$. With each stratum $\Gamma_i(o)$ we associate a unit vector in $l^2(V)$ defined by

$$|\phi_i\rangle = \frac{1}{\sqrt{\kappa_i}} \sum_{\alpha \in \Gamma_i(o)} |\alpha\rangle, \tag{2-5}$$

where, $\kappa_i = |\Gamma_i(o)|$ is called the *i*-th valency of the graph $(\kappa_i := |\{\gamma : \partial(o, \gamma) = i\}| = |\Gamma_i(o)|)$.

2.2 Pseudo-distance regular graphs

Given a vertex $\alpha \in V$ of a graph Γ , consider the stratification (2-4) with respect to α such that $\Gamma_i(\alpha) = \emptyset$ for i > d. Then, we say that Γ is pseudo-distance-regular around vertex α whenever for any $\beta \in \Gamma_k(\alpha)$ and $0 \le k \le d$, the numbers

$$c_k(\beta) := \frac{1}{\kappa(\beta)} \sum_{\gamma \in \Gamma_1(\beta) \cap \Gamma_{k-1}(\alpha)} \kappa(\gamma), \ a_k(\beta) := \frac{1}{\kappa(\beta)} \sum_{\gamma \in \Gamma_1(\beta) \cap \Gamma_k(\alpha)} \kappa(\gamma), \ b_k(\beta) := \frac{1}{\kappa(\beta)} \sum_{\gamma \in \Gamma_1(\beta) \cap \Gamma_{k+1}(\alpha)} \kappa(\gamma)$$
(2-6)

do not depend on the considered vertex $\beta \in \Gamma_k(\alpha)$, but only on the value of k. In such a case, we denote them by c_k , a_k and b_k respectively.

It should be noticed that for regular graphs Γ ($\kappa(\beta) = \kappa \equiv \kappa_1$ for all $\beta \in V$), the numbers c_k, a_k and b_k read as

$$c_k = |\Gamma_1(\beta) \cap \Gamma_{k-1}(\alpha)|, \quad a_k = |\Gamma_1(\beta) \cap \Gamma_k(\alpha)|, \quad b_k = |\Gamma_1(\beta) \cap \Gamma_{k+1}(\alpha)|, \quad (2-7)$$

where we tacitly understand that $\Gamma_{-1}(\alpha) = \emptyset$. The intersection numbers (2-7) and the valencies $\kappa_i = |\Gamma_i(\alpha)|$ satisfy the following obvious conditions

$$a_i + b_i + c_i = \kappa, \quad \kappa_{i-1}b_{i-1} = \kappa_i c_i, \quad i = 1, ..., d,$$

$$\kappa_0 = c_1 = 1, \quad b_0 = \kappa_1 = \kappa, \quad (c_0 = b_d = 0).$$
(2-8)

One should notice that, the definition of pseudo-distance regular graphs together with the Eq.(2-8), imply that in general, the valencies κ_i (the size of the *i*-th stratum) for i = 0, 1, ..., d do not depend on the considered vertex $\beta \in \Gamma_k(\alpha)$, but only on the value of k.

The notion of pseudo-distance regularity has a close relation with the concept of QD type graphs introduced by Obata [17], such that for the adjacency matrices of this type of graphs, one can obtain a quantum decomposition associated with the stratification (2-4) as

$$A = A^+ + A^- + A^0, (2-9)$$

where, the matrices A^+ , A^- and A^0 are defined as follows: for $\beta \in \Gamma_k(\alpha)$, we set

$$A^{+}|\beta\rangle = \sum_{\delta\in\Gamma_{k+1}(\alpha),\delta\sim\beta}|\delta\rangle, \quad A^{-}|\beta\rangle = \sum_{\delta\in\Gamma_{k-1}(\alpha),\delta\sim\beta}|\delta\rangle, \quad A^{0}|\beta\rangle = \sum_{\delta\in\Gamma_{k}(\alpha),\delta\sim\beta}|\delta\rangle, \quad (2-10)$$

Since $\beta \in \Gamma_k(\alpha)$ and $\beta \sim \delta$ then $\delta \in \Gamma_{k-1}(\alpha) \cup \Gamma_k(\alpha) \cup \Gamma_{k+1}(\alpha)$.

It has been shown in Ref. [17] that, if the closed subspace of $l^2(V)$ spanned by $\{|\phi_i\rangle, i = 0, \ldots, d-1\}$ be invariant under the quantum components A^+, A^- and A^0 , then there exist two sequences (called Szegö- Jacobi sequences) $\{\omega_l\}_{l=1}^{\infty}$ and $\{\alpha_l\}_{l=1}^{\infty}$ derived from A such that

$$A^{+}|\phi_{l}\rangle = \sqrt{\omega_{l+1}}|\phi_{l+1}\rangle, \quad l \ge 0 \quad ,$$

$$A^{-}|\phi_{0}\rangle = 0, \quad A^{-}|\phi_{l}\rangle = \sqrt{\omega_{l}}|\phi_{l-1}\rangle, \quad l \ge 1,$$

$$A^{0}|\phi_{l}\rangle = \alpha_{l}|\phi_{l}\rangle, l \ge 0,$$
(2-11)

where $\omega_{l+1} = \frac{\kappa_{l+1}}{\kappa_l} \kappa_-^2(j)$ with $\kappa_-(j) = |\{i \in \Gamma_l(\alpha) : i \sim j\}|$ for $j \in \Gamma_{l+1}(\alpha)$, and $\alpha_l = \kappa_0(j) = |\{i \in V_l; i \sim j\}|$ for $j \in \Gamma_l(\alpha)$, for $l \ge 0$. One can easily check that the coefficients α_l and ω_l are given by

$$\alpha_l = \kappa - b_l - c_l, \qquad \omega_l = b_{l-1}c_l, \quad l = 1, ..., d.$$
(2-12)

By using (2-9) and (2-11), one can obtain

$$A|\phi_l\rangle = \sqrt{\omega_{l+1}}|\phi_{l+1}\rangle + \alpha_l|\phi_l\rangle + \sqrt{\omega_l}|\phi_{l-1}\rangle, \quad l \ge 0.$$
(2-13)

One should notice that, the vectors $|\phi_i\rangle$, i = 0, 1, ..., d-1 form an orthonormal basis for the so called Krylov subspace $K_d(|\phi_0\rangle, A)$ defined as

$$K_d(|\phi_0\rangle, A) = span\{|\phi_0\rangle, A|\phi_0\rangle, \cdots, A^{d-1}|\phi_0\rangle\}.$$
(2-14)

Then it can be shown that [24], the orthonormal basis $|\phi_i\rangle$ are written as

$$|\phi_i\rangle = P_i(A)|\phi_0\rangle,\tag{2-15}$$

where $P_i = a_0 + a_1 A + ... + a_i A^i$ is a polynomial of degree *i* in indeterminate *A* (for more details see for example [19, 24]).

It may be noted that, the pseudo-distance-regularity is a generalization of the notion of distance-regularity which is defined as:

Definition (distance-regular graphs). A pseudo-distance regular graph $\Gamma = (V, E)$ is called distance-regular with diameter d if for all $k \in \{0, 1, ..., d\}$, and $\alpha, \beta \in V$ with $\beta \in \Gamma_k(\alpha)$, the numbers $c_k(\beta)$, $a_k(\beta)$ and $b_k(\beta)$ defined in (2-6) depend only on k but do not depend on the choice of α and β .

Now, it should be noticed that, the stratification of distance-regular graphs will be independent of the choice of the reference vertex (the vertex which stratification is done with respect to it).

2.3 Spectral distribution of the graphs

It is well known that, for any pair $(A, |\phi_0\rangle)$ of a matrix A and a vector $|\phi_0\rangle$, one can assign a measure μ as follows

$$\mu(x) = \langle \phi_0 | E(x) | \phi_0 \rangle, \tag{2-16}$$

where $E(x) = \sum_{i} |u_i\rangle \langle u_i|$ is the operator of projection onto the eigenspace of A corresponding to eigenvalue x, i.e.,

$$A = \int x E(x) dx. \tag{2-17}$$

Then, for any polynomial P(A) we have

$$P(A) = \int P(x)E(x)dx,$$
(2-18)

where for discrete spectrum the above integrals are replaced by summation. Therefore, using the relations (2-16) and (2-18), the expectation value of powers of adjacency matrix A over reference vector $|\phi_0\rangle$ can be written as

$$\langle \phi_0 | A^m | \phi_0 \rangle = \int_R x^m \mu(dx), \quad m = 0, 1, 2, \dots$$
 (2-19)

Obviously, the relation (2-19) implies an isomorphism from the Hilbert space of the stratification onto the closed linear span of the orthogonal polynomials with respect to the measure μ .

From orthonormality of the unit vectors $|\phi_i\rangle$ given in Eq.(2-5) (with $|\phi_0\rangle$ as unit vector assigned to the reference node) we have

$$\delta_{ij} = \langle \phi_i | \phi_j \rangle = \int_R P_i(x) P_j(x) \mu(dx), \qquad (2-20)$$

where, we have used the equation (2-15). Now, by substituting (2-15) in (2-13) and rescaling P_k as $Q_k = \sqrt{\omega_1 \dots \omega_k} P_k$, the spectral distribution μ under question will be characterized by the property of orthonormal polynomials $\{Q_k\}$ defined recurrently by

$$Q_0(x) = 1, \qquad Q_1(x) = x,$$

$$xQ_k(x) = Q_{k+1}(x) + \alpha_k Q_k(x) + \omega_k Q_{k-1}(x), \quad k \ge 1.$$
 (2-21)

If such a spectral distribution is unique, the spectral distribution μ is determined by the identity

$$G_{\mu}(x) = \int_{R} \frac{\mu(dy)}{x - y} = \frac{1}{x - \alpha_0 - \frac{\omega_1 - \omega_2}{x - \alpha_1 - \frac{\omega_1 - \omega_2}{x - \alpha_2 - \frac{\omega_3}{x - \alpha_3 - \cdots}}} = \frac{Q_d^{(1)}(x)}{Q_{d+1}(x)} = \sum_{l=0}^d \frac{\gamma_l}{x - x_l},$$
 (2-22)

where, x_l are the roots of the polynomial $Q_{d+1}(x)$. $G_{\mu}(x)$ is called the Stieltjes/Hilbert transform of spectral distribution μ and polynomials $\{Q_k^{(1)}\}$ are defined recurrently as

$$Q_0^{(1)}(x) = 1, \qquad Q_1^{(1)}(x) = x - \alpha_1,$$

$$xQ_k^{(1)}(x) = Q_{k+1}^{(1)}(x) + \alpha_{k+1}Q_k^{(1)}(x) + \omega_{k+1}Q_{k-1}^{(1)}(x), \quad k \ge 1,$$
(2-23)

respectively. The coefficients γ_l appearing in (2-22) are calculated as

$$\gamma_l := \lim_{x \to x_l} [(x - x_l) G_{\mu}(x)]$$
(2-24)

Now let $G_{\mu}(z)$ is known, then the spectral distribution μ can be determined in terms of $x_l, l = 1, 2, ...$ and Gauss quadrature constants $\gamma_l, l = 1, 2, ...$ as

$$\mu = \sum_{l=0}^{d} \gamma_l \delta(x - x_l) \tag{2-25}$$

(for more details see Refs. [18, 25, 26, 27]).

3 PST of a qudit over antipodes of pseudo distanceregular networks

3.1 State Transference in *d*-dimensional Quantum Systems

A *d*-dimensional quantum system associated with a simple, connected, finite graph G = (V, E)is defined by attaching a *d*-level particle to each vertex of the graph so that with each vertex $i \in V$ one can associate a Hilbert space $\mathcal{H}_i \simeq \mathcal{C}^d$. The Hilbert space associated with G is then given by

$$\mathcal{H}_G = \bigotimes_{i \in V} \mathcal{H}_i = (\mathcal{C}^d)^{\otimes N}, \tag{3-26}$$

where N := |V| denotes the total number of vertices (sites) in G.

Then, the quantum state transfer protocol involves two steps: initialization and evolution. First, a quantum state

$$|\psi\rangle_A = a_0|0\rangle_A + \sum_{\nu=1}^{d-1} a_\nu|\nu\rangle_A \in \mathcal{H}_A$$

(with $a_{\nu} \in \mathcal{C}$ and $\sum_{\nu=0}^{d-1} |a_{\nu}|^2 = 1$) to be transmitted is created. The state of the entire spin system after this step is given by

$$|\psi(t=0)\rangle = |\psi_A\rangle \otimes |0...00_B\rangle = a_0 |0_A\rangle \otimes |0...00_B\rangle + a_1 |1_A\rangle \otimes |0...00_B\rangle + ... + a_{d-1} |(d-1)_A\rangle \otimes |0...00_B\rangle$$
(3-27)

Then, the network couplings are switched on and the whole system is allowed to evolve under $U(t) = e^{-iHt}$ for a fixed time interval, say t_0 . From the fact that $H|0_A\rangle \otimes |0...00_B\rangle = 0$, the final state at time t_0 will be

$$|\psi(t_0)\rangle = a_0|0_A 0...00_B\rangle + \sum_{\nu=1}^{d-1} a_{\nu} \{\sum_{k=1}^N f_{kA}^{(\nu)}(t_0)|0\ldots \underbrace{\nu}_{k-th} 0...0\rangle\},$$
(3-28)

where $f_{kA}^{(\nu)}(t_0) := \langle 0...0 \underbrace{\nu}_{k-th} 0...0 | e^{-iHt_0} | \nu_A 0...0 \rangle$ for k = 1, 2, ..., N; $\nu = 1, ..., d-1$. In order to transfer the state $|\psi_A\rangle$ to the site *B* perfectly (in order to PST is achieved), the following conditions must be fulfilled

$$|f_{AB}^{(\nu)}(t_0)| = 1$$
 for $\nu = 1, 2, ..., d-1$ and some $0 < t_0 < \infty$ (3-29)

which can be interpreted as the signature of perfect communication (or PST) between A and B in time t_0 . The effect of the modulus in (3-29) is that the state (3-28) will be

$$|\psi(t_0)\rangle = a_0|0_A 0...0_B\rangle + \sum_{\nu=1}^{d-1} e^{i\phi_\nu} a_\nu|0_A 0...0\rangle \otimes |\nu\rangle_B$$

so, the state at B, after transmission, will no longer be $|\psi\rangle_A$, but will be of the form

$$a_0|0\rangle + \sum_{\nu=1}^{d-1} e^{i\phi_{\nu}} a_{\nu}|\nu\rangle_B.$$
(3-30)

The phase factors $e^{i\phi_{\nu}}$ for $\nu = 1, 2, ..., d-1$ are independent of a_0, \ldots, a_{d-1} and will thus be known quantities for the graph, which one can correct for with appropriate phase gates.

The model we will consider is a pseudo distance-regular network consisting of N sites labeled by $\{1, 2, ..., N\}$ and diameter D. In Ref. [1], we introduced the PST of a qubit in terms of the SU(2) generators. Let us now consider a state with d levels. First, we prepare the generators for SU(d) systems and thereby introduce the Hamiltonians for a qudit system. The generators of SU(d) group may be conveniently constructed by the elementary matrices of d dimension, $\{e_{pq}|p,q \in \{1,...,d\}\}$. The elementary matrices are given by

$$(e_{pq})_{ij} = \delta_{ip}\delta_{jq}, \ 1 \le i, j \le d; \ e_p := e_{pp}.$$
 (3-31)

which are matrices with one matrix element equal to unity and all others equal to zero. These matrices satisfy the commutation relation

$$[e_{pq}, e_{rs}] = \delta_{sp} e_{rq} - \delta_{qr} e_{ps}$$

There are d(d-1) traceless matrices

$$\lambda_{pq}^{+} = e_{pq} + e_{qp},$$

$$\lambda_{pq}^{-} = \frac{1}{i} (e_{pq} - e_{qp}); \quad 1 \le p < q \le d$$
(3-32)

which are the off-diagonal generators of the SU(d) group. The d-1 additional traceless matrices

$$H_m = \frac{2}{\sqrt{2m(m+1)}} \{\sum_{k=1}^m e_k - me_{m+1}\}; \quad m = 1, 2, ..., d-1$$
(3-33)

are the diagonal generators so that we obtain a total of $d^2 - 1$ generators. SU(2) generators are, for instance, given as $\sigma_x = \lambda_{12}^+ = e_{21} + e_{12}$, $\sigma_y = \lambda_{21}^- = -i(e_{12} - e_{21})$ and $\sigma_z = H_1 = e_1 - e_2$.

We now assume that at time t = 0, the qudit in the first (input) site of the network is prepared in the state $|\psi_{in}\rangle$. We wish to transfer the state to the Nth (output) site of the network with unit efficiency after a well-defined period of time. As regards in the above argument, we choose the standard basis $e_i = |i\rangle$, i = 0, 1, ..., d - 1 for an individual qudit, and assume that initially all particles are in the state $|0\rangle$; i.e., the network is in the state $|\underline{0}\rangle = |0_A 00...00_B\rangle$. Then, we consider the dynamics of the system to be governed by the quantum-mechanical Hamiltonian

$$H_G = \frac{1}{2} \sum_{m=0}^{D} J_m P_m(\frac{1}{2} \sum_{i \sim j} \vec{\lambda}_i \cdot \vec{\lambda}_j + [\kappa_{max} - |E|(\frac{d-1}{d})]I_{d^N}),$$
(3-34)

where, $\vec{\lambda}_i$ is a $d^2 - 1$ dimensional vector with generators of SU(d) as its components acting on the one-site Hilbert space \mathcal{H}_i , J_m is the coupling strength between the reference site 1 and all of the sites belonging to the *m*-th stratum with respect to 1, and P_m 's are polynomials given in (2-15) which are obtained using three term recursion relations (2-21) and the fact that $P_m = \frac{1}{\sqrt{\omega_1 \omega_2 \dots \omega_m}} Q_m$. As it is seen from the Eq. (3-34), the terms of the hamiltonian for $m \ge 1$ are nonlinear functions of $\sum_{i\sim j} \vec{\lambda}_i \cdot \vec{\lambda}_j$.

In the following we note that the term $H_{ij} := \vec{\lambda}_i \cdot \vec{\lambda}_j$ in the hamiltonian (3-34), restricted to the one particle subspace (the subspace of the full Hilbert space spanned by the states with only one site excited), is related to the adjacency matrix of the corresponding graph. To do so, we write H_{ij} as follows

$$H_{ij} = \sum_{1 \le p < q \le d} (\lambda_{pq}^{+(i)} \otimes \lambda_{pq}^{+(j)} + \lambda_{pq}^{-(i)} \otimes \lambda_{pq}^{-(j)}) + \sum_{m=1}^{d-1} H_m^{(i)} \otimes H_m^{(j)}.$$
 (3-35)

Before we proceed, one should notice that we have

$$e_{pq} \otimes e_{rs} = e_{(p-1)d+r,(q-1)d+s}$$
 (3-36)

Then, from the fact that

$$\lambda_{pq}^{+(i)} \otimes \lambda_{pq}^{+(j)} + \lambda_{pq}^{-(i)} \otimes \lambda_{pq}^{-(j)} = 2(e_{pq}^{(i)} \otimes e_{qp}^{(j)} + e_{qp}^{(i)} \otimes e_{pq}^{(j)}),$$

and using the notation

$$(m,n) \equiv m + (n-1)d,$$

one can obtain

$$\sum_{1 \le p < q \le d} (\lambda_{pq}^{+(i)} \otimes \lambda_{pq}^{+(j)} + \lambda_{pq}^{-(i)} \otimes \lambda_{pq}^{-(j)}) = 2 \sum_{1 \le p < q \le d} [e_{(q,p),(p,q)} + e_{(p,q),(q,p)}]$$
(3-37)

Now, we evaluate the term $\sum_{m=1}^{d-1} H_m \otimes H_m$ in (3-35) in terms of the elementary matrices e_{pq} as follows: First we note that

$$H_m \otimes H_m = \frac{2}{m(m+1)} \{ \sum_{p=1}^m \sum_{p'=1}^m e_{(p',p)} - m \sum_{p=1}^m [e_{(m+1,p)} + e_{(p,m+1)}] + m^2 e_{(m+1,m+1)} \}.$$
 (3-38)

The Eq. (3-38) can be rewritten as follows

$$H_m \otimes H_m = \left\{ \frac{2}{m(m+1)} \sum_{p=1}^m e_{(p,p)} + \frac{2m}{m+1} e_{(m+1,m+1)} \right\} + \left\{ \frac{2}{m(m+1)} \sum_{p,p'=1; p \neq p'}^m e_{(p',p)} - \frac{2}{m+1} \sum_{p=1}^m [e_{(m+1,p)} + e_{(p,m+1)}] \right\}.$$
(3-39)

Then, one can show that

$$\sum_{m=1}^{d-1} H_m \otimes H_m = 2 \sum_{p=1}^d e_{(p,p)} - \frac{2}{d} I,$$
(3-40)

for proof see Appendix A.

Therefore, by using (3-37) and (3-40), H_{ij} in (3-35) is written as follows

$$H_{ij} = 2 \sum_{1 \le p < q \le d} [e_{(q,p),(p,q)} + e_{(p,q),(q,p)}] + 2 \sum_{p=1}^{d} e_{(p,p)} - \frac{2}{d}I.$$
 (3-41)

Now, one should notice that the permutation matrix which permutes two qudits, in terms of the elementary basis e_{pq} can be written as

$$P = \sum_{p,q=1}^{d} e_{(p,q),(q,p)} = \sum_{p=1}^{d} e_{(p,p)} + \sum_{1 \le p < q \le d}^{d} [e_{(p,q),(q,p)} + e_{(q,p),(p,q)}].$$
 (3-42)

Then, the Eq.(3-41) takes the following form

$$H_{ij} = 2P_{ij} \otimes I_{d^{N-2}} - \frac{2}{d}I_{d^N}, \qquad (3-43)$$

where, P_{ij} denotes the permutation operator which permutes *i*-th and *j*-th sites and I_{d^N} is $d^N \times d^N$ identity matrix, where N is the number of vertices or sites (N := |V|).

Now, we denote a state in which the *i*-th site has been exited to the level ν by $|\nu_i\rangle \equiv |0....0, \underbrace{\nu}_{i} 0...0\rangle$. Then, the permutation operators P_{ij} only changes this state so that the number and type of excited local states is fixed. This implies that the hamiltonian H_G can be diagonalized in each subspace $S^{(\nu)}$ spanned by the vectors $|\nu_i\rangle$, i = 1, ..., N, for $\nu = 1, ..., d-1$.

We will refer to the states with only one site excited as one particle states and the subspace spanned by these vectors comprise the one-particle sector of the full Hilbert space. Then, the whole one particle subspace S can be written as

$$S = S^{(1)} \oplus S^{(2)} \oplus \ldots \oplus S^{(d-1)}.$$

In the other words, in d^N dimensional Hilbert space, we deal with d-1 one particle subspaces (recall that, each of these subspaces has dimension N). In the case of PST of a qubit, we have only one one-particle subspace of dimension N.

One can easily show that, the operator $\sum_{i\sim j} P_{ij}$ restricted to the one particle subspace $S^{(l)}$, can be related to the adjacency matrix A, as follows

$$\sum_{i \sim j} P_{ij} - \sum_{i=1}^{N} [\kappa_{\max} - \kappa(i)] H_l^{(i)} = A + (|E| - \kappa_{\max}) I_N,$$
(3-44)

where $\kappa_{\max} := \max\{\kappa(i), i = 1, 2, ..., N\}, |E|$ is the number of the edges of the graph, and $H_l^{(i)}$ is the projection operator $I \otimes ... \otimes I \otimes \underbrace{H_l}_i \otimes I \ldots I$ with H_l defined as in (3-33). For regular graphs, where we have $\kappa(i) \equiv \kappa$ for all i = 1, 2, ..., N, the equation (3-44) reads as

$$\sum_{i \sim j} P_{ij} = A + \kappa (\frac{N-2}{2}) I_N, \tag{3-45}$$

in which we have substituted $|E| = \frac{N\kappa}{2}$.

Then, by using (3-43) and (3-44), the hamiltonian in (3-34) restricted to each one particle subspace $S^{(\nu)}$, for $\nu = 1, 2, ..., d - 1$, can be written in terms of the adjacency matrix A, as follows

$$H_G = \frac{1}{2} \sum_{m=0}^{D} J_m P_m \left(\sum_{i \sim j} P_{ij} - \frac{|E|}{d} I_N + [\kappa_{max} - |E|(\frac{d-1}{d})]I_N\right) = \sum_{m=0}^{D} J_m P_m(A).$$
(3-46)

For the purpose of the perfect transference of a qudit, we consider pseudo-distance-regular graphs with $\kappa_D = |\Gamma_D(o)| = 1$, i.e., the last stratum of the graph contains only one site. Then, we impose the constraints that the amplitudes $\langle \phi_i^{(\nu)} | e^{-iHt_0} | \phi_0^{(\nu)} \rangle$ be zero for all i = 0, 1, ..., D-1and $\nu = 1, 2, ..., d-1$, and $\langle \phi_D^{(\nu)} | e^{-iHt_0} | \phi_0^{(\nu)} \rangle = e^{i\theta}$, where θ is an arbitrary phase. Recall that we have

$$|\phi_0^{(\nu)}\rangle = |\nu 0...0\rangle, \ |\phi_i^{(\nu)}\rangle = P_i(A)|\phi_0^{(\nu)}\rangle; \ \nu = 1, 2, ..., d-1.$$

Therefore, the amplitudes $\langle \phi_i^{(\nu)} | e^{-iHt_0} | \phi_0^{(\nu)} \rangle$, for i = 0, 1, ..., D must be evaluated. One can easily show that these amplitudes are independent of the value of ν , i.e., it suffices to evaluate them for one choice of ν , say $\nu = 1$.

As it has been shown in Ref. [1], the above mentioned constraints (by choosing $\nu = 1$, the other choices give the same results), are equivalent to the following ones:

$$\sum_{k=0}^{D} \gamma_k P_i(x_k) e^{-2it_0 \sum_{m=0}^{D} J_m P_m(x_k)} = 0, \quad i = 0, 1, ..., D - 1$$

 $\sum_{k=0} \gamma_k P_i(x_k) e^{-2it_0 \sum_{m=0}^{-} J_m P_m(x_k)} = 0, \quad i = 0, 1, ..., D - 1.$ Denoting $e^{-2it_0 \sum_{m=0}^{D} J_m P_m(x_k)}$ by η_k , the above constraints are rewritten as follows

$$\sum_{k=0}^{D} P_i(x_k)\eta_k\gamma_k = 0, \quad i = 0, 1, ..., D - 1,$$
$$\sum_{k=0}^{D} P_D(x_k)\eta_k\gamma_k = e^{i\theta}.$$
(3-47)

The Eq.(3-47) can be written as

$$\mathbf{P}\begin{pmatrix} \eta_{0}\gamma_{0} \\ \eta_{1}\gamma_{1} \\ \vdots \\ \eta_{D}\gamma_{D} \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ e^{i\theta} \end{pmatrix}, \qquad (3-48)$$

with $P_{ij} = P_i(x_j)$. From the orthogonality relation (2-20) and using (2-25), one can obtain

$$\delta_{ij} = \langle \phi_i | \phi_j \rangle = \sum_{l=0}^{D} P_i(x_l) \gamma_l P_j(x_l) \quad \rightarrow \quad \mathrm{PWP}^t = I \quad \rightarrow \quad \mathrm{PW}^{1/2} (\mathrm{PW}^{1/2})^t = I \quad \rightarrow \quad (\mathrm{PW}^{1/2})^{-1} = (\mathrm{PW}^{1/2})^t \quad \rightarrow \quad \mathrm{P}^{-1} = W\mathrm{P}^t,$$

with $W := diag(\gamma_0, \gamma_1, \ldots, \gamma_D)$. Therefore, P is invertible and the Eq. (3-48) can be rewritten as

$$\begin{pmatrix} \eta_0 \gamma_0 \\ \eta_1 \gamma_1 \\ \vdots \\ \eta_D \gamma_D \end{pmatrix} = \mathbf{P}^{-1} \begin{pmatrix} 0 \\ \vdots \\ 0 \\ e^{i\theta} \end{pmatrix} = W \mathbf{P}^t \begin{pmatrix} 0 \\ \vdots \\ 0 \\ e^{i\theta} \end{pmatrix}.$$
(3-49)

The above equation implies that $\eta_k \gamma_k$ for k = 0, 1, ..., D are the same as the entries in the last column of the matrix $P^{-1} = WP^t$ multiplied with the phase $e^{i\theta}$, i.e., for the purpose of PST, the following equations must be satisfied

$$\eta_k \gamma_k = \gamma_k e^{-2it_0 \sum_{m=0}^{D} J_m P_m(x_k)} = e^{i\theta} (W \mathbf{P}^t)_{kD} \quad , \quad \text{for} \quad k = 0, 1, ..., D.$$
(3-50)

One should notice that, the Eq.(3-50) can be rewritten as

$$(J_0, J_1, \dots, J_D) = -\frac{1}{2t_0} [\theta + (2l_0 + f(0))\pi, \theta + (2l_1 + f(1))\pi, \dots, \theta + (2l_D + f(D))\pi] (WP^t), (3-51)$$

or

$$J_k = -\frac{1}{2t_0} \sum_{m=0}^{D} [\theta + (2l_m + f(m))\pi] (WP^t)_{mk}, \qquad (3-52)$$

where l_k for k = 0, 1, ..., D are integers and f(k) is equal to 0 or 1 (we have used the fact that γ_k and $(WP^t)_{kD}$ are real for k = 0, 1, ..., D, and so we have $\gamma_k = |(WP^t)_{kD}|$). The result (3-52) gives an explicit formula for suitable coupling constants so that PST between the first node $(|\phi_0\rangle)$ and the opposite one $(|\phi_D\rangle)$ can be achieved.

4 Examples of pseudo-distance-regular networks

1. The networks G_n

The networks G_n presented in [28], consist of two balanced binary trees of height n with the 2^n leaves of the left tree identified with the 2^n leaves of the right tree in the simple way shown in Fig. 1 (for n = 2). The number of vertices in G_n is $2^{n+1} + 2^n - 2$. For the purpose of PST over G_n , we prepare the initial state to be transferred, at the left root of the graph and wants to calculate the suitable strength coupling constants so that the probability of the presence of the initial state at the right root be equal to one for some finite time t_0 . One can show that G_n is a pseudo-distance-regular graph with (2n+1) strata, where stratum j consists of 2^j vertices for j = 1, 2, ..., n + 1 and $2^{(2n+1-j)}$ for j = n + 1, ..., 2n + 1. Therefore, its QD parameters are

$$\alpha_i = 0, \quad i = 0, 1, \dots, 2n; \quad \omega_i = 2, \quad i = 1, 2, \dots, 2n.$$

Then one can show that [19, 20] the polynomials $P_i(x)$ are given by

$$P_i(x) = \frac{1}{\sqrt{2^i}} Q_i(x) = U_i(\frac{x}{2\sqrt{2}}), \qquad (4-53)$$

where U_i 's are the Tchebishef polynomial of the second kind. Then, x_l 's (the roots of the Tchebishef polynomial $Q_{2n+1}(x) = \sqrt{2^{2n+1}}U_{2n+1}(\frac{x}{2\sqrt{2}})$) and the coefficients γ_l are given by

$$x_{l} = 2\sqrt{2}\cos\frac{(l+1)\pi}{2(n+1)},$$
$$\gamma_{l} = \frac{(-1)^{l+1}}{n+1}\sin\frac{(l+1)\pi}{2(n+1)}\sin\frac{(2n+1)(l+1)\pi}{2(n+1)}, \quad l = 0, 1, \dots, 2n.$$
(4-54)

Therefore, we have $P_{ij} = U_i(\frac{x_j}{2\sqrt{2}}) = U_i(\cos \frac{\pi(j+1)}{2(n+1)})$ or equivalently

$$P = \begin{pmatrix} 1 & 1 & \dots & 1 \\ U_1(\frac{\pi}{2(n+1)}) & U_1(\cos\frac{\pi}{n+1}) & \dots & U_1(\cos\frac{(2n+1)\pi}{2(n+1)}) \\ \vdots & \vdots & \dots & \vdots \\ U_{2n}(\frac{\pi}{2(n+1)}) & U_{2n}(\cos\frac{\pi}{n+1}) & \dots & U_{2n}(\cos\frac{(2n+1)\pi}{2(n+1)}) \end{pmatrix}.$$
 (4-55)

Furthermore, as it is seen from (4-55), the matrix P is a polynomial transformation [29]. By using the result (3-52), we obtain

$$J_{k} = -\frac{1}{2t_{0}} \sum_{m=0}^{2n} \frac{(-1)^{m+1}}{n+1} \sin \frac{(m+1)\pi}{2(n+1)} \sin \frac{(2n+1)(m+1)\pi}{2(n+1)} [\theta + (2l_{m} + f(m))\pi] U_{k}(\cos \frac{(m+1)\pi}{2(n+1)}),$$
(4-56)

for $k = 0, 1, \dots, 2n$.

In the following we consider the case n = 2 in detail. In this case, we have

$$x_0 = \sqrt{6}, \quad x_1 = \sqrt{2}, \quad x_2 = 0, \quad x_3 = -\sqrt{2}, \quad x_4 = -\sqrt{6};$$

 $\gamma_0 = \gamma_4 = \frac{1}{12}, \quad \gamma_1 = \gamma_3 = \frac{1}{4}, \quad \gamma_2 = \frac{1}{3}.$

Then by using the equations (4-55) and (3-49), we have

$$P = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ \sqrt{3} & 1 & 0 & -1 & -\sqrt{3} \\ 2 & 0 & -1 & 0 & 2 \\ \sqrt{3} & -1 & 0 & 1 & -\sqrt{3} \\ 1 & -1 & 1 & -1 & 1 \end{pmatrix} \rightarrow P^{-1} = WP^{t} = \frac{1}{12} \begin{pmatrix} 1 & \sqrt{3} & 2 & \sqrt{3} & 1 \\ 3 & 3 & 0 & -3 & -3 \\ 4 & 0 & -4 & 0 & 4 \\ 3 & -3 & 0 & 3 & -3 \\ 1 & -\sqrt{3} & 2 & -\sqrt{3} & 1 \end{pmatrix}.$$

Now, using the Eq. (4-56), one can obtain the following suitable coupling constants in order to PST between the most left and the most right nodes be achieved:

$$J_0 = -\frac{\theta + 2\pi/3}{2t_0}, \quad J_1 = J_3 = \frac{\pi}{4\sqrt{3}t_0}, \quad J_2 = -J_4 = -\frac{\pi}{6t_0}$$

2. The modified G_n -type networks

This type of networks consist of two balanced binary trees of height n + 1. For the graphs G_n , set entrance and exit to the left root and the right root of the trees, respectively (see Fig. 2 for n = 2). These networks are pseudo distance regular networks which are characterized by the following intersection array

$$\{b_0, b_1, \dots, b_{2n-1}; c_1, c_2, \dots, c_{2n}\} = \{3, \underbrace{2, 2, \dots, 2}_{n-1}, \underbrace{1, 1, \dots, 1}_{n}; \underbrace{1, 1, \dots, 1}_{n}, \underbrace{2, 2, \dots, 2}_{n-1}, 3\}.$$

Then by using (2-8) and (2-12), one can obtain

$$\kappa \equiv \kappa_1 = 3, \quad \kappa_i = \kappa_{2n-i} = 3 \times 2^{i-1}, \quad i = 1, 2, \dots, n \quad \kappa_{2n} = 1,$$

 $\alpha_i = 0, \quad i \neq n+1, \quad \alpha_{n+1} = 1; \quad \omega_1 = \omega_{2n} = 3, \quad \omega_i = 2, \quad i = 2, \dots, 2n-1.$

Now, by using the QD parameters α_i and ω_i , and the recursion relations (2-21) and (2-23), one can obtain the Stieltjes function for any given n and spectral distribution $\mu(x)$ for the modified G_n -type network. In the following we consider the case n = 2 in detail. In this case, we have

$$\alpha_i = 0, \quad i = 0, 1, 2, 4, \quad \alpha_3 = 1; \quad \omega_1 = \omega_4 = 3, \quad \omega_2 = \omega_3 = 2.$$

Then by using the recursion relations (2-21) and (2-23), one can obtain the stieltjes function as

$$G_{\mu}(x) = \frac{x^4 - 7x^2 - x^3 + 2x + 6}{x(x-3)(x^3 + 2x^2 - 4x - 7)}$$

By using the recursion relations (2-21), we obtain

$$Q_{1}(x) = x, \quad Q_{2}(x) = x^{2} - 3, \quad Q_{3}(x) = x(x^{2} - 5), \quad Q_{4}(x) = x^{4} - x^{3} - 7x^{2} + 5x + 6, \quad Q_{5}(x) = x(x^{4} - x^{3} - 10x^{2} + 5x + 21)$$

$$(4-57)$$

Then, from the fact that, x_l , l = 0, 1, 2, 3, 4 are roots of $Q_5(x)$ (the denominator of $G_{\mu}(x)$), we obtain

$$x_0 = 0, \quad x_1 = 3, \quad x_2 \simeq -2.4728, \quad x_3 \simeq -1.4626, \quad x_4 \simeq 1.9354;$$

 $\gamma_0 = \frac{2}{7}, \quad \gamma_1 = \frac{1}{26}, \quad \gamma_2 \simeq 0.3101, \quad \gamma_3 \simeq 0.1786, \quad \gamma_4 \simeq 0.1872.$

By using the Eq.(2-21) and the fact that $P_{ij} = P_i(x_j) = \frac{1}{\sqrt{\sqrt{\omega_1 \dots \omega_i}}}Q_i(x_j)$, we obtain

$$P \simeq \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & \sqrt{3} & -1.4277 & -0.8445 & 1.1174 \\ -1.3417 & 0 & 1.3930 & -0.3850 & 0.1492 \\ 0 & 2\sqrt{3} & -0.4594 & 0.6974 & 0.4046 \\ 1 & 2 & 0.5572 & -1.4304 & -0.6270 \end{pmatrix} \rightarrow \\ P^{-1} = WP^{t} \simeq \begin{pmatrix} 0.2869 & -0.0155 & -0.3725 & -0.1982 & 0.2133 \\ -0.0303 & -0.0157 & 0.0493 & 0.2495 & 0.0965 \\ 0.2509 & -0.1527 & 0.2961 & -0.0807 & 0.1465 \\ 0.0712 & -0.3078 & -0.1565 & 0.2957 & -0.2813 \\ 0.4213 & 0.4916 & 0.1836 & -0.2664 & -0.1750 \end{pmatrix}$$

Now, using the Eq. (3-52), one can obtain the following suitable coupling constants in order to PST between the most left and the most right nodes be achieved:

$$J_0 = -\frac{\theta + 0.4925\pi}{2t_0}, \quad J_1 = -\frac{0.1838\pi}{2t_0}, \quad J_2 = -\frac{0.0271\pi}{2t_0}, \quad J_3 = -\frac{0.0293\pi}{2t_0}, \quad J_4 = \frac{0.4563\pi}{2t_0}.$$

3. Pseudo-distance-regular network derived from Icosahedron network

This network has 12 nodes with

$${b_0, b_1, b_2; c_1, c_2, c_3} = {5, 2, 1; 1, 2, 5}.$$

Then by using (2-8) and (2-12), one can obtain

$$\kappa = 5, \ \kappa_2 = 5, \ \kappa_3 = 1,$$

$$\alpha_0 = 0, \ \alpha_1 = \alpha_2 = 2, \ \alpha_3 = 0; \ \omega_1 = 5, \ \omega_2 = 4, \ \omega_3 = 5.$$

The stratification with respect to the initial node 1 and the final node 12 produces the same strata (see Fig. 3). Therefore, the PST between the nodes 1 and 12 can be achieved. To do so, by using the recursion relations (2-21), we obtain

$$Q_1(x) = x, \quad Q_2(x) = x^2 - 2x - 5, \quad Q_3(x) = x^3 - 4x^2 - 5x + 10, \quad Q_4(x) = x^4 - 4x^3 - 10x^2 + 20x + 25.$$

(4-58)

Then, from the fact that, x_l , l = 0, 1, 2, 3 are roots of Q_4 , we obtain

$$x_0 = -1, \ x_1 = 5, \ x_2 = \sqrt{5}, \ x_3 = -\sqrt{5}; \ \gamma_0 = \frac{5}{12}, \ \gamma_1 = \frac{1}{12}, \ \gamma_2 = \gamma_3 = \frac{1}{4}$$

By using the Eq.(2-21) and the fact that $P_{ij} = P_i(x_j) = \frac{1}{\sqrt{\sqrt{\omega_1 \dots \omega_i}}}Q_i(x_j)$, we obtain

$$P = \begin{pmatrix} 1 & 1 & 1 & 1 \\ -\frac{1}{\sqrt{5}} & \sqrt{5} & 1 & -1 \\ -\frac{1}{\sqrt{5}} & \sqrt{5} & -1 & 1 \\ 1 & 1 & -1 & -1 \end{pmatrix} \rightarrow (P)^{-1} = WP^{t} = \frac{1}{12} \begin{pmatrix} 5 & -\sqrt{5} & -\sqrt{5} & 5 \\ 1 & \sqrt{5} & \sqrt{5} & 1 \\ 3 & 3 & -3 & -3 \\ 3 & -3 & 3 & -3 \end{pmatrix}.$$

Now, using the Eq. (3-52), we obtain the following coupling constants

$$J_0 = -\frac{2\theta + \pi}{4t_0}, \quad J_1 = J_2 = 0, \quad J_3 = \frac{\pi}{4t_0}.$$

4. 3- simplex fractal with decimation number b = 2

Consider the (n-1)-simplex fractal with decimation number b = 2 [30]-[32] such that all of the n-1 vertices (20...0), (020...0), ..., (0...02) are connected to each other (see Fig. 4 for n = 4). Then, the number of vertices of (n-1)-simplex fractal is $N = C_1^{n-1} + C_2^{n-1} = \frac{n(n-1)}{2}$ such that the degree of each vertex is $\kappa = 2(n-2)$. Also, it can be easily shown that the network has 3 strata with respect to the reference node (200...0). For n = 4, the last stratum contains only one nodes so that the PST can be considered; In this case, the intersection array, size of strata and the QD parameters are given by

$$\{b_0, b_1; c_1, c_2\} = \{4, 1; 1, 1, 4\},\$$

$$\kappa = 4, \quad \kappa_2 = 1,\$$

$$\alpha_0 = 0, \quad \alpha_1 = 2, \quad \alpha_2 = 0; \quad \omega_1 = \omega_2 = 4.$$

Then, by using the recursion relations (2-21), we obtain

$$Q_1(x) = x, \quad Q_2(x) = x^2 - 2x - 4, \quad Q_3(x) = x^3 - 2x^2 - 8x.$$
 (4-59)

Then, from the fact that, x_l , l = 0, 1, 2, 3 are roots of Q_3 , we obtain

$$x_0 = 0$$
, $x_1 = 4$, $x_2 = -2$; $\gamma_0 = \frac{1}{2}$, $\gamma_1 = \frac{1}{6}$, $\gamma_2 = \frac{1}{3}$

By using the Eq.(2-21) and the fact that $P_{ij} = P_i(x_j) = \frac{1}{\sqrt{\omega_1 \dots \omega_i}} Q_i(x_j)$, we obtain

$$P = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 2 & -1 \\ -1 & 1 & 1 \end{pmatrix} \rightarrow (P)^{-1} = WP^{t} = \frac{1}{6} \begin{pmatrix} 3 & 0 & -3 \\ 1 & 2 & 1 \\ 2 & -2 & 2 \end{pmatrix}$$

By using the Eq. (3-52), we obtain the following constants in order to PST be achieved:

$$J_0 = -\frac{2\theta + 5\pi/3}{4t_0}, \quad J_1 = -\frac{\pi}{3t_0}, \quad J_2 = \frac{\pi}{12t_0}.$$

5. Pseudo-distance-regular network derived from Hadamard network with 16 nodes

Consider the pseudo-distance-regular network given in Fig. 5. This network is obtained from the Hadamard network with intersection array $\{4, 3, 2, 1; 1, 2, 3, 4\}$. As Fig. 5 shows, the network is symmetric with respect to the initial and final (horizontal) nodes 1 and 16 $\in \Gamma_4(1)$ and also with respect to the initial and final (vertical) nodes 6 and $11 \in \Gamma_4(6)$. One should notice that stratification of the network with respect to the nodes 1 and 16 produces the same strata. For stratification with respect to the node 1 or 16, we have

$$\kappa = 4, \quad \kappa_2 = 6, \quad \kappa_3 = 4, \quad \kappa_4 = 1,$$

 $\alpha_i = 0, \quad i = 0, \dots, 4; \quad \omega_1 = 4, \quad \omega_2 = 6, \quad \omega_3 = 6, \quad \omega_4 = 4.$

(see Eq. (2-11)). Then, by using the recursion relations (2-21), one can obtain

$$Q_1(x) = x, \ Q_2(x) = x^2 - 4, \ Q_3(x) = x^3 - 10x, \ Q_4(x) = x^4 - 16x^2 + 24, \ Q_5(x) = x^5 - 20x^3 + 64x$$

$$(4-60)$$

Then, from the fact that, x_l , l = 0, 1, 2, 3, 4 are roots of Q_5 , we obtain

$$x_0 = 0$$
, $x_1 = 2$, $x_2 = -2$, $x_3 = 4$, $x_4 = -4$; $\gamma_0 = \frac{3}{8}$, $\gamma_1 = \gamma_2 = \frac{1}{4}$, $\gamma_3 = \gamma_4 = \frac{1}{16}$

Then by using the equations (4-55) and (3-49), we have

$$P = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & -1 & 2 & -2 \\ -\frac{\sqrt{6}}{3} & 0 & 0 & \sqrt{6} & \sqrt{6} \\ 0 & -1 & 1 & 2 & -2 \\ 1 & -1 & -1 & 1 & 1 \end{pmatrix} \rightarrow P^{-1} = WP^{t} = \frac{1}{16} \begin{pmatrix} 6 & 0 & -2\sqrt{6} & 0 & 6 \\ 4 & 4 & 0 & -4 & -4 \\ 4 & -4 & 0 & 4 & -4 \\ 1 & 2 & \sqrt{6} & 2 & 1 \\ 1 & -2 & \sqrt{6} & -2 & 1 \end{pmatrix}.$$

Now, using the Eq. (3-52), we obtain the following constants:

$$J_0 = -\frac{\theta + 5\pi/8}{2t_0}, \quad J_1 = J_3 = -\frac{\pi}{8t_0}, \quad J_2 = -\frac{3\pi}{8\sqrt{6}t_0}, \quad J_4 = \frac{3\pi}{16t_0}$$

5 Conclusion

PST of an arbitrary *d*-level quantum state over antipodes of pseudo distance-regular networks, was investigated. By using the spectral analysis techniques and algebraic structures of pseudo distance-regular graphs an explicit formula for coupling constants in the Hamiltonians was given so that state of a particular qudit initially encoded on one site can be evolved freely to the opposite site without any dynamical control.

Appendix A

Proof of the Eq.(3-40):

By using the Eq. (3-39), we have

$$\sum_{m=1}^{d-1} H_m \otimes H_m = \sum_{m=1}^{d-1} \left\{ \frac{2}{m(m+1)} \sum_{p=1}^m e_{(p,p)} + \frac{2m}{m+1} e_{(m+1,m+1)} \right\} + \sum_{m=1}^{d-1} \left\{ \frac{2}{m(m+1)} \sum_{p,p'=1; p \neq p'}^m e_{(p',p)} - \frac{2}{m+1} \sum_{p=1}^m [e_{(m+1,p)} + e_{(p,m+1)}] \right\}.$$
(A-i)

We evaluate the first sum in the above equation, the second one can be evaluated similarly.

$$\begin{split} &\sum_{m=1}^{d-1} \left\{ \frac{2}{m(m+1)} \sum_{p=1}^{m} e_{(p,p)} + \frac{2m}{m+1} e_{(m+1,m+1)} \right\} = \sum_{m=1}^{d-1} \frac{2}{m(m+1)} \sum_{p=1}^{m} e_{(p,p)} + \sum_{m=1}^{d-1} \frac{2m}{m+1} e_{(m+1,m+1)} \\ &= \sum_{p=1}^{1} e_{(p-1)d+p} + \frac{1}{3} \sum_{p=1}^{2} e_{(p-1)d+p} + \ldots + \frac{2}{2(d-1)} \sum_{p=1}^{d-1} e_{(p-1)d+p} + e_{d+2} + \frac{4}{3} e_{2d+3} + \ldots + \frac{2(d-1)}{d} e_d = e_1 \sum_{m=1}^{d-1} \frac{2}{m(m+1)} + e_{d+2} \sum_{m=2}^{d-1} \frac{2}{m(m+1)} + \ldots + e_{d-1} \frac{2}{d(d-1)} + e_{d+2} + \frac{4}{3} e_{2d+3} + \ldots + \frac{2(d-1)}{d} e_d = e_1 \sum_{\alpha=1}^{d} \left[\frac{2(\alpha-1)}{\alpha} + 2(\frac{1}{\alpha} - \frac{1}{d}) \right] e_{(\alpha-1)d+\alpha} = 2(1-1/d)(e_1 + e_{d+2} + \ldots + e_{d^2}) = 2(1-1/d) \sum_{p=1}^{d} e_{(p-1)d+p}, \end{split}$$
where, we have used the identity $\sum_{m=\alpha}^{d-1} \frac{2}{m(m+1)} = \sum_{m=\alpha}^{d-1} (\frac{1}{m} - \frac{1}{m+1}) = 2(\frac{1}{\alpha} - \frac{1}{d}).$

The second sum in (A-i) can be evaluated as

$$\sum_{m=1}^{d-1} \left\{ \frac{2}{m(m+1)} \sum_{p,p'=1; p \neq p'}^{m} e_{(p',p)} - \frac{2}{m+1} \sum_{p=1}^{m} [e_{(m+1,p)} + e_{(p,m+1)}] \right\} = -\frac{2}{d} \sum_{p,p'=1; p \neq p'}^{d} e_{(p-1)d+p'} e$$

Therefore, we obtain

$$\sum_{m=1}^{d-1} H_m \otimes H_m = 2 \sum_{p=1}^d e_{(p-1)d+p} - \frac{2}{d} \sum_{p,p'=1}^d e_{(p-1)d+p'} = 2 \sum_{p=1}^d e_{(p-1)d+p} - \frac{2}{d} I.$$

References

- [1] M. A. Jafarizadeh and R. Sufiani, (2008), Phys. Rev. A 77, 022315.
- [2] D. Bouwmeester, A. Ekert, and A. Zeilinger, (2000), The Physics of Quantum Information (Springer-Verlag, Berlin).

- [3] S. Bose, (2003), Phys. Rev. Lett. 91, 207901.
- [4] V. Subrahmanyam, (2003), arXiv: quant-ph/0307135.
- [5] T. Shi, Ying Li, Z. Song, and C. Sun, (2004), arXiv: quant-ph/ 0408152.
- [6] T. J. Osborne and N. Linden, (2004), Phys. Rev. A 69, 052315.
- [7] M. Christandl, N. Datta, A. Ekert, and A. J. Landahl, (2004), Phys. Rev. Lett. 92, 187902.
- [8] M. Christandl, N. Datta, T. C. Dorlas, A. Ekert, A. Kay and A. J. Landahl, (2005), Phys. Rev. A 71, 032312.
- [9] D. Burgarth and S. Bose, (2005), Phys. Rev. A 71, 052315.
- [10] D. Burgarth and S. Bose, (2005), New J. Phys. 7, 135.
- [11] M. H. Yung and S. Bose, (2005), Phys. Rev. A 71, 032310.
- [12] M. H. Yung, (2006), Phys. Rev. A 74, 030303.
- [13] I. Affleck and J.B. Marston, Phys. Rev. B 37, 11538 (1988).
- [14] J.B. Marston and I. Affleck, Phys. Rev. B 39, 11538 (1989).
- [15] C. Honerkamp and W. Hofstetter, Phys. Rev. Lett. 92, 170403 (2004).
- [16] A. Bayat and V. Karimipour, arXiv: quant-ph/0612144.
- [17] N. Obata, (2004), Quantum Probabilistic Approach to Spectral Analysis of Star Graphs, Interdisciplinary Information Sciences, Vol. 10, 41-52.
- [18] A. Hora, and N. Obata, (2003), Fundamental Problems in Quantum Physics, World Scientific, 284.
- [19] M. A. Jafarizadeh and S. Salimi, (2006), J. Phys. A : Math. Gen. 39, 1-29.

- [20] M. A. Jafarizadeh, S. Salimi, (2007), Annals of physics, Vol. 322 1005-1033.
- [21] M. A. Jafarizadeh, R. Sufiani, (2007), Physica A, 381, 116-142.
- [22] M. A. Jafarizadeh and R. Sufiani, (2007), International Journal of Quantum Information Vol. 5, No. 4, 575-596.
- [23] M. A. Jafarizadeh, R. Sufiani and S. Jafarizadeh, (2007), J. Phys. A: Math. Theor. 40, 4949-4972.
- [24] M. A. Jafarizadeh, R. Sufiani, S. Salimi and S. Jafarizadeh, Eur. Phys. J. B 59, 199-216.
- [25] J. A. Shohat, and J. D. Tamarkin, (1943), The Problem of Moments, American Mathematical Society, Providence, RI.
- [26] T. S. Chihara, (1978), An Introduction to Orthogonal Polynomials, Gordon and Breach, Science Publishers Inc.
- [27] A. Hora, and N. Obata, (2002), Quantum Information V, World Scientific, Singapore.
- [28] A. Childs, E. Farhi, and S. Gutmann, (2002), Quantum Inf. Process. 1, 35.
- [29] M. Puschel and J. M. F. Moura, (2003), SIAM Journal of Computing, vol. 32, no. 5, pp. 12801316.
- [30] M. A. Jafarizadeh, R. Sufiani and S. Jafarizadeh, arXiv: cond-mat/07072570.
- [31] M. A. Jafarizadeh, Physica A 287 (2000) 1-25.
- [32] M. A. Jafarizadeha, Europian Physical journal B, vol. 4 (1998) 103.

Figure Captions

Figure-1: Shows the network G_2 .

Figure-2: shows the modified G_2 -type network.

Figure-3: Shows the pseudo-distance-regular network derived from Icosahedron network.

Figure-4: Shows the 3- simplex fractal with decimation number b = 2.

Figure-5: Shows the pseudo-distance-regular network derived from Hadamard network with 16 nodes.









