CHARACTERS OF PRIME DEGREE

EDITH ADAN-BANTE

ABSTRACT. Let G be a finite nilpotent group, χ and ψ be irreducible complex characters of G of prime degree. Assume that $\chi(1)=p$. Then either the product $\chi\psi$ is a multiple of an irreducible character or $\chi\psi$ is the linear combination of at least $\frac{p+1}{2}$ distinct irreducible characters.

1. Introduction

Let G be a finite group and $\chi, \psi \in \operatorname{Irr}(G)$ be irreducible complex characters of G. We can check that the product of $\chi\psi$, where $\chi\psi(g)=\chi(g)\psi(g)$ for all g in G, is a character and so it can be expressed as a linear combination of irreducible characters. Let $\eta(\chi\psi)$ be the number of distinct irreducible constituents of the product $\chi\psi$.

Fix a prime number p. Assume that G is a nilpotent group and $\chi, \psi \in Irr(G)$ are characters of degree p. In this note, we study the product $\chi\psi$.

Theorem A. Let G be a finite nilpotent group, χ and ψ be irreducible complex characters of prime degree. Assume that $\chi(1) = p$. Then one of the following holds:

- (i) $\chi\psi$ is the sum of p^2 distinct linear characters.
- (ii) $\chi \psi$ is the sum of p distinct linear characters of G and of p-1 distinct irreducible characters of G with degree p.
- (iii) all the irreducible constituents of $\chi\psi$ are of degree p. Also, either $\chi\psi$ is a multiple of an irreducible character, or it has at least $\frac{p+1}{2}$ distinct irreducible constituents, and at most p distinct irreducible constituents, i.e.

either
$$\eta(\chi\psi) = 1$$
 or $\frac{p+1}{2} \le \eta(\chi\psi) \le p$.

(iv) $\chi\psi$ is an irreducible character.

It is proved in Theorem A of [1] that given any prime p, any p-group P, any faithful characters $\chi, \psi \in \operatorname{Irr}(P)$, either the product $\chi\psi$ is a multiple of an irreducible, or $\chi\psi$ is the linear combination of at least $\frac{p+1}{2}$ distinct irreducible characters, i.e. either $\eta(\chi\psi)=1$ or $\eta(\chi\psi)\geq \frac{p+1}{2}$. It is proved in [4] that given any prime p and any integer n>0, there exists a p-group P and characters $\varphi, \gamma \in \operatorname{Irr}(P)$ such that $\eta(\varphi\gamma)=n$. Thus without the hypothesis that the characters in Theorem A of [1] are faithful, the result may not hold. In this note, we are proving that if the characters have "small" degree then the values that $\eta(\chi\psi)$ can take have the same constraint as if they were faithful.

Date: 2008.

¹⁹⁹¹ Mathematics Subject Classification. 20c15.

 $Key\ words\ and\ phrases.$ Characters of prime degree, nilpotent groups, p-groups, complex characters, products of characters.

2. Proofs

We are going to use the notation of [5]. In addition, we denote by $\operatorname{Lin}(G) = \{\chi \in \operatorname{Irr}(G) \mid \chi(1) = 1\}$ the set of linear characters, and by $\operatorname{Irr}(G \bmod N) = \{\chi \in \operatorname{Irr}(G) \mid \operatorname{Ker}(\chi) \geq N\}$ the set of irreducible characters of G that contain in their kernel the subgroup N. Also, denote by $\overline{\chi}$ the complex conjugate of χ , i.e. $\overline{\chi}(g) = \overline{\chi(g)}$ for all g in G.

Lemma 2.1. Let G be a finite group and $\chi, \psi \in \text{Irr}(G)$. Let $\alpha_1, \alpha_2, \ldots, \alpha_n$, for some n > 0, be the distinct irreducible constituents of the product $\chi \psi$ and a_1, a_2, \ldots, a_n be the unique positive integers such that

$$\chi\psi = \sum_{i=1}^{n} a_i \alpha_i.$$

If $\alpha_1(1) = 1$, then $\psi \overline{\alpha_1} = \overline{\chi}$. Hence the distinct irreducible constituents of the character $\chi \overline{\chi}$ are 1_G , $\overline{\alpha_1}\alpha_2$, $\overline{\alpha_1}\alpha_2$, ..., $\overline{\alpha_1}\alpha_n$, and

$$\chi \overline{\chi} = a_1 1_G + \sum_{i=2}^n a_i (\overline{\alpha_1} \alpha_i).$$

Proof. See Lemma 4.1 of [3].

Lemma 2.2. Let G be a finite p-group for some prime p and $\chi \in Irr(G)$ be a character of degree p. Then one of the following holds:

- (i) $\chi \overline{\chi}$ is the sum of p^2 distinct linear characters.
- (ii) $\chi \overline{\chi}$ is the sum of p distinct linear characters of G and of p-1 distinct irreducible characters of G with degree p.

Proof. See Lemma 5.1 of [2].
$$\Box$$

Lemma 2.3. Let G be a finite p-group, for some prime p, and $\chi, \psi \in \text{Irr}(G)$ be characters of degree p. Then either $\eta(\chi\psi) = 1$ or $\eta(\chi\psi) \geq \frac{p+1}{2}$.

Proof. Assume that the lemma is false. Let G and $\chi, \psi \in \operatorname{Irr}(G)$ be a counterexample of the statement, i.e. $\chi(1) = \psi(1) = p$ and $1 < \eta(\chi\psi) < \frac{p+1}{2}$.

Working with the group $G/(\text{Ker}(\chi) \cap \text{Ker}(\psi))$, by induction on the order of G, we may assume that $\text{Ker}(\chi) \cap \text{Ker}(\psi) = \{1\}$. Set $n = \eta(\chi\psi)$. Let $\theta_i \in \text{Irr}(G)$, for i = 1, ..., n, be the distinct irreducible constituents of $\chi\psi$. Set

(2.4)
$$\chi \psi = \sum_{i=1}^{n} m_i \theta_i$$

where $m_i > 0$ is the multiplicity of θ_i in $\chi \psi$.

If $\chi\psi$ has a linear constituent, then by Lemma 2.1 and Lemma 2.2 we have that $\eta(\chi\psi) \geq p$. If $\chi\psi$ has an irreducible constituent of degree p^2 , then $\chi\psi \in \operatorname{Irr}(G)$ and so $\eta(\chi\psi) = 1$. Thus we may assume that $\theta_i(1) = p$ for $i = 1, \ldots, n$.

Since G is a p-group, there must exist a subgroup H and a linear character ξ of H such that $\xi^G = \chi$. Then $|G:H| = \chi(1) = p$ and thus H is a normal subgroup. By Clifford theory we have then

$$\chi_H = \sum_{i=1}^p \xi_i$$

for some $\xi_1 = \xi, \dots, \xi_p$ distinct linear characters of H.

Claim 2.6. H is an abelian group.

Proof. Suppose that $\psi_H \in \operatorname{Irr}(H)$. Since $(\xi \psi_H)^G = \chi \psi$ by Exercise 5.3 of [5], and $\xi \psi_H \in \operatorname{Irr}(H)$, it follows that either $\xi \psi_H$ induces irreducibly, and thus $\eta(\chi \psi) = 1$, or $\xi \psi_H$ extends to G and thus $(\xi \psi_H)^G$ is the sum of the p distinct extensions of $\xi \psi_H$, i.e. $\eta(\chi \psi) = p$. Therefore $\psi_H \notin \operatorname{Irr}(G)$ and since H is normal in G of index p and $\psi(1) = p$, ψ is induced from some $\tau \in \operatorname{Lin}(H)$.

Since H is normal of index p, both ξ and τ are linear characters and thus $\operatorname{Ker}(\xi) \cap \operatorname{Ker}(\tau) \geq [H, H]$. Observe that $\operatorname{core}_G(\operatorname{Ker}(\xi) \cap \operatorname{Ker}(\tau)) = \operatorname{core}_G(\operatorname{Ker}(\xi)) \cap \operatorname{core}_G(\operatorname{Ker}(\tau)) = \operatorname{Ker}(\chi) \cap \operatorname{Ker}(\psi)$. Since H is a normal subgroup of G, so is [H, H] and thus $\{1\} = \operatorname{Ker}(\chi) \cap \operatorname{Ker}(\psi) \geq [H, H]$. Therefore H is abelian.

By the previous claim, observe that also ψ is induced by some linear character τ of H and thus

$$\psi_H = \sum_{i=1}^p \tau_i$$

for some $\tau_1 = \tau, \dots, \tau_p$ distinct linear characters of H. Observe also that the center of both χ and ψ is contained in H.

Claim 2.8.
$$Z(G) = Z(\chi) = Z(\psi)$$
.

Proof. Suppose that $\mathbf{Z}(\chi) \neq \mathbf{Z}(\psi)$. Set $U = \mathbf{Z}(\chi) \cap \mathbf{Z}(\psi)$. Either U is properly contained in $\mathbf{Z}(\chi)$, or it is properly contained in $\mathbf{Z}(\psi)$. We may assume that $U < \mathbf{Z}(\psi)$ and thus we may find a subgroup $T \leq \mathbf{Z}(\psi)$ such that T/U is chief factor of G. Since H is abelian, $\mathbf{Z}(\psi) < H$ and $\tau^G = \psi$, then $\psi_T = p\tau_T$ and so $(\tau_i)_T = \tau_T$ for $i = 1, \ldots, p$. Because $\xi^G = \chi$, $\xi \in \text{Lin}(H)$ and $T \not\leq \mathbf{Z}(\chi)$, the stabilizer of ξ_T is H. Thus the stabilizer of $\xi_T\tau_T$ in G is H. By Clifford theory we have that $\xi\tau_i \in \text{Lin}(H)$ induces irreducibly and $\xi\tau_i$ are distinct characters for $i = 1, \ldots, p$. By (2.7) we have that $\chi\psi = (\xi\psi_H)^G = (\xi(\tau_1 + \cdots + \tau_p))^G = (\xi\tau_1)^G + \cdots + (\xi\tau_p)^G$, and thus $\eta(\chi\psi) = p$. We conclude that such T can not exist and so $\mathbf{Z}(\chi) = \mathbf{Z}(\psi)$.

Given any $z \in Z(\chi)$ and $g \in G$, we have $z^g \cong z \pmod{\ker(\chi)}$ since $Z(G/\ker(\chi)) = Z(\chi)/\ker(\chi)$. Hence $[z,g] = z^{-1}z^g$ lies in $\ker(\chi)$. This same z lies in $Z(\psi) = Z(\chi)$. Hence [z,g] also lies in $\ker(\psi)$. Therefore $[z,g] \in \ker(\chi) \cap \ker(\psi) = 1$ for every $z \in Z(\chi) = Z(\psi)$ and every $g \in G$. This implies that $Z(\chi) = Z(\psi) = Z(G)$.

Set $Z = \mathbf{Z}(G)$. Since Z is the center of G, $\xi^G = \chi$ and $\tau^G = \psi$, we have

(2.9)
$$\chi_Z = p\xi_Z \text{ and } \psi_Z = p\tau_Z.$$

Because $\chi_Z \psi_Z = p^2 \xi_Z \tau_Z$, (2.4) implies that

$$(2.10) (\theta_i)_Z = p\xi_Z \tau_Z$$

for all $i = 1, \ldots, n$.

Let Y/Z be a chief factor of G with $Y \leq H$. Since Z is the center of G and $Z = \mathbf{Z}(\chi)$, the set $\text{Lin}(Y \mid \xi_Z)$ of all extensions of ξ_Z to linear characters is $\{(\xi_1)_Y = \xi_Y, (\xi_2)_Y, \dots, (\xi_p)_Y\}$ and it is a single G-conjugacy class. By Clifford theory we have that

(2.11)
$$\chi_Y = \sum_{i=1}^p (\xi_i)_Y.$$

Since H is the stabilizer of τ_Y in G and $\psi(1) = p$, as before we have that the set $\text{Lin}(Y \mid \tau_z) = \{(\tau_1)_Y = \tau_Y, (\tau_2)_Y, \dots, (\tau_p)_Y\}$ and

(2.12)
$$\psi_Y = \sum_{i=1}^p (\tau_i)_Y.$$

Claim 2.13. The stabilizer $\{g \in G \mid (\xi_Y \tau_Y)^g = \xi_Y \tau_Y \}$ of $\xi_Y \tau_Y \in \text{Lin}(Y)$ in G is H.

Proof. Assume notation (2.4). Suppose $\xi_Y \tau_Y$ is a G-invariant character. Since |Y:Z|=p and $\xi_Y \tau_Y$ is an extension of $\xi_Z \tau_Z$, it follows then that all the extensions of $\xi_Z \tau_Z$ to Y are G-invariant. Thus by (2.4) and (2.10), given any i, there exists some extension $v_i \in \text{Lin}(Y)$ of $\xi_Z \tau_Z$ such that $(\theta_i)_Y = p v_i$. Thus $(\chi \psi)_Y = (\sum_{i=1}^n m_i \theta_i)_Y = \sum_{i=1}^n m_i (v_i)_Y = \sum_{i=1}^n m_i p v_i$ has at most $n < \frac{p+1}{2}$ distinct irreducible constituents. On the other hand, by (2.11) and (2.12) we have

$$(\chi \psi)_Y = \chi_Y \psi_Y = (\sum_{i=1}^p (\xi_i)_Y)(\sum_{j=1}^p (\tau_j)_Y) = p \sum_{j=1}^p \xi_Y(\tau_j)_Y,$$

and so $(\chi \psi)_Y$ has p distinct irreducible constituents. That is a contradiction and thus $G_{\xi_Y \tau_Y} = H$.

By Clifford theory and the previous claim, we have that for each i = 1, ..., n, there exists a unique character $\sigma_i \in \text{Lin}(H \mid \xi_Y \tau_Y)$ such that

$$\theta_i = (\sigma_i)^G.$$

If Y = H then $|G: Z| = |G: H||H: Z| = p^2$. Since $\chi(1) = \psi(1) = p$, by Corollary 2.30 of [5] we have that χ and ψ vanish outside Z. Since $\theta_i(1) = p$ for all i and $|G: Z| = |G: \mathbf{Z}(\theta_i)| = p^2$, it follows that there exists a unique irreducible character lying above $\xi_Z \tau_Z$ and thus $\eta(\chi \psi) = 1$.

2.15. Fix a subgroup $X \leq H$ of G such that X/Y is a chief factor of G. Let α , $\beta \in \text{Lin}(X)$ be the linear characters such that

$$\xi_X = \alpha \text{ and } \tau_X = \beta.$$

Since σ_i lies above $\xi_Y \tau_Y \in \text{Lin}(Y)$ for all i and X/Y is a chief factor of a p-group, there is some $\delta_i \in \text{Irr}(X \mod Y)$ such that

$$(2.16) (\sigma_i)_X = \delta_i \alpha \beta.$$

Claim 2.17. The subgroup [X,G] generates Y = [X,G]Z modulo Z.

Proof. Working with the group $\bar{G} = G/\operatorname{Ker}(\chi)$, using the same argument as in the proof of Claim 3.26 of [1], we have that $[\bar{X}, \bar{G}]$ generates $\bar{Y} = [\bar{X}, \bar{G}]\bar{Z}$ modulo \bar{Z} . Since $Z = \mathbf{Z}(\chi)$, we have that $\operatorname{Ker}(\chi) \leq Z$. Thus $\bar{Z} = Z/\operatorname{Ker}(\chi)$ and the claim follows.

2.18. Observe that G/H is cyclic of order p. So we may choose $g \in G$ such that the distinct cosets of H in G are H, Hg, Hg^2 , ..., Hg^{p-1} .

Since $\chi = \xi^G$ and $\xi_X = \alpha$, it follows from 2.15 that

$$\chi_X = \alpha + \alpha^g + \dots + \alpha^{g^{p-1}} = \sum_{i=0}^{p-1} \alpha^{g^i}.$$

Similarly, we have that

$$\psi_X = \beta + \beta^g + \dots + \beta^{g^{p-1}} = \sum_{j=0}^{p-1} \beta^{g^j}.$$

Combining the two previous equations we have that

(2.19)
$$\chi_X \psi_X = (\sum_{j=0}^{p-1} \alpha^{g^j})) (\sum_{j=0}^{p-1} \beta^{g^j}) = \sum_{i=0}^{p-1} \sum_{j=0}^{p-1} \alpha^{g^i} \beta^{g^j}.$$

By (2.4) and (2.16), we have that

(2.20)
$$(\chi \psi)_X = (\sum_{i=1}^n m_i \theta_i)_X = \sum_{i=1}^n m_i [\sum_{j=0}^{p-1} (\delta_i \alpha \beta)^{g^j}].$$

Claim 2.21. Let $g \in G$ be as in 2.18. For each i = 0, 1, ..., p - 1, there exist $j \in \{0, 1, ..., p - 1\}$ and $\delta_{g^i} \in \text{Lin}(X \mod Y)$ such that

(2.22)
$$\alpha \beta^{g^i} = (\alpha \beta)^{g^j} \delta_{g^i}.$$

Also $|\{\delta_{q^i} \mid i = 0, 1, 2, \dots, p - 1\}| \le n$.

Proof. See Proof of Claim 3.30 of [1].

Claim 2.23. Let $g \in G$ be as in 2.18. Then there exist three distinct integers $i, j, k \in \{0, 1, 2, ..., p-1\}$, and some $\delta \in \operatorname{Irr}(X \operatorname{mod} Y)$, such that

$$\alpha \beta^{g^i} = (\alpha \beta)^{g^r} \delta,$$

$$\alpha \beta^{g^j} = (\alpha \beta)^{g^s} \delta,$$

$$\alpha \beta^{g^k} = (\alpha \beta)^{g^t} \delta,$$

for some $r, s, t \in \{0, 1, 2, \dots, p-1\}$.

Proof. See Proof of Claim 3.34 of [1].

Claim 2.24. We can choose the element g in 2.18 such that one of the following holds:

(i) There exists some j = 2, ..., p-1 such that

$$\alpha \beta^g = (\alpha \beta)^{g^r},$$

$$\alpha \beta^{g^j} = (\alpha \beta)^{g^s},$$

for some $r, s \in \{0, 1, ..., p-1\}$ with $r \neq 1$.

(ii) There exist j and k such that 1 < j < k < p, and

$$\begin{array}{rcl} \alpha\beta^g & = & (\alpha\beta)^{g^r}\delta, \\ \alpha\beta^{g^j} & = & (\alpha\beta)^{g^s}\delta, \\ \alpha\beta^{g^k} & = & (\alpha\beta)^{g^t}\delta, \end{array}$$

for some $\delta \in \operatorname{Irr}(X \operatorname{mod} Y)$ and some $r, s, t \in \{0, 1, \dots, p-1\}$ with $r \neq 1$.

Proof. See Proof of Claim 3.35 of [1].

Let g be as in Claim 2.24. Since X/Y is cyclic of order p, we may choose $x \in X$ such that X = Y < x >. Since H is abelian, we have [X, H] = 1. Suppose that $[x, g^{-1}] \in Z$. Then x centralizes both g^{-1} and H modulo Z. Hence $xZ \in \mathbf{Z}(G/Z)$ and so $[x, G] \leq Z$. Since Y/Z is a chief section of the p-group G, we have that $[Y, G] \leq Z$ and so $[< x > Y, G] = [X, G] \leq Z$ which is false by Claim 2.17. Hence $[x, g^{-1}] \in Y \setminus Z$ and so

(2.25)
$$Y = Z < y > \text{ is generated over } Z \text{ by } y = [x, g^{-1}].$$

Since $[Y,G] \leq Z$ we have that $z=[y,g^{-1}] \in Z$. If z=1, then G=H < g > centralizes Y=Z < y >, since H centralizes Y < X by 2.15, and G centralizes Z. This is impossible because $Z=\mathbf{Z}(G) < Y$. Thus

(2.26)
$$z = [y, g^{-1}]$$
 is a non-trivial element of Z .

By (2.25) we have $y=[x,g^{-1}]=x^{-1}x^{g^{-1}}$. By (2.26) we have $z=[y,g^{-1}]=y^{-1}y^{g^{-1}}$. Finally $z^{g^{-1}}=z$ since $z\in Z$. Since $X=Z< x,y>\leq H$ is abelian, it follows that

(2.27)
$$z^{g^{-j}} = z, \ y^{g^{-j}} = yz^j \text{ and } x^{g^{-j}} = xy^j z^{\binom{j}{2}},$$

for any integer $j=0,1,\ldots,p-1$. Because $g^{-p}\in H$ centralizes X by 2.15, we have $z^p=1$ and $y^pz^{\binom{p}{2}}=1$.

Observe that the statement is true for $p \leq 3$ since then $\frac{p+1}{2} \leq 2$. Thus we may assume that p is odd. Hence p divides $\binom{p}{2} = \frac{p(p-1)}{2}$ and $z^{\binom{p}{2}} = 1$. Therefore

$$(2.28) y^p = z^p = 1.$$

It follows that y^i , z^i and $z^{\binom{i}{2}}$ depend only on the residue of i modulo p, for any integer $i \geq 0$.

2.29. Observe that $\operatorname{Ker}(\xi_Z) \cap \operatorname{Ker}(\tau_Z) \leq \operatorname{Ker}(\chi) \cap \operatorname{Ker}(\psi) = 1$ implies that z is not in both $\operatorname{Ker}(\xi_Z)$ and $\operatorname{Ker}(\tau_Z)$. Without loss of generality we may assume that $\tau_Z(z) \neq 1$. Since β is an extension of τ_Z , we may assume that $\beta(z) \neq 1$.

Claim 2.30. $\xi_Z \tau_Z(z)$ is primitive p-th root of unit.

Proof. Suppose that $(\xi_Z \tau_Z)(z) = 1$. Then $(\xi_Z \tau_Z)([y, g^{-1}]) = 1$ and so $(\xi_Z \tau_Z)^g(y) = (\xi_Z \tau_Z)(y)$. Since H is abelian, |G:H| = p, θ_i lies above $\xi_Z \tau_Z$ for all i, and $g \in G \setminus H$, it follows that $Y = \langle y, \mathbf{Z}(G) \rangle$ is contained in $\mathbf{Z}(\theta_i)$. That is contradiction with Claim 2.13. Thus $(\xi_Z \tau_Z)(z) \neq 1$. Since z is of order p and $\xi_Z \tau_Z$ is a linear character, the claim follows.

Claim 2.31. Suppose that

(2.32)
$$\alpha \beta^g = (\alpha \beta)^{g^r} \delta,$$

and

(2.33)
$$\alpha \beta^{g^j} = (\alpha \beta)^{g^s} \delta,$$

for some $j \in \{0, 1, ..., p-1\}$, $j \neq 1$, some $\delta \in Irr(X \mod Y)$, and some $r, s \in \{0, 1, ..., p-1\}$. Then

(2.34)
$$\delta(x) = \beta(z)^{hj(r-1)},$$

where $2h \equiv 1 \mod p$.

Proof. By Claim 2.30 and the same argument as in the proof of Claim 3.40 of [1], the statement follow. \Box

Suppose that Claim 2.24 (ii) holds. Then by Claim 2.31 we have that $\delta(x) = \beta(z)^{hj(r-1)}$ and $\delta(x) = \beta(z)^{hk(r-1)}$. Since $\beta(z) = \tau_Z(z)$ is a primitive p-th root of unit by 2.29, we have that $hj(r-1) \equiv hk(r-1) \mod p$. Since $r \not\equiv 1 \mod p$ and $2h \equiv 1 \mod p$, we have that $k \equiv j \mod p$, a contradiction. Thus Claim 2.24 (i) must hold.

We apply now Claim 2.31 with $\delta = 1$. Thus $1 = \delta(x) = \beta(z)^{hj(r-1)}$. Therefore $hj(r-1) \equiv 0 \mod p$. Since $2h \equiv 1 \mod p$, either $j \equiv 0 \mod p$ or $r-1 \equiv 0 \mod p$. Neither is possible. That is our final contradiction and Lemma 2.3 is proved. \square

Proof of Theorem A. Since G is a nilpotent group, G is the direct product $G_1 \times G_2$ of its Sylow p-subgroup G_1 and its Hall p'-subgroup G_2 . We can write then $\chi = \chi_1 \times \chi_2$ and $\psi = \psi_1 \times \psi_2$ for some characters $\chi_1, \psi_1 \in \operatorname{Irr}(G_1)$ and some characters $\chi_2, \psi_2 \in \operatorname{Irr}(G_2)$. Since $\chi(1) = p$, we have that $\chi_2(1) = 1$ and thus $\chi_2\psi_2 \in \operatorname{Irr}(G_2)$. If $\psi(1) \neq p$, since $\psi(1)$ is a prime number, we have that $\psi_1(1) = 1$ and thus $\chi_1\psi_1$ is an irreducible. Therefore $\chi\psi \in \operatorname{Irr}(G)$ and (iv) holds. We may assume then that $\psi(1) = p$ and thus $\psi_2(1) = 1$. Then $\chi_2\psi_2$ is a linear character and so we may assume that G is a p-group.

If $\chi\psi$ has a linear constituent, by Lemma 2.1 and Lemma 2.2 we have that (i) or (ii) holds. So we may assume that all the irreducible constituents of $\chi\psi$ are of degree at least p. If $\chi\psi$ has an irreducible constituent of degree p^2 , then $\chi\psi\in {\rm Irr}(G)$ and (iv) holds. We may assume then that all the irreducible constituents of $\chi\psi$ have degree p. Since $\chi\psi(1)=p^2$, it follows that $\eta(\chi\psi)\leq p$. By Lemma 2.3 we have that either $\eta(\chi\psi)=1$ or $\eta(\chi\psi)\geq \frac{p+1}{2}$, and so (iii) holds.

Examples. Fix a prime p > 2

- (i) Let E be an extraspecial group of order p^3 and $\phi \in Irr(E)$ of degree p. We can check that the product $\phi \overline{\phi}$ is the sum of all the linear characters of E.
- (ii) In the proof of Proposition 6.1 of [2], an example is constructed of a p-group G and a character $\chi \in Irr(G)$ such that $\chi \overline{\chi}$ is the sum of p distinct linear characters and of p-1 distinct irreducible characters of degree p.
- (iii) Given an extraspecial group E of order p^3 , where p > 2, and $\phi \in \operatorname{Irr}(E)$ a character of degree p, we can check that $\phi\phi$ is a multiple of an irreducible. In Proposition 6.1 of [1], an example is provided of a p-group G and a character $\chi \in \operatorname{Irr}(G)$ such that $\eta(\chi\chi) = \frac{p+1}{2}$. In [6], an example is provided of a p-group P and two faithful characters $\delta, \epsilon \in \operatorname{Irr}(P)$ of degree p such that $\eta(\delta\epsilon) = p 1$.

Let G be the wreath product of a group of order p with itself. Thus G has a normal abelian subgroup of N of order p^p and index p. Let $\lambda \in \text{Lin}(N)$. We can check that $\chi = \lambda^G$ and $\psi = (\lambda^2)^G$ are irreducible characters of degree p and $\chi \psi$ is the sum of p distinct irreducible characters of degree p.

We wonder if there exists a p-group P with characters $\chi, \psi \in \operatorname{Irr}(P)$ of degree p such that $\frac{p+1}{2} < \eta(\chi\psi) < p-1$.

(iv) Let Q be a p-group and $\kappa \in Irr(Q)$ be a character of degree p. Set $P = Q \times Q$, $\chi = \kappa \times 1_G$ and $\psi = 1_G \times \kappa$. Observe that χ, ψ and $\chi \psi$ are irreducible characters of P.

References

1. E. Adan-Bante, Products of characters and finite p-groups, J. of Algebra, 277 (2004) 236-255.

- E. Adan-Bante, Products of characters and finite p-groups II, Arch. Math. 82 No 4 (2004), 289-297.
- E. Adan-Bante, Products of characters and derived length II, J. Group Theory 8 (2005), 453-459
- 4. E. Adan-Bante, Products of characters and restriction of characters, preprint.
- 5. I.M.Isaacs, Character Theory of Finite Groups. New York-San Francisco–London: Academic Press 1976
- 6. M. Loukaki, A. Moreto, On the number of constituents of products of characters , Algebra Colloq. 14, no.2, 207-208,(2007).

DEPARTMENT OF MATHEMATICAL SCIENCE, NORTHERN ILLINOIS UNIVERSITY, WATSON HALL 320 DEKALB, IL 60115-2888, USA

 $E\text{-}mail\ address\colon \texttt{EdithAdan@illinoisalumni.org}$