

Resonant interactions of nonlinear water waves in a finite basin

Elena Kartashova[‡], Sergey Nazarenko, Oleksii Rudenko[†]

[†]Weizmann Institute of Science, Rehovot, Israel

^{*}Mathematics Institute, University of Warwick, Coventry CV4-7AL, UK

[‡]RISC, J. Kepler University, Linz, Austria

e-mail: lena@risc.uni-linz.ac.at

We study exact four-wave resonances among gravity water waves in a square box with periodic boundary conditions. We show that these resonant quartets are linked with each other by shared Fourier modes in such a way that they form independent clusters. These clusters can be formed by two types of quartets: (1) *angle-resonances* which cannot directly cascade energy but which can redistribute it among the initially excited modes and (2) *scale-resonances* which are much more rare but which are the only ones that can transfer energy between different scales. We find such resonant quartets and their clusters numerically on the set of 1000 x 1000 modes, classify and quantify them and discuss consequences of the obtained cluster structure for the wavefield evolution. Finite box effects and associated resonant interaction among discrete wave modes appear to be important in most numerical and laboratory experiments on the deep water gravity waves, and our work is aimed at aiding the interpretation of the experimental and numerical data.

Contents

Introduction	1
Construction of q-classes	2
Scale-resonances	3
Collinear quartets	3
Non-collinear quartets	4
Tridents	4
Non-tridents	4
Clusters	5
Angle-resonances	6
Dynamics	7
The wave field	7
A quartet	7
Summary and Discussion of Results	8
Acknowledgments	8
References	9

INTRODUCTION

Weakly nonlinear systems of random waves are usually studied in the framework of wave turbulence theory (WTT) (for introduction to WTT see book [1]). Besides weak nonlinearity and phase randomness, this statistical description is based on the infinite-box limit. This approach yields wave kinetic equations for the wave spectrum which have important stationary solutions, Kolmogorov-Zakharov (KZ) spectra [1]. Importance of KZ spectra is in that they correspond to a constant flux

of energy in Fourier space and, therefore, they are analogous to the Kolmogorov energy-cascade spectrum in hydrodynamic turbulence. WTT approach can also study evolution of higher momenta of wave amplitudes and even their probability density function, and, therefore, examine conditions for deviation from Gaussianity and onset of intermittency [2, 3, 4, 5]. Significant effort has been done in the past to test WTT and its predictions numerically [3, 6, 7, 8, 9] as well as experimentally [10, 11, 12, 13]. It has been noted however that in both the numerical and the laboratory experiments the domain boundaries typically play a very important role and the infinite-box limit assumed by WTT is not achieved. It is important to understand that WTT takes the infinite-box limit before the weak-nonlinearity limit, which physically means that a lot of modes interact simultaneously if they are in *quasi-resonance*, i.e. satisfy the following conditions

$$\begin{cases} |\omega(\mathbf{k}_1) \pm \omega(\mathbf{k}_2) \pm \dots \pm \omega(\mathbf{k}_s)| < \Omega, \\ \mathbf{k}_1 \pm \mathbf{k}_2 \pm \dots \pm \mathbf{k}_s = 0 \end{cases} \quad (1)$$

with some resonance broadening $\Omega > 0$ which is a monotonically increasing function of nonlinearity (mean wave amplitude). Here \mathbf{k} and $\omega(\mathbf{k})$ are wave vector and dispersion function correspondingly and a wave is regarded in a general form $\exp i(\mathbf{k}\mathbf{x} - \omega t)$. WTT is supposed to work when the resonance broadening Ω is greater than the spacing δ_ω between the adjacent wave modes

$$\Omega > (\partial\omega/\partial k)2\pi/L \quad (2)$$

where L is the box size. For some types of waves, for example for the capillary water waves, this condition is easy to satisfy, and therefore to achieve WTT regime [14]. However, this condition is often violated for some other types of waves, in particular for the surface gravity waves which will be the main object of this Letter. This

occurs in numerical simulations due to limitations on the numerical resolution [7, 9] and in laboratory experiments - due to an insufficient basin size [10, 11]. In this case, the Fourier space discreteness (which is due to a finite box size) leads to significant depletion of the number of wave resonances with respect to the infinite box limit. In turn, this results in a slowdown of the energy cascade through the k -space with respective steepening of the wave spectra [10, 15]. In addition, the wavenumber grid will cause the wave spectra to be anisotropic in this case.

What happens when the condition (2) is so badly violated that only waves which are in exact resonance (i.e. $\Omega = 0$) can interact? In this case, the mechanism of the wave phase randomization based on many quasi-resonant waves interacting simultaneously will be absent and, therefore, one should expect less random and more coherent behavior. In [16] it was shown that in many wave systems, resonantly interacting waves in finite domains are partitioned into small independent clusters in Fourier space, such that there cannot be an energy flux between different clusters. In particular, in [17] some examples of wave systems were given in which no resonances exist (capillary water waves, $\omega = |\mathbf{k}|^{3/2}$), infinite number of resonances exists (oceanic planetary waves, $\omega = |\mathbf{k}|^{-1}$). Both of these examples are three-wave systems [i.e. $s = 3$ in (1)]. In the present paper we will concentrate on finding resonances for the deep water gravity waves, which is a four-wave system.

The problem of computing exact resonances in a confined laboratory experiment is highly non-trivial because the wavenumbers are integers and (1) is a system of Diophantine equations on many integer variables in large powers. Computational time of solving this system by a simple enumeration of possibilities in this case grows exponentially with each variable and the size of spectral domain under consideration. A specially developed q -class method [18, 19, 20] has allowed to accelerate the computation and find all the resonances among waves in large spectral domains, in a matter of minutes. In this Letter we use q -class method to construct resonant wave clusters formed by 4-wave resonances among water gravity waves covered by the kinematic resonance conditions in the form:

$$\begin{cases} |\mathbf{k}_1|^{1/2} + |\mathbf{k}_2|^{1/2} = |\mathbf{k}_3|^{1/2} + |\mathbf{k}_4|^{1/2}, \\ \mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4 \end{cases} \quad (3)$$

with $\mathbf{k}_i = (m_i, n_i)$ and integer $|m_i|, |n_i| \leq 1000$. Our main aim is to understand how anisotropic resonance clusters influence the general dynamics of the complete wave field.

CONSTRUCTION OF q -CLASSES

We adopt the general definition of a q -class given in [18] for the dispersion function $\omega = |\mathbf{k}|^{1/2}$ in following way. Consider the set of algebraic numbers $R = \pm k^{1/4}$. Any such number k has a unique representation

$$k = \gamma q^{1/4}, \gamma \in \mathbb{Z}$$

where q is a product

$$q = p_1^{e_1} p_2^{e_2} \dots p_n^{e_n},$$

while p_1, \dots, p_n are all different primes and the powers $e_1, \dots, e_n \in \mathbb{N}$ are all smaller than 4. Then the set of numbers from R having the same q is called a q -class Cl_q . The number q is called a class index. For a number $k = \gamma q^{1/4}$, γ is called the weight of k . For instance, a wave vector $\mathbf{k} = (160, 40)$ belongs to the q -class with $q = 1700$. Obviously, for any two numbers k_1, k_2 belonging to the same q -class, all their linear combinations with integer coefficients belong to the same class q .

It can be shown that Sys.(3) have two general types of solutions:

Type I: All 4 wave vectors belong to the same class Cl_q , in this case first equation of the Sys.(3) can be rewritten as

$$\gamma_1 \sqrt[4]{q} + \gamma_2 \sqrt[4]{q} = \gamma_3 \sqrt[4]{q} + \gamma_4 \sqrt[4]{q} \quad (4)$$

with integer $\gamma_1, \gamma_2, \gamma_3, \gamma_4$.

and

Type II: All 4 wave vectors belong to two different classes Cl_{q_1}, Cl_{q_2} ; in this case first equation of the Sys.(3) can be rewritten as

$$\gamma_1 \sqrt[4]{q_1} + \gamma_2 \sqrt[4]{q_2} = \gamma_3 \sqrt[4]{q_1} + \gamma_4 \sqrt[4]{q_2} \quad (5)$$

with integer γ_1, γ_2 .

Notice that (5) *is not identity* in the initial variables m_i, n_i because an integer can have several different presentations as a sum of two squares, for instance, 4-tuple $\{-1, 4\}, \{2, -5\}, \{-4, 1\}, \{5, -2\}$ is an example of II-type solution, with all weights $\gamma_i = 1$ and q -class indexes $q_1 = 17, q_2 = 29$.

The I- and II-type of solutions describe substantially different energy exchanges in the k -space. The II-type resonances are called *angle-resonances* [23] and consist of wavevectors with pairwise equal lengths, i.e. $|\mathbf{k}_1| = |\mathbf{k}_3|$ and $|\mathbf{k}_2| = |\mathbf{k}_4|$ or $|\mathbf{k}_1| = |\mathbf{k}_4|$ and $|\mathbf{k}_2| = |\mathbf{k}_3|$. Thus, these resonances do not transfer energy outside of the initial range of $|\mathbf{k}|$ and, therefore, cannot provide an energy cascade mechanism. However, these resonances can redistribute energy among the initial wavenumbers, in both

the direction $\mathbf{k}/|\mathbf{k}|$ and the scale $|\mathbf{k}|$. Since the initial support of energy in $|\mathbf{k}|$ cannot change, the II-type resonances alone would form a finite dimensional system, and it would be reasonable to expect a relaxation of such a system to a thermodynamic Rayleigh-Jeans distribution determined by the initial values of the motion integrals (the energy and the waveaction in this case). Note that such a thermalization could happen only among the resonant wavenumbers, and many wavenumbers which are initially excited but not in resonance would not evolve at all. (Such an absence of the evolution was called "frozen turbulence" in [6] where the capillary waves were studied for which there are no exact resonances).

On the other hand, the I-type resonances are called *scale-resonances* [23], and they can generate new wavelengths, - this corresponds to 3 or 4 different weights γ in (4) [23]. Thus, they are the only kind of resonances that can transfer energy outside of the range of initial $|\mathbf{k}|$.

SCALE-RESONANCES

We have studied a cluster structure of scale-resonances in the spectral domain $|m|, |n| \leq 1000$. In our computational domain we have found 230464 such resonances, among them 213760 collinear (i.e. all 4 wave vectors are collinear) and only 16704 (7.25%) non-collinear resonances. However, the nonlinear interaction coefficient in collinear quartets is equal to zero and, therefore, they have no dynamical significance [24]. On the other hand, for mathematical completeness we will consider these solutions too, for in some 4-wave resonance system the interaction coefficients in collinear quartets are not necessarily equal zero, of course.




Figure 1: Color online. Structure of collinear resonances in the spectral domain $|\mathbf{k}| \leq 100$ (left panel) and of non-collinear resonances in the same domain (right panel)

In the Table I the structure of all clusters is presented while in the Table II - the structure of non-collinear is given; cluster length is the number of quartets belonging to one cluster.

Cluster length	Number of clusters	Cluster length	Number of clusters
4	43136	352	3
8	1256	368	6
48	452	416	1
56	184	512	1
64	20	592	2
96	14	800	1
112	10	1024	1
256	7	1152	1
304	2	1376	1

Table I: Clustering in the entire set of quartets in the domain 1000x1000

Cluster length	Number of clusters	Cluster length	Number of clusters
8	1312	80	18
16	48	96	6
24	8	128	8
48	14	272	2
64	4	368	2

Table II: Clustering in the *non-collinear* subset of quartets in the domain 1000x1000.

In the next sections, on all the Figures, the structure of resonance clusters in the *spectral space* is shown as follows: each wave vector is presented as a node of integer lattice and nodes belonging to one solution are connected by lines. In Fig. 1 structure of collinear and non-collinear quartets is shown in a smaller domain $\sqrt{m_i^2 + n_i^2} \leq 100$.

Collinear quartets

First of all, let us make an important remark. If a 4-tuple $\{\{m_1, n_1\}, \{m_2, n_2\}, \{m_3, n_3\}, \{m_4, n_4\}\}$ consists of all collinear wave vectors, then the ratio $|m_i|/|n_i|$ is the same for all 4 wave vectors. Let us assume that $m_i, n_i \neq 0$, then

$$0 \neq |n_i|/|m_i| = c, \quad \forall i = 1, 2, 3, 4, \quad (6)$$

where c is an arbitrary finite non-zero constant and

$$|\mathbf{k}_i|^{1/2} = (m_i^2 + n_i^2)^{1/4} = m_i^{1/2}(1 + c^2)^{1/4}, \quad (7)$$

and Sys.(3) takes the form

$$\begin{cases} m_1^{1/2} + m_2^{1/2} = m_3^{1/2} + m_4^{1/2}, \\ m_1 + m_2 = m_3 + m_4, \end{cases} \Rightarrow \begin{cases} (m_1 m_2)^{1/2} = (m_3 m_4)^{1/2} \\ \Rightarrow |m_1| = |m_3 m_4|/|m_2|. \end{cases}$$




Figure 2: Color online. First non-axial trident

Now we can compute m_1 taking arbitrary integer m_2, m_3, m_4 provided that $|m_3 m_4|$ is divisible on $|m_2|$, and keeping in mind that $n_i = c m_i$, we can find all collinear solutions with $c \neq 0$. Obviously, a rational number c defines a line in spectra space and not all the lines are allowed. Case $c = 0$ corresponds to the solutions lying on the axes X ($n_i = c m_i$) and Y ($m_i = c n_i$), i.e. with all $m_i = 0$ or $n_i = 0$ correspondingly, e.g. 4-tuple $\{\{4, 0\}, \{-49, 0\}, \{-36, 0\}, \{-9, 0\}\}$. Parametrization of the resonances in this case was first given in [24]. As we have already mentioned before, these quartets are dynamically irrelevant because there is no nonlinear interaction within these quartets [24].

Non-collinear quartets

Tridents

Non-collinear scale-resonances have been studied first in [9, 15] in the spectral domain $\sqrt{m_i^2 + n_i^2} \leq 100$ and a special type of quartets named *tridents* have been singled out. By definition, a wave quartet is called a trident if:

- 1) There exist two vectors among four in a quartet, say \mathbf{k}_1 and \mathbf{k}_2 , such that $\mathbf{k}_1 \uparrow \downarrow \mathbf{k}_2$, thus they satisfy $(\mathbf{k}_1 \cdot \mathbf{k}_2) = -k_1 k_2$;
- 2) Two other vectors in the quartet, \mathbf{k}_3 and \mathbf{k}_4 , have the same length: $k_3 = k_4$;
- 3) \mathbf{k}_3 and \mathbf{k}_4 are equally inclined to \mathbf{k}_1 , thus $(\mathbf{k}_1 \cdot \mathbf{k}_3) = (\mathbf{k}_1 \cdot \mathbf{k}_4)$.

The following presentation for a trident quartet has been suggested in [9, 15],

$$\mathbf{k}_1 = (a, 0), \mathbf{k}_2 = (-b, 0), \mathbf{k}_3 = (c, d), \mathbf{k}_4 = (c, -d), \quad (8)$$

and two-parametric series of solutions has been written out:

$$\begin{cases} a = (s^2 + t^2 + st)^2, & b = (s^2 + t^2 - st)^2, \\ c = 2st(s^2 + t^2), & d = s^4 - t^4, \end{cases} \quad (9)$$

with arbitrary integer s, t . It is easy to check that vectors $\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4$ belong to the same class Cl_1 , with weights $\gamma_1 = s^2 + t^2 + st, \gamma_2 = s^2 + t^2 - st, \gamma_3 = \gamma_4 = s^2 + t^2$, and obviously $\gamma_1 + \gamma_2 = \gamma_3 + \gamma_4, \forall s, t \in \mathbb{Z}$.

Parametrization (8) corresponds to the tridents oriented along the X-axis with its vectors \mathbf{k}_1 and \mathbf{k}_2 and, therefore, we will call them *axial* tridents. There exist also *non-axial* tridents, for instance, the quartet $\{\{49, 49\}, \{-9, -9\}, \{5, 35\}, \{35, 5\}\}$. As mentioned in [9, 15], all the non-axial tridents can be obtained from the axial ones, (8), via a rotation by angles with rational values of cosine combined with respective re-scaling (to obtain an integer-valued solution out of rational-valued ones).

Cluster length	Number of clusters
8	1320
16	48
48	40
96	10
144	2
176	2

Table III: Clustering in the subset of *tridents* in the domain 1000x1000

In the computation domain $|m|, |n| \leq 1000$ we have found 13888 non-axial tridents, the first non-axial trident is shown on the Fig. 2. Among 13888, 13504 tridents have no vectors on any axis and 384 tridents have just one vector on an axis (for instance, $\{\{180, 135\}, \{0, 64\}, \{120, 119\}, \{60, 80\}\}$). The total amount of all possible tridents is 14848, thus only 960 are axial (6.5% of the total number). The data on tridents' clustering are given in the Table III.

Non-tridents

The study of the resonance solution set in the spectral domain $|m_i|, |n_i| \leq 1000$ shows that not all non-collinear cascading quartets are *tridents*. They are called further *non-tridents*, e.g. a quartet $\{\{990, 180\}, \{128, 256\}, \{718, 236\}, \{400, 200\}\}$ is a non-trident quartet.

The overall number of tridents is 14848 while the number of non-tridents is 1856. Notice that the first non-trident quartet $\{\{180, 135\}, \{0, 64\}, \{120, 119\}, \{60, 80\}\}$ lies in the spectral domain $|\mathbf{k}| \leq 225$. This means, that if we are interested only in the large-scale quartets, say, quartets with $|\mathbf{k}| \leq 100$, the complete set of scale-resonances consists of 1728 quartets, among them - 1632 collinear quartets and 96 tridents, but no non-tridents yet.




Figure 3: Color online. First non-trident

Clusters

In the previous Section we have shown that there are three different types of scale-resonances: collinear quartets, tridents and non-tridents. Before studying the structure of clusters formed by these resonances let us notice the following simple but important fact. Suppose a quartet

$$\{\tilde{\mathbf{k}}_1, \tilde{\mathbf{k}}_2, \tilde{\mathbf{k}}_3, \tilde{\mathbf{k}}_4\} \quad (10)$$

is a solution of (3), then each permutation of indexes $1 \leftrightarrow 2, 3 \leftrightarrow 4, 1 \leftrightarrow 3, 1 \leftrightarrow 4, 2 \leftrightarrow 3, \dots$ will generate a new solution of (10), in general case, all together 8 different *symmetry generated solutions*. Of course, in some particular cases, when some of the vectors belonging to a quartet coincide, the overall number of symmetry generated solutions can be smaller. For instance, the quartet $\{\{0, -54\}, \{0, 294\}, \{90, 120\}, \{-90, 120\}\}$ is part of the cluster of 8 symmetry generated solutions, while the quartet $\{\{17, 31\}, \{-153, -279\}, \{-68, -124\}, \{-68, -124\}\}$ belongs to the cluster of 4 such solutions. On the Fig. 4 the smallest tridents cluster is shown, with and without multi-edges. Graphical presentation of a cluster as a graph with multi-edges is of course mathematically correct but would make the pictures of bigger clusters somewhat nebulous (see Fig. 4, left panel). Therefore, further on we omit multi-edges in our graphical presentations as it is shown in Fig. 4, right panel.

There are also clusters formed by different types of scale-resonances, for instance, clusters containing tridents and non-tridents (Fig. 5, left panel) or collinear and non-collinear quartets (Fig. 5, right panel)

One of the most important characteristic of the resonances structure is the *wavevector multiplicity* (introduced in [20]), which describes how many times a given wavevector is a part of some solution. In the Table IV the wavevectors multiplicities are given, for non-collinear quartets. It turned out that 91% of all these wavevectors (6720 from overall amount 7384) have multiplicity 8, i.e.





Figure 4: Color on line. The smallest tridents cluster formed by 8 symmetry generated solutions, two presentation forms are given: with multi-edges (left panel) and without multi-edges (right panel)

Figure 5: Color on line. **Left panel:** A shortest cluster formed by both tridents and non-tridents; cluster length is 64, among them 48 tridents and 16 non-tridents. **Right panel:** A shortest cluster formed by both collinear and non-collinear quartets; cluster length is 48, among them 32 collinear and 16 non-collinear quartets. No multi-edges are shown.

they are parts of symmetry generated group of solution. This fact simplifies drastically the computations of the dynamical system (13) corresponding to such a cluster because interaction coefficient, Z , will be the same for all symmetry generated solutions.

Multiplicity	Amount of vectors
8	6720
16	424
24	192
32	36
40	8
48	4

Table IV: Wavevectors multiplicity computed for the non-collinear quartets in the spectral domain $|m_i|, |n_i| \leq 1000$.

As can be seen from the Fig. 4, the dynamics of a cluster consisting of 8 connected quartets is defined by only 4 different wave-frequencies! In the Fig. 6 the cluster of 32 connected quartets is shown, with all together only 7 different wave-frequencies.




Figure 6: Example of a non-trident cluster of length 32, no multi-edges are shown

Obviously, the cluster structure defines the form of dynamical system corresponding to the cluster. The main goal of this detailed clusters study is of course a hope to construct an isomorphism (i.e. one-to-one correspondence) between a cluster and a dynamical system. In [21] this construction have been presented for an arbitrary 3-wave resonance system, with triads as primary elements of the planar graph. Its implementation in Mathematica is given in [22], where also interaction coefficients similar to Z in (13) are computed. To construct this isomorphism two following facts have been used: 1) in a 3-wave resonance system only *scale-resonances* do exist, and 2) if we add an arbitrary triad to a cluster, in general we always add some new wave-frequencies as well (the only exception is identified in [21]). A 4-wave resonance system does not possess these nice properties, on the contrary - there exist scale- and angle-resonances, and most of the quartets are parts of a symmetry generated solutions set. It would be a challenge to develop some general approach to construct dynamical systems for resonance cluster in an arbitrary 4-wave system.

ANGLE-RESONANCES

Until now we have studied the structure of the scale-resonances which are described by (4) and generate new wavenumbers, thereby providing the energy cascade mechanism. The angle-resonances represented by (5) do not generate new wavenumbers but can redistribute energy among the modes which were excited initially (or which are forced externally). As we have mentioned earlier, if taken on their own, these resonances would lead to the thermal equilibrium distribution of the energy and the waveaction among the (initially excited) resonant waves. The important fact, however, is that the scale- and the angle-resonances are not independent and can form a *mixed cluster* containing both types of this resonances [23]. The energy cascade mechanism in such a

mixed cluster is presented schematically in Fig.7 where the quadrangles S_1 and S_2 denote scale-resonances so that 4-tuples $(V_{1,1}, \dots, V_{1,4})$ and $(V_{2,1}, \dots, V_{2,4})$ represent scale-resonances and squares $A_1, \dots, A_i, \dots, A_n$ represent angle-resonances.




Figure 7: Color on line. Schematic presentation of mixed resonance cascade

If a wave takes a part simultaneously in angle- and scale-resonances, corresponding nodes of S_1, S_2 and A_j are connected by (red) dashed arrows so that $V_{1,1} = V_{n,2}$ and $V_{n,1} = V_{2,3}$. We have found many examples of such mixed clusters in our solution set, for instance:

$$S_1 = \{ \{-64, -16\}, \{784, 196\}, \{144, 36\}, \{576, 144\} \},$$

$$A_n = \{ \{-64, -16\}, \{4, 16\}, \{-64, 16\}, \{4, -16\} \},$$

$$\text{with } V_{1,1} = V_{n,2} = (-64, -16),$$

$$S_2 = \{ \{-49, -196\}, \{4, 16\}, \{-36, -144\}, \{-9, -36\} \},$$

which is further on connected with an angle-resonance

$$A_{\bar{n}} = \{ \{-49, -196\}, \{784, 196\}, \{-49, 196\}, \{784, -196\} \},$$

(not shown in Fig.7) *via* a node-vector $(-49, -196)$.

The energy flux over scales due to the mixed cluster is weak: one wave can participate in a few dozen of angle-resonances (see Fig. 8), which means that only a small part of its energy will go to a scale-resonance. Moreover, even this weak energy cascade may terminate at finite wavenumber if it happens to move along a *finite* cluster which terminates before reaching the dissipation range (or before reaching the range of high frequencies where the nonlinearity gets large enough for the quasi-resonances to take over the energy flux from the exact resonances).

The number of angle-resonances in the spectral domain $|m_i|, |n_i| \leq 1000$ is of order of $6 \cdot 10^8$ while the number of scale-resonances in the same domain is of order of $8 \cdot 10^5$, among them less than $2 \cdot 10^4$ are non-collinear and do play a role in the energy exchange among the modes within the quartets (see next the Section).




Figure 8: Color on line. Wave (64,0) takes part in 2 scale-resonances, both *non-tridents*. The upper number in the circle is m , the lower is n , and (red) thick lines drawn between vectors on the same side of the Eqs.(3). Wave (119,120) takes part in 1 scale-resonance and in 12 angle-resonances.

Another important characteristic of the resonances structure is the multiplicity of a vector, which describes how many times a given vector is a part of some solution. The multiplicity histogram for angle-resonances is shown




Figure 9: The multiplicities histogram for angle-resonances.

in Fig.9. On the axis X the multiplicity of a vector is shown and on the axis Y the number of vectors with a given multiplicity. This has been cut off - multiplicities go very high, indeed the vector (1000,1000) takes part in 11075 solutions. All this indicates that the angle-resonances play an important role in the overall dynamics of the wave field.

DYNAMICS

The wave field

Once the clusters are found, one can consider an evolution of amplitudes of waves that belong to each individual cluster by considering a respective reduction of the dynamical equation. For the gravity wave case, the appro-

priate dynamical equation is Zakharov equation for the complex amplitude $a_{\mathbf{k}}(t)$ corresponding to the \mathbf{k} -mode,

$$i \frac{da_{\mathbf{k}}}{dt} = \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{\mathbf{k} + \mathbf{k}_1 = \mathbf{k}_2 + \mathbf{k}_3} T_{\mathbf{k}_2, \mathbf{k}_3}^{\mathbf{k}, \mathbf{k}_1} a_{\mathbf{k}_1}^* a_{\mathbf{k}_2} a_{\mathbf{k}_3} e^{i(\omega_{\mathbf{k}} + \omega_{\mathbf{k}_1} - \omega_{\mathbf{k}_2} - \omega_{\mathbf{k}_3})t}, \quad (11)$$

where $T_{\mathbf{k}_2, \mathbf{k}_3}^{\mathbf{k}, \mathbf{k}_1} \equiv T(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$ is an interaction coefficient for gravity water waves which can be found in [25].

For very weak waves, amplitudes $a_{\mathbf{k}}(t)$ vary in time much slower than the linear oscillations. The factor $e^{i(\omega_{\mathbf{k}} + \omega_{\mathbf{k}_1} - \omega_{\mathbf{k}_2} - \omega_{\mathbf{k}_3})t}$ on RHS will rapidly oscillate for most waves except for those in an exact resonance for which this factor is 1. Thus, only the resonant modes will give a contribution to the dynamics in this case and the oscillating contributions of the non-resonant terms will average out in time to zero and will not give any contribution to the cumulative change of $a_{\mathbf{k}}(t)$. Leaving only the resonant terms, we have

$$i \frac{da_{\mathbf{k}}}{dt} = \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^R T_{\mathbf{k}_2, \mathbf{k}_3}^{\mathbf{k}, \mathbf{k}_1} a_{\mathbf{k}_1}^* a_{\mathbf{k}_2} a_{\mathbf{k}_3}, \quad (12)$$

where \sum^R means a summation only over $\mathbf{k}_1, \mathbf{k}_2$ and \mathbf{k}_3 which are in resonance with \mathbf{k} . Obviously, we should consider \mathbf{k} 's from the same cluster only (i.e. solve the problem for one cluster at a time). Note that the fast timescale of the linear dynamics completely disappeared from this equation. Thus, paradoxically, the dynamics of very weak waves in finite boxes is strongly nonlinear: it is more nonlinear than in WTT which works for larger amplitudes and where quasi-resonances ensure phase randomness. This explains the fact found in the three-wave example that even relatively large clusters often exhibit a periodic or quasi-periodic behavior [26].

A study of the system (12) is possible analytically for small clusters (perhaps even integrating the system in some lucky cases) and numerically for large clusters, which is an interesting subject for future research. Some properties, however, can already be seen in the example of a single quartet which has been studied before [27]. Let us briefly discuss these properties.

A quartet

The dynamical system describing slowly changing amplitudes of a quartet has the form [27]:

$$\begin{cases} i\dot{a}_1 = 2Za_2^*a_3a_4 \\ i\dot{a}_2 = 2Za_1^*a_3a_4 \\ i\dot{a}_3 = 2Za_4^*a_1a_2 \\ i\dot{a}_4 = 2Za_3^*a_1a_2 \end{cases} \quad (13)$$

and $Z = T_{\mathbf{k}_2, \mathbf{k}_3}^{\mathbf{k}, \mathbf{k}_1}$. The mathematical analysis of the system (13) can be performed similar to what has been done in [28] for an integrable 3-wave system of resonantly interacting planetary waves, though computations of the modulus of elliptic integral are more involved and variety of different dynamical scenarios is substantially richer (see [27] for details). The general answer can be given in terms of Jacobean elliptic functions.

Obviously, the quartets with interaction coefficient $Z = 0$ do not influence the general dynamics of the wave field at the corresponding time scale. As it was shown in [24], for all *collinear* quartets $Z \equiv 0$, and therefore they can be excluded from consideration.

On the other hand, in [29] so-called "degenerate quartets" (*tridents* in our terminology) have been studied and it was established numerically that they have strictly periodic behavior: after appropriate translation in the horizontal plane, two snapshots of the free-surface taken at $t = 0$ and $t = T$ are identical.

As to exact the resonances, in a general situation and for some specific initial conditions, the system (13) "does not exhibit strict periodicity" in numerical simulations [27]. Thus, general formulae for the solutions of (13) have to be studied in more details in order to distinguish periodic and non-periodic dynamics of an arbitrary quartet. On the other hand, as our main interest in this Letter is the large-scale dynamics, all scale-resonances with a non-zero interaction coefficient are tridents and therefore demonstrate a periodic time behavior. Of course, if a trident is involved into a cluster with some other quartets then one should be cautious about the predictions obtained for an isolated quartet.

SUMMARY AND DISCUSSION OF RESULTS

- In this Letter, we studied properties of deep water gravity waves bounded by a square periodic box. At very small wave amplitudes, when the nonlinear resonance broadening is less than the k -space spacing, WTT fails and only the waves which are in *exact* four-wave resonance can interact. This situation appears to be typical for all existing numerical simulations and laboratory experiments. Indeed, as mentioned in [9], to overcome the wavenumber discreteness associated with the final box one needs at least 10000 x 10000 numerical resolution which is presently unavailable for this type of problems. On the other hand, as shown in [10], even in a 10m x 6m laboratory flume the finite-box effects are very strong. Thus, to understand the wave behavior in laboratory experiments and in numerical simulations it is crucial to study exact resonances among discrete wave modes, which was the focus of the present Letter.

- We found numerically all resonant quartets on the set of 1000 x 1000 modes making use of the q -class method originally developed in [18]. We found that all resonant quartets separate into noninteracting with each other clusters. Each cluster may consist of two types of quartets: scale- and angle-resonances. The angle-resonances cannot transfer energy to any k -modes which are not already present in the system. They cannot carry an energy flux through scales and their main role is to thermalize the initially excited modes. The scale-resonances are much more rare than the angle ones and yet their role is important because they are the only resonances that can transfer energy between different scales. Most of the scale-resonances, but not all, are of the *trident* type, for which a partial parametrization can be written out explicitly. If one is interested in large-scale modes only, say 100 x 100 domain, then the tridents are the only scale-resonances, which is very fortunate due to the available parametrization.

- Even though the angle-resonances cannot cascade energy, they are important for the overall cascade process because they are involved in the same wave clusters with the scale-resonances. One wave mode may typically be connected simultaneously to many angle modes and only one (or several) scale modes. Thus, one can split large clusters into "reservoirs", each formed by a large number of angle quartets in quasi-thermal equilibrium, and which are connected with each other by sparse links formed by scale quartets. This structure suggests a significant energy cascade slowdown and anisotropy with respect to the infinite-box limit. Further study is needed to examine the structure of such large clusters and possible energy cascade routes from the region of excitation at low wavenumbers to the dissipative large k range.

- We also discussed consequences of the cluster structures for the dynamics, and argued that one should expect a less random and more regular behavior in the case of very low amplitude waves with respect to larger (but still weak) waves described by WTT. More study is needed in future both analytically, for small clusters, and numerically, for large clusters. A particularly interesting question to answer in this case is about any possible universal mechanisms of transition between the regular dynamics to chaos and possible coexistence of the regular and chaotic motions.

Acknowledgments

Acknowledgements. E.K. acknowledges the support of the Austrian Science Foundation (FWF) under project P20164-N18 "Discrete resonances in nonlinear wave systems". O.R. and S.N. acknowledge the support of the Transnational Access Programme at RISC-Linz, funded

by European Commission Framework 6 Programme for Integrated Infrastructures Initiatives under the project SCIENCE (Contract No. 026133).

-
- [1] V.E. Zakharov, V.S. L'vov and G. Falkovich. **Kolmogorov Spectra of Turbulence**. Series in Nonlinear Dynamics, Springer (1992)
- [2] Y. Lvov and S.V. Nazarenko, *Phys. Rev. E* **69**: 066608 (2004)
- [3] Y. Choi, Y. Lvov, S.V. Nazarenko and B. Pokorni. *Physics Letters A*: **339** (3-5): 361 (2004)
- [4] Y. Choi, Y. Lvov and S.V. Nazarenko. *Physics Letters A*: **332** (3-4): 230 (2004)
- [5] A.C. Newell, S.V. Nazarenko and L. Biven, *Physica D* **152-153**: 520 (2001)
- [6] A.N. Pushkarev and V.E. Zakharov, *Physica D* **135** (1-2): 98 (2000)
- [7] V.E. Zakharov, A.O. Korotkevich, A.N. Pushkarev and A.I. Dyachenko, *JETP Lett.* **82** (8): 487 (2005)
- [8] S. Annenkov and V. Shrira, *Phys. Rev. Lett* **96**: 204501 (2006)
- [9] Y.V. Lvov, S. Nazarenko and B. Pokorni. *Physica D* **218**: 24 (2006)
- [10] P. Denissenko, S. Lukaschuk and S. Nazarenko, *Phys. Rev. Lett.* **99**: 014501 (2007)
- [11] E. Falcon, C. Laroche and S. Fauve. *Phys. Rev. Lett.* **98**: 094503 (2007)
- [12] G. Kolmakov, A. Levchenko, M. Braznikov, L. Mezhov-Deglin, A. Slichenko and P. McClintock. *Phys. Rev. Lett.* **93**: 74501 (2004)
- [13] M. Braznikov, G. Kolmakov, A. Levchenko and L. Mezhov-Deglin. *Europhysics Lett.* **58**: 510 (2002)
- [14] V. M. Eguiluz, M. T. Levinsen and P. Alstrom. *Europhys. Lett.* **58**: 517 (2002)
- [15] S. Nazarenko. *J. Stat. Mech.* LO2002 (2006)
- [16] E. Kartashova. *Phys. Rev. Lett.* **72**: 2013 (1994)
- [17] E. Kartashova. *AMS Transl.* **182** (2): 95 (1998)
- [18] E. Kartashova. *J. Low Temp. Phys.* **145** (1-4): 286 (2006)
- [19] E. Kartashova and A. Kartashov. *Int. J. Mod. Phys. C* **17**(11): 1579 (2006)
- [20] E. Kartashova and A. Kartashov. *Comm. Comp. Phys.* **2** (4): 783 (2007)
- [21] E. Kartashova and G. Mayrhofer. *Physica A: Stat. Mech. Appl.* **385**: 527 (2007)
- [22] E. Kartashova, C. Raab, Ch. Feurer, G. Mayrhofer and W. Schreiner. In Ch. Kharif, E. Pelinovsky (Eds): **Extreme Waves in the Ocean**, Springer (to appear). E-print: <http://arxiv.org/abs/0706.3789> (2008)
- [23] E. Kartashova. *Phys. Rev. Lett.* **98** (21): 214502 (2007)
- [24] A.I. Dyachenko, Y.V. Lvov and V.E. Zakharov, *Physica D* **87** (1-4): 233 (1995)
- [25] V.P. Krasitskii. *J. Fluid Mech.* **272**: 1 (1994)
- [26] E. Kartashova and V.S. L'vov. Submitted to *Phys. Rev. Lett.* E-print arXiv:0801.3374v1 (2008)
- [27] M. Stiassnie and L. Shemer. *Wave motion* **41**: 307 (2005)
- [28] E. Kartashova and V.S. L'vov. *Phys. Rev. Lett.* **98** (19): 198501 (2007)
- [29] L. Shemer and M. Stiassnie. In Y.Toba, H.Mitsuyasu, Reidel F.D. (Eds): **The ocean surface**, p.51-57. Dordrecht, Holland (1985)