

On the Global Structure of Hopf Hypersurfaces in a Complex Space form.

A.A.Borisenko.

Abstract. It is known that a tube over a Kähler submanifold in a complex space form is a Hopf hypersurface. In some sense the reverse statement is true: a connected compact generic immersed C^{2n-1} regular Hopf hypersurface in the complex projective space is a tube over an irreducible algebraic variety. In the complex hyperbolic space a connected compact generic immersed C^{2n-1} regular Hopf hypersurface is a geodesic hypersphere.

Introduction.

A natural class of real hypersurfaces in a complex space form $\overline{M}(c)$ of constant holomorphic curvature $4c$ is the class of Hopf hypersurfaces. For a unit normal vector ξ of a hypersurface M the vector $J\xi$ is a tangent vector to M , where J is the complex structure of the complex space form $\overline{M}(c)$.

Definition A hypersurface $M \subset \overline{M}(c)$ is called a Hopf hypersurface if the vector $J\xi$ is a principal direction at every point of M .

Y.Maeda [11] proved that for Hopf hypersurfaces in the n -dimensional complex projective space \mathbf{CP}^n the corresponding principal curvature in the direction $J\xi$ is constant. It is known that a tube over a Kähler submanifold in a complex projective space is a Hopf hypersurface. T.E. Cecil and P.J. Ryan studied the local and global structure of Hopf hypersurfaces with constant rank of the focal map Φ_r .

Let M be an embedded hypersurface of $\overline{M}(c)$ of the regularity class C^2 . Let NM be the normal bundle of M with projection $p : NM \rightarrow M$ and let BM be the unit normal bundle. For $\xi \in NM$ let $F(\xi)$ be the point in $\overline{M}(c)$ reached by traversing a distance $|\xi|$ along the geodesic in $\overline{M}(c)$ originating at $x = p(\xi)$ with the initial tangent vector ξ .

A point $P \in \overline{M}(c)$ is called a focal point of multiplicity $\nu > 0$ of (M, x) if $P = F(\xi)$ and the Jacobian of the map F has nullity ν at ξ .

Definition The tube of radius r over M is the image of the map $\Phi_r : BM \rightarrow \overline{M}(c)$ given by $\Phi_r(\xi) = F(r\xi)$, $\xi \in BM$.

T.E. Cecil and P.J. Ryan had proved the following result:

Lemma 1. [1] Let M be a connected, orientable Hopf hypersurface of \mathbf{CP}^n with corresponding constant principal curvature $\mu = 2 \cot 2r$. Suppose the map Φ_r has constant rank q on M . Then q is even and every point $x_0 \in M$ has a neighbourhood U such that $\Phi_r(U)$ is an embedded complex $q/2$ -dimensional submanifold of \mathbf{CP}^n .

We remark that, in Lemma 1 and Lemma 13 below, C^3 regularity is enough. From Lemmas 1 and 13 we obtain that Hopf hypersurface with Φ_r of constant rank is an analytical hypersurface. It follows from this fact that $\Phi_r(U)$ is a complex submanifold and parametrizations functions of $\Phi_r(U)$ satisfy an elliptic system of the PDE's with analytical coefficients. From C^2 regularity of $\Phi_r(U)$ we obtain that $\Phi_r(U)$ is analytic.

The global version of Lemma 1 has the following form [1]:

Let M be a connected compact embedded real Hopf hypersurface in \mathbf{CP}^n with corresponding constant principal curvature $\mu = 2 \cot 2r$. Suppose the map Φ_r has constant rank q on M . Then Φ_r factors through a holomorphic immersion of the complex $q/2$ -dimensional manifold M/T_0 into \mathbf{CP}^n , where T_0 are $(2n - q - 1)$ -dimensional spheres, the leaves of the distribution

$$T_0(x) = \{y \in T_x M, (\Phi_r)_*(y) = 0\}.$$

Acknowledgements. The author began to study Hopf hypersurfaces when he was a visiting professor at the University of Valencia (Spain) from January 1997 to June 1997. The author would like to thank Vicente Miquel and Yury Nikolaevsky for closely reading a preliminary version of this paper and for their remarks. The author would like to thank the referee for valuable remarks.

1. The main results

The following theorem gives a complete description of the global structure of Hopf hypersurfaces in complex space forms.

Let M be an immersed regular hypersurface in a regular manifold N . Suppose that for a point $P \in N$ of self-intersection the linear span of the tangent hyperplanes to the branches of M coincides with tangent space $T_P N$ of the ambient manifold. This point is called a generic point of self-intersection. If every point of self-intersection of the hypersurface M is a generic point of self-intersection then the hypersurface M is called a generic immersed hypersurface.

Theorem. 1. *Let M be a C^{2n-1} regular compact generic immersed orientable Hopf hypersurface in the complex projective space $\mathbf{C}P^n$ ($n \geq 2$). Then M is a tube over an irreducible algebraic variety.*

Corollary *Let M be a C^{2n-1} regular connected compact embedded Hopf hypersurface in the complex projective space $\mathbf{C}P^n$ ($n \geq 2$). Then M is a tube over an irreducible algebraic variety.*

The following are some standard examples of Hopf hypersurfaces in $\mathbf{C}P^n$ of constant holomorphic curvature 4.

1. A geodesic hypersphere M is the set of points at a fixed distance $r < \frac{\pi}{2}$ from a point $P \in \mathbf{C}P^n$. It is obvious that M is also the tube of radius $\frac{\pi}{2} - r$ over the hyperplane $\mathbf{C}P^{n-1} \subset \mathbf{C}P^n$ dual to the point P .
2. A tube over a totally geodesic $\mathbf{C}P^k$ ($1 \leq k \leq n - 1$).
3. A tube over a totally geodesic real projective space RP^n and over a complex quadric $Q^{n-1} = \{(z_0, \dots, z_n) \in \mathbf{C}P^n : z_0^2 + z_1^2 + \dots + z_n^2 = 0\}$.

A tube of small radius r over a closed irreducible algebraic manifold in $\mathbf{C}P^n$ is an analytic Hopf hypersurface. But let $f = x_0^6 x_3^2 + x_1^3 x_2^5 = 0$ be the algebraic variety M in $\mathbf{C}P^3$. The point $P(1, 0, 0, 0)$ is a singular point ($\text{grad } f/P = 0$). In any neighbourhood of the point P the normal curvatures at smooth points vary from $-\infty$ to $+\infty$. From Lemma 12 below it follows that normal curvatures of the tube of any radius r tend to $+\infty$. It follows that the tube of any radius r has regularity less than $C^{1,1}$.

V. Miquel had proved the following theorem:

Theorem(V. Miquel [13]) *Let M be a connected compact embedded Hopf hypersurface in $\mathbf{C}P^n$ contained in a geodesic ball of radius $R < \frac{\pi}{2}$.*

Suppose that

1. M has constant mean curvature H ;
2. The principal curvature μ in the direction $J\xi$ satisfies the inequality

$$\mu \geq 2 \cot \left(2 \text{arc cot} \left[\frac{(2n-1)H - \mu}{2n-2} \right] \right).$$

Then M is a geodesic hypersphere.

We prove the following theorem.

Theorem. 2. *Let M be a C^{2n-1} regular connected compact generic immersed orientable Hopf hypersurface in the complex projective space $\mathbf{C}P^n$ ($n \geq 2$) contained in a geodesic ball of radius $R < \frac{\pi}{2}$. Then M is a geodesic hypersphere.*

Let $\mathbf{C}H^n$ be the complex hyperbolic space of constant holomorphic curvature -4 . We prove the following theorem.

Theorem. 3. *Let M be a connected compact generic immersed orientable C^{2n-1} regular Hopf hypersurface in the complex hyperbolic space $\mathbf{C}H^n$ ($n \geq 2$). Then the Hopf hypersurface M is a geodesic hypersphere.*

2. Lemmas

Lemma 2. *(Y. Maeda, [11]) Let M be a connected Hopf hypersurface in the complex projective space $\mathbf{C}P^n$. Then the principal curvature μ of M in the direction $J\xi$ is constant.*

Let A_ξ be the shape operator of M .

Lemma 3. *(T.E. Cecil, P.J. Ryan [1]) Suppose $J\xi$ is an eigenvector of A_ξ with an eigenvalue μ . Then we have:*

- a) $(F_*)_{r\xi}(X, 0) = 0$ if $\lambda = \cot r$ is an eigenvalue of A_ξ and X is a vector in the eigenspace T_λ corresponding to the eigenvalue λ .
- b) $(F_*)_{r\xi}(J\xi, 0) = 0$ if $\mu = 2\cot 2r$.
- c) $(F_*)_{r\xi}(X, V) \neq 0$ except as determined by (a) and (b).

Now, let M be a real hypersurface of a complex space form $\overline{M}^n(c)$ of constant holomorphic curvature $4c$ and let ξ be a unit normal field on M . If $X \in T_P M$, $P \in M$, then one has a decomposition

$$JX = \phi X + f(X)\xi$$

into the tangent and normal components respectively. So, ϕ is a $(1, 1)$ -tensor field and f is a 1-form. Then they satisfy

$$\phi^2 X = -X + f(X)U, \quad \phi U = 0, \quad f(\phi X) = 0$$

for any vector field X tangent to M , where $U = -J\xi$. Moreover, we have

$$g(\phi X, Y) + g(X, \phi Y) = 0, \quad f(X) = g(X, U);$$

$$g(\phi X, \phi Y) = g(X, Y) - f(X)f(Y)$$

with g the metric tensor in $\overline{M}^n(c)$. We denote by A the shape operator on $T_P M$ associated with ξ .

Lemma 4. *1.([9]) Let M be a Hopf hypersurface in $\overline{M}^n(c)$. Then we have*

$$a) \quad -2c\phi = \mu(\phi A + A\phi) - 2A\phi A;$$

$$b) \quad X\mu = (U\mu)f(X)$$

and

$$(U\mu)g((\phi A + A\phi)X, Y) = 0,$$

where μ is the principal curvature in the direction $U = -J\xi$, X, Y are vectors tangent to M , and $U\mu$ is the derivative of the function μ in the direction U . Moreover, if $\phi A + A\phi = 0$ then

$$cg(X, \phi Y) = -g(\phi AX, AY) = g(A\phi X, AY),$$

$$cg(\phi X, \phi X) = -g(A\phi X, A\phi X)$$

and so $c \leq 0$.

2.([11]) Let M be a Hopf hypersurface in $\mathbf{C}P^n$. If $X \in T_\alpha \subset T_P M$, then

$$JX \in T_{\mu\alpha+2/2\alpha-\mu} \subset T_P M,$$

where T_α is an eigenspace corresponding to a principal curvature α .

It follows from the equation (a) of the first part of the lemma that α cannot be equal to μ or to $\mu/2$.

Definition Let A be a subset of a metric space X . Let $\delta(A)$ denote the diameter of A , and let

$$\delta^p(A) = [\delta(A)]^p \text{ for } p > 0,$$

$$\delta^0(A) = \begin{cases} 1, & \text{if } A \neq \emptyset; \\ 0 & \text{if } A = \emptyset. \end{cases}$$

For $p \geq 0$ and $\varepsilon > 0$ define.

$$H_\varepsilon^p(A) = \inf \left\{ \sum_{i=1}^{\infty} \delta^p(A_n) : A \subset \cup A_n \text{ and } \delta(A_n) < \varepsilon \right\};$$

$$H^p(A) = \lim_{\varepsilon \rightarrow 0^+} H_\varepsilon^p(A) = \sup H_\varepsilon^p(A).$$

We call H^p the Hausdorff p -measure.

Lemma 5. (H. Federer, [4]) If $m > \nu \geq 0$ and $k \geq 1$ are integers, A is an open subset of R^m , $B \subset A$, Y is a normed vector space and $f : A \rightarrow Y$ is a map of class C^k such that

$$\text{Dim } \text{im } f_*(x) \leq \nu \quad \text{for } x \in B,$$

then

$$H^{\nu+(m-\nu)/k}[f(B)] = 0.$$

Definition Let Ω be a complex manifold. A set $A \subset \Omega$ is called an analytic set in Ω if for each point $a \in \Omega$, there exist a neighbourhood U of a and functions f_1, \dots, f_N holomorphic in U such that $A \cap U = Z_{f_1} \cap \dots \cap Z_{f_k} \cap U$, where Z_f is the set of zeros of a holomorphic function f .

A point a of an analytic set A is called a regular point if there exists a neighbourhood U of a in Ω such that $A \cap U$ is a complex submanifold of U . The complex dimension of $A \cap U$ is then called the dimension of A at the point a and is denoted by $\dim_a A$. The set of all regular points of an analytic set is denoted by $\text{reg } A$. Its complement $A \setminus \text{reg } A$ is denoted by $\text{sng } A$. The set $\text{sng } A$ is called the set of singular points of the set A . It can be shown by induction on the dimension of the manifold Ω that $\text{sng } A$ is nowhere dense and closed. This allows us to define the dimension of A at any point a of A as

$$\dim_a A = \lim_{z \rightarrow a} \dim_z A \quad (z \in \text{reg } A).$$

The set A is called purely p -dimensional if $\dim_z A = p$ for all $z \in A$ [2], [3].

Lemma 6. (B. Shiffman, [16]) Let E be a closed subset of a complex manifold Ω and let A be a purely q -dimensional analytic subset of $\Omega \setminus E$. If $H^{2q-1}(E) = 0$ then the closure \overline{A} of the set A in Ω is a purely q -dimensional analytic subset of Ω .

Definition(D.Mumford, [14]) Let $U \subset \mathbf{C}^n$ be an open set. A closed subset $X \subset U$ is a $*$ -analytic subset of U if X can be decomposed

$$X = X^{(r)} \cup X^{(r-1)} \cup \dots \cup X^{(0)},$$

where for all i , $X^{(i)}$ is an i -dimensional complex submanifold of U and $\overline{X^{(i)}} \subset X^{(i)} \cup X^{(i-1)} \dots \cup X^{(0)}$. If $X^{(r)} \neq \emptyset$, then r is called the dimension of X .

An analytic set is always $*$ -analytic [14].

Lemma 7. (Chow's Theorem, [14]) If $X \subset \mathbf{C}P^n$ is a closed $*$ -analytic subset, then X is a finite union of algebraic varieties.

Lemma 8. [3] An analytic set A in a complex manifold Σ is irreducible if and only if the set $\text{reg } A$ is connected.

Let $X \subset \mathbf{C}P^n$ denote a closed irreducible algebraic variety of dimension l which may have singularities and let $X_e \subset X$ denote the non-empty open subset of its smooth points. For the definitions of irreducible singular and smooth points see [14]. Define

$$V'_X = \left\{ (x, y) \in \mathbf{C}P^n \times \mathbf{C}\check{P}^n \mid x \in X_e \text{ and } y \text{ is tangent hyperplane at } x \right\},$$

where $\mathbf{C}\check{P}^n$ is the dual complex projective space.

The closure V_X of V'_X on Zariski topology in $\mathbf{C}P^n \times \mathbf{C}\check{P}^n$ is called the tangent hyperplane bundle of X . It is a closed irreducible algebraic variety of dimension $(n-1)$. The first projection maps V_X onto X

$$\pi_1: V_X \rightarrow X, \quad (x, y) \rightarrow x.$$

Consider now the second projection

$$\pi_2: V_X \rightarrow \mathbf{C}\check{P}^n, \quad (x, y) \rightarrow y.$$

Its image $\check{X} = \pi_2(V_X)$ is a closed irreducible variety of $\mathbf{C}\check{P}^n$ of dimension at most $(n-1)$, the dual variety of X [9].

Lemma 9. (*Duality Theorem*) [6], [10] *The tangent hyperplane bundles of an closed irreducible algebraic variety X and its dual variety \check{X} coincide: WE have $V_{\check{X}} = V_X$ and hence $\check{\check{X}} = X$.*

Let $\mathbf{C}P^n$ be the complex projective space with standard Fubini-Study metric. To a hyperplane $L \subset \mathbf{C}P^n$ passing through a point $x \in \mathbf{C}P^n$ we associate the point $y \in \mathbf{C}P^n$ representing the complex line in \mathbf{C}^{n+1} orthogonal to L . Then the distance $\rho(x, y)$ is equal to $\pi/2$. One can identify $\mathbf{C}\check{P}^n$ with $\mathbf{C}P^n$ in this way and consider \check{X} as a subset in $\mathbf{C}P^n$.

It is possible to define a tube over an closed irreducible algebraic variety $X \subset \mathbf{C}P^n$ which may have singularities. Let $(x, y) \in V_X \subset \mathbf{C}P^n \times \mathbf{C}\check{P}^n = \mathbf{C}P^n \times \mathbf{C}P^n$, $x \in X$, $y \in \check{X}$, and let $L(x, y)$ be a complex projective line through $x, y \in \mathbf{C}P^n$. Then $L(x, y)$ is a totally geodesic two-dimensional sphere in $\mathbf{C}P^n$ of curvature 4, the distance $\rho(x, y)$ is equal to $\pi/2$, and x and y are poles of the sphere $L(x, y)$. The set of points of $L(x, y)$ at a distance r from the point x is a circle $S_r(x, y)$ with the center x . The union

$$S_r = \bigcup_{(x,y) \in V_X} S_r(x, y)$$

is called the tube of radius r over X . The set S_r is the tube of radius $\frac{\pi}{2} - r$ over the dual variety \check{X} .

If all the points of X are regular this definition coincides with one above.

The set of points $\text{sng } V_X \subset V_X$ such that $(x, y) \in \text{sng } V_X$ if $x \in \text{sng } X$ or $y \in \text{sng } \check{X}$ is a closed algebraic subvariety of V_X , $\text{reg } V_X = V_X \setminus \text{sng } V_X$ is an open set of V_X in the Zariski topology.

Let $X \subset \mathbf{C}P^n$ be a closed irreducible algebraic variety and let x_0 be a Zariski open set in X . Then the closure of x_0 in the classical topology is X [14].

Let us take the Segre map

$$\sigma: \mathbf{C}P^n \times \mathbf{C}\check{P}^n \rightarrow \mathbf{C}P^{(n+1)^2-1}.$$

Then $\sigma(V_X)$ is a closed irreducible algebraic variety in $\mathbf{C}P^{(n+1)^2-1}$ and the set $\text{reg } V_X$ is an open set of V_X in the Zariski topology.

As corollary we obtain the following result

Lemma 10. *The closure of the set $\text{reg } V_X \subset \mathbf{C}P^n \times \mathbf{C}P^n$ in the standard topology coincides with the tangent bundle V_X .*

Therefore the tube over X is the closure of the set

$$\bigcup_{(x,y) \in \text{Reg } V_X} S_r(x, y)$$

Lemma 11. [5] Let X be a compact topological space. Suppose A is a closed subset such that $X \setminus A$ is a smooth n -dimensional orientable manifold without boundary. Then

$$H_q(X, A) \simeq H^{n-q}(X \setminus A),$$

where H_i, H^i are homology and cohomology groups.

Lemma 12. [1] Suppose $J\xi$ is an eigenvector of the shape operator A_ξ of a Hopf hypersurface M in the complex projective space, with the corresponding eigenvalue $2 \cot 2\Theta$, $0 < \Theta < \frac{\pi}{2}$. Suppose $J\xi, X_2, \dots, X_n$ is a basis of principal vectors of A_ξ with $A_\xi X_j = \cot \Theta_j X_j$, $2 \leq j \leq n$, $0 < \Theta_j < \pi$; $\frac{\partial}{\partial t_j}$ ($2 \leq j \leq k$) are normal vectors. Then the shape operator A_r of the tube Φ_r is given in terms of its principal vectors by

- (a) $A_r \left(\frac{\partial}{\partial t_j} \right) = -\cot r \left(\frac{\partial}{\partial t_j} \right)$, $2 \leq j \leq k$;
- (b) $A_r (X_j, 0) = \cot (\Theta_j - r) (X_j, 0)$, $2 \leq j \leq n$;
- (c) $A_r (J\xi, 0) = \cot (2(\Theta - r)) (J\xi, 0)$.

For a complex hyperbolic space \mathbf{CH}^n the following analog of Lemma 1 holds:

Lemma 13. [13] Let M be an orientable Hopf hypersurface of \mathbf{CH}^n such that the principal curvature μ in the direction $J\xi$ is constant and equal to $\mu = 2 \coth 2r$. Suppose that Φ_r has constant rank q on M . Then for every point $x_0 \in M$ there exists an open neighbourhood U of x_0 such that $\Phi_r U$ is a $q/2$ -dimensional complex submanifold embedded in \mathbf{CH}^n .

Lemma 14. [15] Let Ω be a Hermitian complex manifold with exact fundamental form $\omega = d\gamma$. Let A be an analytical q -dimensional set with boundary $\partial A \subset \Omega$ such that $A \cup \partial A$ is compact.

Then

$$H^{2q}(A) \leq \frac{1}{q} (\max_{\partial A} |\gamma|) H^{2q-1}(\partial A),$$

where $H^{2q}(A), H^{2q-1}(\partial A)$ are Hausdorff measures, and

$$|\gamma|(z) = \max \{ |\gamma(v)| : v \in T_z \Omega, |v| = 1 \}.$$

Lemma 15. [8] Let M be a Hopf hypersurface of a complex space form $\overline{M}^n(c)$ ($c \neq 0$). If U is an eigenvector of A , then the principal curvature $\mu = g(AU, U)$ is constant.

3. Proofs of the Theorems

Let M_s be the set of points of M such that $\text{rank}(\Phi_r)_*(M_s) = s$, $F_s = \Phi_r(M_s)$, $F = \Phi_r(M)$. From Lemma 4 we obtain that if $X \in T_\alpha \subset T_P M$ where T_α is the eigenspace corresponding to the principal curvature $\alpha = \cot r$, then $JX \in T_\alpha$. Hence s is even and if $s < 2q$, then $s \leq 2q - 2$.

Let

$$E = \bigcup_{s < 2q} F_s \cup F_0$$

$$F_0 = \{x \in F : x = \Phi_r(L_1) = \Phi_r(L_2), L_1 \neq L_2 \subset M, \text{rank}(\Phi_r)_*(P_1) = \text{rank}(\Phi_r)_*(P_2) = 2q\},$$

for $P_i \in L_i$, where L_i are leaves of the distribution $\text{Ker}(\Phi_r)_*$.

Proof of the theorem 1. Let M be a compact Hopf hypersurface in \mathbf{CP}^n . This means that the vector $J\xi$ is a principal direction of M , where ξ is the unit normal vector and J is the complex structure in \mathbf{CP}^n . From Lemma 2 it follows that the corresponding principal curvature μ is constant, $\mu = 2 \cot 2r$. Let $2q$ be the maximal rank of $(\Phi_r)_*$ on M . Let $P \in M$ be a point such that $\text{rank}(\Phi_r)_*(P) = 2q$ and let M_{2q} be the corresponding connected component of M such that $P \in M_{2q}$ and for $Q \in M_{2q}$ $\text{rank}(\Phi_r)_*(Q) = 2q$. Set $F_{2q} = \Phi_r(M_{2q})$, $\tilde{F} = F_{2q} \cap (\mathbf{CP}^n \setminus E)$. From Lemma 1 we obtain that \tilde{F} is a purely analytic set, $\dim_z \tilde{F} = q$, $z \in \tilde{F}$.

Locally F_0 is a transversal intersection of two complex submanifolds of dimension q . Hence F_0 is an analytic set of real dimension $\leq 2q - 2$. Then its Hausdorff measure

$$H^{2q-1}(F_0) = 0.$$

Now apply Lemma 5 to the set $E_1 = \bigcup_{s < 2q} F_s$ and the map Φ_r . Then $\nu \leq 2q - 2$.

If the class of regularity of M is greater or equal to $2(n - q + 1)$ then the class of regularity of Φ_r is $k \geq 2(n - q + 1) - 1$ and

$$\nu + \frac{2n - 1 - \nu}{k} \leq 2q - 2 + \frac{2n - 1}{k} \leq 2q - 1,$$

for $k \geq 2n - 1$. From Lemma 5 we have $H^{2q-1}(E_1) = 0$ and so $H^{2q-1}(E) = 0$. From Lemma 6 we obtain that the closure of \tilde{F} is a purely q -dimensional analytic subset of $\mathbf{C}P^n$. Since any analytic subset is $*$ -analytic we get from Chow's Theorem (Lemma 7) that $\text{cl } \tilde{F} \subset \mathbf{C}P^n$ is a finite union of algebraic varieties. An analytic set A is an irreducible if and only if the set $\text{reg } A$ is connected. From Lemma 8 it follows that $\text{cl } \tilde{F}$ is irreducible as analytic set and we obtain that $\text{cl } \tilde{F} = X$ is an irreducible algebraic variety.

Let S_r be a tube over $X = \text{cl } \tilde{F}$. From Lemma 10 we have $S_r \subset M$ and $S_r = \text{cl } M_{2q}$. We will prove that $\text{cl } M_{2q} = M$. Suppose that $\text{cl } M_{2q} \neq M$. Then in every neighbourhood of a point $P \in \partial M_{2q}$ there exist points $Q \in M \setminus \text{cl } M_{2q}$. Let $P \in \partial M_{2q}$. Then $P \in S_r(x, y)$ such that $x \in \text{sng } X$, $y \in \text{sng } \check{X}$. Then

$$\partial M_{2q} = \bigcup_{x \in \text{sng } X, y \in \text{sng } \check{X}} S_r(x, y).$$

Otherwise some neighbourhood of P belongs to $\text{cl } M_{2q}$ and $P \in \text{int } \text{cl } M_{2q}$. The set of points

$$\text{sng}(X, \check{X}) = \text{sng } X \times \mathbf{C}P^n \cap \mathbf{C}P^n \times \text{sng } \check{X} \subset V_X \subset \mathbf{C}P^n \times \mathbf{C}P^n$$

is a closed algebraic subvariety of V_X . The dimension of $\text{sng}(X, \check{X}) \leq n - 2$ because the dimension of V_X is equal to $n - 1$. The set ∂M_{2q} is a fiber bundle over $\text{sng}(X, \check{X})$ with the circle S^1 as a leaf. The real dimension of $\text{sng}(X, \check{X})$ is $\leq 2(n - 2)$ whence

$$H_{2n-3}(\text{sng}(X, \check{X}), \mathbf{Z}) = 0.$$

For $E = \partial M_{2q}$, $B = \text{sng}(X, \check{X})$, $F = S^1$ the exact Thom-Gysin sequence has the form [17]

$$\begin{aligned} H_{2n-1}(\text{sng}(X, \check{X}), \mathbf{Z}) &\rightarrow H_{2n-3}(\text{sng}(X, \check{X}), \mathbf{Z}) \rightarrow \\ &\rightarrow H_{2n-2}(\partial M_{2q}, \mathbf{Z}) \rightarrow H_{2n-2}(\text{sng}(X, \check{X}), \mathbf{Z}), \\ 0 &\rightarrow 0 \rightarrow H_{2n-2}(\partial M_{2q}, \mathbf{Z}) \rightarrow 0. \end{aligned}$$

We obtain

$$H_{2n-2}(\partial M_{2q}, \mathbf{Z}) = 0.$$

Next, we apply Lemma 11 with $X = M$, $A = \partial M_{2q}$. Then

$$H_{2n-1}(M, \partial M_{2q}) = H^0(M \setminus \partial M_{2q}).$$

But $M \setminus \partial M_{2q}$ has $m > 1$ connected components and

$$H^0(M \setminus \partial M_{2q}, \mathbf{Z}) = \bigoplus_{i=1}^m \mathbf{Z}$$

is the direct sum of m copies of \mathbf{Z} [17].

For the pair $(M, \partial M_{2q})$ the exact homology sequence has the following form

$$\begin{aligned} H_{2n-1}(\partial M_{2q}, \mathbf{Z}) &\rightarrow H_{2n-1}(M, \mathbf{Z}) \rightarrow H_{2n-1}(M, \partial M_{2q}, \mathbf{Z}) \rightarrow \\ &\rightarrow H_{2n-2}(\partial M_{2q}, \mathbf{Z}); \end{aligned}$$

$$H_{2n-1}(\partial M_{2q}, \mathbf{Z}) = H_{2n-2}(\partial M_{2q}, \mathbf{Z}) = 0; \quad H_{2n-1}(M, \mathbf{Z}) = \mathbf{Z}.$$

It follows that $H_{2n-1}(M, \partial M_{2q}, \mathbf{Z}) = \mathbf{Z}$. This contradicts to above result. Thus $\text{cl } M_{2q} = M$ and M is a tube over the irreducible algebraic variety $\text{cl } \tilde{F} = X$.

Proof of the theorem 2. Let S be the hypersphere of the minimal radius r_0 such that the hypersurface M is contained in the ball D with boundary $\partial D = S$. Let P be a point of tangency of M and S . Let ξ be the inward unit normal vector at the point P . Then the principal curvature in the direction $J\xi$ is $\mu = 2\cot 2\rho \geq 2\cot 2r_0$, and so $\rho \leq r_0 < \pi/2$. Another principal curvature $k_i = \cot \Theta_i$ at the point P satisfies the conditions $\cot \Theta_i \geq \cot r_0$, where $2\cot 2r_0, \cot r_0$ are principal curvatures of the hypersphere S . Then $\Theta_i \leq r_0$. Let $r = \rho - \pi/2$. From Lemma 12 we obtain that the principal curvatures of the tube Φ_r over M are equal to

$$(k_i)_r = \text{tg}(\rho - \Theta_i) \leq \text{tg}(r_0 - \Theta_i) < \infty.$$

Hence $\text{rank}(\Phi_r)_*(P) = 2(n-1)$ and from Theorem 1 we get that $\Phi_r(M) = \text{cl } \tilde{F} = X$ is an irreducible hypersurface of degree d . Let X_k be a sequence of smooth algebraic hypersurfaces such that $\lim X_k = X$, $\text{degree } X_k = d$ [7], and let \check{X}, \check{X}_k be dual algebraic varieties. Then

$$M = \Phi_{\frac{\pi}{2}-r}(X) = \Phi_r(\check{X})$$

and from Lemma 9 we get that $\check{X} = \lim \check{X}_k$. From the above for $\Phi_{\frac{\pi}{2}-r}(X_k) = M_k$,

$$\lim M_k = M.$$

For large k , M_k is contained in the balls D_k of radius $R < \pi/2$ and M_k does not intersect complex projective space $x_0 = 0$.

Let $f = 0$ be the equation of the algebraic hypersurface X_n where f is a homogeneous polynomial, $\text{grad } f \neq 0$. By Bezou Theorem [15] the system of equations

$$x_0 = 0, \quad f = 0, \quad f_{x_0} = 0$$

has a nontrivial solution for n is ≥ 3 and degree of the polynomial $f \geq 2$. This means that M_k intersects the hyperplane $x_0 = 0$. It follows that f is a linear function and the X_k are hyperplanes, M_k are hyperspheres. Then the hypersurface M is a geodesic hypersphere too.

For $n = 2$ the equation of the tube has the following parametric form

$$z_j = x_j \cos r + \sin r \frac{\frac{\partial f}{\partial x_j}}{|\text{grad } f|} e^{it};$$

x_j are coordinates of points of the algebraic variety, $0 \leq t \leq 2\pi$; $0 \leq r \leq \frac{\pi}{2}$, r is radius of the tube Φ_r ; $j = 0, 1, 2$.

From the real point of view X is a compact two-dimensional manifold.

Denote

$$g_1 = |x_0 \cos r|, \quad g_2 = \left| \frac{\frac{\partial f}{\partial x_0}}{|\text{grad } f|} \sin r \right|,$$

If the degree of the polynomial f is ≥ 2 the zero sets of these regular functions on the manifold X are non empty on the manifold X . Hence there exists a point $P \in X$ such that $g_1 = g_2 = \rho$. Then $z_0 = \rho (e^{i\alpha} + e^{i(\beta+t)})$. Moreover, if $t = \alpha - \beta - \pi$ then $z_0 = 0$.

This means that M_k intersects the hyperplane $x_0 = 0$.

Thus f is a linear function and M_k and M are geodesic hyperspheres as in the case $n \geq 3$.

Proof of the theorem 3. Let S be the hypersphere of the minimal radius r_0 such that the hypersurface M is contained in the ball D with boundary S . Let P_0 be a point of tangency of M and S . Let ξ be the inward unit normal vector of M at the point P_0 . From Lemma 15 it follows that the principal curvature μ in the direction $J\xi$ is constant. At the point P_0 this curvature satisfies the inequality $\mu \geq 2\coth 2r_0$ and $\mu = 2\coth 2r$. We now follow the proof of Theorem 1, using Lemma 13 instead of

Lemma 1. Consider the map Φ_r . For a Hopf hypersurface $\text{rank}(\Phi_r)_*$ is always even. This follows from Lemma 4.

Suppose $2q$ is the maximal rank of $(\Phi_r)_*$ at the points of M . Let $P \in M$ be a point such that $\text{rank}(\Phi_r)_*(P) = 2q$ and M_{2q} is the connected component of M such that for $Q \in M_{2q}$ $\text{rank}(\Phi_r)_*(Q) = 2q$. As in the proof of Theorem 1, set

$$F = \Phi_r(M), \quad F_{2q} = \Phi_r(M_{2q}), \quad F_s = \Phi_r(M_s),$$

$$E = F_0 \bigcup_{s < 2q} F_s; \quad \tilde{F} = F_{2q} \cap \mathbf{C}H^n \setminus E.$$

We obtain that $\text{cl} \tilde{F} = X$ is a compact analytic set in $\mathbf{C}H^n$ with boundary $\partial X \subset E$. The Hausdorff measure $H^{2q-1}(\partial X) = 0$. From Lemma 14 it follows that $H^{2q}(X)$ is equal to 0. This is possible only if $q = 0$ and X is a point. Then M is a tube over a point and M is a geodesic hypersphere.

References.

- [1] *T.E. Cecil, P.J. Ryan, " Focal sets and real hypersurfaces in complex projective space"*, Trans. Amer. Math. Soc., 269 (1982) 481–499
- [2] *E.M. Chirka, " Complex analytic sets"*, Nauka, Moscow, 1985
- [3] *E.M. Chirka, Complex analytic sets. Encyclopaedia of Mathematical Sciences. Several complex variables I*, A.G.Vitushkin (ed), 7 Springer–Verlag, 1990
- [4] *H. Federer, Geometric measure theory*, Springer–Verlag, 1969
- [5] *A.T. Fomenko, D.B. Fuchs, Homotopic Topology* Nauka, Moscow, 1989
- [6] *J. Harris, Algebraic Geometry*, Springer–Verlag, 1996
- [7] *W.V.D. Hodge and D.Pedoe, Methods of algebraic geometry*, Cambridge, 1 (1947)
- [8] *U-Hang Ki, Young Jin Suh On real hupersurfaces of a complex space form*, Math. J. Okayama Univ. , 32 (1990) , 207-221
- [9] *M. Kon, Pseudo–Einstein real hypersurfaces in complex space forms*, J. Dif. Geometry , 14 (1979) , 339-354
- [10] *K. Lamotke, The topology of complex projective varieties after S. Lefschetz*, Topology , 20 (1981) , 15–51
- [11] *Y. Maeda, On real hypersurfaces of a complex projective space*, J. Math. Soc. Japan. , 28 (1976) , 529–540
- [12] *V. Miquel, Compact Hopf hypersurfaces of constant mean curvature in complex space forms*, Annals of Global Analysis and Geometry , 12 (1994) , 211–218
- [13] *S. Montiel, Real hypersurfaces of a complex hyperbolic space*, J. Math. Soc. Japan , 37 (1985) , 515–535
- [14] *D. Mumford, Algebraic geometry. I, Grundlehren der mathematischen Wissen- schaften , Springer–Verlag, Berlin–Heidelberg–New York , 221 (1976) , 189*
- [15] *I.R. Shafarevich, Basis of algebraic geometry*, Springer Verlag, 1977
- [16] *B. Shiffman, On the removal of singularities of analytic sets* Michig. Math. J., 15 (1968) , 111–120
- [17] *E. Spanier, Algebraic Topology*, Mc GRAW-HILL Book Company, New York – London, 1966