# <span id="page-0-0"></span>The Calderón Projection: New Definition and Applications

Bernhelm Booß-Bavnbek, Matthias Lesch, and Chaofeng Zhu

ABSTRACT. We consider an arbitrary linear elliptic first–order differential operator A with smooth coefficients acting between sections of complex vector bundles  $E, F$  over a compact smooth manifold M with smooth boundary  $\Sigma$ . We describe the analytic and topological properties of  $A$  in a collar neighborhood U of  $\Sigma$  and analyze various ways of writing  $A \upharpoonright U$  in product form. We discuss the sectorial projections of the corresponding tangential operator, construct various invertible doubles of A by suitable local boundary conditions, obtain Poisson type operators with different mapping properties, and provide a canonical construction of the Calderón projection. We apply our construction to generalize the Cobordism Theorem and to determine sufficient conditions for continuous variation of the Calderón projection and of well-posed selfadjoint Fredholm extensions under continuous variation of the data.

### CONTENTS



<sup>2000</sup> Mathematics Subject Classification. Primary 58J32; Secondary 35J67, 58J50, 57Q20.

Key words and phrases. Calderón projection, Cauchy data spaces, cobordism theorem, continuous variation of operators and boundary conditions, elliptic differential operator, ellipticity with parameter, Lagrangian subspaces, regular boundary value problem, sectorial projection, selfadjoint Fredholm extension, Sobolev spaces, symplectic functional analysis.

This work was supported by the network "Mathematical Physics and Partial Differential Equations" of the Danish Agency for Science, Technology and Innovation. The second named author was partially supported by Sonderforschungsbereich/Transregio "Symmetries and Universality in Mesoscopic Systems" (Bochum–Duisburg/Essen–Köln–Warszawa) and the Hausdorff Center for Mathematics (Bonn). The third named author was partially supported by FANEDD 200215, 973 Program of MOST, Fok Ying Tung Edu. Funds 91002, LPMC of MOE of China, and Nankai University.



2 BERNHELM BOOSS-BAVNBEK, MATTHIAS LESCH, AND CHAOFENG ZHU

#### LIST OF FIGURES

<span id="page-1-0"></span>

# 1. Introduction

<span id="page-1-1"></span>This paper is about basic analytical properties of elliptic operators on compact manifolds with smooth boundary. Our main achievements are (i) to develop the basic elliptic analysis in full generality, and not only for the generic case of operators of Dirac type in product metrics (i.e., we assume neither constant coefficients in normal direction nor symmetry of the tangential operator); (ii) to establish the cobordism invariance of the index in greatest generality; and (iii) to prove the continuity of the Calderón projection and of related families of global elliptic boundary value problems under parameter variation. We take our point of departure in the following observations.

<span id="page-2-0"></span>1.1. Dirac operator *folklore*. Most analysis of geometrical and physical problems involving a Dirac operator  $A$  on a compact manifold  $M$  with smooth boundary  $\Sigma$  acting on sections of a (complex) bundle E seems to rely on quite a few basic facts which are part of the shared folklore of people working in this field of global analysis (e.g., see Booß–Bavnbek and Wojciechowski [[BBW93](#page-56-1)] for properties (WiUCP), (InvDoub) and (Cob), and Nicolaescu [[Nic97](#page-57-0), Appendix] and Booß–Bavnbek, Lesch and Phillips [[BBLP05](#page-56-2)] for property (Param)):

- WiUCP: the weak inner unique continuation property (also called *weak* UCP to the boundary), i.e., there are no nontrivial elements in the null space ker A vanishing at the boundary  $\Sigma$  of M;
- **InvDoub:** the existence of a suitable elliptic invertible continuation  $\widetilde{A}$  of  $A$ . acting on sections of a vector bundle over the closed double or another suitable closed manifold  $M$  which contains  $M$  as submanifold; this yields a Poisson type operator  $K_{+}$  which maps sections over the boundary into sections over M; and a precise Calderón projection  $C_+$ , i.e., an idempotent mapping of sections over the boundary onto the Cauchy data space which consists of the traces at the boundary of elements in the nullspace of A (possibly in a scale of Sobolev spaces);
- Cob: the existence of a selfadjoint regular Fredholm extension of any total (formally selfadjoint) Dirac operator  $A$  in the underlying  $L^2$ -space with domain given by a pseudodifferential boundary condition; that implies the vanishing of the signature of the associated quadratic form, induced by the leading symbol in normal direction at the boundary; moreover, that actually is equivalent to the Cobordism Theorem asserting a canonical splitting of the induced tangential operator  $B = B^+ \oplus B^-$  over  $\Sigma$  with  $ind B<sup>+</sup> = 0;$
- Param: the continuous dependence of a family of operators, their associated Calderón projections, and of any family of well-posed (elliptic) boundary value problems on continuous or smooth variation of the coefficients.

<span id="page-2-1"></span>1.2. In search of generalization. With the renewed interest in geometrically defined elliptic operators of first order of general type, arising, e.g., from perturbations of Dirac operators, we ask to what extent the preceding list can be generalized to arbitrary linear elliptic differential operators with smooth coefficients. It is hoped that the results of this paper can serve as guidelines for similar constructions and results for hypo- and sub-elliptic operators where the symbolic calculus is not fully available.

There are immediate limits for generalization of some of the mentioned features by counter examples: UCP, even weak inner UCP may not hold for arbitrary elliptic systems of first order, see indications in that direction in Plis  $[Pli61, Corollary 1]$  $[Pli61, Corollary 1]$  $[Pli61, Corollary 1]$ , p. 610] and the first-order Alinhac type counterexample to strong UCP [Bär99, Example, p. 184]. Moreover, from just looking at the deficiency indices, we see that the formally selfadjoint operator  $i\frac{d}{dx}$  on the positive line does not admit a selfadjoint extension. This example is instructive because, quite opposite to the half-infinite domain, on a bounded one-dimensional interval any system of first-order differential equations satisfies property (Cob) by a deformation argument.

We go through the list.

*Property (WIUCP).* It seems that the precise domain of validity is unknown. The local stability of weak inner UCP has been obtained by Booß–Bavnbek and Zhu in [[BBZ05](#page-57-2), Lemma 3.2]. In spite of the local definition of UCP, the property (WiUCP) has a threefold global geometric meaning: (i) there are no ghost solutions, i.e., each section  $u \in \Gamma^\infty(M; E)$  belonging to the null space ker A over the manifold M has a non-trivial trace  $u \nvert \Sigma$  at the boundary; (ii) equivalently, the maximal extension  $A_{\text{max}} = (A_{\text{min}}^t)^*$  is surjective in  $L^2(M, E)$  for densely defined closed minimal  $A_{\min}$ :  $\mathscr{D}(A_{\min}) = H_0^1(M, E) \to L^2(M, E)$ ; and (iii), as noted by Booß–Bavnbek and Furutani in [[BBF98](#page-56-3), Section 3.3] and in various follow-ups, it seems that assuming weak inner UCP of  $A$  and  $A<sup>t</sup>$  is mandatory for obtaining the continuity of Cauchy data spaces and the continuous change of the Calderón projection under variation of the coefficients, i.e., property (Param).

*Property (InvDoub).* Different approaches are available: one approach ( $[BBW93$  $[BBW93$ , Chapter 9]) has been the gluing of A and its formal adjoint  $A<sup>t</sup>$  to an invertible elliptic operator  $\tilde{A}$  over the closed double  $\tilde{M}$ . This construction is explicit, if the metric structures underlying the Dirac operator's definition are product near the boundary. In the selfadjoint case, it yields at once the Lagrangian property of the Cauchy data space in the symplectic Hilbert space  $L^2(\Sigma, E_{\Sigma})$  of square integrable sections in  $E_{\Sigma} := E[\Sigma]$ . Then (Cob) follows.

Property (InvDoub) generalizes to the non-product case for operators of Dirac type and, as a matter of fact, for any elliptic operator satisfying weak inner UCP under the somewhat restrictive condition that the tangential operator has selfadjoint leading symbol. Here the trick is that this condition permits the prolongation of the given operator to a slightly larger manifold  $M'$  with boundary reaching constant coefficients in normal direction close to the new boundary and maintaining UCP under the prolongation (as well as formal selfadjointness of the coefficients, if present at the old boundary). This is explained in the Appendix.

But what can be done for general elliptic operators? A very general and elegant construction of the Calderón projection was given by Hörmander in  $[H\ddot{o}r85, Theo$ rem 20.1.3] on the symbol level. Unfortunately, he obtains only an almost projection (up to smoothing operators) which limits its applicability in our context.

In this paper, we shall exploit another general definition of the Calderón projection which is due to Seeley [[See69](#page-57-4), Theorem 1 and Appendix, Lemma]. Seeley's construction provides a precise projection, not only an approximate one, and does not require UCP. First he replaces A by an invertible operator  $A_1$  by adding the projection onto the finite–dimensional space of inner solutions. Then he extends the operator  $D:=\begin{pmatrix} 0 & A_1^t \ A_1 & 0 \end{pmatrix}$ ) to the closed double  $\widetilde{M}^{\prime}$  of a slightly extended manifold  $M'$  with boundary. In general, such a prolongation may, however, destroy weak inner UCP even when UCP was established on the original manifold. Seeley constructs on the symbol level (and by adding a suitable correction term) an elliptic extension  $\tilde{D}$  of D over the whole of the closed manifold  $\tilde{M}'$  which is always invertible. He shows that  $\tilde{D}$  provides the wanted Poisson operator and a truly pseudodifferential Calderón projection  $\mathscr{P}_+$  along the original boundary  $\Sigma$ . If the tangential operator is formally selfadjoint then  $\mathscr{P}_+$  has selfadjoint leading symbol and can be replaced by the orthogonal projection which is also pseudodifferential and has the same symbol (and may be denoted by the same letter  $\mathscr{P}_+$ ). In this way, the choices in the *construction of the invertible double* are removed totally, as the operation of the resulting  $\mathscr{P}_+$  is concerned. This makes  $\mathscr{P}_+$  a good candidate for property (Cob).

However, Seeley's general construction, similarly the recent Grubb [[Gru09](#page-57-5), Section 11.1], in difference to the simple gluing in the case of Dirac type operators of [[BBW93](#page-56-1), Chapter 9], does not immediately lead to the Lagrangian property of im  $\mathscr{P}_+$ . Moreover and more seriously, when working with curves of elliptic problems Seeley's construction does not give a hint under what conditions the Calderón projections vary continuously when varying a parameter. There are too many choices involved in Seeley's construction.

<span id="page-4-0"></span>1.3. Our present approach. This motivates our present approach (inspired by Himpel, Kirk and Lesch [[HKL04](#page-57-6)]), namely the construction of the invertible double as a canonically given local boundary problem for the double D exactly on the original manifold  $M$ , without any choices, prolongations etc. This leaves us with full control of the UCP situation; leads directly to the wanted Fredholm Lagrangian property (Cob); and, moreover and here most decisively, provides explicit formulas for treating the parameter dependence in property (Param).

This program is opened in Section [2](#page-5-0) by explaining our basic choice of product structures near the boundary for the sake of comprehensible analysis, even if the original geometric structures are non-product; moreover, for the convenience of the reader and for fixing our notation we summarize a few basic facts about regular boundary conditions.

To begin with, we do not assume selfadjoint leading symbol of the tangential operator  $B_0$  nor constant coefficients  $B_x = B_0$  along an inward coordinate x. Most of our estimates depend on the single fact that  $B_0 - \lambda$  is parameter dependent elliptic for  $\lambda$  in a conic neighborhood of iR in the sense of Shubin [[Shu80](#page-57-7), Section II.9]. More precisely, we depend on the related concept of sectorial spectral projections introduced in 1970 by Burak [[Bur70](#page-57-8)] and recently further developed in Ponge [[Pon06](#page-57-9), Section 3] as part of the current upsurge of interest in spectral properties of non-selfadjoint elliptic operators. Because of our interest in the continuous dependence of this kind of generalized positive spectral projections on the input data we found it necessary to develop the concept of sectorial projections once again from scratch. This is done in Section [3](#page-11-0) where we develop an abstract Hilbert space framework for the concept of sectorial projections and apply it to tangential operators perceived as parametric elliptic operators.

In Section [4](#page-19-0) we provide the construction and the relevant properties of the invertible double yielded by a local elliptic boundary value problem, induced by fixing an invertible bundle homomorphism  $T$  over  $\Sigma$ .

In Section [5](#page-23-0) we establish suitable Sobolev regularity of the inverse operator which leads to the definition and basic properties of Poisson operator and Calderón projection, both definitions made dependent of the choice of the above mentioned homomorphism  $T$ . We shall show that the range of the Calderón projection does not depend on the choice of  $T$  and is, in fact, equal to the Cauchy data space. That yields the relation between the *canonical* Calderón projection defined as orthogonal projection onto the Cauchy data space, and our *relative* Calderón projections, which depend on T. However, it is not the canonical, but only the relative definition that establishes the Lagrangian property of the Cauchy data space and its continuous dependence of the coefficients for general elliptic differential operators of first order.

That is the subject of the two closing sections of this paper, which present the fruits of the analysis endeavour of Sections 1-5.

Property (Cob). In Section [6,](#page-30-0) we give a first application of our construction of the Calderón projection: we give our reading of Ralston  $[Ral70]$  $[Ral70]$  $[Ral70]$  and infer that the arguments of this 1970 paper establish the following findings for any formally selfadjoint differential operator  $A$  over a compact manifold  $M$  with smooth boundary Σ:

- the existence of a selfadjoint Fredholm extension  $A_P$  given by a pseudodifferential boundary condition P;
- the vanishing of the signature of  $i\omega$  on the space  $V(B_0)$  of eigensections to purely imaginary eigenvalues of the tangential operator  $B_0$  of A over the boundary (or on ker  $B_0$  in the case that  $B_0$  is formally selfadjoint); here  $i\omega$  denotes the form induced by the Green form of A on the symplectic

von Neumann space  $\beta(A) := \mathscr{D}(A_{\text{max}})/\mathscr{D}(A_{\text{min}})$ , i.e., the leading symbol of A over the boundary  $\Sigma$  in normal direction;

• and, equivalently, but seemingly never recognized by people working in global analysis, the General Cobordism Theorem, stating that the index of any elliptic differential operator  $B^+$  over a closed manifold  $\Sigma$  must vanish. if  $B^+$  can be written as the left lower corner of a formally selfadjoint tangential operator over  $\Sigma$  induced by an elliptic formally selfadjoint A on a smooth compact manifold M with  $\partial M = \Sigma$ .

Property (Param). In Section [7,](#page-41-1) as a second application of our construction of the Calderón projection, we establish that property in great generality. Roughly speaking, we let an operator family  $(A_z)$  and the family  $(A_z^t)$  of formally adjoint operators vary continuously in the operator norm from  $L_1^2(M)$  to  $L^2(M)$  and assume that the leading symbol  $(J_0(A_z))$  of  $(A_z)$  over the boundary  $\Sigma$  in normal direction also varies continuously in the  $L^2_{1/2}(\Sigma)$  operator norm with z running in a parameter space Z. We assume for all  $A_z$  and  $A_z^t$  property (WiUCP) or, almost equivalently, that the dimensions of the spaces of "ghost solutions" without trace at the boundary remain constant under the variation. Then in Theorems [7.9](#page-46-2) and [7.2a](#page-42-0) we show that the inverse of the "invertible double" and, under slightly sharpened continuity, the Poisson operator in respective operator norms vary continuously; and so does the resolvent of a family  $(A_{zP_z})$  of well-posed Fredholm extensions of now formally selfadjoint  $(A_z)$  with orthogonal pseudodifferential projections  $(P_z)$ varying in  $L^2_{1/2}(\Sigma)$  operator norm.

Unfortunately, we can neither prove nor disprove the continuous variation of the Calderón projection in the same generality. However, if the leading symbol of the tangential operator is selfadjoint, we can prove the continuous variation of the sectorial projection (Proposition  $7.15$ ) and so (Corollary  $7.4$ ) of the Calderón projections by our correction formula [\(5.31\)](#page-28-0) and Theorem [7.2b](#page-42-0). Our Proposition [7.13](#page-48-1) shows that the difficulties for proving continuous variation of the sectorial projection disappear also for non-selfadjoint leading symbol, if the variation is of order  $< 1$ .

In Appendix A we discuss various special cases with emphasis on constant coefficients in normal direction in a collar around the boundary.

The main results of this paper have been announced in [[BBL09](#page-56-4)].

# <span id="page-5-0"></span>2. Elliptic differential operators of first order on manifolds with boundary

<span id="page-5-1"></span>2.1. Product form and metric structures near the boundary. We shall begin with a basic observation: Dirac operators emerge from a Riemannian structure on the manifold and a Hermitian metric on the vector bundle (together with Clifford multiplication and a connection). Talking about a general differential operator it is in our view very misleading to pretend that the operator will depend on metrics and such. All we need is the operator and an  $L^2$ -structure on the sections. The latter basically only requires a density (take dvol in the Riemannian case) on the manifold and a metric on the bundle. In this paper, we prefer to choose metrics and such as simple as possible and push all complications into the operator.

The message is this: we can always work in the product case and do have to worry only about the operator. In detail:

Let M be a compact manifold with boundary and  $\pi : E \to M$  a vector bundle. Given a Hermitian metric h on E and a Riemannian metric g on M we can form the Hilbert space  $L^2(M, E; g, h)$  which is the completion of  $\Gamma_0^{\infty}(M \setminus \partial M; E)$  with regard to the scalar product

<span id="page-6-2"></span>(2.1) 
$$
\langle u, v \rangle_{g,h} := \int_M h(u(x), v(x)) \, \mathrm{dvol}_g(x).
$$

The space of smooth sections of the vector bundle E over M is denoted by  $\Gamma^{\infty}(M; E)$ ; the corresponding space of smooth compactly supported sections is denoted by  $\Gamma_0^{\infty}(M;E)$ . Given another Riemannian metric  $g_1$  and another Hermitian metric  $h_1$ on E there is a smooth positive function  $\rho \in C^{\infty}(M)$  such that

$$
(2.2) \t\t dvol_{g_1} = \varrho \, dvol_g,
$$

and there is a unique smooth section  $\theta \in \Gamma^\infty(M; \text{End } E)$  such that for  $x \in M, \xi, \eta \in$  $E_x$  we have

<span id="page-6-0"></span>(2.3) 
$$
h_{1,x}(\xi,\eta) = h_x(\theta(x)\xi,\eta), \qquad h_x(\theta(x)\xi,\eta) = h_x(\xi,\theta(x)\eta).
$$

With regard to h the operator  $\theta(x)$  is selfadjoint and positive definite, thus with regard to *n* the operator  $\sigma(x)$  is senadjoint and positive definite, thus we may form  $\sqrt{\theta}$  which is again a smooth selfadjoint and positive definite section of (End E, h). It is clear that [\(2.3\)](#page-6-0) determines  $\theta(x)$  uniquely and the claimed smoothness of  $x \mapsto \theta(x)$  can be checked easily in local coordinates. In sum, we find for  $u, v \in \Gamma^{\infty}(M; E)$ 

(2.4)  
\n
$$
\langle u, v \rangle_{g_1, h_1} = \int_M h_1(u(x), v(x)) \, dvol_{g_1}(x)
$$
\n
$$
= \int_M h(\theta(x)u(x), v(x)) \varrho(x) \, dvol_g(x)
$$
\n
$$
= \langle \sqrt{\varrho \theta} u, \sqrt{\varrho \theta} v \rangle_{g, h}.
$$

Thus we arrive at

<span id="page-6-1"></span>Lemma 2.1. The map

$$
\Psi : \Gamma^\infty(M; E) \longrightarrow \Gamma^\infty(M; E), \quad u \longmapsto \sqrt{\varrho \theta}^{-1} u
$$

extends to an isometry from  $L^2(M,E;g,h)$  onto  $L^2(M,E;g_1,h_1)$ .

Now assume that we are given a differential operator A in  $L^2(M,E;g,h)$  of first order. It may be a Dirac operator which is constructed from the metrics  $q$ and  $h$ . q and  $h$  may be wildly non–product near the boundary. Suppose there are metrics  $g_1$ ,  $h_1$  which we like more, e.g., product near the boundary. Then consider the differential operator  $\Psi A \Psi^{-1}$  in  $L^2(M, E; g_1, h_1)$ .  $\Psi A \Psi^{-1}$  is still a differential operator and since  $\Psi$  is unitary all spectral properties are preserved.

Let us be even more specific and choose a neighborhood U of  $\partial M =: \Sigma$  and a diffeomorphism  $\phi: U \to [0,\varepsilon) \times \Sigma$  with  $\phi \upharpoonright \Sigma = \text{id}_{\Sigma}$ . Furthermore, we choose a metric  $g_1$  on M such that

<span id="page-6-3"></span>(2.5) 
$$
\phi_* g_1 = dx^2 \oplus g_{\Sigma},
$$

$$
g_{\Sigma} := g | \Sigma,
$$

is a product metric which induces the same metric on the boundary as  $g$ . Here  $x$ denotes the normal inward coordinate near the boundary in the metric  $q_1$ .

 $\phi$  is covered by a bundle isomorphism  $\mathscr{F} : E \restriction U \to [0,\varepsilon) \times E_{\Sigma}, E_{\Sigma} := E \restriction \Sigma$ , i.e., we have the commutative diagram

(2.6) 
$$
E\upharpoonright U \xrightarrow{\mathscr{F}} [0,\varepsilon) \times E_{\Sigma}
$$

$$
\downarrow^{\pi} \qquad \qquad \downarrow^{\text{id} \times \pi}
$$

$$
U \xrightarrow{\phi} [0,\varepsilon) \times \Sigma.
$$

Likewise, we may now choose a metric  $h_1$  on E such that  $h_1(x) := \mathscr{F}_* h_1\{x\} \times \Sigma$  $h_1 \, \Sigma = h \, \Sigma =: h_\Sigma$  is independent of  $x \in [0, \varepsilon)$ . The mappings  $\mathscr{F}$  and  $\phi$  induce a map

(2.7) 
$$
\Psi_1: \Gamma^\infty(U;E) \longrightarrow C^\infty([0,\varepsilon),\Gamma^\infty(E_\Sigma)) \nf \longmapsto \left(x \mapsto (p \mapsto \mathscr{F}(f(\phi^{-1}(x,p))))\right),
$$

which extends to a unitary isomorphism  $L^2(U,E;g_1,h_1) \to L^2([0,\varepsilon],L^2(\Sigma,E_{\Sigma};g))$  $\Sigma, h(\Sigma)$ ). On  $\Sigma$  and  $E_{\Sigma}$  we have the fixed metrics  $g_{\Sigma}$  respectively  $h_{\Sigma}$  and we will suppress the reference to them in the notation.

Together with the unitary isomorphism  $\Psi$  of Lemma [2.1](#page-6-1) we obtain the claimed isometry

<span id="page-7-4"></span>(2.8) 
$$
\Phi := \Psi_1 \circ \Psi : L^2(U, E; g, h) \longrightarrow L^2([0, \varepsilon], L^2(E_{\Sigma})).
$$

Now  $\Phi A \Phi^{-1}$  is a first order differential operator in the product Hilbert space  $L^2([0, \varepsilon)) \otimes L^2(\Sigma, E_{\Sigma}; g_{\Sigma}, h_{\Sigma})$  and hence it takes the form

<span id="page-7-3"></span>(2.9) 
$$
D := \Phi A \Phi^{-1} =: J\Big(\frac{d}{dx} + B\Big)
$$

with a bundle endomorphism  $J \in C^{\infty}([0,\varepsilon), \Gamma^{\infty}(\Sigma; \text{End } E_{\Sigma}))$  and a smooth family of first order differential operators  $B \in C^{\infty}([0,\varepsilon), \text{Diff}^{1}(\Sigma; E_{\Sigma}))$ ; Diff<sup>1</sup> $(\Sigma; E_{\Sigma})$ denoting the space of first order differential operators acting on sections of  $E_{\Sigma}$ . For the moment we consider here only the smooth case, but so far one can replace "smooth" by "continuous" or "Lipschitz" or whatever.

Let us repeat: now all metric structures are product near the boundary and we do not have to worry about them. If we start, e.g., with a Dirac operator A on a Riemannian manifold with non–product metric the 'non–product situation' is reflected in the varying coefficients of D. From now on we have to worry only about those varying coefficients and nothing else.

After these somewhat pedagogical remarks we are ready to formulate the general set–up of the paper.

<span id="page-7-0"></span>2.2. The general set–up. We are going to fix some notation which will be used throughout the paper. Assume that the following data are given:

- a compact smooth Riemannian manifold  $(M, g)$  with smooth boundary  $\Sigma := \partial M$ .
- Hermitian vector bundles  $(E, h^E), (F, h^F)$ ,
- a first order elliptic differential operator

(2.10) 
$$
A: \Gamma^{\infty}(M; E) \longrightarrow \Gamma^{\infty}(M; F).
$$

•  $A^t: \Gamma^\infty(M; F) \longrightarrow \Gamma^\infty(M; E)$  denotes the formal adjoint of A with respect to the metrics  $g, h^E, h^F$ .

For further reference we record Green's formula for A.

<span id="page-7-2"></span>**Lemma 2.2.** Let  $\nu \in \Gamma^\infty(\Sigma; TM\vert \Sigma)$  be the outward normal vector field. Then we have with the notation of [\(2.1\)](#page-6-2) and [\(2.5\)](#page-6-3) for  $u \in \Gamma^\infty(M; E)$ ,  $v \in \Gamma^\infty(M; F)$ 

<span id="page-7-1"></span>(2.11) 
$$
\langle Au, v \rangle_{g,h} - \langle u, A^t v \rangle_{g,h}
$$

$$
= \frac{1}{i} \int_{\Sigma} h_{\Sigma} (\sigma_A^1(\nu^b) u | \Sigma, v | \Sigma) dvol_{g_{\Sigma}} = -\langle J(0) u | \Sigma, v | \Sigma \rangle_{g_{\Sigma}, h_{\Sigma}},
$$

where  $i := \sqrt{-1}$ ,  $\nu^b$  denotes the cotangent vector field corresponding to  $\nu$  in the metric g, and  $\sigma_A^1$  denotes the leading symbol of A.

Note that  $\phi_* \nu = -\frac{d}{dx}$ . Recall also from the previous Subsection [2.1](#page-5-1) that by construction all transformations are trivial on the boundary, that is,

(2.12) 
$$
\phi \upharpoonright \Sigma = \text{id}, \quad \mathscr{F} \upharpoonright E_{\Sigma} = \text{id}, \n(\Psi_1 f) \upharpoonright \Sigma = f \upharpoonright \Sigma, \quad (\Psi f) \upharpoonright \Sigma = f \upharpoonright \Sigma, \n(\Phi f) \upharpoonright \Sigma = f \upharpoonright \Sigma.
$$

We consider A as an unbounded operator between the Sobolev (and Hilbert) spaces

(2.13) 
$$
L_s^2(M, E; g, h^E), \quad L_s^2(M, F; g, h^F), \quad s \ge 0.
$$

If A acts as an unbounded operator between the Hilbert spaces  $H_1, H_2$ , we denote by A<sup>∗</sup> its functional analytic adjoint. For 0th order operators and for elliptic operators on closed manifolds the distinction between formal adjoint and (true) adjoint does not really matter; so in this case we use both notations interchangeably.

The closure of  $A \rvert \Gamma_0^{\infty}(M \setminus \Sigma; E)$  in  $L^2$  is denoted by  $A_{\min}$  and we put

(2.14) 
$$
\mathscr{D}(A_{\max}) := \{ f \in L^2 \mid Af \in L^2 \}
$$

the domain of an unbounded operator T will always be denoted by  $\mathscr{D}(T)$ . As ex-plained in Subsection [2.1](#page-5-1) there exists a collar  $U \approx [0, \varepsilon) \times \Sigma$  and linear isomorphisms

,

.

<span id="page-8-1"></span>(2.15) 
$$
\Phi^G : \Gamma^\infty(U;G) \longrightarrow C^\infty([0,\varepsilon),\Gamma^\infty(G_\Sigma)), \quad G = E, F,
$$

which extend to isometries

<span id="page-8-2"></span>(2.16) 
$$
L^2(U, G; g, h^G) \longrightarrow L^2([0, \varepsilon], L^2(\Sigma, G_\Sigma; g_\Sigma, h^{G_\Sigma}), \quad G = E, F,
$$

where  $g_{\Sigma} = g \restriction \Sigma, G_{\Sigma} := G \restriction \Sigma$  and  $h^{G_{\Sigma}} = h^G \restriction G_{\Sigma}$ . Now we consider

<span id="page-8-5"></span>
$$
(2.17) \tD := \Phi^F A(\Phi^E)^{-1} : C^\infty([0,\varepsilon), \Gamma^\infty(E_\Sigma)) \longrightarrow C^\infty([0,\varepsilon), \Gamma^\infty(F_\Sigma)).
$$

Since A is a first order elliptic differential operator we find

<span id="page-8-3"></span>(2.18) 
$$
D = J_x \left( \frac{d}{dx} + B_x \right),
$$

where  $J_x \in \text{Hom}(E_\Sigma, F_\Sigma)$ ,  $0 \le x \le \varepsilon$ , is a smooth family of bundle homomorphisms and  $(B_x)_{0 \le x \le \varepsilon}$  is a smooth family of first order elliptic differential operators between sections of  $E_{\Sigma}$ . Note that in view of [\(2.11\)](#page-7-1)  $J_0$  equals  $i\sigma_A^1(\nu^b)$ , where  $\nu = -\frac{d}{dx}$  is the outward normal vector field.

To avoid an inflation of parentheses we will most often use the notation  $B_x, J_x$ instead of  $B(x)$ ,  $J(x)$  etc. Only to avoid double subscripts we will write  $B(x)$ ,  $J(x)$ in subscripts.

Since  $\Phi^E, \Phi^F$  are unitary (cf. [\(2.15\)](#page-8-1), [\(2.16\)](#page-8-2)) we have  $D^t = \Phi^E A^t (\Phi^F)^{-1}$  and hence

<span id="page-8-4"></span>(2.19) 
$$
-D^{t} = J_{x}^{t} \frac{d}{dx} - B_{x}^{t} J_{x}^{t} + (J_{x}')^{t}
$$

$$
= J_{x}^{t} \left(\frac{d}{dx} - (J_{x}')^{-1} B_{x}^{t} J_{x}^{t}\right) + (J_{x}')^{t}
$$

<span id="page-8-6"></span>If A is formally selfadjoint, we have the relations

(2.20) 
$$
J^t = -J, \quad JB = J' - B^t J
$$

 $($ ' denotes differentiation by x).

<span id="page-8-0"></span><sup>1</sup>For simplicity we content ourselves with Sobolev spaces of nonnegative order. On a manifold with boundary Sobolev spaces of negative order are a nuisance, although with some care they could be dealt with here, cf. [[LM72](#page-57-11)].

Alternatively, we may choose the following normal form in a collar of the boundary:

<span id="page-9-1"></span>(2.21) 
$$
D = J_x \left( \frac{d}{dx} + B_x \right) + \frac{1}{2} J'_x,
$$

In this normalization,  $A = A^t$  implies the relations

$$
(2.22)\t\t\t Jt = -J, \quad JB = -BtJ.
$$

The normal form  $(2.21)$  determines  $J$  and  $B$  uniquely.

Returning to general (not necessarily formally selfadjoint) A we remark that the ellipticity of  $A$  and hence of  $D$  imposes various restrictions. The obvious ones are that  $J_x$  is invertible and that  $B_x$  is elliptic for all x. What's more, ellipticity of D means that for  $\lambda \in \mathbb{R}, \xi \in T_p^*\Sigma, (\lambda, \xi) \neq (0, 0)$ , the operator

<span id="page-9-4"></span>
$$
(2.23)\t\t\t i\lambda + \sigma_{B(x)}^1(p,\xi)
$$

is invertible for all  $(x, p) \in [0, \varepsilon) \times \Sigma$ . Here,  $\sigma_{B(x)}^1$  denotes the leading symbol of  $B_x$ . In other words, for  $\xi \in T_p^* \Sigma \setminus \{0\}$  the endomorphism  $\sigma_{B(x)}^1(p,\xi) \in \text{End}(E_p)$ has no eigenvalues on the imaginary axis  $i\mathbb{R}.$ 

<span id="page-9-0"></span>2.3. Regular boundary conditions. For the convenience of the reader and to fix some notation we briefly summarize a few basic facts about boundary conditions for  $A$ . Standard references are [[BBW93,](#page-56-1) Hör85, [LM72,](#page-57-11) [See69](#page-57-4)]. We will adopt the point of view of the elementary functional analytic presentation [[BL01](#page-57-12)]. However, we try to be as self-contained as possible.

It is well-known that the trace map

(2.24) 
$$
\varrho: \Gamma_0^{\infty}(M; E) \longrightarrow \Gamma^{\infty}(\Sigma; E), \quad f \longmapsto f \upharpoonright \Sigma
$$

extends by continuity to a bounded linear map between Sobolev spaces

<span id="page-9-5"></span>
$$
(2.25) \tL_s^2(M, E) \longrightarrow L_{s-1/2}^2(\Sigma, E_\Sigma), \t s > 1/2.
$$

For the domain of  $A_{\text{max}}$  this can be pushed a bit further. Namely, for  $s \geq 0$  the trace map extends by continuity to a bounded linear map

<span id="page-9-2"></span>(2.26) 
$$
\mathscr{D}(A_{\max,s}) \longrightarrow L_{s-1/2}^2(\Sigma, E_{\Sigma}), \quad s \ge 0,
$$

that is, there is a constant  $C_s$ , such that for  $f \in L_s^2(M, E)$  with  $Af \in L_s^2(M, E)$ 

<span id="page-9-3"></span>
$$
(2.27) \t\t\t \|gf\|_{s-1/2} \leq C_s(\|f\|_s + \|Af\|_s) \t (s \geq 0).
$$

Here  $||f||_s$  denotes the Sobolev norm of order s. Furthermore, norms of operators from  $L_s^2$  to  $L_{s'}^2$  will be denoted by  $\|\cdot\|_{s,s'}$ , and  $\|\cdot\|_{\infty}$  denotes the sup-norm of a function.

The proof of  $(2.26)$ ,  $(2.27)$  in [[BBW93](#page-56-1), Theorem 13.8 and Corollary 13.9] simplifies [[LM72](#page-57-11)] for operators of Dirac type but remains valid for any elliptic differential operator of first order, cf. also [[BL01](#page-57-12), Lemma 6.1].

**Definition 2.3.** (a) Let  $CL^0(\Sigma; E_{\Sigma}, G)$  denote the space of *classical pseudodiffer*ential operators of order 0, acting from sections of  $E_{\Sigma}$  to sections of another smooth Hermitian vector bundle  $G$  over  $\Sigma$ .

(b) Let  $P \in CL^0(\Sigma; E_\Sigma, G)$ . We denote by  $A_P$  the operator A acting on the domain

(2.28) 
$$
\mathscr{D}(A_P) := \left\{ f \in L_1^2(M, E) \middle| P(\varrho f) = 0 \right\},
$$

and by  $A_{\text{max},P}$  the operator A acting on the domain

(2.29) 
$$
\mathscr{D}(A_{\max,P}) := \{ f \in L^2(M,E) \mid Af \in L^2(M,F), P(\varrho f) = 0 \}.
$$

(c) The boundary condition P for A is called regular if  $A_{\text{max},P} = A_P$ , i.e., if  $f, Af \in L^2, P(\varrho f) = 0$  already implies that  $f \in L^2_1(M, E)$ .

(d) The boundary condition P is called *strongly regular* if  $f \in L^2$ ,  $Af \in L^2_k$ ,  $P(\varrho f) =$ 0 already implies  $f \in L^2_{k+1}(M, E)$ ,  $k \geq 0$ .

[\(2.27\)](#page-9-3) shows that  $\mathscr{D}(A_P)$  is in any case a closed subspace of  $L^2_1(M,E)$ .

<span id="page-10-2"></span>**Proposition 2.4.** Let P be regular for A. Then  $A_P$  is a closed semi-Fredholm operator with finite–dimensional kernel.

PROOF. Let  $(f_n) \subset \mathscr{D}(A_P)$  be a sequence with  $f_n \to f$  and  $Af_n \to g$  $L^2(M, F)$ . Then  $Af = g$  weakly, hence  $f \in \mathscr{D}(A_{\max}) = \mathscr{D}(A_{\max,0})$  and in view of [\(2.26\)](#page-9-2), [\(2.27\)](#page-9-3) we have  $P(\varrho f) = 0$  and the regularity of P implies  $f \in L_1^2(M, E)$ , thus  $f \in \mathscr{D}(A_P)$ .

Hence  $A_P$  is closed and thus  $\mathscr{D}(A_P)$  is complete in the graph norm. In view of the Closed Graph Theorem the previous argument shows that the inclusion  $\iota : \mathscr{D}(A_P) \hookrightarrow L_1^2(M, E)$  is bounded.  $\iota$  is thus an injective bounded linear map from the Hilbert space  $\mathscr{D}(A_P)$  (equipped with the graph norm) onto a closed subspace of  $L_1^2(M, E)$ ; the closedness is also a consequence of the argument at the beginning of this proof. Consequently, on  $\mathscr{D}(A_P)$  the graph norm and the  $L_1^2$ -norm are equivalent. I.e., for  $f \in \mathcal{D}(A_P)$  we have

(2.30) 
$$
\frac{1}{C}||f||_1 \le ||f||_0 + ||Af||_0 \le C||f||_1.
$$

Since the inclusion  $L_1^2(M, E) \hookrightarrow L^2(M, E)$  is compact, the inclusion  $\mathscr{D}(A_P) \hookrightarrow$  $L^2(M, E)$  is compact, too. Consequently,  $A_P$  is a semi-Fredholm operator with finite–dimensional kernel.

### <span id="page-10-1"></span>Remark 2.5.

(1)  $P =$  Id is strongly regular and its domain  $\mathscr{D}(A_P) = L_{1,0}^2(M, E)$  equals the closure of  $\Gamma_0^{\infty}(M \setminus \Sigma; E)$  in  $L_1^2(M, E)$ . This is seen by induction. Namely, if  $f \in L^2, Af = g \in L^2$  $f \in L^2, Af = g \in L^2$  $f \in L^2, Af = g \in L^2$  and  $gf = 0$  we may extend<sup>2</sup> f by 0 to obtain  $\tilde{f}, \tilde{A}\tilde{f} \in L^2_{loc}$ and hence  $f \in L_1^2$ .

For the induction we first note that similarly as in [[BL01](#page-57-12), Cor. 2.14] one shows that to the map

$$
(2.31) \qquad \varrho^{(k+1)}: L^2_{k+1}(M, E) \longrightarrow \bigoplus_{j=0}^k L^2_{k-j+1/2}(\Sigma, E_{\Sigma}), \quad f \longmapsto (\varrho A^j f)_{j=0}^k
$$

there exists a continuous linear right–inverse

(2.32) 
$$
e^{(k+1)} : \bigoplus_{j=0}^{k} L_{k-j+1/2}^2(\Sigma, E_{\Sigma}) \longrightarrow L_{k+1}^2(M, E).
$$

To complete the induction consider  $f \in L^2$ ,  $Af \in L^2_k$ ,  $gf = 0$ . By induction we may assume  $f \in L^2_k$ . Put

(2.33) 
$$
f_1 := f - e^{(k+1)}(0, \varrho Af, \dots, \varrho A^k f).
$$

Then  $f - f_1 \in L^2_{k+1}$ ,  $f_1 \in L^2_k$  and  $\varrho A^j f_1 = 0, j = 0, \ldots, k$ . Hence we may extend f by 0 to obtain  $\hat{f} \in L^2$  with  $\hat{A}^j \hat{f} \in L^2$ ,  $j = 0, \ldots, k+1$ . From local elliptic regularity we infer  $f \in L^2_{k+1,loc}$  and thus  $f \in L^2_{k+1}$ .

 $(\text{im } A_{\text{Id}})^{\perp} = \{ f \in L^2(M, F) \mid A^t f = 0 \}$  which is known to be (or see Propo-sition [5.12](#page-29-0) and Theorem [6.1](#page-30-2) below) infinite–dimensional if dim  $M > 1$ . Hence we cannot expect regularity to imply that  $A_P$  is Fredholm. However, if P and the dual boundary condition for  $A^t$  are regular then  $A_P$  is Fredholm.

<span id="page-10-0"></span><sup>&</sup>lt;sup>2</sup>We think of M as being a subset of an open manifold  $\widetilde{M}$  to which A can be extended as an elliptic operator.

(2) Regular boundary conditions are closely related to the well-posed boundary conditions of Seeley [[See69,](#page-57-4) [BBW93](#page-56-1)]. One of the main results in [[BL01](#page-57-12)] states that for symmetric Dirac operators and symmetric boundary conditions (given by operators P with closed range) regularity and well–posedness are equivalent.

(3) It is well-known that if P has closed range then P may be replaced by an orthogonal projection with the same kernel. In this setting the dual boundary condition can easily be computed:

**Proposition 2.6.** Let  $P \in CL^0(\Sigma; E_{\Sigma})$ ,  $P = P^2 = P^*$ . Then

(2.34) 
$$
(A_P)^* = A^t_{\max,(\text{Id}-P)J_0^t}.
$$

PROOF. This follows easily from Green's formula Lemma [2.2.](#page-7-2)

We recall from  $[H\ddot{\sigma}r85,$  Definition 20.1.1] (see also  $[BBW93,$  $[BBW93,$  $[BBW93,$  Remark 18.2d]):

**Definition 2.7.** Let  $P \in CL^0(\Sigma; E_{\Sigma}, G)$ . We say that P defines a local elliptic boundary condition for our operator  $A$  (or, equivalently, we say  $P$  satisfies the Šapiro-Lopatinski condition for A), if and only if the leading symbol  $\sigma_P^0$  of P maps the space  $M_{y,\zeta}^+$  isomorphically onto the fibre  $G_y$  for each point  $y \in \Sigma$  and each cotangent vector  $\zeta \in T^*_y(\Sigma)$ ,  $\zeta \neq 0$ . Here  $M^+_{y,\zeta}$  denotes the space of boundary values of bounded solutions u on the positive real line of the linear system  $\frac{d}{dt}u$  +  $\sigma_{B(0)}^1(y,\zeta)u=0$  of ordinary differential equations.

<span id="page-11-2"></span>**Remark 2.8.** Note that a solution of the ordinary differential equation  $\frac{d}{dt}u$  +  $\sigma_{B(0)}^1(y,\zeta)u=0$  is bounded if and only if the initial value  $u_0$  belongs to the range of the positive spectral projection  $P_+(\sigma_{B(0)}^1(y,\zeta))$  (cf. Section [3](#page-11-0) below) of the matrix  $\sigma_{B(0)}^1(y,\zeta)$ . Hence  $M^+_{y,\zeta} = \text{im } P_+(\sigma_{B(0)}^1(y,\zeta))$  and local ellipticity means that  $\sigma_P^0$  maps im  $P_+(\sigma_{B(0)}^1(y,\zeta))$  isomorphically onto  $G_y$ .

We obtain from  $[H\ddot{\sigma}r85,$  Theorem 20.1.2, Theorem 20.1.8] (differently also along the lines of [[BBW93](#page-56-1), Theorem 19.6]):

<span id="page-11-3"></span>**Proposition 2.9.** Any P satisfying the Šapiro-Lopatinskii condition for A is strongly regular and the corresponding  $A_P$  is a Fredholm operator.

# 3. Sectorial projections of an elliptic operator

<span id="page-11-1"></span><span id="page-11-0"></span>3.1. Parameter dependent ellipticity. Regarding properties of the tangential operator  $B_0$  on  $\Sigma$ , it is natural to distinguish three situations of increasing generality:

- (i)  $B_0$  is formally selfadjoint,
- (ii)  $B_0 B_0^t$  is an operator of order zero, and
- (iii)  $B_0$  is the tangential operator of an elliptic operator over the whole manifold M.

Whereas (i) implies that the spectrum spec( $B_0$ ) of  $B_0$  is contained in the real axis and (ii) that for all  $p \in \Sigma$  and  $\xi \in T_p^* \Sigma$  the leading symbol  $\sigma_{B(0)}^1(p,\xi) \in \text{End}(E_p)$ is selfadjoint, the general case (iii) a priori only implies that  $\sigma_{B(0)}^1(p,\xi)$  has no eigenvalues on the imaginary axis iR for all  $p \in \Sigma$  and  $\xi \in T_p^* \Sigma \setminus \{0\}$ , as explained above after [\(2.23\)](#page-9-4).

One may ask, what consequences can be drawn from the general property (iii) for the spectrum of  $B_0$ ? A first answer is Proposition [3.3](#page-12-0) below. In fact, (iii) contains more information than just that the leading symbol  $\sigma_{B(0)}^1(p,\xi)$  has no eigenvalues on  $i\mathbb{R}$ .

For the convenience of the reader let us briefly recall the notion of a (pseudo) differential operator with parameter, cf. Shubin [[Shu80](#page-57-7), Section II.9].

Let  $\Lambda \subset \mathbb{C}$  be an open conic subset, i.e.,  $z \in \Lambda$ ,  $r > 0 \Rightarrow rz \in \Lambda$ . For an open subset  $U \subset \mathbb{R}^n$  let  $S^m(U, \mathbb{R}^n \times \Lambda)$  denote the space of smooth functions

$$
a: U \times \mathbb{R}^n \times \Lambda \longrightarrow \mathbb{C}, \quad (x, \xi, \lambda) \longmapsto a(x, \xi, \lambda),
$$

such that for multi-indices  $\alpha, \beta \in \mathbb{Z}_+^n$ ,  $\gamma \in \mathbb{Z}_+^2$  and each compact subset  $K \subset U$  we have

$$
\left|\partial_x^{\alpha}\partial_{\xi}^{\beta}\partial_{\lambda}^{\gamma}a(x,\xi,\lambda)\right| \leq C_K\big(1+|\xi|+|\lambda|\big)^{m-|\beta|-|\gamma|}.
$$

We emphasize that  $\partial_\lambda^\gamma$  denotes real partial derivatives – we do not require holomorphicity in  $\lambda$ .

In other words,  $S^m(U, \mathbb{R}^n \times \Lambda)$  are the symbols of Hörmander type (1,0).

We shall call a symbol  $a \in S^m(U, \mathbb{R}^n \times \Lambda)$  classical if it has an asymptotic expansion

$$
(3.1) \t\t a \sim \sum_{j=0}^{\infty} a_{m-j},
$$

where  $a_{m-j} \in S^{m-j}(U, \mathbb{R}^n \times \Lambda)$  with homogeneity

$$
a_{m-j}(r\xi, r\lambda) = r^{m-j} a_{m-j}(\xi, \lambda) \text{ for } r \ge 1, \ |\xi|^2 + |\lambda|^2 \ge 1.
$$

We denote the classical symbols by  $\mathrm{CS}^m(U, \mathbb{R}^n \times \Lambda) \subset S^m(U, \mathbb{R}^n \times \Lambda)$ .

**Definition 3.1.** Let  $E_{\Sigma}$  be a complex vector bundle of finite fibre dimension N over a smooth closed manifold  $\Sigma$  and let  $\Lambda \subset \mathbb{C}$  be open and conic. A *classical* pseudodifferential operator of order m with parameter  $\lambda \in \Lambda$  is a family  $B(\lambda) \in$  $CL^{m}(\Sigma; E_{\Sigma}), \lambda \in \Lambda$ , such that locally  $B(\lambda)$  is given by

$$
B(\lambda)u(x) = (2\pi)^{-n} \int_{\mathbb{R}^n} \int_U e^{i\langle x-y,\xi\rangle} b(x,\xi,\lambda)u(y)dyd\xi
$$

with b an  $N \times N$  matrix of functions belonging to  $\mathbb{CS}^m(U, \mathbb{R}^n \times \Lambda)$ .

**Remark 3.2.** (1) A *pseudodifferential operator with parameter* is more than just a map from Λ to the space of pseudodifferential operators.

(2) Our definition of a pseudodifferential operator with parameter is slightly different from that of Shubin [[Shu80](#page-57-7), Section II.9]; however, the main results of loc. cit. do also hold for this class of operators.

The leading symbol of a classical pseudodifferential operator  $B$  of order  $m$  with parameter is now a smooth function  $\sigma_B^m(x,\xi,\lambda)$  on  $T^*\Sigma \times \Lambda \setminus \{(x,0,0) \mid x \in \Sigma\}$ which is homogeneous in the following sense

$$
\sigma_B^m(x, r\xi, r\lambda) = r^m \sigma_B^m(x, \xi, \lambda) \text{ for } (\xi, \lambda) \neq (0, 0), r > 0.
$$

Parameter dependent ellipticity is defined as invertibility of this homogeneous leading symbol. The basic example of a pseudodifferential operator with parameter is the resolvent of an elliptic differential operator.

<span id="page-12-0"></span>**Proposition 3.3.** Let  $\Sigma$  be a closed manifold and let  $B \in \text{Diff}^1(\Sigma; E_{\Sigma})$  be a first order differential operator. Let  $\Lambda \subset \mathbb{C}$  be an open conic subset such that  $B-\lambda, \lambda \in \Lambda$ , is parameter dependent elliptic, i.e., for each  $(p, \xi) \in T^*\Sigma$ ,  $\xi \neq 0$ , and each  $\lambda \in \Lambda$ the homomorphism

<span id="page-12-1"></span>
$$
\sigma_B^1(p,\xi) - \lambda : E_p \longrightarrow E_p
$$

is invertible. Then there exists  $R > 0$  such that  $B - \lambda$  is invertible for  $\lambda \in \Lambda, |\lambda| \geq$ R, and we have

$$
||(B - \lambda)^{-1}||_{s, s+\alpha} \le C_{\alpha} |\lambda|^{-1+\alpha}
$$

for such  $\lambda$  and  $0 \leq \alpha \leq 1$ .



<span id="page-13-1"></span>FIGURE 1. Construction of a closed cone  $\Lambda$  such that  $B_0 - \lambda$  is elliptic with parameter  $\lambda \in \Lambda.$ 

For the proof see [[Shu80](#page-57-7), Theorem 9.3]. In our situation Proposition [3.3](#page-12-0) has the following consequences:

**Proposition 3.4.** Let  $\Sigma$  be a closed manifold and let

$$
D = J_x \left(\frac{d}{dx} + B_x\right)
$$

be a first order elliptic differential operator on the collar  $[0, \varepsilon) \times \Sigma$ . Then

- (a)  $B_0 \lambda$  is parameter dependent elliptic in an open conic neighborhood  $\Lambda$ of the imaginary axis  $i\mathbb{R}$ .
- (b)  $B_0$  is an operator with compact resolvent, spec  $B_0$  consists of a discrete set of eigenvalues of finite multiplicity. At most finitely many eigenvalues lie on the imaginary axis iR.

For an eigenvalue  $\lambda$  even the *generalized eigenspace*  $\bigcup_N \text{ker}(B_0 - \lambda)^N$  is finitedimensional; note that  $B_0$  is not necessarily selfadjoint.

PROOF. From the ellipticity of D we infer that  $\sigma_{B(0)}^1(p,\xi) - it$  is invertible for  $(p, \xi, t) \in T^* \Sigma \times \mathbb{R}, (\xi, t) \neq (0, 0)$ . Since

$$
\bigcup_{(p,\xi)\in T^*\Sigma, \, |\xi|=1} \operatorname{spec} \sigma^1_{B(0)}(p,\xi)
$$

is bounded in  $\mathbb C$  and in view of the homogeneity we find an angle  $\vartheta > 0$  such that

spec 
$$
\sigma_{B(0)}^1(p,\xi) \cap \Lambda = \emptyset
$$
.

Here  $\Lambda$  is as in Figure [1.](#page-13-1)

(a) now follows from the previous proposition. Since  $B_0$  is elliptic, its spectrum is either discrete or equals  $\mathbb C$ . The previous lemma implies that  $B_0 - \lambda$  is invertible for  $\lambda \in \Lambda$  large enough. Hence we conclude that spec  $B_0$  is discrete and that (b)  $holds.$ 

<span id="page-13-0"></span>3.2. Sectorial operators: abstract Hilbert space framework. We shall now discuss the positive respectively negative sectorial spectral projections of an elliptic differential operator B of first order on a closed manifold  $\Sigma$ . We start with a purely functional analytic discussion.

<span id="page-14-7"></span>3.2.1. Idempotents in a Hilbert space. Let us briefly summarize some facts about (not necessarily bounded) idempotents in a Hilbert space. A densely defined operator P in the Hilbert space H is an *idempotent* if  $P \circ P = P$ ; as an identity between unbounded operators  $P \circ P = P$  means im  $P \subset \mathcal{D}(P)$  and  $P(Px) = Px$ for  $x \in \mathcal{D}(P)$ .

Given subspaces  $U, V \subset H$  with

<span id="page-14-0"></span>
$$
(3.3) \tU \cap V = \{0\},\
$$

<span id="page-14-1"></span>
$$
(3.4) \t\t U + V
$$
 dense in H,

the projection  $P_{U,V}$  along U onto V is an (not necessarily bounded) idempotent and every idempotent P in H is of this form with  $\mathscr{D}(P_{U,V}) = U + V, U = \text{ker } P$  and  $V = \operatorname{im} P.$ 

It is easy to see that  $P^*_{U,V} = P_{V^{\perp},U^{\perp}}$  is also an (not necessarily densely defined) idempotent. Thus  $P_{U,V}$  is closable iff  $U^{\perp}+V^{\perp}$  is dense or, equivalently  $\overline{U}\cap\overline{V}=\{0\}.$ In that case, the closure of  $P_{U,V}$  is  $P_{U,V} = P_{\overline{U},\overline{V}}$ . Consequently,  $P_{U,V}$  is a closed operator if and only if  $U, V$  are closed subspaces of  $H$ .

<span id="page-14-4"></span>**Lemma 3.5.** (a) Let  $P_{U,V}$  be an idempotent in the Hilbert space H, where  $U, V$ are closed subspaces satisfying  $(3.3)$ ,  $(3.4)$  above. Then  $P_{U,V}$  is bounded if and only if  $U + V = H$ .

(b) Let  $P = P_{U,V}$  be a bounded idempotent in the Hilbert space H. Then  $P + \text{Id} - P^*$ is an invertible operator.

Denote by  $P_{\text{ort}}$  the orthogonalization of P, i.e.,  $P_{\text{ort}} = P_{V^{\perp} \cdot V}$  is the orthogonal projection onto im P. Then we have

<span id="page-14-3"></span>(3.5) 
$$
P_{\text{ort}} = P(P + \text{Id} - P^*)^{-1},
$$

<span id="page-14-2"></span>(3.6) 
$$
(P^*)_{\text{ort}} = (P + \text{Id} - P^*)^{-1}P.
$$

PROOF. (a) is a consequence of the Closed Graph Theorem.

(b) By (a) U, V are closed subspaces of H satisfying  $U \cap V = \{0\}$ ,  $U \oplus V = H$ . Then  $\overline{\text{Id}} - P^* = P_{U^{\perp}, V^{\perp}}$ . Since bounded idempotents are bounded below  $P \upharpoonright U^{\perp}$  maps  $U^{\perp} = \ker P^{\perp}$  bijectively onto V and Id  $-P^*$  maps  $U = \ker(\text{Id} - P^*)^{\perp}$  bijectively onto  $V^{\perp}$ . Hence  $P + \text{Id} - P^*$  is invertible. Moreover, this description gives (P + Id  $-P^*$ )<sup>-1</sup> explicitly: given  $v \in V = \text{im } P$  then  $(P + \text{Id} - P^*)^{-1}v$  is the unique element  $\xi \in U^{\perp}$  with  $P\xi = v$  and thus  $P(P + \text{Id} - P^*)^{-1}v = v$ . Furthermore, if  $v \in V^{\perp}$  then  $(P + \text{Id} - P^{*})^{-1}v$  is the unique element  $\eta \in U = \ker P$  with  $(\text{Id} - P^*)\eta = v$ . This proves  $P(P + \text{Id} - P^*)^{-1} = P_{V^{\perp}, V} = P_{\text{ort}}$ . The equality [\(3.6\)](#page-14-2) is proved similarly.

Alternatively, one may apply [\(3.5\)](#page-14-3) to  $P^*$  to find  $(P^*)_{\text{ort}} = P^*(P^* + \text{Id} - P)^{-1}$ . Then [\(3.6\)](#page-14-2) follows from  $(P + Id - P^*)P^* = PP^* = P(P^* + Id - P)$ .

Our construction of  $P_{\text{ort}}$  is a slight modification of the construction given by M. Birman and A. Solomyak and disseminated in [[BBW93](#page-56-1), Lemma 12.8].

Lemma [3.5](#page-14-4) (a) shows that unbounded idempotents in a Hilbert space are abundant. See also Example [3.13](#page-17-1) below.

3.2.2. The semigroups  $Q_{\pm}(x)$  of a sectorial operator. In this subsection let H be a separable Hilbert space and  $B$  a closed operator in  $H$ .

<span id="page-14-5"></span>**Definition 3.6.** We call  $B$  a *weakly sectorial* operator if

- (1) B has compact resolvent.
- (2) There exists a closed conic neighborhood  $\Lambda$  of iR such that spec  $B \cap \Lambda$  is finite and

(3.7) 
$$
\|(B - \lambda)^{-1}\| = \mathcal{O}(|\lambda|^{-\alpha}), \quad |\lambda| \to \infty, \lambda \in \Lambda,
$$

<span id="page-14-6"></span>for some  $0 < \alpha \leq 1$ .



<span id="page-15-0"></span>FIGURE 2. The contours  $\Gamma_{\pm}$  in the plane defining the semigroups  $Q_{\pm}$ .

If  $\alpha = 1$  then we call B a sectorial operator.

We fix a weakly sectorial operator  $B$  in the sense of Definition [3.6.](#page-14-5)

**Convention 3.7.** (a)  $c > 0$  is chosen large enough such that (3.8) spec  $B \cap \{z \in \mathbb{C} \mid |z| = c\} = \emptyset$ ,

> and such that  $\{z \in \mathbb{C} \mid |z| = c\}$  contains all eigenvalues on the imaginary axis.

(b) We specify two complementary contours  $\Gamma_{+}$  in the plane as sketched in Figure [2](#page-15-0) with  $\Gamma_+$  encircling, up to finitely many exceptions, the eigenvalues of B with real part  $\geq 0$  and  $\Gamma_-\$ encircling the remaining eigenvalues. Of course, for this to be possible  $c$  has to be large enough.

# <span id="page-15-3"></span>Definition 3.8.

(3.9) 
$$
Q_{+}(x) := \frac{1}{2\pi i} \int_{\Gamma_{+}} e^{-\lambda x} (\lambda - B)^{-1} d\lambda, \qquad x > 0,
$$

<span id="page-15-1"></span>(3.10) = Id + 
$$
\frac{1}{2\pi i}
$$
 $\int_{\Gamma_+} e^{-\lambda x} \lambda^{-1} B(\lambda - B)^{-1} d\lambda$ ,

(3.11) 
$$
Q_{-}(x) := \frac{1}{2\pi i} \int_{\Gamma_{-}} e^{-\lambda x} (\lambda - B)^{-1} d\lambda, \qquad x < 0,
$$

<span id="page-15-2"></span>(3.12) 
$$
= \frac{1}{2\pi i} \int_{\Gamma_{-}} e^{-\lambda x} \lambda^{-1} B(\lambda - B)^{-1} d\lambda.
$$

When the dependence on B matters we will write  $Q_{\pm}(x, B)$ .

Formulas [\(3.10\)](#page-15-1), [\(3.12\)](#page-15-2) are obtained by adding and subtracting  $\lambda^{-1}$  inside the integral and taking into account that 0 lies inside  $\Gamma_+$  but outside  $\Gamma_-$ .

 $Q_{+}(x)$  are certainly bounded operators for  $x > 0$   $(x < 0)$ . Heuristically,  $Q_{+}(0)$ should be the positive/negative sectorial spectral projection of  $B$ , obtained from holomorphic functional calculus. However,  $Q_{\pm}(0)$  is not defined everywhere. To avoid ambiguities, we shall keep to the following two rigorous definitions instead of dealing directly with  $Q_{\pm}(0)$ .

**Definition 3.9.** We put  $\mathscr{D}(P_{+,0}) := \{ \xi \in H \mid \lim_{x \to 0+} Q_+(x) \xi \text{ exists} \}$  and  $P_{+,0} \xi :=$  $\lim_{x\to 0+} Q_+(x)\xi$  for  $\xi \in \mathscr{D}(P_{+,0})$ .  $P_{-,0}$  is defined analogously using  $Q_-(x)$ .



<span id="page-16-0"></span>FIGURE 3. A two-component contour  $\Gamma$ , separating an inner sector around the real axis where all eigenvalues of  $B_0$  show up, from two outer sectors which totally belong to the resolvent set of  $B_0$ .

[\(3.10\)](#page-15-1), [\(3.12\)](#page-15-2) and the estimate [\(3.7\)](#page-14-6) imply that  $\mathscr{D}(B) \subset \mathscr{D}(P_{+,0})$  and for  $\xi \in \mathscr{D}(B)$  we have

(3.13) 
$$
P_{+,0}\xi = \xi + \frac{1}{2\pi i} \int_{\Gamma_+} \lambda^{-1} (\lambda - B)^{-1} d\lambda \ (B\xi)
$$

(3.14) 
$$
P_{-,0}\xi = \frac{1}{2\pi i} \int_{\Gamma_-} \lambda^{-1} (\lambda - B)^{-1} d\lambda (B\xi),
$$

thus  $P_{\pm,0}$  is densely defined  $(\mathscr{D}(B)$  is indeed a core for  $P_{\pm,0}$ ). Note that  $Q_{\pm}(x,B)^*$  =  $Q_{\pm}(x, B^*)$  (cf. Prop. [3.11\)](#page-16-1), hence the densely defined operator  $P_{\pm,0}(B^*)$  is contained in  $P_{\pm,0}(B)^*$ . Thus  $P_{\pm,0}$  is closable:

<span id="page-16-2"></span>**Definition 3.10.** The closure of  $P_{\pm,0}$  will be called the *positive/negative sectorial* spectral projection  $P_{\pm}$  of B.

<span id="page-16-1"></span>**Proposition 3.11.** For  $x, y > 0$  we have

- (a)  $Q_+(x, B)^* = Q_+(x, B^*), Q_-(-x, B)^* = Q_-(-x, B^*).$ (b)  $Q_+(x)Q_+(y) = Q_+(x+y)$ . (c)  $Q_+$  is differentiable and  $\frac{dQ_+}{dx}(x) = -BQ_+(x)$ . (d)  $Q_+(x)Q_-(-y) = Q_-(-y)Q_+(x) = 0.$
- (e)  $P_+Q_+(x) \subset Q_+(x)P_+, P_+Q_-(-x) = 0.$

PROOF. The proof is straightforward and analogous to the proof in [[Pon06](#page-57-9)] of the fact that  $P_+$  is an idempotent.

Corollary 3.12.  $P_{\pm}$  are complementary, i.e.,  $P_{+} = \text{Id} - P_{-}$ , (possibly unbounded) idempotents in H.

PROOF. Since  $\mathscr{D}(B)$  is a core for  $P_{\pm}$  it suffices to check that for  $\xi \in \mathscr{D}(B)$  we have  $P_{\pm}^2 \xi = P_{\pm} \xi$  and  $(P_+ + P_-) \xi = \xi$ .

If  $\xi \in \mathcal{D}(B)$  then using Proposition [3.11](#page-16-1) we find

(3.15) 
$$
Q_{+}(x)P_{+}\xi = \lim_{y \to 0+} Q_{+}(x)Q_{+}(y)\xi = \lim_{y \to 0+} Q_{+}(x+y)\xi = Q_{+}(x)\xi,
$$

hence  $P_+\xi \in \mathcal{D}(P_{+,0}) \subset \mathcal{D}(P_+)$  and  $P_+^2\xi = P_+\xi$ .

Secondly, we take  $a \xi \in \mathcal{D}(B)$  and find

$$
(P_{+} + P_{-})\xi = \xi + \frac{1}{2\pi i} \int_{\Gamma} \lambda^{-1} (\lambda - B)^{-1} d\lambda (B\xi),
$$

where  $\Gamma$  is chosen as in Figure [3.](#page-16-0) Pushing the radius of the circle arches to  $\infty$  shows  $(P_+ + P_-)\xi = \xi.$ 

The fact that the sectorial projections are a priori unbounded operators may seem strange. The following example shows that the phenomenon really occurs:

<span id="page-17-1"></span>**Example 3.13.** Let  $D$  be a discrete selfadjoint positive definite operator in  $H$ . I.e., there is an orthonormal basis  $(e_n)_{n\in\mathbb{N}}$  of H such that  $De_n = \lambda_n e_n$ , where  $0 < \lambda_1 \leq \lambda_2 \leq \ldots \to \infty$ .

Pick a parameter  $0 \le \alpha \le 1$  and define the operator B in  $H \oplus H$  as follows:

(3.16) 
$$
\mathscr{D}(B) := \{ (u, v) \in H \oplus H \mid v \in \mathscr{D}(D), Du - D^{2-\alpha}v \in H \},\newline B(u, v) := (Du - D^{2-\alpha}v, -Dv).
$$

One immediately checks that for  $\lambda \notin \text{spec } D \cup - \text{spec } D$  the resolvent of B is given by

<span id="page-17-2"></span>
$$
(3.17) \ (B - \lambda)^{-1}(\xi, \eta) = ((D - \lambda)^{-1}\xi - 2(D - \lambda)^{-1}D^{2-\alpha}(D + \lambda)^{-1}\eta, -(D + \lambda)^{-1}\eta).
$$

Because of  $0 \le \alpha \le 1$  the resolvent is indeed bounded. Furthermore, [\(3.17\)](#page-17-2) shows that outside a conic neighborhood of the real axis, equivalently in a conic neighborhood of the  $i\mathbb{R}$ , we have an estimate

(3.18) 
$$
\|(B - \lambda)^{-1}\| = \mathcal{O}(|\lambda|^{-\alpha}), \quad |\lambda| \to \infty.
$$

Hence, if  $0 < \alpha \leq 1$  then B is a weakly sectorial operator in the sense of Definition [3.6,](#page-14-5) spec B = spec D ∪ – spec D and the positive/negative spectral subspaces of B are given by

(3.19) 
$$
\lim_{x \to a} P_{+}(B) = H \oplus 0,
$$

$$
\ker P_{+}(B) = \{(u, D^{\alpha - 1}u) \mid u \in H\} = \text{Graph}(D^{\alpha - 1}).
$$

Consequently, if  $0 < \alpha < 1$  then im  $P_{+}(B) \oplus \ker P_{+}(B)$  is not closed and hence the positive sectorial projection  $P_+(B)$  is not bounded.

We leave it as an intriguing problem to find an example of a sectorial operator with decay rate  $\alpha = 1$  in [\(3.7\)](#page-14-6) such that  $P_+$  is unbounded.

## 3.3. Sectorial operators: parametric elliptic differential operators.

<span id="page-17-0"></span>3.3.1. The geometric situation. We return to our geometric situation and consider the tangential operator B (previously denoted by  $B(0)$  or  $B_0$ , for convenience we omit (0) as long as we do not need  $B(x)$ ) of an elliptic differential operator A on a compact manifold with boundary, cf. Section [2,](#page-5-0) in particular [\(2.18\)](#page-8-3).

Then it is known that the positive sectorial projection is bounded:

<span id="page-17-3"></span>**Theorem 3.14.** Let  $B$  be a first order elliptic differential operator on the closed manifold  $\Sigma$ . Furthermore, assume that  $B-\lambda$  is parametric elliptic in an open conic neighborhood  $\Lambda$  of i $\mathbb R$ . Then the positive/negative sectorial projections  $P_{+}$  of B are pseudodifferential operators of order 0. In particular  $P_{\pm}$  acts as bounded operator in each Sobolev space  $L_s^2(\Sigma, E_{\Sigma})$ .

The proof is an adaption of the classical complex power construction of Seeley [[See67](#page-57-13)]. See Burak [[Bur70](#page-57-8)], Wojciechowski [[Woj85](#page-57-14)], and recently Ponge [[Pon06](#page-57-9)].

We also note that it follows from Proposition [3.3](#page-12-0) that  $B$  is a sectorial operator in the sense of Definition [3.6.](#page-14-5) Also, recall from [\(3.2\)](#page-12-1) the resolvent estimate:

<span id="page-18-1"></span>For all  $s \in \mathbb{R}$ ,  $0 \leq \alpha \leq 1$ , we have

(3.20) 
$$
\sup_{\lambda \in \Gamma_{\pm}} |\lambda|^{1-\alpha} ||(\lambda - B)^{-1}||_{s, s+\alpha} \leq C(s, \alpha),
$$

where  $\|\cdot\|_{s,s+\alpha}$  denotes the operator norm between the Sobolev spaces  $L_s^2(\Sigma, E_{\Sigma})$ and  $L^2_{s+\alpha}(\Sigma, E_{\Sigma})$ , see also the following remark.

Here and in the following we shall denote the closed interval  $[0, \infty)$  by  $\mathbb{R}_+$ . Similarly  $\mathbb{Z}_+ := \{0, 1, 2, 3, \ldots\}.$ 

Remark 3.15. (1) We recall that (cf. e.g. [[BL01](#page-57-12), Cor. 2.20])

$$
(3.21) \qquad L_s^2(\mathbb{R}_+ \times \Sigma, E_\Sigma) = L_s^2(\mathbb{R}_+, L^2(\Sigma, E_\Sigma)) \cap L^2(\mathbb{R}_+, L_s^2(\Sigma, E_\Sigma)), \quad s \ge 0.
$$

<span id="page-18-2"></span>In particular, if  $s \in \mathbb{Z}_+$  then a Sobolev norm for  $L_s^2(\mathbb{R}_+ \times \Sigma, E_{\Sigma})$  is given by

$$
(3.22) \quad \|f\|_{L_s^2(\mathbb{R}_+\times\Sigma, E_\Sigma)}^2 = \int_0^\infty \|\partial_x^s f(x)\|_{L^2(\Sigma, E_\Sigma)}^2 + \|(\mathrm{Id} + |B|)^s f(x)\|_{L^2(\Sigma, E_\Sigma)}^2 dx.
$$

Since the spaces  $L_s^2(\ldots)$  have the interpolation property ([**[Tay96](#page-57-15)**, Sec. 4.2], [**[BL01](#page-57-12)**, Sec. 2) for  $s \geq 0$ , it will be sufficient in most cases to deal with integer  $s \in \mathbb{Z}_{+}$ .

(2) Note that since B is elliptic, the Sobolev norms on sections of  $E_{\Sigma}$  can be defined using  $B$ , i.e.,

(3.23) 
$$
\|\xi\|_{L_s^2(\Sigma, E_\Sigma)}^2 = \|(Id + |B|)^s \xi\|_{L^2(\Sigma, E_\Sigma)}^2.
$$

(3) Whenever it is clear from the context whether we are taking norms of sections over  $\mathbb{R}_+ \times \Sigma$  or over  $\Sigma$  we will, as before, denote Sobolev norms of order s by a subscript s.

3.3.2. Mapping properties of  $Q_+$ . The following Proposition will be useful for the study of the mapping properties of the invertible double and of the remainder terms in the construction of the Poisson operator and the Calderón projection, see Subsections [5.2](#page-24-0) and [5.3.](#page-27-0) Proposition [3.16](#page-18-0) establishes a weak convergence of  $Q_{+}(x) \rightarrow P_{+}, x \rightarrow 0+,$  in compensation for the generally not valid convergence in the operator norm.

<span id="page-18-0"></span>**Proposition 3.16.** Let  $\varphi \in C_0^{\infty}(\mathbb{R}_+), m \in \mathbb{Z}_+$ .

(a) For  $s \in \mathbb{R}$  the operator

(3.24) 
$$
\mathrm{id}_{\mathbb{R}_+}^m \varphi Q_+ : \xi \longmapsto (x \mapsto x^m \varphi(x) Q_+(x) \xi)
$$

maps  $L_s^2(\Sigma, E_{\Sigma})$  continuously to  $L^2_{\text{comp}}(\mathbb{R}_+, L^2_{s+m+1/2}(\Sigma, E_{\Sigma}))$ .

(b) For  $s \geq -1/2$  it maps continuously to  $L^2_{s+m+1/2,\text{comp}}(\mathbb{R}_+ \times \Sigma, E_{\Sigma}).$ 

PROOF. Let us first prove the claim (b). It is fairly easy to see that  $\mathrm{id}_{\mathbb{R}_+}^m \varphi Q_+$  maps  $L_s^2(\Sigma, E_{\Sigma})$  continuously to  $L^2_{\text{comp}}(\mathbb{R}_+, L_{s'}^2(\Sigma, E_{\Sigma}))$  for some s'. Thus once we have proved that the range of  $\mathrm{id}_{\mathbb{R}_+}^{m} \varphi Q_+$  is contained in the space  $L^2_{\text{comp}}(\mathbb{R}_+, L^2_{s+m+1/2}(\Sigma, E_{\Sigma}))$  the continuity will follow from the Closed Graph Theorem.

Furthermore, since  $\mathrm{id}_{\mathbb{R}_+}^m \varphi Q_+$  commutes with B it suffices to prove the claim for s large enough: namely, we pick a  $\lambda_0$  in the resolvent set of B. Then for arbitrary  $s \geq -1/2$  we choose k large enough such that the claim holds for  $s + k$ . The claim for s now follows from the identity

(3.25) id<sup>m</sup> R<sup>+</sup> ϕQ+|L 2 <sup>s</sup> = (λ<sup>0</sup> − B) k id<sup>m</sup> R<sup>+</sup> ϕQ+|L 2 s+k (λ<sup>0</sup> − B) −k |L 2 s .

Finally, by complex interpolation (cf. e.g. [[Tay96](#page-57-15), Sec. 4.2]) it suffices therefore to consider  $s = n + 1/2$ ,  $n \in \mathbb{Z}_+$ . Now pick  $\xi \in L^2_{n+1/2}(\Sigma, E_{\Sigma})$  and

put  $f(x) := x^m \varphi(x) Q_+(x) \xi$ . It is straightforward to check that f is smooth on  $(0, \infty) \times \Sigma$ . From

(3.26) 
$$
\left(\frac{d}{dx} + B\right)^j f(x) = \left(\left(\frac{d}{dx}\right)^j x^m \varphi(x)\right) Q_+(x) \xi
$$

we infer by the boundedness of  $P_+(B)$  (according to Theorem [3.14\)](#page-17-3) that

<span id="page-19-1"></span>
$$
\begin{aligned} \left(\frac{d}{dx} + B\right)^j f\big|_{x=0} \\ &= \begin{cases} 0, & j = 0, \dots, m-1, \\ \left(\frac{d}{dx}\right)^j \big|_{x=0} x^m \varphi(x) P_+(B) \xi \in L^2_{n+1/2}(\Sigma, E_\Sigma), & j \ge m, \end{cases} \\ &\in L^2_{s+m+1/2-j-1/2}(\Sigma, E_\Sigma). \end{aligned}
$$

From  $(3.27)$  and (an obvious adaption of) [[BL01](#page-57-12), Cor. 2.17] (cf. also Remark [2.5\)](#page-10-1) we infer that  $f \in L^2_{s+m+1/2}(\mathbb{R}_+ \times \Sigma, E_{\Sigma})$ . Hence (b) is proved.

For  $s \ge -1/2$  the claim (a) follows from (b). For arbitrary s we again conjugate by  $(\lambda_0 - B)^k$  as above and we reach the conclusion.

Remark 3.17. The claim of the previous proposition also follows by applying more sophisticated pseudodifferential techniques (cf. Grubb [[Gru96](#page-57-16), Thm. 2.5.7]). Our proof only uses the basic trace results for Sobolev spaces, the ellipticity of B, and the boundedness of the positive sectorial projection on Sobolev spaces. The previous proposition can therefore be generalized to situations where pseudodifferential techniques are not necessarily available. An abstract version is as follows (see also Subsection [5.1](#page-23-1) where scales of Hilbert spaces are recalled to some extent):

**Proposition 3.18.** Let B be a sectorial operator in a Hilbert space. Let  $H_s :=$  $\mathscr{D}((B^*B)^{s/2}), s \geq 0$ , be the scale of Hilbert spaces of  $B^*B$  and  $\widetilde{H}_s := \mathscr{D}((BB^*)^{s/2})$ be the scale of Hilbert spaces of  $BB^*$ . For negative s the spaces  $H_s$  and  $\widetilde{H}_s$  are defined by duality (cf. [[BL01](#page-57-12), Sec. 2.A]). Furthermore, put for  $s > 0$ 

$$
\mathscr{H}_{s}(\mathbb{R}_{+},H_{\bullet}):=\bigcap_{0\leq t\leq s}L^{2}_{s}(\mathbb{R}_{+},H_{s-t})
$$

(cf.  $[\text{BL01}, \text{Sec. 2}, \text{Prop. 2.10}]$  $[\text{BL01}, \text{Sec. 2}, \text{Prop. 2.10}]$  $[\text{BL01}, \text{Sec. 2}, \text{Prop. 2.10}]$  for other descriptions).

Assume that the positive sectorial projection  $P_+$  of B maps  $H_s$  continuously to  $H<sub>s</sub>$  for all s.

Let  $\varphi \in C_0^{\infty}(\mathbb{R}_+), m \in \mathbb{Z}_+$ . Then

(a) For  $s \in \mathbb{R}$  the operator

(3.28) 
$$
\mathrm{id}_{\mathbb{R}_+}^m \varphi Q_+ : \xi \longmapsto (x \mapsto x^m \varphi(x) Q_+(x) \xi)
$$

maps  $\mathscr{H}_s(\mathbb{R}_+, H_{\bullet})$  continuously to  $L^2_{\text{comp}}(\mathbb{R}_+, \widetilde{H}_{s+m+1/2}).$ (b) For  $s \geq -1/2$  it maps continuously to  $\mathscr{H}_{s+m+1/2,\text{comp}}(\mathbb{R}_+,\widetilde{H}_\bullet)$ .

For elliptic pseudodifferential operators the distinction between  $H_s$  and  $H_s$ is, of course, unnecessary. For general unbounded operators, however, we cannot expect  $H_s$  to be equal to  $\widetilde{H}_s$ .

#### 4. The invertible double

<span id="page-19-0"></span>We return to the set–up described at the beginning of Subsection [2.2](#page-7-0) and give a construction of the invertible double of a general first order elliptic differential operator.

<span id="page-20-0"></span>**4.1. The construction of**  $\tilde{A}_{P(T)}$ . We introduce the operator

(4.1) 
$$
\widetilde{A} := A \oplus (-A^t) : \Gamma^\infty(M; E \oplus F) \longrightarrow \Gamma^\infty(M; F \oplus E).
$$

We are going to consider a special class of boundary conditions for  $\widetilde{A}$ :

<span id="page-20-3"></span>**Definition 4.1.** Let  $T \in CL^0(\Sigma; E_{\Sigma}, F_{\Sigma})$  be a classical pseudodifferential operator of order 0, acting from sections of  $E_{\Sigma}$  to sections of  $F_{\Sigma}$ . We put

(4.2) 
$$
P(T) = (-T \text{ Id}) \in \text{CL}^{0}(\Sigma; E_{\Sigma} \oplus F_{\Sigma}, F_{\Sigma}).
$$

Viewed as an operator in  $L^2_s(\Sigma, E_\Sigma \oplus F_\Sigma)$  the operator  $P(T)$  has closed range which equals

(4.3) 
$$
\operatorname{im} P(T) = L_s^2(\Sigma, F_\Sigma) \subset L_s^2(\Sigma, E_\Sigma \oplus F_\Sigma).
$$

Since this is a closed subspace of  $L_s^2(\Sigma, E_\Sigma \oplus F_\Sigma)$ , the boundary condition for  $\widetilde{A}$ given by  $P(T)$  can be realized by a pseudodifferential orthogonal projection, as noted in Remark [2.5.](#page-10-1)3.

To be more specific we recall that the realization  $A_{P(T)}$  of A with respect to the boundary condition  $P(T)$  has domain

<span id="page-20-4"></span>(4.4) 
$$
\mathscr{D}(\widetilde{A}_{P(T)}) := \left\{ \begin{pmatrix} f_+ \\ f_- \end{pmatrix} \in L_1^2(M, E \oplus F) \middle| \varrho f_- = T \varrho f_+ \right\}.
$$

<span id="page-20-2"></span>**Lemma 4.2.** If T is invertible, the dual of the boundary condition  $P(T)$  for  $\widetilde{A}$  is  $P(-J_0^{-1}(T^t)^{-1}J_0^t),$ 

i.e.,

(4.5) 
$$
\left(\widetilde{A}_{P(T)}\right)^{*} = \widetilde{A}_{\max,P(-J_{0}^{-1}(T^{t})^{-1}J_{0}^{t})}^{*}.
$$

PROOF. Let  $f = \begin{pmatrix} f_+ \\ f_- \end{pmatrix} \in \mathscr{D}(\widetilde{A}_{P(T)})$  and  $g = \begin{pmatrix} g_+ \\ g_- \end{pmatrix} \in \mathscr{D}((\widetilde{A}_{P(T)})^*)$ . Note that  $g_+ \in L^2(M, F), g_- \in L^2(M, E)$ . Green's formula Lemma [2.2](#page-7-2) yields

(4.6)  
\n
$$
0 = \langle Af, g \rangle - \langle f, A^t g \rangle
$$
\n
$$
= \langle Af_+, g_+ \rangle - \langle f_+, A^t g_+ \rangle - \langle A^t f_-, g_- \rangle + \langle f_-, Ag_- \rangle
$$
\n
$$
= -\langle J_0 \varrho f_+, \varrho g_+ \rangle - \langle \varrho f_-, J_0 \varrho g_- \rangle
$$
\n
$$
= -\langle \varrho f_+, J_0^t \varrho g_+ + T^t J_0 \varrho g_- \rangle.
$$

This holds for all  $f \in \mathscr{D}(\tilde{A}_{P(T)})$  if and only if  $J_0^t \varrho g_+ + T^t J_0 \varrho g_- = 0$  and we reach the conclusion.

# <span id="page-20-1"></span>4.2. The local ellipticity of  $P(T)$  for  $\tilde{A}$ .

 $\sim$   $\sim$   $\sim$   $\sim$ 

<span id="page-20-5"></span>**Proposition 4.3.** Let T be an invertible bundle homomorphism from  $E_{\Sigma}$  to  $F_{\Sigma}$ with  $J_0^t T > 0$ . Then the boundary condition defined by  $P(T)$  satisfies the Sapiro-Lopatinski $\check{i}$  condition for  $\check{A}$ .

## <span id="page-20-6"></span>Remark 4.4.

(1) Obvious candidates for T with  $J_0^t T > 0$  are  $J_0$  and  $(J_0^t)^{-1}$ . We can in addition choose T to be unitary by putting  $T := (J_0 J_0^t)^{-1/2} J_0$ .

(2) Note that if  $J_0^t T$  is positive definite then it is in particular selfadjoint and hence the dual condition for  $\tilde{A}^t$  (cf. Lemma [4.2\)](#page-20-2) is given by

<span id="page-20-7"></span>(4.7) 
$$
T^{\text{dual}} = -J_0^{-1} (T^t)^{-1} J_0^t = -(T^t J_0)^{-1} J_0^t = -(J_0^t T)^{-1} J_0^t = -T^{-1}.
$$

In particular we find that fulfilling the assumption  $J_0^t T > 0$  of the preceding proposition implies that the boundary condition for  $\tilde{A}^t$  defined by  $P(T^{\text{dual}})$  also satisfies the Šapiro-Lopatinski condition. To see this we recall from  $(2.19)$  that the front bundle endomorphism  $J_0(A^t)$  of  $A^t$  is given by  $J_0(A^t) = -J_0^t$ , so

(4.8) 
$$
(-J_0^t)^t T^{\text{dual}} = J_0 T^{-1} = J_0 (J_0^t T)^{-1} J_0^t > 0.
$$

Proof. We refer to Remark [2.8](#page-11-2) and use the language of idempotents. During this proof, for an endomorphism b of a finite–dimensional vector space,  $P_+(b)$ will denote the spectral projection corresponding to a closed contour encircling all eigenvalues  $\lambda$  with Re  $\lambda \geq 0$  (respectively  $\lt 0$ ), cf. Section [3.](#page-11-0)

From  $(2.18)$ ,  $(2.19)$  we see that the tangential operator of A has leading symbol  $b_0 \oplus -(J_0^t)^{-1}b_0^*J_0^t$ ,  $b_0 := \sigma_{B(0)}^1$ . Consequently the positive spectral projection of  $b_0 \oplus -(J_0^t)^{-1}b_0^*J_0^t$  is given by  $P_+(b_0) \oplus (J_0^t)^{-1}P_-(B^t)J_0^t$ . In each  $y \in \Sigma$  and  $\zeta \in$  $T^*_y(E_\Sigma)$ ,  $\zeta \neq 0$ , we consider the Sapiro-Lopatinskii mapping from im  $P_+(b_0(y,\zeta)) \oplus$  $(\tilde{J}_0^t)^{-1}$  im  $P_-(b_0(y,\zeta)^*)$  to  $F_y$ , given by  $\sigma_{P(T)}^0 = (-T \text{ Id})$ :

$$
\text{im } P_+(b_0(y,\zeta)) \oplus (J_0^t)^{-1} \text{ im } P_-(b_0(y,\zeta)^*) \longrightarrow F_y
$$
  

$$
(e_+, (J_0^t)^{-1}e_-) \longmapsto -Te_+ + (J_0^t)^{-1}e_-.
$$

Multiplying by  $J_0^t$  we see that this map is bijective if and only if the map

<span id="page-21-1"></span>(4.9) 
$$
E_y = \operatorname{im} P_+(b_0(y,\zeta)) \oplus \operatorname{im} P_-(b_0(y,\zeta)^*) \longrightarrow E_y
$$

$$
(e_+,e_-) \longmapsto -J_0^t T e_+ + e_-
$$

is bijective. To explain why  $E_y = \lim P_+(b_0(y,\zeta)) \oplus \lim P_-(b_0(y,\zeta)^*)$  we note that  $\lim P_+(b_0(y,\zeta))^{\perp} = \ker P_+(b_0(y,\zeta))^* = \ker P_+(b_0(y,\zeta)^*) = \lim P_-(b_0(y,\zeta)^*)$ , so the sum on the left of [\(4.9\)](#page-21-1) is indeed an orthogonal decomposition (cf. also Lemma [3.5\)](#page-14-4).

Since the dimensions on the left and on the right side of [\(4.9\)](#page-21-1) coincide it suffices to show that the map in [\(4.9\)](#page-21-1) is injective: so let  $-J_0^tTe_+ + e_- = 0, e_+ \in$  $\lim P_+(b_0(y,\zeta)), e_-\in \lim P_-(b_0(y,\zeta))^* = \lim P_+(b_0(y,\zeta))^{\perp}.$  Taking scalar product with  $e_+$  we find  $0 = -\langle J_0^t T e_+, e_+ \rangle$ . This implies, since by assumption  $J_0^t T > 0$ , that  $e_{+} = 0$ . But then  $e_{-} = 0$  as well.

<span id="page-21-0"></span>**4.3. The solution space** ker  $\tilde{A}_{P(T)}$ . Next we indicate why the boundary conditions of Definition [4.1](#page-20-3) are significant. Before doing that we recall the various solution spaces and Cauchy data spaces associated to A.

Definition 4.5. (a) Put

$$
Z^{s}(A) := \{ f \in L_s^2(M, E) \mid Af = 0 \}, \quad s \ge 0.
$$

 $Z^s(A^t) \subset L^2_s(M, F)$  is defined analogously. For brevity we often write

(4.10) 
$$
Z_{+}^{s} := Z^{s}(A), \quad Z_{-}^{s} := Z^{s}(A^{t}).
$$

It follows from [\(2.25\)](#page-9-5) and [\(2.26\)](#page-9-2) that the trace map sends  $Z_{\pm}^{s}$  continuously to  $L_{s-1/2}^2(\Sigma, E_\Sigma)$  respectively  $L_{s-1/2}^2(\Sigma, F_\Sigma), s \geq 0$ .

(b) We define the Cauchy data spaces by

(4.11) 
$$
N_{\pm}^{s} := \varrho(Z_{\pm}^{s+1/2}), \quad s \ge -1/2, N_{\pm} := N_{\pm}^{0}.
$$

(c) Finally let

<span id="page-21-2"></span>(4.12) 
$$
Z_{+,0}(A) := \{ f \in L_1^2(M,E) \mid Af = 0, \varrho f = 0 \}
$$

denote the space of all *inner solutions*. It is the finite–dimensional kernel of  $A_{\text{Id}}$  (cf. Proposition [2.4\)](#page-10-2).  $Z_{-,0}(A) := Z_{+,0}(A^t)$  denotes the corresponding kernel of  $A_{\text{Id}}^t$ . (d) We say that A has the weak inner unique continuation property (UCP) if  $Z_{+,0}(A) = \{0\}.$ 

<span id="page-22-1"></span>**Proposition 4.6.** Let  $T$  be as in Definition [4.1](#page-20-3). Then there is a canonical inclusion

$$
Z_{+,0}(A) \oplus Z_{-,0}(A) \subset \ker \overline{A}_{P(T)}.
$$

If, in addition,  $J_0^t T$  is positive definite, then the inclusion is an equality.

PROOF. If  $f_{\pm} \in Z_{\pm,0}$  then  $\binom{f_{+}}{f_{-}} = f \in \ker \widetilde{A}_{P(T)}$  since  $\varrho f_{-} = 0 = T \varrho f_{+}$ , cf.  $(4.4).$  $(4.4).$ 

Now assume that  $J_0^t T$  is positive definite and let  $\begin{pmatrix} f_+ \\ f_- \end{pmatrix} = f \in \text{ker } \widetilde{A}_{P(T)}$ . Then certainly  $f_{\pm} \in Z_{\pm}^1$  and  $\varrho f_{-} = T \varrho f_{+}$ . Since  $J_0^t T$  is nonnegative and invertible, the operator  $W := (J_0^t T)^{1/2}$  exists and is invertible. Now Green's formula Lemma [2.2](#page-7-2) yields

(4.13)  
\n
$$
||W \varrho f_{+}||^{2} = \langle \varrho f_{+}, J_{0}^{t} T \varrho f_{+} \rangle
$$
\n
$$
= \langle J_{0} \varrho f_{+}, \varrho f_{-} \rangle
$$
\n
$$
= -\langle Af_{+}, f_{-} \rangle + \langle f_{+}, A^{t} f_{-} \rangle
$$
\n
$$
= 0,
$$

and since W is invertible we find  $\rho f_ - = 0$ . Thus  $\rho f_ + = T^{-1} \rho f_ - = 0$  and hence  $f_{\pm} \in Z_{\pm,0}.$ 

<span id="page-22-0"></span>4.4. The main result. Recall from Proposition [2.9](#page-11-3) that by checking the Sapiro-Lopatinskii condition we have not only proved regularity of  $P(T)$  for  $\tilde{A}$ , but also that  $A_{P(T)}$  is a Fredholm operator. Let us summarize the results of Proposition [2.4,](#page-10-2) Proposition [2.9,](#page-11-3) Proposition [4.3,](#page-20-5) Remark [4.4,](#page-20-6) and Proposition [4.6:](#page-22-1)

<span id="page-22-3"></span>Theorem 4.7. Let M be a compact Riemannian manifold with boundary and

 $A: \Gamma^{\infty}(M; E) \longrightarrow \Gamma^{\infty}(M; F)$ 

a first order elliptic differential operator. Write

$$
D = \Phi^F A (\Phi^E)^{-1} =: J_x \left( \frac{d}{dx} + B_x \right)
$$

as in [\(2.16\)](#page-8-2), [\(2.17\)](#page-8-5) for suitable isometries  $\Phi^E$ ,  $\Phi^F$ .

Let T be an invertible bundle homomorphism from  $E_{\Sigma}$  to  $F_{\Sigma}$  and consider the boundary condition for  $\tilde{A} := A \oplus (-A^t)$  given by

$$
(4.14) \t\t P(T) = (-T \t\t Id).
$$

Assume furthermore that

<span id="page-22-2"></span> $(4.15)$  $t_0^t T$  is positive definite, in particular selfadjoint.

Then

(a)  $P(T)$  is strongly regular for  $\widetilde{A} := A \oplus (-A^t)$ .

(b) The dual condition is given by  $P(T^{dual}) = P(-T^{-1})$ . It is strongly regular for  $\widetilde{A}^t.$ 

(c) The operator  $A_{P(T)}$  is Fredholm with compact resolvent and

$$
\ker \tilde{A}_{P(T)} = Z_{+,0}(A) \oplus Z_{-,0}(A),
$$
  
coker  $\tilde{A}_{P(T)} \simeq Z_{-,0}(A) \oplus Z_{+,0}(A).$ 

(d) Finally, if A and  $A^t$  satisfy weak inner UCP then  $\widetilde{A}_{P(T)}$  is invertible. Moreover in this case the inverse  $(\widetilde{A}_{P(T)})^{-1}$  maps  $L_s^2(M, F \oplus E)$  continuously to  $L_{s+1}^2(M, E \oplus E)$ F) for all  $s \geq 0$ .

Proof. We only have to comment on the very last statement. As in the proof of Proposition [2.4](#page-10-2) we infer from the strong regularity that on  $\{f \in L^2_k \mid$  $P(T)(\varrho f) = 0$  we have estimates

(4.16) 
$$
\frac{1}{C}||f||_k \leq ||f||_{k-1} + ||\widetilde{A}f||_{k-1} \leq C||f||_k.
$$

Hence  $\widetilde{A}_{P(T)}^{-1}$  maps  $L_k^2$  continuously to  $L_{k+1}^2$ ,  $k \in \mathbb{Z}_+$ . Now the claim follows from complex interpolation.

Remark 4.8. We emphasize that the condition [\(4.15\)](#page-22-2) holds for

(4.17) 
$$
T \in \left\{J_0, (J_0^t)^{-1}, (J_0J_0^t)^{-1/2}J_0\right\}.
$$

# <span id="page-23-5"></span>5. Calderón projection from the invertible double

<span id="page-23-1"></span><span id="page-23-0"></span>5.1. Sobolev scale. Next we recall the purely functional analytic notion of the Sobolev scale of an operator (cf.  $[LM72]$  $[LM72]$  $[LM72]$ ,  $[BL01]$  $[BL01]$  $[BL01]$ ). For the moment let D be a closed operator in the Hilbert space H. For  $s \in \mathbb{R}$  let  $H_s(D)$  be the completion of

(5.1) 
$$
H_{\infty}(D) := \bigcap_{s \geq 0} \mathcal{D}((D^*D)^{s/2})
$$

with respect to the scalar product

(5.2) 
$$
\langle x, y \rangle_s := \langle (\mathrm{Id} + D^* D)^s x, y \rangle.
$$

Obviously  $H_1(D) = \mathscr{D}(D)$  and the scalar product  $\langle ., . \rangle_0$  extends to a perfect pairing between  $H_s(D)$  and  $H_{-s}(D)$ . Furthermore, the spaces  $H_s(D)$  have the interpolation property, that is for  $s < t$  and  $0 \le \Theta \le 1$  we have

<span id="page-23-2"></span>(5.3) 
$$
H_{\Theta t + (1 - \Theta)s}(D) = [H_s(D), H_t(D)]_{\Theta}
$$

in the sense of complex interpolation theory (cf. e.g. [[Tay96](#page-57-15), Sec. 4.2]).

Note that D induces bounded linear maps  $H_s(D) \to H_{s-1}(D^*)$ . We will mostly use the case  $|s| \leq 1$ .

Since  $H_{-1}(D)$  is canonically (C–anti–)isomorphic to the dual of  $H_1(D)$  it fol-lows from [\(5.3\)](#page-23-2) that  $H_s(D), |s| \leq 1$ , depends only on the spaces  $H_0(D)$  and  $H_1(D)$ ; it does not depend on the particular operator  $D$  generating the scale. This independence, of course, is not true for  $|s| > 1$ .

If the condition [\(4.15\)](#page-22-2)  $(J_0^t T > 0)$  is fulfilled, then in view of [\(4.4\)](#page-20-4) it is appro-priate to put<sup>[3](#page-23-3)</sup>

(5.4) 
$$
L_{s,-T^{-1}}^2(M, F \oplus E) := H_s((\widetilde{A}_{P(T)})^*), -1 \leq s \leq 1.
$$

<span id="page-23-4"></span>Obviously, for  $1/2 < s \leq 1$  we have by  $(4.7)$ 

(5.5) 
$$
L_{s,-T^{-1}}^2(M, F \oplus E) = \left\{ \begin{pmatrix} f_+ \\ f_- \end{pmatrix} \in L_s^2(M, E \oplus F) \ \middle| \ \varrho f_- = -T^{-1} \varrho f_+ \right\}.
$$

The latter definition makes sense also for  $s > 1$  but the equality [\(5.4\)](#page-23-4) is limited to  $|s| \leq 1$ .

We have by construction  $L^2_{s,T} \subset L^2_s$ , hence  $\varrho$  induces bounded linear maps

<span id="page-23-7"></span>(5.6) 
$$
L_{s,-T^{-1}}^2(M, F \oplus E) \longrightarrow L_{s-1/2}^2(\Sigma, F_{\Sigma}),
$$

$$
\begin{pmatrix} f \\ g \end{pmatrix} \longmapsto f \upharpoonright \Sigma, \qquad 1/2 < s \le 1.
$$

Denote by  $\varrho^*$  the  $L^2$ -dual of  $\varrho$ . I.e.,  $\varrho^*$  is a bounded linear map

<span id="page-23-6"></span>
$$
(5.7) \t\t p^*: L_s^2(\Sigma, F_{\Sigma}) \longrightarrow L_{s-1/2, -T^{-1}}^2(M, F \oplus E), -1/2 \le s < 0,
$$

<span id="page-23-3"></span><sup>3</sup>Lateron the following considerations will always be used for the dual boundary condition  $T<sup>dual</sup> = -T<sup>-1</sup>$ , see [\(4.7\)](#page-20-7). Therefore we present them already here for  $-T<sup>-1</sup>$  instead of T.

such that for  $\xi \in L^2_s(\Sigma, F_\Sigma)$  and  $f \in L^2_{-s+1/2,-T^{-1}}(M, F \oplus E)$  we have  $\langle \varrho^* \xi, f \rangle =$  $\langle \xi, \rho f \rangle$ .

<span id="page-24-0"></span>5.2. Induced Poisson type operators and inverses. We use the notation of the previous Section [4.](#page-19-0) Throughout the whole section we assume [\(4.15\)](#page-22-2)

<span id="page-24-1"></span>(5.8) 
$$
J_0^t
$$
 is positive definite, in particular selfadjoint

and additionally

<span id="page-24-4"></span>(5.9) 
$$
[J_0^t T, B_0^t] \text{ is of order } 0.
$$

<span id="page-24-6"></span>**Remark 5.1.**  $[J_0^t T, B_0^t] = 0$  for the choice  $T = (J_0^t)^{-1}$ .

If  $A = A<sup>t</sup>$  and  $B<sub>0</sub> - B<sup>t</sup><sub>0</sub>$  is of order 0 then  $[J<sub>0</sub><sup>t</sup>T, B<sub>0</sub><sup>t</sup>]$  is of order 0 for all three choices of  $T$  in  $(4.17)$ .

Recall from Remark [4.4](#page-20-6) that condition [\(5.8\)](#page-24-1) implies that the dual boundary condition for  $\widetilde{A}^t$  is then given by  $-T^{-1}$ .

According to Theorem [4.7](#page-22-3) the boundary condition  $P(T)$  is regular for  $\widetilde{A}$  and

<span id="page-24-5"></span>(5.10) 
$$
\ker \widetilde{A}_{P(T)} = Z_{+,0}(A) \oplus Z_{-,0}(A) = Z_{+,0}(\widetilde{A}),
$$

$$
\operatorname{coker} \widetilde{A}_{P(T)} \simeq Z_{-,0}(A) \oplus Z_{+,0}(A) = Z_{-,0}(\widetilde{A}).
$$

The orthogonal projections onto  $Z_{+,0}(A)$ ,  $Z_{-,0}(A)$  are denoted by  $P_{Z_{+,0}(\widetilde{A})}, P_{Z_{-,0}(\widetilde{A})}$ respectively.

In a collar of the boundary we expand  $A$  in the following form, omitting the explicit reference to  $\Phi^E, \Phi^F$  etc. (cf. [\(2.17\)](#page-8-5), [\(2.18\)](#page-8-3)),

<span id="page-24-2"></span>(5.11) 
$$
A = J_0 \left(\frac{d}{dx} + B_0\right) + C_1 x + C_0
$$

with a first order differential operator  $C_1$  and x-independent bundle morphism  $C_0 \in \Gamma^\infty(\Sigma; \text{Hom}(E_\Sigma, F_\Sigma))$ . Here and in the sequel, by slight abuse of notation, x will also denote the operator of multiplication by the function  $x \mapsto x$ .

To see [\(5.11\)](#page-24-2) we expand J and B near  $x = 0$ 

(5.12) 
$$
J_x = J_0 + J_x^{(1)}x = J_0 + J_0'x + J_x^{(2)}x^2
$$

$$
B_x = B_0 + B_x^{(1)}x.
$$

Noting that  $\left[\frac{d}{dx}, x\right] = 1$  we find

(5.13)  
\n
$$
J_x \frac{d}{dx} = J_0 \frac{d}{dx} + J_x^{(1)} x \frac{d}{dx}
$$
\n
$$
= J_0 \frac{d}{dx} + (J_x^{(1)} \frac{d}{dx}) x - J_x^{(1)}
$$
\n
$$
= J_0 \frac{d}{dx} + (J_x^{(1)} \frac{d}{dx} - J_x^{(2)}) x - J_0'
$$
\n
$$
J_x B_x = J_0 B_0 + (J_0 B_x^{(1)} + J_x^{(1)} B_x) x,
$$

thus

(5.14) 
$$
C_1 = J_x^{(1)} \frac{d}{dx} - J_x^{(2)} + J_0 B_x^{(1)} + J_x^{(1)} B_x,
$$

$$
C_0 = -J'_0.
$$

<span id="page-24-3"></span>The formal adjoint can be written similarly as

(5.15) 
$$
A^t = \left(-\frac{d}{dx} + B_0^t\right)J_0^t + \widetilde{C}_1x + \widetilde{C}_0,
$$

where again  $\widetilde{C}_1$  is a first order differential operator and  $\widetilde{C}_0$  is an x-independent bundle morphism. More precisely,

(5.16) 
$$
\widetilde{C}_1 = C_1^t + \frac{1}{x} ([x, C_1^t] - [x, C_1^t]_{|x=0}),
$$

$$
\widetilde{C}_0 = C_0^t + [x, C_1^t]_{|x=0} = -(J_0')^t + (J_0')^t = 0,
$$

thus  $C_0$  in fact vanishes. Note that in [\(5.11\)](#page-24-2), [\(5.15\)](#page-24-3) x is intentionally on the right of  $C_1$ . From this it also becomes clear that  $\tilde{C}_0 = 0$  because in the expansion of  $A^t = \left(-\frac{d}{dx} + B_x^t\right)J_x^t$  near  $x = 0$  the commutator  $[x, \frac{d}{dx}]$  does not show up.

<span id="page-25-3"></span>**Remark 5.2.** We note that if the tangential operator  $B_0$  has a selfadjoint leading symbol we may replace  $B_0$  by  $\frac{1}{2}(B_0 + B_0^t)$  and still write  $A, A^t$  in the form  $(5.11), (5.15)$  $(5.11), (5.15)$  $(5.11), (5.15)$ . This changes, of course,  $C_0$  and  $C_0$ .

We fix a real number  $c > 0$  as in Section [3,](#page-11-0) Convention [3.7](#page-0-0) and consider the corresponding family of operators  $Q_{+}(x)$  of Definition [3.8.](#page-15-3) Then we define the following operators mapping (distributional) sections of  $E[\Sigma]$  into (distributional) sections of  $E[\mathbb{R}_+ \times \Sigma]$ :

<span id="page-25-2"></span>(5.17) 
$$
(R\xi)(x) := \begin{pmatrix} Q_+(x)\xi \\ Q_-(-x)^*\xi \end{pmatrix}, \quad R_T\xi := \begin{pmatrix} \mathrm{Id} & 0 \\ 0 & -T \end{pmatrix} R\xi.
$$

 $R_T$  will allow us to study the regularity properties of the (generalized) inverse of  $A_{P(T)}$  (cf. [\(5.27\)](#page-27-1) below) and of the Poisson operator (cf. Definition [5.9](#page-28-1) below). The Poisson operator is a map sending sections on the boundary into the kernel of A in the interior. R and  $R_T$  do almost have this property. For the constant coefficient operator  $A = J_0 \left( \frac{d}{dx} + B_0 \right)$  one has indeed  $\widetilde{A}Q_+ = 0$  by Proposition [3.11](#page-16-1) a,c. Even in the constant coefficient case  $\widetilde{A}R$  is not necessarily 0 but, thanks to  $(5.9)$ , small in a certain sense. This will become clear below. Note that  $R_T$  does not map into the domain of  $A_{P(T)}$ . Its role will become transparent in formula [\(5.21\)](#page-26-0) below.

For a cut–off function  $\varphi \in C_0^{\infty}(\mathbb{R}_+)$  we consider  $\varphi R_T$  as an operator from sections of  $E_{\Sigma} = E \Sigma$  to sections of  $E \oplus F$  over M; note that the range of  $\varphi R_T$ consists of sections vanishing outside a collar of  $\Sigma$ . From Proposition [3.16](#page-18-0) we infer that  $\varphi R_T$  maps  $L_s^2(\Sigma, E_\Sigma)$  continuously to  $L_{s+1/2, \text{comp}}^2(M, E \oplus F)$ ,  $s \ge -1/2$ .

<span id="page-25-0"></span>To calculate  $\widetilde{A}\varphi R_T$  we proceed by component:

(5.18) 
$$
A\varphi Q_{+}(x)\xi = \left((C_1x + C_0)\varphi(x) + J_0\varphi'(x)\right)Q_{+}(x)\xi
$$

and

<span id="page-25-1"></span>(5.19) 
$$
A^{t}\varphi TQ_{-}(-x)^{*}\xi = \left( (\widetilde{C}_{1}xT + \widetilde{C}_{0}T + [B_{0}^{t}, J_{0}^{t}T])\varphi(x) - J_{0}^{t}\varphi'(x) \right)Q_{-}(-x)^{*}\xi.
$$

The mapping properties of the right hand sides with respect to Sobolev spaces can be deduced from Proposition [3.16.](#page-18-0)

**Definition 5.3.** We write  $S(A,T)\xi$  for the differential expression  $\tilde{A}$  applied to  $\varphi R_T \xi$ .

**Remark 5.4.** The "differential expression" is emphasized here since  $\varphi R_T$  does not map into the domain of  $A_{P(T)}$ . However, by duality (cf. Subsection [5.1\)](#page-23-1)  $A_{P(T)}$ may also be viewed as a bounded operator  $L^2(M, E \oplus F) \longrightarrow L_{-1, -T^{-1}}^2(M, F \oplus E)$ . This should be viewed as applying  $\tilde{A}$  in the distributional sense.

The distinction between  $S(A,T)$  and  $\overline{A}_{P(T)}$  acting on  $L^2(M, E \oplus F)$  is crucial. The difference between the two operators is (see [\(5.21\)](#page-26-0) below)  $\varrho^* J_0(P_+ + P_-^*)$ .

 $S(A, T)$  will allow us to control the error (in terms of regularity, not in terms of size) between the approximate Poisson operator constructed from  $R_T$  and the true Poisson operator.

 $S(A, T)$  also depends on the choice of c in Convention [3.7,](#page-0-0) but this will be suppressed in the notation.  $S(A, T)$  is a  $2 \times 1$  column consisting (up to sign) of the right hand sides of [\(5.18\)](#page-25-0) and [\(5.19\)](#page-25-1).

We single out an immediate but important consequence of Proposition [3.16:](#page-18-0)

#### <span id="page-26-1"></span>Proposition 5.5.

 $S(A,T)$  maps  $L_s^2(\Sigma, E_{\Sigma})$  continuously to  $L_{s+1/2, comp}^2(M, F \oplus E)$ ,  $s \ge -1/2$ .

From the mapping properties of  $\varphi R_T$  we conclude in particular that for  $\xi \in$  $L_s^2(\Sigma, E_{\Sigma}), s \geq -1/2$ , we have  $\varphi R_T \xi \in L^2(M, E \oplus F) = H_0(\widetilde{A}_{P(T)})$ . Hence we may apply  $\widetilde{A}_{P(T)} \in \mathcal{B}(L^2(M, E \oplus F), L^2_{-1,-T^{-1}}(M, F \oplus E))$  (cf. Subsection [5.1\)](#page-23-1) to  $\varphi R_T$ . Recall that this is defined by duality and the result will be different from the differential expression  $\widetilde{A}$  applied to  $\varphi R_T \xi$ . Indeed for  $f \in \mathscr{D}((\widetilde{A}_{P(T)})^*)$  $L^2_{1,-T^{-1}}(M, F \oplus E)$  we find using Green's formula, [\(5.8\)](#page-24-1) and the boundary condition  $\varrho_{-}f = -T^{-1}\varrho_{+}f; \, \varrho_{\pm}f := \varrho(f_{\pm})$ :

(5.20)  
\n
$$
\langle \varphi R_T \xi, (\widetilde{A}_{P(T)})^* f \rangle
$$
\n
$$
= \langle J_0 P_+ \xi, \varrho_+ f \rangle - \langle J_0^t T P_-^* \xi, \varrho_- f \rangle + \langle S(A, T) \xi, f \rangle
$$
\n
$$
= \langle J_0 (P_+ + P_-^*) \xi, \varrho f \rangle + \langle S(A, T) \xi, f \rangle
$$
\n
$$
= \langle \varrho^* J_0 (P_+ + P_-^*) + S(A, T) \rangle \xi, f \rangle.
$$

Here  $P_{\pm}$  denote the positive/negative sectorial spectral projections of  $B_0$  in the sense of Definition [3.10.](#page-16-2) Recall  $P_{+}(B_0^t) = P_{+}(B_0)^*$ .

<span id="page-26-0"></span>Thus as an identity in  $H_{-1}((\tilde{A}_{P(T)})^*) = L_{-1, -T^{-1}}^2(M, F \oplus E)$  we arrive at

(5.21) 
$$
\widetilde{A}_{P(T)} \varphi R_T \xi = ( \varrho^* J_0 (P_+ + P_-^*) + S(A, T) ) \xi.
$$

It is important to note that, even if  $\xi$  has better regularity than  $L_{-1/2}^2$ , this is just an identity in  $H_{-1}((\widetilde{A}_{P(T)})^*)$  since  $\varphi R_T \xi$  does not fulfill the boundary condition for  $A_{P(T)}$ . The boundary condition plays no role as long as we view  $\varphi R_T \xi$  as an element of  $L^2$ .

With some care we can now basically proceed as in  $[HKL04, Sec. 3.2 and 3.3]$  $[HKL04, Sec. 3.2 and 3.3]$  $[HKL04, Sec. 3.2 and 3.3]$ : Firstly we note that by Lemma [3.5](#page-14-4)  $P_+ + P_-^*$  is invertible. Secondly we introduce the (Hilbert space) pseudo–inverse  $G$  of  $A_{P(T)}$ , namely,

(5.22) 
$$
\widetilde{G}f := \begin{cases} \widetilde{A}_{P(T)}^{-1}f, & f \in \operatorname{im} \widetilde{A}_{P(T)}, \\ 0, & f \in \operatorname{im} \left( \widetilde{A}_{P(T)} \right)^{\perp}, \end{cases}
$$

taking account of the possible absence of weak unique continuation. Here  $\widetilde{A}_{P(T)}^{-1}f$ denotes the inverse image in  $\left(\ker \widetilde{A}_{P(T)}\right)^{\perp}$  of f under  $\widetilde{A}_{P(T)}$ .  $L_{\text{P}}$ 

$$
\mathbf{Re}(\mathbf{r})
$$

(5.23) 
$$
U: L^2(M, E \oplus F) \longrightarrow L^2(M, F \oplus E)
$$

be the partial isometry which sends  $Z_{+,0}(\tilde{A})$  onto  $Z_{-,0}(\tilde{A})$  by interchanging the summands in  $(5.10)$  and which is zero on the orthogonal complement. Then

(5.24) 
$$
P_{Z_{+,0}(\widetilde{A})} = U^*U, \quad P_{Z_{-,0}(\widetilde{A})} = UU^*,
$$

and

(5.25) 
$$
\widetilde{G} = (\mathrm{Id} - U^*U)(\widetilde{A}_{P(T)} + U)^{-1}.
$$

We recall that  $Z_{+,0}(\widetilde{A}), Z_{-,0}(\widetilde{A})$  are finite–dimensional and consist of sections which are smooth up to the boundary (cf. Remark [2.5](#page-10-1) (1)). Hence  $U$  and  $U^*$  are smoothing operators. Furthermore, from  $\varrho U = 0$ ,  $\varrho U^* = 0$  we immediately obtain

<span id="page-27-3"></span>(5.26) 
$$
U^* \varrho^* = 0, \quad U \varrho^* = 0.
$$

By construction ker  $A_{P(T)} \subset H_{\infty}(A_{P(T)})$ . Hence  $A_{P(T)} + U$  induces invertible bounded linear maps from  $H_s(\widetilde{A}_{P(T)})$  onto  $H_{s-1}((\widetilde{A}_{P(T)})^*)$ . Consequently,  $\widetilde{G}$  induces bounded linear maps from  $H_s((\widetilde{A}_{P(T)})^*)$  onto  $H_{s+1}(\widetilde{A}_{P(T)})$ . Together with the mapping properties of  $\varrho^*$  we conclude that  $\tilde{G}\varrho^*$  maps  $L_s^2(\Sigma, F_\Sigma)$  continuously to  $L^2_{s+1/2}(M, E \oplus F)$  for  $-1/2 \leq s < 0$ .

 $G\varrho^*$  is the main building block of the Poisson operator (Definition [5.9](#page-28-1) below) which should act at least on  $L^2$ . Therefore we have to improve the bound on s, which is now straightforward:

To apply the pseudo–inverse  $\tilde{G}$  to [\(5.21\)](#page-26-0) it is enough that (5.21) is an identity in  $H_{-1}$ . Hence we find

<span id="page-27-1"></span>(5.27) 
$$
\widetilde{G}\varrho^* = \left( (\text{Id} - P_{Z_{+,0}(\widetilde{A})}) \varphi R_T - \widetilde{G}S(A,T) \right) (J_0(P_+ + P_-^*))^{-1}.
$$

<span id="page-27-2"></span>**Theorem 5.6.** For  $-1/2 \leq s \leq 1/2$  the operator  $\tilde{G}\varrho^*$  maps  $L^2_s(\Sigma, F_{\Sigma})$  continuously to  $L_{s+1/2}^2(M, E \oplus F)$ .

PROOF. This follows immediately from Proposition [3.16,](#page-18-0) [\(5.18\)](#page-25-0), [\(5.19\)](#page-25-1), [\(5.27\)](#page-27-1), and Proposition [5.5.](#page-26-1)

<span id="page-27-0"></span>5.3. The Calderón projection. From the invertible double the construction of the Calderón projection is straightforward. During the whole subsection we assume that the conditions [\(5.8\)](#page-24-1) and [\(5.9\)](#page-24-4) are fulfilled.

**Definition 5.7.** For a section  $f = \begin{pmatrix} f_+ \\ f_- \end{pmatrix}$  of  $E \oplus F$  we recall the notation  $\varrho_{\pm}(f) :=$  $\varrho(f_\pm)$  and  $r_\pm(f) := f_\pm$ . Furthermore, we put for sections f, g of E, F

.

(5.28) 
$$
e_{+}(f) := \begin{pmatrix} f \\ 0 \end{pmatrix}, e_{-}(g) := \begin{pmatrix} 0 \\ g \end{pmatrix}
$$

<span id="page-27-4"></span>**Proposition 5.8.**  $\widetilde{G}\varrho^*$  maps  $L_s^2(\Sigma, F_\Sigma)$  to  $Z_+^{s+1/2} \oplus Z_-^{s+1/2}, -1/2 \le s \le 1/2$ .

PROOF. In view of Theorem [5.6](#page-27-2) it remains to be shown that  $\tilde{G}\varrho^*$  maps into the kernel of  $A$ .

Let  $f \in L_s^2(\Sigma, F_\Sigma), -1/2 \le s \le 1/2$ , and a test function  $\varphi \in \Gamma_0^{\infty}(M \setminus \Sigma; F \oplus E)$ be given. In view of [\(5.7\)](#page-23-6) we choose a real number s' with  $-1/2 \leq s' < 0, s' \leq$ s. Then, by [\(5.7\)](#page-23-6), we have  $\varrho^* f \in L^2_{s'-1/2,-T^{-1}}(M, F \oplus E)$  and thus  $\widetilde{G}\varrho^* f \in$  $H_{s'+1/2}(\tilde{A}_{P(T)})$ ; note  $s'+1/2 \geq 0$ . Since  $\varphi$  has compact support away from  $\Sigma$ we certainly have  $\varphi \in H_{\infty}((\tilde{A}_{P(T)})^*i)$ . Hence  $\langle \tilde{A}\tilde{G}\varrho^*f, \varphi \rangle = \langle \tilde{G}\varrho^*f, (\tilde{A}_{P(T)})^*\varphi \rangle$ . Viewing the rhs as the dual pairing between  $H_{s'+1/2}(A_{P(T)})$  and  $H_{-s'-1/2}(A_{P(T)})$ we may also move  $\widetilde{G}$  to the right and find  $\langle \widetilde{A}\widetilde{G}\varrho^*f, \varphi \rangle = \langle \varrho^*f, \widetilde{G}^*(\widetilde{A}_{P(T)})^*\varphi \rangle$ , cf. Subsection [5.1.](#page-23-1) By construction of the generalized inverse we have, note again that  $\varphi \in H_{\infty}((\widetilde{A}_{P(T)})^*) \subset L^2(M, F \oplus E),$ 

(5.29) 
$$
\widetilde{G}^*(\widetilde{A}_{P(T)})^*\varphi = (I - U^*U)\varphi.
$$

However,  $U^*U\varphi \in H_\infty(\widetilde{A}_{P(T)})$  and  $\varrho U^*U\varphi = 0$ , cf. [\(5.26\)](#page-27-3). Thus  $\langle \varrho^*f, (\mathrm{Id} - U^*U)\varphi \rangle =$  $\langle f, \varrho \varphi \rangle = 0.$ 

This calculation shows that if  $\widetilde{A}$  is applied in the weak sense to  $\widetilde{G}\varrho^*f$  one gets  $\widetilde{G}\varrho^*f \in Z^{s+1/2} \oplus Z^{s+1/2}$ 0. But then  $\widetilde{G}\varrho^* f \in Z^{s+1/2}_+ \oplus Z^{s+1/2}_ \frac{s+1}{2}$ .

# <span id="page-28-1"></span>Definition 5.9.

(1) Define the Poisson operator by

$$
K_{\pm} := \pm r_{\pm} \widetilde{G} \varrho^* J_0.
$$

 $K_{\pm}$  maps  $L_s^2(\Sigma, E_{\Sigma})$  continuously to  $L_{s+1/2}^2(M, E)$   $(L_{s+1/2}^2(M, F))$  for  $-1/2 \leq s \leq 1/2$ .

(2)

$$
C_+ := \varrho_+ K_+, \quad C_- := T^{-1} \varrho_- K_-.
$$

 $C_{+}$  is called the Calderón projection of A. Recall that T is the operator defining the boundary condition for  $\widetilde{A}$ .

 $K_{\pm}$  and  $C_{\pm}$  depend on the pair  $(A, T)$ .

We summarize the result of the construction before Theorem [5.6,](#page-27-2) cf. in particular [\(5.27\)](#page-27-1):

<span id="page-28-3"></span>**Proposition 5.10.** Let  $\varphi \in C_0^{\infty}(\mathbb{R}_+)$  and let  $R_T$  be defined as in [\(5.17\)](#page-25-2). Furthermore, let  $P_+ := P_+(B_0), P_- := P_-(B_0)$  be the positive respectively negative sectorial spectral projections of  $B_0$  as introduced in Definition [3.10](#page-16-2), cf. Theorem [3.14](#page-17-3).

<span id="page-28-2"></span>Then the Poisson operators are given by

(5.30) 
$$
K_{\pm} = \pm r_{\pm} \Big( (\text{Id} - P_{Z_{+,0}(\widetilde{A})}) \varphi R_T - \widetilde{G}S(A,T) \Big) (P_+ + P_-^*)^{-1},
$$

and the Calderón projections are given by (see also  $(5.6)$  and  $(5.7)$ )

<span id="page-28-0"></span>(5.31) 
$$
C_{+} = (P_{+} - \rho_{+}\tilde{G}S(A,T))(P_{+} + P_{-}^{*})^{-1}
$$

$$
C_{-} = (P_{-} + T^{-1}\tilde{G}S(A,T))(P_{+} + P_{-}^{*})^{-1}.
$$

PROOF. The Theorem follows immediately from  $(5.27)$ .

**Remark 5.11.** Note that in the formula for  $K_{+}$  in [\(5.30\)](#page-28-2)  $R_T$  can be replaced by R (cf.  $(5.17)$ ), hence the first summand for  $K_{+}$  is independent of T.

Note that "our" Calderón projection differs from  $P_{+,\text{ort}} = P_+(P_+ + P_-^*)^{-1}$ (cf. Lemma [3.5,](#page-14-4) [\(3.5\)](#page-14-3)) by an operator which regularizes by at least one Sobolev order. So our construction of the invertible double yields naturally a version of the orthogonalized Calderón projection; if  $T = (J_0^t)^{-1}$  then  $C_+$  is indeed an orthogonal projection, see the next Proposition.

This is in contrast to the *classical* Calderón projection  $\mathscr{P}_+$  of Seeley [[See69](#page-57-4)] which is a pseudodifferential operator of order 0 whose leading symbol coincides with that of  $P_+$ . Hence our  $C_+$  differs from the orthogonalized Calderón projection  $\mathscr{P}_{+}$ <sub>ort</sub> by an operator which regularizes by at least one Sobolev order. With some more work one can indeed show that  $C_+$  is a pseudodifferential operator which differs from  $\mathscr{P}_{+,\text{ort}}$  by an operator of order -1.

If  $B_0 = B_0^t$ , or more generally if  $B_0$  has a selfadjoint leading symbol, then Proposition [5.10](#page-28-3) shows in particular the well–known fact (cf. [[BBW93](#page-56-1), Corollary 14.3]) that  $C_+ - P_+(B_0)$  is an operator of order -1.

Our approach also reproves a stronger result in the product situation: namely assume that  $A = J_0(\frac{d}{dx} + B_0)$  in a collar of the boundary with  $B_0 = B_0^t$ . Using the following modified version<sup>[4](#page-29-1)</sup> of  $R$ 

(5.32) 
$$
(R'\xi)(x) := \begin{pmatrix} Q_+(x)\xi \\ (J_0^t T)^{-1}Q_-(-x)^*(J_0^t T)\xi \end{pmatrix}, \quad R'_T\xi := \begin{pmatrix} \mathrm{Id} & 0 \\ 0 & -T \end{pmatrix} R'\xi,
$$

one then has  $\widetilde{A}R_T \xi = 0$  and hence  $S(A, T)\xi$  is supported in supp  $\varphi'$  away from the boundary. Then it is not difficult to see that  $S(A, T)$  is smoothing (cf. [**[HKL04](#page-57-6)**, Prop 3.15 and Eq. (3.38)]) and thus  $C_{+} - P_{+}(B_{0})$  is a smoothing operator.

This result was proved by G. Grubb [[Gru99](#page-57-17), Prop. 4.1]. Before it was shown by S. Scott [[Sco95](#page-57-18), Prop. 2.2] for selfadjoint Dirac operators on spin manifolds in the case ker  $B_0 = 0$ .

In general,  $C_+$  and  $P_+(B_0)$  belong to different connected components of the Grassmannian of pseudodifferential projections with fixed leading symbol even for symmetric  $B_0$ , see [[BBW93](#page-56-1), Remark 22.25]. For operators of Dirac type, however,  $C_+$  can be continuously deformed into a finite range perturbation of  $P_+(B_0)$  in the  $L^2$  and the  $L^2_{1/2}$  operator topology, see [[Nic97](#page-57-0), Corollary C.3].

<span id="page-29-0"></span>**Proposition 5.12.**  $C_{\pm}$  are idempotents with  $C_{+} + C_{-} =$  Id and

$$
C_{+}(L_s^2) = N_+^s,
$$
  
\n
$$
C_{-}(L_s^2) = T^{-1}N_-^s,
$$
  $-1/2 \le s \le 1/2.$ 

Furthermore if  $T := (J_0^t)^{-1}$  then  $C_{\pm}^* = C_{\pm}$ , i.e.,  $C_{\pm}$  act as orthogonal projections on  $L^2$ . In that case  $C_{\pm} L_0^2$  are  $L^2$  extensions of pseudodifferential projections.

<span id="page-29-2"></span>Remark 5.13. In view of the previous Proposition and Theorem [4.7](#page-22-3) we can always construct  $A_{P(T)}$  in such a way that  $C_{\pm}$  are orthogonal projections (as mentioned, by choosing  $T := (J_0^t)^{-1}$ . However, even if A is symmetric it may happen that  $A_{P(T)}$  is not selfadjoint.

If  $A = A^t$  and if  $T = J_0(-J_0^2)^{1/2}$  satisfies [\(5.9\)](#page-24-4) we may construct a selfadjoint extension of  $\widetilde{A}$  at the cost of a non-orthogonal Calderón projection.

Only if  $J_0^2 = -Id$  and  $T = (J_0^t)^{-1}$  then  $\tilde{A}_{P(T)}$  is selfadjoint and  $C_{\pm}$  are orthogonal projections.

PROOF OF PROPOSITION [5.12.](#page-29-0) We already know from Proposition [5.8](#page-27-4) that

$$
C_{+}(L_s^2) \subset N_+^s,
$$
  

$$
C_{-}(L_s^2) \subset T^{-1}N_-^s.
$$

We show

(i) 
$$
N_+^s \cap T^{-1}N_-^s = \{0\},
$$
  
(ii)  $C_+ + C_- = \text{Id}.$ 

This easily implies the first claim.

(i) Let  $\xi \in N^s_+ \cap T^{-1}N^s_-$ . Then there are  $f \in Z^{s+1/2}_+$ ,  $g \in Z^{s+1/2}_-$  with  $\varrho f =$  $\xi = T^{-1} \varrho g$ . Then

(5.33) 
$$
\begin{pmatrix} f \ g \end{pmatrix} \in \ker \widetilde{A}_{P(T)} = Z_{+,0}(\widetilde{A}).
$$

Since elements of  $Z_{+,0}(\widetilde{A})$  vanish on the boundary we infer  $\xi = 0$ .

<span id="page-29-1"></span><sup>&</sup>lt;sup>4</sup>The whole discussion after [\(5.17\)](#page-25-2) can be carried out with this modified  $R'$ , too.

(ii) Let  $\xi \in L^2(\Sigma, E_\Sigma), -1/2 \leq s \leq 1/2$ , and  $f \in \mathscr{D}((\tilde{A}_{P(T)})^*)$ . Then  $\varrho_+ f =$  $-T\varrho-f$  (cf. Remark [4.4\)](#page-20-6) and exploiting the selfadjointness of  $J_0^tT$  we obtain

(5.34)  
\n
$$
\langle (C_+ + C_-)\xi, J_0^t \varrho_+ f \rangle
$$
\n
$$
= \langle \varrho_+ \widetilde{G} \varrho^* J_0 \xi - T^{-1} \varrho_- \widetilde{G} \varrho^* J_0 \xi, J_0^t \varrho_+ f \rangle
$$
\n
$$
= \langle \varrho_+ \widetilde{G} \varrho^* J_0 \xi, J_0^t \varrho_+ f \rangle - \langle \varrho_- \widetilde{G} \varrho^* J_0 \xi, J_0 J_0^{-1} (T^{-1})^t J_0^t \varrho_+ f \rangle
$$
\n
$$
= \langle (\varrho_+ \oplus \varrho_-) \widetilde{G} \varrho^* J_0 \xi, (J_0^t \oplus J_0) (\varrho_+ f \oplus \varrho_- f) \rangle
$$
\n
$$
= \langle \widetilde{G} \varrho^* J_0 \xi, (\widetilde{A}_{P(T)})^* f \rangle
$$
\n
$$
= \langle \varrho^* J_0 \xi, f \rangle = \langle \xi, J_0^t \varrho f \rangle.
$$

This proves (ii).

Finally, let  $T = (J_0^t)^{-1}$  and pick  $\xi \in N^0_+, \eta \in T^{-1}N^0_-\text{. Choose } f \in Z^{1/2}_+$  with  $\rho_{+}f = \xi$  and  $g \in Z_{-}^{1/2}$  with  $T^{-1}\rho_{+}g = \eta$ . Then Green's formula Lemma [2.2](#page-7-2) gives (5.35)  $\langle \xi, \eta \rangle = \langle J_0 \varrho_+ f, \varrho_+ g \rangle = -\langle A f, g \rangle + \langle f, A^t g \rangle = 0.$ 

Hence  $N^0_+ \perp T^{-1} N^0_-$  and we are done.

To prove the pseudodifferential property we recall from [[See69](#page-57-4), Appendix] Seeley's construction of the Calderón projection which always yields a pseudodifferential projection  $\mathscr{P}_{+}^{s}$  onto  $N_{+}^{s}$ . By orthogonalization, cf. Lemma [3.5,](#page-14-4) we obtain an orthogonal pseudodifferential projection onto  $N_{+}^{s}$  which must coincide with  $C_{+}$ for  $s = 0$ .

# 6. The General Cobordism Theorem

<span id="page-30-0"></span>In the preceding sections, we gave a new definition of the Calderón projection. We achieved a canonical construction, free of extensions and other choices, and in greatest generality. Our main goal with the new definition was a construction which admits to follow precisely a continuous variation of the coefficients of an elliptic differential operator up to the induced variation of the Calderon projection. We shall return to this application below in Section [7.](#page-41-1)

An added bonus of our construction of the Calderón projection is that it leads immediately, and somewhat surprisingly, to a simple proof and a wide generalization of the classical Cobordism Theorem. We shall now give five different formulations of the Cobordism Theorem and show that the first claim (I), expressed in the language of symplectic functional analysis, follows immediately from our construction of the Calderón projection, and that the four other definitely non–trivial claims (II)-(V) are easily derived from the first claim. Put differently, we shall show that the Cobordism Theorem and its various generalizations and reformulations are an almost immediate consequence of our construction of the Calderón projection.

In all of this section we shall assume that our first order elliptic operator A over the smooth compact manifold M with boundary  $\Sigma$  is formally selfadjoint, i.e.,  $A = A<sup>t</sup>$ . For convenience we shall write  $B_0 := B(0)$  and  $J_0 := J(0)$  as in the previous sections.

<span id="page-30-1"></span>6.1. The General Cobordism Theorem. The main result of this section is

<span id="page-30-2"></span>**Theorem 6.1** (The General Cobordism Theorem). Let  $A : \Gamma^\infty(M; E) \to \Gamma^\infty(M; E)$ be a first order formally selfadjoint elliptic differential operator on a smooth compact manifold M with boundary acting between sections of the vector bundle E. We assume that [\(5.9\)](#page-24-4) is satisfied<sup>[5](#page-30-3)</sup> by  $T = J_0(-J_0^2)^{-1/2}$ 

<span id="page-30-3"></span><sup>&</sup>lt;sup>5</sup>As noted in Remark [5.1](#page-24-6) this is the case if  $B_0 - B_0^t$  is of order 0.

#### Then we have the following results:

(I) Let  $C_{\pm}$  denote the Calderón projections introduced in Definition [5.9](#page-28-1), constructed from the invertible double with  $T \in \{(J_0^t)^{-1}, J_0, J_0(-{J_0}^2)^{-1/2}\}\$ . Then the range of  $C_+$  is a Lagrangian subspace of the strongly symplectic Hilbert space  $(L^2(\Sigma, E_{\Sigma}), -J_0)$ . Note that im  $C_+$  is independent of T. Moreover, im  $C_-$  is also Lagrangian, if  $T := J_0(-J_0^2)^{-1/2}$ .

(II) We have sign i $P_0J_0 \upharpoonright W_0 = 0$ . Here  $W_0$  denotes the (finite-dimensional) sum of the generalized eigenspaces of  $B_0$  corresponding to purely imaginary eigenvalues and  $P_0$  denotes the orthogonal projection onto  $W_0$ ; in general  $J_0$  will not map  $W_0$  to itself.

If  $B_0 = B_0^t$ , then  $J_0$  anticommutes with  $B_0$  and we have sign  $iJ_0$  ker  $B_0 = 0$ .

(III) Under the same additional assumption, i.e., for  $B_0 = B_0^t$ , the tangential operator  $B_0$  is odd with respect to the grading given by the unitary operator  $\alpha$  :=  $iJ_0(-J_0^2)^{-1/2}$  and hence splits into matrix form  $B_0 = \begin{pmatrix} 0 & B^{-1} \\ B^{-1} & 0 \end{pmatrix}$  $B^+$  0  $\Big)$  with respect to the  $\pm 1$ –eigenspaces of  $\alpha$ . The index of  $B^+$ : ker( $\alpha - 1$ )  $\longrightarrow$  ker( $\alpha + 1$ ) vanishes.

(IV) While we do not know whether  $C_+$  is a pseudodifferential operator for  $T = J_0 (J_0^2)^{-1/2}$ , we can prove the following:

There exists a pseudodifferential projection P over  $\Sigma$  such that ker P is a Lagrangian subspace of the strongly symplectic Hilbert space  $(L^2(\Sigma, E_{\Sigma}), -J_0)$ , and (ker P, im  $C_+$ ) is a Fredholm pair of closed subspaces of  $L^2(\Sigma, E_{\Sigma})$ .

(V) There exists a selfadjoint pseudodifferential Fredholm extension  $A_P$ .

In the following, we shall first give our view of elements of symplectic functional analysis. Based on that, we shall prove the preceding theorem as indicated above, i.e., we shall prove  $(I)$  directly from our new construction of the Calderón projection, then the implications  $(I) \implies (II) \implies (III)$ , and then  $(II) \implies (IV)$ ,  $(V)$ .

To us, our order of proving Theorem [6.1](#page-30-2) is the most simple and natural, beginning with and footing on claim (I). However, at the end of this section we shall explain that one can reverse the order of the proof. In particular, we shall show that  $(V)$  was essentially proved by Ralston in 1970 in [[Ral70](#page-57-10)] and that  $(I)$  can be derived from (V) independently.

Remark 6.2. (1) An alternative reading of (III) is the following: the index of any elliptic first order differential operator  $C$  with smooth coefficients over a smooth closed manifold  $\Sigma$  must vanish, if the block operator  $B := \begin{pmatrix} 0 & C^t \\ C & 0 \end{pmatrix}$  $C = 0$  can be written as the tangential operator of an elliptic formally selfadjoint operator A on a smooth compact manifold M with  $\partial M = \Sigma$ . That is a new generalization of the illustrious Cobordism Theorem for Dirac operators on spin manifolds which played a decisive role for the first proof of the Atiyah-Singer Index Theorem (1963). Since then it was slightly generalized and an impressive variety of different proofs were given. Our point here is to show that the Cobordism Theorem has nothing to do with the specific form of Dirac type operators but generalizes to all elliptic formally selfadjoint differential operators of first order.

(2) To some extent, our approach is motivated by [[BL01](#page-57-12), Section 1.C], and is similar to C. Frey's [[Fre05](#page-57-19), Section 3.4]. The main difference between our approach and Frey's is that we reduce the arguments a bit more to purely algebraic reasoning. That permits us to get through also in the general case, not assuming Dirac type, i.e., not assuming constant coefficients near the boundary in normal direction - as Frey does and all the other literature on cobordism invariance.

(3) From a geometric point of view, the assumption of constant coefficients may appear sufficiently general for many applications, justified for index problems by  $K$ –theory and homotopy invariance of the index, and sufficiently challenging for constantly attracting ever new and more simple and more ingenious proofs of the cobordism invariance of the index over the last 50 years. For a recent example and summary of the highlights we refer to Braverman [[Bra02](#page-57-20)]. However, from an analysis point of view, it is not natural to assume constant coefficients near the boundary. Moreover, it seems more than timely that the pilot work by Ralston, admitting general coefficients and so showing a way for our General Cobordism Theorem, is taken into account in the analysis and topology literature.

<span id="page-32-0"></span>6.2. Elements of symplectic functional analysis. Let us briefly summarize the basic set–up of symplectic functional analysis (see, e.g., [[BBF98](#page-56-3)], [[BBZ04](#page-57-21)] or [[KL04](#page-57-22)]).

6.2.1. Basic definitions. Let  $H$  be a real or complex Hilbert space. A weakly symplectic form on H is a non–degenerate Hermitian sesqui–linear form  $\omega$  on H. Sesqui–linear means, following the tradition in mathematical physics,  $\omega(\lambda x, y)$  =  $\overline{\lambda}\omega(x,y), \omega(x,\lambda y) = \lambda \omega(x,y)$ , and Hermitian means  $\omega(y,x) = -\overline{\omega(x,y)}$  for  $x, y \in$ H. Finally, non-degeneracy means that the map  $H \ni x \mapsto \omega(x, \cdot) \in H^*$  is an injective continuous linear map; it then follows that the range of this map is dense.

The pair  $(H, \omega)$  is called a (complex) weakly symplectic Hilbert space. Since  $\omega$ is continuous there is a unique skew–symmetric injective map  $\gamma \in \mathscr{B}(H)$  such that

(6.1) 
$$
\omega(x, y) = \langle \gamma x, y \rangle, \quad x, y \in H.
$$

Here as before,  $\mathscr{B}(H)$  denotes the space of bounded endomorphisms of H. For a subspace  $\lambda \subset H$  we have

(6.2) 
$$
\lambda^{\omega} := \{ x \in H \mid \omega(x, y) = 0 \text{ for } y \in \lambda \} = (\gamma \lambda)^{\perp},
$$

hence  $\lambda$  is Lagrangian (i.e.  $\lambda^{\omega} = \lambda$ ) if and only if  $\lambda = (\gamma \lambda)^{\perp}$ .

The pair  $(H, \omega)$  is called *strongly* symplectic, if the injective operator with dense range  $H \ni x \mapsto \omega(x, \cdot) \in H^*$  is in fact surjective and hence has a bounded inverse. Equivalently, the skew–symmetric operator  $\gamma$  which implements  $\omega$  is invertible.

In a strongly symplectic Hilbert space we may choose an equivalent scalar product  $\langle \cdot, \cdot \rangle_{\gamma}$  such that the operator which implements  $\omega$  with respect to  $\langle \cdot, \cdot \rangle_{\gamma}$  is the *unitary reflection*  $\gamma(-\gamma^2)^{-1/2}$ : namely

<span id="page-32-1"></span>(6.3) 
$$
\omega(x, y) = \langle \gamma x, y \rangle = \langle (-\gamma^2)^{1/2} \gamma (-\gamma^2)^{-1/2} x, y \rangle = \langle \gamma (-\gamma^2)^{-1/2} x, y \rangle_{\gamma}
$$

with  $\langle \xi, \eta \rangle_{\gamma} := \langle (-\gamma^2)^{1/2} \xi, \eta \rangle$ .

The scalar product  $\langle \cdot, \cdot \rangle_{\gamma}$  is equivalent to  $\langle \cdot, \cdot \rangle$  in the sense that there is a constant  $C$  such that

(6.4) 
$$
C^{-1}\langle x,x\rangle \leq \langle x,x\rangle_{\gamma} \leq C\langle x,x\rangle, \quad x \in H.
$$

In view of [\(6.3\)](#page-32-1) the operator which implements  $\omega$  with respect to  $\langle \cdot, \cdot \rangle_{\gamma}$  is the unitary reflection  $\gamma(-\gamma^2)^{-1/2}$ .

Finally, we comment on isomorphisms: let  $R : (H_1, \omega_1) \longrightarrow (H_2, \omega_2)$  be an invertible bounded linear map which is symplectic but not necessarily isometric. If  $\omega_j(\cdot, \cdot) = \langle \gamma_j \cdot, \cdot \rangle, j = 1, 2$  with skew–symmetric  $\gamma_j$  then  $\gamma_1 = R^* \gamma_2 R$ , in particular  $(H_1, \omega_1)$  is strongly symplectic if and only if  $(H_2, \omega_2)$  is.

<span id="page-32-2"></span>Remark 6.3. For simplicity, we assume from now on that all vector spaces, Hilbert spaces, and symplectic spaces are complex. Note, however, that our definition of symplectic Hilbert space does not require the existence of a Lagrangian subspace (as opposed to e.g.  $[KL04, Sec. 6]$  $[KL04, Sec. 6]$  $[KL04, Sec. 6]$ . Indeed, it might be that there are no Lagrangian subspaces at all. Take, e.g.,  $H := \mathbb{C}$  and  $\omega(x, y) := i\overline{x}y$ ; see also [**[KL04](#page-57-22)**, Rem. 6.13].

34 BERNHELM BOOSS-BAVNBEK, MATTHIAS LESCH, AND CHAOFENG ZHU

6.2.2. Algebraic observations. Dealing with elliptic problems on manifolds with boundary naturally leads to symplectic Hilbert spaces, see below Paragraph [6.2.4.](#page-34-0) However, keeping the arguments on a purely algebraic level where possible may make proofs more transparent. Moreover, Sobolev chains of symplectic Hilbert spaces are equipped with a variety of non–equivalent norms but compatible symplectic forms. For such applications it is nice to establish results independent of topological choices.

So, let  $(H, \omega)$  be a symplectic vector space; i.e., no boundedness of the symplectic form is assumed.

First we note that the quadratic form  $x \mapsto i\omega(x, x)$  has well–defined signature, if  $H$  is finite–dimensional. In that case we have

<span id="page-33-2"></span>(6.5) sign  $i\omega = 0 \iff \exists \lambda \subset H$  Lagrangian subspace,

cf. Remark [6.7.](#page-34-1)1.

Next, we recall a simple algebraic observation, taken from [[BBZ04](#page-57-21), Lemma 1.2]. Here, and in the proposition further below, the point is to establish the Lagrangian property for isotropic subspaces via a purely algebraic Fredholm pair property, i.e., finite–dimensional intersection and finite codimension of sum; no closedness is assumed. The algebraic Fredholm pair property can be considered as the natural generalization of the well-known definition of Lagrangian subspaces in finite dimension as isotropic subspaces of maximal dimension.

<span id="page-33-4"></span>**Lemma 6.4.** Let  $(H, \omega)$  be a symplectic vector space with transversal subspaces  $\lambda, \mu$ , i.e.,  $\lambda + \mu = H$ ,  $\lambda \cap \mu = 0$ . If  $\lambda, \mu$  are isotropic subspaces, then they are Lagrangian subspaces.

<span id="page-33-0"></span>PROOF. From linear algebra we have

$$
\lambda^{\omega} \cap \mu^{\omega} = (\lambda + \mu)^{\omega} = \{0\},
$$
  
since  $\lambda + \mu = H$ . From  
(6.6)  

$$
\lambda \subset \lambda^{\omega}, \mu \subset \mu^{\omega}
$$
  
we get  

$$
(6.7) \qquad H = \lambda^{\omega} \oplus \mu^{\omega}.
$$
  
Since  $H = \lambda \oplus \mu$  we conclude from (6.6) and (6.7) that  $\lambda$ 

Since  $H = \lambda \oplus \mu$  we conclude from [\(6.6\)](#page-33-0) and [\(6.7\)](#page-33-1) that  $\lambda = \lambda^{\omega}$  and  $\mu = \mu^{\omega}$ .

<span id="page-33-1"></span>The preceding result can be generalized:

**Proposition 6.5.** Let  $(H, \omega)$  be a symplectic vector space with isotropic subspaces  $\lambda, \mu$ . If  $(\lambda, \mu)$  forms an algebraic Fredholm pair with  $\text{ind}(\lambda, \mu) \geq 0$ , then  $\lambda$  and  $\mu$ are Lagrangian subspaces of H and we have

 $\text{ind}(\lambda, \mu) = 0, \ (\lambda + \mu)^{\omega} = \lambda \cap \mu, \text{ and } (\lambda + \mu)^{\omega \omega} = \lambda + \mu.$ 

For the proof we refer to [[BBZ04](#page-57-21), Proposition 1.13a].

6.2.3. Symplectic reduction. We recall a lemma on symplectic reduction from [[KL04](#page-57-22), Prop 6.12] (see also [[BBZ09](#page-57-23), Proposition 2.2] for a generalization to weakly symplectic Hilbert spaces):

<span id="page-33-3"></span>**Lemma 6.6.** Let  $(H, \omega)$  be a strongly symplectic Hilbert space,  $\lambda \subset H$  a Lagrangian subspace and  $W \subset H$  a closed co-isotropic subspace. Assume that  $\lambda + W^{\omega}$  is closed. Then the form

 $\widetilde{\omega}(x + W^{\omega}, y + W^{\omega}) := \omega(x, y), \quad x, y \in W$ 

is a strongly symplectic form on  $W/W^{\omega}$ . Moreover, the symplectic reduction of  $\lambda$ by W

(6.8) 
$$
\text{Red}_W(\lambda) := ((\lambda + W^{\omega}) \cap W) / W^{\omega} \subset W / W^{\omega}
$$

is a Lagrangian subspace of  $W/W^{\omega}$ .

<span id="page-34-1"></span>**Remark 6.7.** (1) Without the assumption  $\lambda + W^{\omega}$  closed  $\tilde{\omega}$  will still be a nondegenerate sesqui-linear form on the quotient  $W/W^{\omega}$ . However, in that case it might be that there are no Lagrangian subspaces at all as pointed out in Remark [6.3.](#page-32-2) If the form  $\tilde{\omega}$  is written as  $\tilde{\omega}(\xi, \eta) = \langle \tilde{\gamma}\xi, \eta \rangle$  with a suitable scalar product and a skew–symmetric unitary operator  $\tilde{\gamma}$ , then the existence of a Lagrangian subspace is equivalent to the fact that the  $\pm i$  eigenspaces of  $\tilde{\gamma}$  have the same dimension, corresponding to [\(6.5\)](#page-33-2).

(2) In [[KL04](#page-57-22), Prop. 6.12] the Lemma was formulated under seemingly more restrictive assumptions. Let us give an equivalent formulation of the Lemma which clarifies the link to loc. cit. and which will be useful below:

<span id="page-34-2"></span>**Lemma 6.8.** Let  $(H, \omega)$ ,  $\lambda \subset H$  and  $W \subset H$  be as in Lemma [6.6](#page-33-3). Let  $\gamma$  be the invertible skew–symmetric operator in H such that  $\omega(\cdot, \cdot) = \langle \gamma \cdot, \cdot \rangle$ .

Let  $W_0 \subset W$  be a closed subspace such that  $W = W_0 \oplus W^{\omega}$  (the sum is not necessarily orthogonal). Furthermore, let  $P_0$  denote the orthogonal projection onto  $W_0$  and let  $Q_0$  denote the projection along  $W^{\omega}$  onto  $W_0$ .

Then  $\omega \restriction W_0 = \langle P_0 \gamma, \cdot \rangle$  is a strongly symplectic form on  $W_0$ ,  $Q_0(\lambda \cap W)$  is a Lagrangian subspace of  $(W_0, \omega)$  and the quotient map  $\pi : W_0 \longrightarrow W/W^{\omega}$  is a bounded invertible symplectic linear operator.

Proof. Let us briefly sketch the proof of this and of the previous lemma. First as remarked at the beginning of this subsection we may choose a scalar product such that the corresponding  $\gamma$  is unitary. Consider first  $W_0 = (W^{\omega})^{\perp} = W \cap \gamma W$ . [[KL04](#page-57-22), Prop. 6.12] and its proof show that  $(W_0, \omega)$  is symplectic and that  $Q_0 = P_0$ maps  $\lambda \cap W$  onto a Lagrangian subspace of  $W_0$ . Furthermore, since  $W_0 = (W^{\omega})^{\perp}$  in this case, the quotient map  $\pi W_0 : (W_0, \omega) \to (W/W^{\omega}, \tilde{\omega})$  is a unitary symplectic isomorphism. This proves that  $(W/W^{\omega}, \tilde{\omega})$  is indeed a strongly symplectic Hilbert space and that  $\text{Red}_W(\lambda)$  is a Lagrangian subspace. Hence Lemma [6.6](#page-33-3) is proved.

To prove Lemma [6.8](#page-34-2) for an arbitrary closed subspace  $\widetilde{W}_0 \subset W$  with  $W =$  $\widetilde{W}_0 \oplus W^{\omega}$  we only have to note that we have the following commutative diagram

(6.9) 
$$
Q_{\widetilde{w}_0} \bigcup_{W_0}^{W_0} P_{w_0} \longrightarrow \widetilde{\pi}
$$

$$
W_0 \longrightarrow \widetilde{w}_0 W/W^{\omega}
$$

where  $\pi, \tilde{\pi}$  denote the quotient map  $W \longrightarrow W/W^{\omega}$  restricted to  $W_0, \widetilde{W}_0$  respectively,  $P_{W_0}$  denotes the orthogonal projection onto  $W_0$  and  $Q_{\widetilde{W}_0}$  denotes the projection along  $W^{\omega}$  onto  $\widetilde{W}_0$ .  $\pi, \widetilde{\pi}$  are symplectic bounded invertible maps. From this all remaining claims follow all remaining claims follow.

<span id="page-34-0"></span>6.2.4. The von-Neumann quotient of all natural boundary values. We recall the basic findings about selfadjoint extensions and the relation between Fredholm Lagrangian pairs in the von-Neumann quotient  $\mathscr{D}(A_{\text{max}})/\mathscr{D}(A_{\text{min}}) =: \beta(A)$  and in  $L^2(\Sigma,E_{\Sigma})$ :

Let  $A_m$  be a closed symmetric operator with domain  $\mathscr{D}_m$  in a Hilbert space H. Following von Neumann and the Russian tradition of M. Krein, Vishik, and Birman, the operator  $A_m$  defines a (strongly) symplectic Hilbert space  $\beta(A_m) := \mathscr{D}_{\text{max}}/\mathscr{D}_m$ of natural boundary values. Here  $\mathscr{D}_{\text{max}}$  denotes the domain of  $A_m^*$ . The Hilbert space structure on  $\beta(A_m)$  is given by the graph scalar product, and the symplectic form is given by Green's form

(6.10) 
$$
\omega(x + f, y + g) := \langle Ax, y \rangle - \langle x, Ay \rangle \text{ for } x, y \in \mathscr{D}_{\text{max}},
$$

independent of the choice of  $f, g \in \mathscr{D}_m$  (see our Equation [\(2.11\)](#page-7-1) and [[BBF98](#page-56-3), Section 3]). It is well known that there is a one-to-one correspondence between

• domains  $\mathscr{D}_m \subset \mathscr{D} \subset \mathscr{D}_{\text{max}}$  which yield a selfadjoint operator  $A_{\mathscr{D}} := A_m^* \mathcal{D}$ 

• and the Lagrangian subspaces  $\lambda$  of  $\beta(A_m)$ 

by

$$
\mathscr{D} \mapsto \lambda := \mathscr{D}/\mathscr{D}_m \quad \text{ and } \quad \lambda \mapsto \mathscr{D} := \{x \in H \mid x + \mathscr{D}_m \in \lambda\}.
$$

In our situation, we set  $H := L^2(M, E)$  and consider  $A_m = A_{\min} = A \, \otimes \, (A_{\min})$ with  $\mathscr{D}_m = \mathscr{D}(A_{\min}) = L^2_{1,0}(M,E)$ , the closure of  $\Gamma_0^{\infty}(M \setminus \Sigma; E)$  in  $L^2_1(M, E)$  like in Remark [2.5.](#page-10-1) Then  $\beta(A_m)$  naturally becomes a subspace of  $L^2_{-1/2}(\Sigma, E_{\Sigma})$ . From now on we will use this identification and view  $\beta(A_m)$  as a subspace of  $L^2_{-1/2}(\Sigma, E_{\Sigma})$ .

To discuss selfadjoint extensions by boundary conditions given by pseudodifferential projections, it is helpful to consider two other symplectic spaces: the strongly symplectic Hilbert space  $L^2(\Sigma, E_{\Sigma})$  with symplectic form induced by  $-J_0$  and the weakly symplectic Hilbert space  $L_1^2(M, E)/L_{1,0}^2(M, E)$  with symplectic form induced by Green's form, or, equivalently, by  $-J_0$ , as well. Here we identify the quotient with the subspace  $L_{1/2}^2(\Sigma, E_{\Sigma}) \subset L^2(\Sigma, E_{\Sigma})$  of sections over the boundary, but with different scalar product  $\langle \cdot, \cdot \rangle_{L^2_{1/2}(\Sigma, E_{\Sigma})}$ , hence  $J_0 = (\text{Id} + |B|)^{-1} J_0$ . Note that  $J_0$  is not invertible for dim  $M > 1$ , as explained in [[BBZ04](#page-57-21), Remark 1.6b].

The relations between the Lagrangian subspaces of these three different symplectic spaces are somewhat delicate because neither  $L^2(\Sigma, E_{\Sigma}) \subset \beta(A_m)$  nor  $L^2(\Sigma, E_{\Sigma}) \supset \beta(A_m)$ . We may, however, recall a very general result from [[BBFO01](#page-56-5), Theorem 1.2a]:

Let  $\beta$  and L be strongly symplectic Hilbert spaces with symplectic forms  $\omega_{\beta}$ and  $\omega_L$ , respectively. Let

<span id="page-35-0"></span>(6.11) 
$$
\beta = \beta_- \oplus \beta_+ \quad \text{and} \quad L = L_- \oplus L_+
$$

be direct sum decompositions by transversal (not necessarily orthogonal) pairs of Lagrangian subspaces. We assume that there exist continuous, injective mappings

<span id="page-35-1"></span>(6.12) 
$$
i_-: \beta_- \longrightarrow L_-
$$
 and  $i_+: L_+ \longrightarrow \beta_+$ 

with dense images and which are compatible with the symplectic structures, i.e.,

<span id="page-35-2"></span>(6.13) 
$$
\omega_L(i_-(x),a)=\omega_\beta(x,i_+(a)) \text{ for all } a\in L_+ \text{ and } x\in \beta_-.
$$

Let  $\lambda_0$  be a fixed Lagrangian subspace of  $\beta$ . We consider the Fredholm La*grangian Grassmannian* of  $\lambda_0$ 

 $\mathscr{FL}_{\lambda_0}(\beta) := \{ \mu \subset \beta \mid \mu \text{ Lagrangian subspace and } (\mu, \lambda_0) \text{ Fredholm pair} \}.$ 

The topology of  $\mathscr{FL}_{\lambda_0}(\beta)$  is defined by the operator norm of the orthogonal projections onto the Lagrangian subspaces.

<span id="page-35-3"></span>Theorem 6.9 (Booß-Bavnbek, Furutani, Otsuki). Under the assumptions [\(6.11\)](#page-35-0),  $(6.12)$ , and  $(6.13)$ , we have a natural continuous mapping

$$
\tau: \mathscr{FL}_{\beta-}(\beta) \longrightarrow \mathscr{FL}_{L-}(L), \quad \mu \longmapsto \mu \cap L,
$$

where  $\beta$  and L are identified with subspaces of  $\beta_+ \oplus L_-$ .

The following splitting lemmata are of independent interest. Note that we do not claim a direct sum decomposition of  $\beta(A)$  into Lagrangian subspaces for now. Later, this will be a consequence of our Theorem [6.1.](#page-30-2) See also the recent [[BBC08](#page-56-6), Section 1. In the following lemma, we could use our  $C_+$  instead of using Seeley's Calderón projection  $\mathscr{P}_+$ . All these direct sum decompositions of  $\beta(A)$  are different. but equally valid, see Lemma [6.11](#page-36-0) and Remark [6.12.](#page-36-1)

Lemma 6.10. Let A be an elliptic formally selfadjoint first order differential operator on a compact smooth manifold M with smooth boundary  $\Sigma$  and let  $\mathscr{P}_+$ denote Seeley's corresponding (pseudodifferential) Calderón projection. Then the space  $\beta(A) := \mathscr{D}(A_{\text{max}})/\mathscr{D}(A_{\text{min}})$  can be described explicitly as the direct sum

(6.14) 
$$
\beta(A) = \operatorname{im}(\mathscr{P}_{+,-1/2}) \oplus \operatorname{im}(\operatorname{Id} - \mathscr{P}_{+,1/2}),
$$

where  $\mathscr{P}_{+,s}$  denotes the extension/restriction of the pseudodifferential  $\mathscr{P}$  to  $L_s^2(\Sigma,E_{\Sigma}).$ 

PROOF. (1) First we show the inclusion im( $\mathscr{P}_{+,-1/2}$ ) ⊂  $\beta(A)$ . Let f belong to  $\text{im}(\mathscr{P}_{+,-1/2})$ . Then there exists  $f_1 \in L^2_{-1/2}(\Sigma, E_{\Sigma})$  with  $f = r_+ \tilde{G} \rho^* J_0 f_1$ , where  $\tilde{G}$ denotes Seeley's 'inverse on the double' (which in difference to our  $\tilde{G}$  is not canonically defined) and  $\rho^*$  Seeley's dual of the trace (which, once again, in difference to our  $\rho^*$  is neither canonically defined). We observe that  $g := \tilde{G}\rho^* J_0 f_1 \in \mathscr{D}(\tilde{A}_{\text{max}})$ with  $\tilde{A}q=0$ . Note that this is Seeley's  $\tilde{A}$  which is neither canonical and therefore not suitable for discussing the parameter dependence, but has the advantage of delivering a pseudodifferential Calderón projection. So  $f = \rho r_+ g$  with  $r_{+}g \in \mathscr{D}(A_{\max}).$  Hence  $f \in \beta(A)$ .

(2) Next we observe that  $L^2_{1/2}(\Sigma, E_{\Sigma}) \subset \beta(A)$  since  $L^2_1(M, E) \subset \mathscr{D}(A_{\max})$ .

(3) Together with argument (1) this implies

<span id="page-36-2"></span>
$$
\operatorname{im}(\mathscr{P}_{+,-1/2}) \oplus \operatorname{im}(\operatorname{Id} - \mathscr{P}_{+,1/2}) \subset \beta(A).
$$

(4) To show the equality, we notice  $\beta(A) \subset L^2_{-1/2}(\Sigma, E_{\Sigma})$ . Applying Seeley's result

$$
(6.15) \t\t\t\t\t\t\mathscr{P}_{+,s} + \mathscr{P}_{-,s} = \text{Id}
$$

for  $s = -1/2$  we can write each  $f \in \beta(A)$  in the form

$$
f = f_1 + f_2
$$
, where  $f_1 \in \text{im}(\mathscr{P}_{+,-1/2})$  and  $f_2 \in \text{im}(\mathscr{P}_{-,-1/2})$ .

By (1),  $f_1 \in \beta(A)$ , so  $f_2 = f - f_1 \in \beta(A)$ , i.e., there exists a  $g \in \mathscr{D}(A_{\max})$  such that  $f_2 = \rho g$ . Note  $\mathscr{P}_+ f_2 = 0$  by the splitting [\(6.15\)](#page-36-2). Applying one version of Gårding's inequality (see, e.g., [[BBW93](#page-56-1), Chapter 18])

$$
(6.16) \t ||g||_{L_1^2(M,E)} \leq C(||g||_{L^2(M,E)} + ||Ag||_{L^2(M,E)} + ||\mathscr{P}_+\rho g||_{L_{1/2}^2(\Sigma,E_{\Sigma})}),
$$

we obtain  $g \in L_1^2(M, E)$  and so  $f_2 = \rho g \in L_1^2_{1/2}(\Sigma, E_{\Sigma})$ , i.e.,  $f_2 \in \text{im}(\mathscr{P}_{-,1/2})$ .  $\square$ 

<span id="page-36-0"></span>**Lemma 6.11.** Let  $P, Q$  be two pseudodifferential projections with the same leading symbol. Then

(6.17) 
$$
\operatorname{im}(P_{-1/2}) \oplus \operatorname{im}(\operatorname{Id} - P_{1/2}) = \operatorname{im}(Q_{-1/2}) \oplus \operatorname{im}(\operatorname{Id} - Q_{1/2}).
$$

PROOF. So, let  $f \in \text{im}(P_{-1/2}) \oplus \text{im}(\text{Id} - P_{1/2})$ . Then there are  $\varphi \in L^2_{-1/2}(M, E)$ ,  $\psi \in L^2_{1/2}(M,E)$  such that

$$
f = P\varphi + (I - P)\psi = Q\varphi + (I - Q)\psi + \underbrace{(P - Q)(\varphi - \psi)}_{=:h}.
$$

By assumption  $P - Q$  is a pseudodifferential operator of order  $\leq -1$ , hence  $h \in$  $L_{1/2}^2(M, E)$  and thus  $f = Q(\varphi + h) + (I - Q)(\psi + h)$  with  $\varphi + h \in L_{-1/2}^2(M, E)$ and  $\psi + h \in L^2_{-1/2}(M, E)$ , proving the claim.

<span id="page-36-1"></span>Remark 6.12. By combining the two preceding lemmata we obtain the useful formula

<span id="page-36-3"></span>(6.18) 
$$
\beta(A) = \text{im}(P_{-1/2}) \oplus \text{im}((\text{Id} - P)_{1/2})
$$

for all pseudodifferential projections P with  $\sigma_0(P) = \sigma_0(\mathscr{P}_+)$  where  $\mathscr{P}_+$  denotes Seeley's (pseudodifferential) Calderón projection.

Equation [\(6.18\)](#page-36-3) generalizes a previous result in [[BBF99](#page-56-7), Proposition 7.15] obtained for the spectral Atiyah-Patodi-Singer projection  $P := P_{>}(B_0)$ . There, the von Neumann space  $\beta(A)$  was expressed as the direct sum of

- the  $L^2_{1/2}(\Sigma, E_{\Sigma})$ -closure of the linear span of the negative eigenspaces of the tangential operator  $B_0$  with
- the  $L^2_{-1/2}(\Sigma, E_{\Sigma})$ -closure of the linear span of the nonnegative eigenspaces of  $B_0$

in the special case that the operator  $A$  is of Dirac type in product metric near the boundary (in particular, that A has a formally selfadjoint tangential operator  $B_0$ and constant coefficients in normal direction near the boundary).

<span id="page-37-0"></span>**6.3. Proof of Theorem [6.1.](#page-30-2)** As announced above, we prove the five claims successively.

6.3.1. Application of the new construction of the Calderón projection.

PROOF OF (I). We recall that  $(u, v) \mapsto \omega(u, v) := \langle -J_0u, v \rangle$  is a symplectic form for the Hilbert space  $(L^2(\Sigma, E_{\Sigma}), \langle \cdot, \cdot \rangle)$ . It is strong since  $J_0$  is a bundle isomorphism. Then the range  $\text{im}(C_+) = N^0_+$  is an isotropic subspace because of Green's formula [\(A.13\)](#page-55-0), applied to the kernel of the formally selfadjoint operator A. Here we use  $T := J_0 \left(-J_0^2\right)^{-1/2}$  to construct  $C_{\pm}$ , see Remark [5.13.](#page-29-2) Then also  $\text{im}(C_{-}) = T^{-1}(N_{+}^{0})$  is an isotropic subspace since the chosen T is clearly symplectic. By Proposition [5.12,](#page-29-0) we have  $C_+ + C_- = \text{Id}$ , so  $N_+^0$  and  $T^{-1}(N_+^0)$  make a pair of transversal isotropic subspaces of  $L^2(\Sigma, E_{\Sigma})$ . Then (I) follows by applying Lemma  $6.4.$ 

**Remark 6.13.** We notice that the splitting  $\mathscr{P}_+ + \mathscr{P}_- =$  Id in Seeley, [[See66](#page-57-24), Lemma 5] does not provide two transversal Lagrangian subspaces but only an isotropic range of  $\mathscr{P}_+$  with the preceding argument in the case of symmetric A. The problem is that even for symmetric A Seeley's continuation into a collar and further over the double does not preserve symmetry in general.

6.3.2. Stability arguments. Before deriving  $(II)$  we shall address stability aspects of the issue.

We see at once that any formally selfadjoint operator of the form

$$
A := -j_t \frac{d}{dt} - \frac{1}{2} (\frac{d}{dt} j_t) - b_t, \quad j_t \text{ invertible}
$$

on the unit interval  $t \in [0,1]$  admits selfadjoint boundary conditions. The symplectic form on the space of boundary values is given by  $J := j_0 \oplus (-j_1)$  with respect to the reversed orientation at the ends of the interval. By continuity, we have sign  $j_0 = \text{sign } j_1$ . So, sign  $iJ = 0$ . (Note that there is no tangential operator in the 1-dimensional case).

In higher dimension, a similar continuity argument does not work in general.

The following lemma yields the stability of the signature of the almost complex form  $J_t$  on the nullspace ker  $B_t$  under variation of the parameter t. It can be considered as an index stability statement (and certainly can be proved also that way instead of the proof given below).

**Lemma 6.14.** Let  $(H, \langle \cdot, \cdot \rangle)$  be a Hilbert space and  $(S_t)$  and  $(J_t)$  be two continuous families of bounded invertible operators on H,  $t \in [0,1]$ . Assume that all  $S_t$  are positive definite. Let  $(B_t)$  be a continuous family of closed Fredholm operators. We assume that all  $iJ_t$  and  $B_t$  are selfadjoint with respect to the new metric  $\langle x, y \rangle_t :=$  $\langle S_t x, y \rangle$ . Moreover, we assume that all  $J_t$  have bounded inverses, and  $J_t B_t = -B_t J_t$ for all t. Then we have

(6.19) 
$$
\operatorname{sign}(iJ_t|\ker B_t) = \text{constant}.
$$

PROOF. We divide the proof into three steps.

Step 1. We can assume that  $S_t = Id$ . Indeed, denote by  $A^{*_t}$  the adjoint operator of A with respect to the scalar product induced by  $S_t$ . Then we have

$$
\langle S_t x, A^{*_t} y \rangle = \langle S_t A x, y \rangle.
$$

So  $A^{*_t} = S_t^{-1}A^*S_t$ . Set

$$
J'_t := S_t^{\frac{1}{2}} J_t S_t^{-\frac{1}{2}}, \quad B'_t := S_t^{\frac{1}{2}} B_t S_t^{-\frac{1}{2}}.
$$

Then  $iJ'_t$  and  $B'_t$  are selfadjoint,  $J'_t B'_t = -B'_t J'_t$ , and

$$
sign(iJ_t \restriction \ker B_t) = sign(iJ'_t \restriction \ker B'_t).
$$

So we can assume  $S_t = Id$ .

Step 2. We reduce to the finite–dimensional case: For each  $t \in [0,1]$ , there is a small  $\varepsilon > 0$  such that  $[-\varepsilon, \varepsilon] \cap \sigma(B_t) \subset \{0\}$ . Then for s close to  $t, \pm \varepsilon \notin \sigma(B_s)$ . Let  $D_{\varepsilon} := \{ z \in \mathbb{C} \mid |z| < \varepsilon \}.$  Define

$$
P_s := -\frac{1}{2\pi i} \int_{\partial D_\varepsilon} (B_s - z \operatorname{Id})^{-1} dz.
$$

Then  $P_s$  is a continuous family of orthogonal projections of finite rank, and  $P_sJ_sP_s$ ,  $P_s B_s P_s : \text{im } P_s \to \text{im } P_s$  satisfy our assumptions, and  $P_t B_t P_t = 0$ . Step 3. Since in the finite–dimensional case  $sign(iJ_t)$  is constant, it suffices to prove

the following

*Claim.* Let H be finite–dimensional. Then we have  $sign(iJ_t) = sign(iJ_t)$  ker  $B_t$ .

In fact, let  $V_t$  denote the orthogonal complement of ker  $B_t$ . Then both ker  $B_t$ and  $V_t$  are invariant subspaces of  $J_t$ . Since on  $V_t$ ,  $B_t$  is invertible, and since we have  $-J_t = B_t^{-1} J_t B_t$ , we have  $sign(iJ_t | V_t) = 0$ . Hence our claim follows.

6.3.3. Proof of  $(II)$ ,  $(III)$ . We now proceed to show the General Cobordism Theorem (II).

We exploit the formal selfadjointness of A and choose, in a collar of the boundary, the normal form

$$
A = J_x \left(\frac{d}{dx} + B_x\right) + \frac{1}{2}J'_x
$$

of Equation [\(2.21\)](#page-9-1) with the relations  $J^* = -J$ ,  $JB = -B^tJ$ . The relation  $JB =$  $-B<sup>t</sup>J$  has consequences for the positive/negative sectorial spectral subspaces with regard to the natural symplectic structure on  $L^2(\Sigma, E_{\Sigma})$ , see the proof of Lemma [6.15](#page-38-0) below.

Similarly to Figure [1](#page-13-1) we now choose contours  $\Gamma_{\leq}$ ,  $\Gamma_{\geq}$  and  $\Gamma_0$  as follows (see Figure [4\)](#page-39-0):  $\Gamma$  encircles all eigenvalues in the left half plane,  $\Gamma$  encircles all eigenvalues in the right half plane, and  $\Gamma_0$  encircles all eigenvalues on the imaginary axis iR. The corresponding spectral projections are denoted by  $P_<(B_0), P_>(B_0)$  and  $P_0(B_0)$ . In view of Theorem [3.14,](#page-17-3) these are pseudodifferential operators of order 0, and hence bounded, and as closed idempotents they do have closed range (see Paragraph [3.2.1\)](#page-14-7).  $P_0(B_0)$  is of finite–rank and hence a smoothing operator with range being the sum of the generalized eigenspaces of  $B<sub>0</sub>$  to imaginary eigenvalues. We therefore have a direct sum decomposition

<span id="page-38-1"></span>
$$
(6.20) \tL2(\Sigma, E_{\Sigma}) = \text{im } P_{<}(B_0) \oplus \text{im } P_0(B_0) \oplus \text{im } P_{>}(B_0) =: W_{<} \oplus W_0 \oplus W_{>}.
$$

In particular,  $(W_>, W_<)$  is a Fredholm pair of closed subspaces of  $L^2(\Sigma, E_{\Sigma})$ . Recall that the closedness of  $W<sub>lt</sub>, W<sub>gt</sub>$  and the finite codimension of  $W<sub>lt</sub> + W<sub>gt</sub>$  in  $L<sup>2</sup>(\Sigma, E<sub>Σ</sub>)$ imply that  $W< +W>$  is closed, see [[BBF99](#page-56-7), Remark A.1].

<span id="page-38-0"></span>**Lemma 6.15.**  $W_>, W_>,$  are isotropic subspaces with  $W_>^{\omega} = W_0 \oplus W_>$  and  $W_<^{\omega} =$  $W_{< \oplus W_0$ .



<span id="page-39-0"></span>FIGURE 4. Three contours encircling all eigenvalues in the right half plane, on the imaginary axis, and all eigenvalues in the left half plane, respectively.

PROOF. This is a consequence of the relation  $J_0B_0 = -B_0^tJ_0$  which implies for  $\xi \in L_1^2(\Sigma, E_{\Sigma})$  (cf. Subsection [3.2\)](#page-13-0)

(6.21)  
\n
$$
J_0 P_{>}(B_0) \xi = \frac{1}{2\pi i} \int_{\Gamma_{>}} \lambda^{-1} J_0 (\lambda - B_0)^{-1} d\lambda B_0 \xi
$$
\n
$$
= \frac{1}{2\pi i} \int_{\Gamma_{>}} \lambda^{-1} (\lambda + B_0^t)^{-1} d\lambda B_0^t J_0 \xi
$$
\n
$$
= P_{<}(B_0^t) J_0 \xi.
$$

Then

$$
W_{>}^{\omega} = (J_0 W_{>} )^{\perp} = (J_0 \text{ im } P_{>} (B_0))^{\perp} = \text{ im } P_{<} (B_0^t)^{\perp}
$$
  
= ker  $P_{<} (B_0^t)^* = \text{ker } P_{<} (B_0) = W_0 \oplus W_{>}.$ 

The other claim is proved analogously.  $\Box$ 

Lemma [6.15](#page-38-0) is now the key to the proofs of the remaining implications in this paragraph. Before proceeding, we note:

**Lemma 6.16.** Let  $P$  and  $Q$  be bounded idempotents in a Banach space  $H$ , and  $P - Q$  compact. Then the pair (ker P, im Q) is Fredholm.

PROOF. Consider the operator

$$
R := QP + (\text{Id} - P)(\text{Id} - Q) = Q(Q + P - Q) + (\text{Id} - Q + Q - P)(\text{Id} - Q)
$$
  
= Q + Q(P - Q) + \text{Id} - Q + (Q - P)(\text{Id} - Q)  
= \text{Id} + Q(P - Q) + (Q - P)(\text{Id} - Q).

Since  $P - Q$  is compact, R is Fredholm. Since  $(\ker P \cap \dim Q) \subset \ker R$  and  $(\ker P +$  $\dim Q$ )  $\supset$  im R, we have dim(ker  $P \cap \mathrm{im} Q$ )  $< +\infty$  and  $\dim(H/(\ker P + \mathrm{im} Q))$   $< +\infty$ . Thus the pair (ker  $P$ , im  $Q$ ) is algebraically Fredholm, and hence Fredholm, since the spaces ker  $P$ , im  $Q$  are closed.

PROOF OF  $(I) \implies (II)$ . Let  $\mathscr{P}_+$  denote Seeley's corresponding (pseudodifferential) Calderón projection. Then we have im  $C_+ = \text{im } \mathscr{P}_+$ . Since the difference

between  $\mathscr{P}_+$  and  $P_>(B_0)$  is of order -1, we have that  $(\text{im }\mathscr{P}_+, W_<)$  is a Fredholm pair by the preceding argument (and finite–dimensional perturbation). Hence im  $\mathscr{P}_+ \oplus W_{\leq 0}$  is a closed subspace.

By (I), im $\mathcal{P}_+$  is a Lagrangian subspace. Applying Lemma [6.8](#page-34-2) to the coisotropic subspace  $W^{\omega}_{\leq} = W_0 \oplus W_{\leq}$  we obtain that

$$
\text{Red}_{W_{\leq}^{\omega}}(\text{im }\mathscr{P}_{+})\simeq \pi(\text{Red}_{W_{\leq}^{\omega}}(\text{im }\mathscr{P}_{+}))\subset W_{0}
$$

is a Lagrangian subspace of

$$
W^{\omega}_{\leq}/W^{\omega\omega}_{\leq} = (W_{\leq} + W_0)/W_{\leq} \simeq W_0.
$$

So, the finite–dimensional symplectic Hilbert space  $(W_0,\langle iP_0J_0 \cdot,\cdot \rangle)$  has a Lagrangian subspace. Therefore

$$
\operatorname{sign} i P_0 J_0 \upharpoonright W_0 = 0. \qquad \qquad \Box
$$

We assume  $B_0 = B_0^t$  and put  $\alpha = -iJ_0(-J_0^2)^{-1/2}$ . Then the special case of claim (II) of Theorem [6.1](#page-30-2) follows. Moreover, we have under that assumption:

PROOF OF  $(II) \Leftrightarrow (III)$ . We follow the conventional lines and refer, e.g., to **[[BBW93](#page-56-1)**, Theorem 21.5] with the immediate modifications: Note that  $\alpha$  is a grading which thanks to  $B_0 = B_0^t$  and  $B_0 J_0 = -B_0^t J_0$  anticommutes with  $B_0$ , making  $B_0$  an odd operator. More precisely, let  $(E_{\Sigma})^{\pm}$  denote the positive (negative) eigenspace of  $\alpha$ . Under the direct sum decomposition  $E_{\Sigma} = (E_{\Sigma})^+ \oplus (E_{\Sigma})^-$ , the operator  $B_0$  takes the form  $B_0 = \begin{pmatrix} 0 & B^- \\ B^+ & 0 \end{pmatrix}$  $B^+$  0 ), where  $B^- := (B^+)^t$  and  $B^+$ : ker( $\alpha - 1$ ) → ker( $\alpha + 1$ ). Then we have ker  $B_0 = \ker B^+ \oplus \ker B^-$ , and the positive (negative) eigenspace of  $iJ_0|_{\text{ker }B_0}$  is ker  $B^{\pm}$ . That proves

$$
\operatorname{sign}(iJ_0|_{\ker B_0}) = \operatorname{ind} B^+.
$$

6.3.4. Proof of  $(II) \implies (IV), (V)$ . To show that  $(IV)$  and  $(V)$  are trivial consequences of (I), one is tempted to set  $P := C_+$ . However, in Proposition [5.12](#page-29-0) we have established that  $C_+$  is the  $L^2$  extension of a pseudodifferential projection only for the boundary operator  $T := (J_0^t)^{-1}$ , contrary to our assumption  $T := J_0(-J_0^2)^{-1/2}$  in (I) for achieving that ker  $C_+ = \text{im } C_-$  becomes a Lagrangian subspace of  $L^2(\Sigma, E_{\Sigma})$ . To prove that  $C_+$  is also pseudodifferential for the last choice of T would require applying more advanced elliptic boundary value theory. Instead of that, we give a simple construction of the wanted  $P$  as a perturbation of the positive spectral projection  $P_>(B_0)$  by a suitable finite–rank operator, and let simple symplectic analysis do the remainder of the work:

PROOF OF  $(II) \implies (IV), (V)$ . The vanishing of the signature sign  $iP_0J_0$  $W_0$  on the finite–dimensional space  $W_0$  implies the existence of a transversal pair of Lagrangian subspaces  $(\lambda, \mu) \subset W_0$ . The pair  $(W_+ + \lambda, W_- + \mu)$  is a transversal pair of Lagrangian subspaces of  $L^2(\Sigma, E_{\Sigma})$ . Denote by P the projection of  $L^2(\Sigma, E_{\Sigma})$ onto  $W$ <sub>></sub> +  $\lambda$ . Then P is a zeroth order pseudodifferential operator, and P –  $\mathscr{P}_+$ is of −1 order. Then (ker P, im  $\mathcal{P}_+$ ) is a Fredhom pair, and ker  $P = W_{<} + \mu$  is a Lagrangian subspace of  $L^2(\Sigma, E_{\Sigma})$ . Since im  $C_+ = \text{im }\mathscr{P}_+$ , (IV) follows.

Now we consider (V). By Remark [6.12](#page-36-1) we have  $\beta(A) = \text{im } P_{-1/2} \oplus \text{im}(\text{Id} - P_{1/2})$ . Clearly im  $P_{-1/2}$  and im(Id  $-P_{1/2}$ ) are isotropic subspaces of  $\beta(A)$ . By Lemma [6.4,](#page-33-4) they are Lagrangian subspaces. Then the extension  $A_P$  is a selfadjoint extension. Fredholmness follows from leading symbol consideration.

<span id="page-41-0"></span>6.4. Alternative routes to the General Cobordism Theorem. We de-scribe alternative routes to prove Theorem [6.1.](#page-30-2) In the present context  $(I)$  is an immediate consequence of our Calderón construction. Therefore, we began the proof of the General Cobordism Theorem with a proof of (I).

One true alternative is to begin with a proof of  $(V)$ : That claim was proved in [[Ral70](#page-57-10), Theorem I] for bounded regions M in Euclidean space. However, Ralston's arguments fully generalize and can be summarized in the following way (in our present notation):

OUTLINES OF A PROOF OF  $(V)$ . (1) First we notice the pointwise vanishing of the signature of the form  $iJ_0\upharpoonright p$  on the fibre  $E_p$  for each  $p \in \Sigma$ . This can be obtained by deforming  $J_0 \upharpoonright p$  into a strict anti-involution  $\widetilde{J_0} \upharpoonright p$  with  $(\widetilde{J_0} \upharpoonright p)^2$ − Id and exploiting the anti-commutative relation [\(2.20\)](#page-8-6) with the elliptic symbol. Consequently, the fibre dimension of the bundle E must be even. That permits the first trick, namely to split  $E = E_+ \oplus E_-$  and to show that there exists a well-posed symmetric Fredholm extension given by the graph of a pseudodifferential elliptic operator  $P : \Gamma^{\infty}(\Sigma; E_+ \restriction \Sigma) \to \Gamma^{\infty}(\Sigma; E_- \restriction \Sigma)$ .

(2) Next we show that the deficiency indices  $\kappa_1$ ,  $\kappa_2$  of A are finite and that their difference is equal to the index ind  $P$ .

(3) Then we show that  $-\operatorname{ind} P = \operatorname{sign} -i\omega \restriction \operatorname{vN}(P)$  where  $\omega$  denotes the Green form and  $vN(P)$  denotes a suitably defined subspace of the von Neumann quotient  $\mathscr{D}(A_{\text{max}})/\mathscr{D}(A_{\text{min}}) =: \beta(A).$ 

(4) Finally we show that ind P vanishes, and hence that  $A_P$  can be extended to a selfadjoint  $A_{\tilde{P}}$  with domain given by a pseudodifferential projection and preserving the Fredholm property. the Fredholm property.

Note that we can deduce  $(I)$  from  $(V)$  in the following way, and independently of the delicacies of our Calderón construction. We firstly show that we have  $(V) \implies$ (II).

PROOF OF  $(V) \implies (II)$ . Since  $A_P$  is a selfadjoint Fredholm operator, by [[BBF98](#page-56-3), Proposition 3.5], im  $\mathscr{P}_{+,-1/2}$  is a Lagrangian subspace of  $\beta(A)$ .

By Remark [6.12](#page-36-1) and Equation [\(6.20\)](#page-38-1), we have

$$
(6.22) \qquad \beta(A) = \text{im } P_{< 1/2}(B_0) \oplus \text{im } P_0(B_0) \oplus \text{im } P_{> -1/2}(B_0).
$$

Using the same method as in the proof of  $(I) \implies (II)$  and applying Lemma [6.8](#page-34-2) to the co-isotropic subspace im  $(P_{\leq,1/2}(B_0))^{\omega'} = \text{im } P_0(B_0) \oplus \text{im } P_{\leq,1/2}(B_0)$ , we obtain that

$$
\mathrm{Red}_{\left(P_{<,1/2}(B_0)\right)^\omega}(\mathrm{im}\, \mathscr{P}_{+,-1/2})\simeq \pi(\mathrm{Red}_{\left(P_{<,1/2}(B_0)\right)^\omega}(\mathrm{im}\, \mathscr{P}_{+,-1/2}))\subset W_0
$$

is a Lagrangian subspace of  $W_0$ . So, the finite–dimensional symplectic Hilbert space  $(W_0, \langle iP_0 J_0 \cdot, \cdot \rangle)$  has a Lagrangian subspace. Therefore

$$
\operatorname{sign} i P_0 J_0 \upharpoonright W_0 = 0. \qquad \qquad \Box
$$

PROOF OF  $(V) \implies (I)$ . By the above proofs we have  $L^2(\Sigma, E_{\Sigma}) = \text{im } P \oplus$  $\text{im}(\text{Id}-P)$  and  $\beta(A) = \text{im }P_{-1/2} \oplus \text{im}(\text{Id}-P_{1/2})$ . By Theorem [6.9,](#page-35-3)  $\text{im } \mathscr{P}_{+,-1/2} \cap$  $L^2(\Sigma, E_{\Sigma}) = \text{im } \mathscr{P}_+ = \text{im } C_+$  is a Lagrangian subspace of  $L^2(\Sigma, E_{\Sigma})$ .

### 7. Parameter dependence

<span id="page-41-1"></span>In this section we discuss the continuous dependence of the Calderón projection and the Poisson operator on the input data. That is, given a first order elliptic differential operator  $A \in \text{Diff}^1(M; E, F)$  and  $T \in \text{Diff}^0(\Sigma; E_\Sigma, F_\Sigma), \Sigma = \partial M$  (cf. Definition [4.1\)](#page-20-3), we want to have criteria to ensure that  $(A, T) \rightarrow K_+ = K_+(A, T)$ respectively  $(A, T) \mapsto C_+ = C_+(A, T)$  is continuous in an appropriate sense.

Therefore we first introduce various metrics on the spaces of pairs  $(A, T)$ . Re-ferring to [\(5.11\)](#page-24-2) and [\(5.15\)](#page-24-3) we consider  $J_0, B_0, C_0, C_1$  as functions of A.

<span id="page-42-5"></span>**Definition 7.1.** (a) Let  $\mathscr{V}(M;E,F)$  denote the subspace of  $\text{Diff}^1(M;E,F)$   $\times$  $\text{Diff}^{0}(\Sigma;E_{\Sigma},F_{\Sigma})$  consisting of those  $(A,T)$  for which  $[B_{0}^{t},J_{0}^{t}T]$  is of order 0 (cf. [\(5.9\)](#page-24-4)). By  $\mathscr{E}(M;E,F)$  we denote the subspace of  $\mathscr{V}(M;E,F)$  consisting of those pairs  $(A, T)$  where

 $(1)$  *A* is elliptic

 $(2)$  T is invertible and satisfies  $(5.8)$ .

Finally, we denote by  $\mathscr{E}_{\text{LCP}}(M; E, F)$  the subspace of  $\mathscr{E}(M; E, F)$  consisting of pairs  $(A, T)$  where A and  $A<sup>t</sup>$  satisfy weak inner UCP.

(b) On the linear space  $\mathcal{V}(M;E,F)$  we introduce two norms:

<span id="page-42-2"></span>
$$
(7.1) \t N_0(A,T) := \|A\|_{1,0} + \|A^t\|_{1,0} + \|T\|_{1/2,1/2},
$$

and

<span id="page-42-1"></span>(7.2) 
$$
N_1(A, T) := ||B_0||_{1,0} + ||B_0^t||_{1,0} + ||[B_0^t, J_0^t T]||_0 + ||T||_0 + ||J_0||_0 + ||C_1||_{1,0} + ||C_0||_0 + ||\widetilde{C}_1||_{1,0} + ||\widetilde{C}_0||_0.
$$

Except for  $C_1$ ,  $\tilde{C}_1$  the norms in [\(7.2\)](#page-42-1) are the mapping norms between Sobolev spaces over Σ while the norms for  $C_1$ ,  $\tilde{C}_1$  are mapping norms between Sobolev spaces over the collar  $[0, \varepsilon) \times \Sigma$ .

(c) We equip the space  $\mathscr{E}(M;E,F)$  with the metric  $d_0$  induced by the metric  $N_0$ of [\(7.1\)](#page-42-2). I.e.

(7.3) 
$$
d_0((A,T),(A',T')) := N_0(A-A',T-T').
$$

The norms  $N_0, N_1$  induce metrics on subspaces of  $\mathcal{V}(M; E, F)$ , in particular on  $\mathscr{E}(M;E,F)$ .

To study the dependence of  $K_+$  and  $C_+$  on  $(A, T)$  the formulas [\(5.30\)](#page-28-2) and [\(5.31\)](#page-28-0) in Proposition [5.10](#page-28-3) are crucial.

To illustrate this let us consider a map  $Z \ni z \mapsto (A(z), T(z)) \in \mathscr{E}(M; E, F)$ from a metric space Z to  $\mathscr{E}(M; E, F)$ . To conclude that the corresponding map  $z \mapsto K_+(A(z), T(z)) \in \mathscr{B}(L^2(\Sigma, E_{\Sigma}), L^2_s(M, E))$  is continuous for some fixed  $0 \leq$  $s \leq 1/2$  it suffices to show the continuity of

<span id="page-42-3"></span>
$$
(1) \ \ z \mapsto \varphi R_{T(z)}(A(z)) \in \mathscr{B}(L^2(\Sigma,E_{\Sigma}),L^2_s(M,E\oplus F)),
$$

<span id="page-42-4"></span> $(2)$   $z \mapsto \varphi N_{T(z)}(A(z)) \in \mathscr{B}(L^2(\Sigma, E_{\Sigma}), L_s(M, E \oplus F)),$ <br>  $(2)$   $z \mapsto S(A(z), T(z)) \in \mathscr{B}(L^2(\Sigma, E_{\Sigma}), L^2(M, F \oplus E)),$ 

(3)  $z \mapsto \widetilde{G}(A(z), T(z)) \in \mathscr{B}(L^2(M, F \oplus E), L^2_1(M, E \oplus F)).$ 

For the continuity of  $z \mapsto C_+(A(z), T(z)) \in \mathscr{B}(L^2(\Sigma, E_{\Sigma}))$  [\(1\)](#page-42-3) has to be replaced by the continuity of the map

$$
(1') z \mapsto P_+(B_0(A(z))) \in \mathscr{B}(L^2(\Sigma, E_{\Sigma})).
$$

The continuity of these maps is by no means necessary for ensuring the continuous dependence of  $K_{+}, C_{+}$ . In order to keep the presentation reasonable in size we estimate generously - we are not striving for optimality here.

Let us now state the main result of this section.

We define the *strong metric* on the space  $\mathscr{E}(M;E,F)$  by

$$
(7.4) \t dstr((A, T), (A', T')) := N0(A - A', T - T') + N1(A - A', T - T').
$$

Note that by complex interpolation  $||T - T'||_{s,s} \leq d_{str}((A,T), (A',T'))$  for all  $0 \le s \le 1/2$ .

<span id="page-42-0"></span>Theorem 7.2. (a) The map

$$
(\mathscr{E}_{\text{UCP}}(M;E,F),d_{str}) \longrightarrow \mathscr{B}(L^2(\Sigma,E_{\Sigma}),L^2_s(M,E))
$$

sending  $(A, T)$  to the Poisson operator  $K_+(A, T)$  is continuous for  $0 \le s < 1/2$ . (b) Let  $(A(z), T(z))_{z\in Z}$  be a continuous family in  $(\mathscr{E}_{\text{UCP}}(M; E, F), d_{str})$  parametrized by a metric space Z. Assume that the corresponding family

$$
z \mapsto P_+(B_0(A(z))) \in \mathscr{B}(L_s^2(\Sigma, E_{\Sigma}))
$$

of positive spectral projections of the tangential operator is continuous for some fixed  $s \in [-1/2, 1/2]$ . Then the map

$$
Z \longrightarrow \mathscr{B}(L^2_s(\Sigma, E_{\Sigma}))
$$

sending z to the Calderón projection  $C_{+}(A(z), T(z))$  is continuous.

**Remark 7.3.** (1) Of course, analogous statements hold for  $K_-, C_-$ . We leave it as an intriguing problem whether the statement about  $K_{+}$  still holds for  $s = 1/2$ . This would be more natural since  $K_{+}$  maps  $L^{2}$  to  $L_{1/2}^{2}$ .

(2) The fact that the continuous dependence of  $P_+$  in (b) has to be assumed is not very satisfactory. The point here is *not* that the construction of  $P_+$  requires a spectral cut. Suppose a spectral cut for  $B_0 = B_0(A(z_0))$  is chosen. Then  $P_+$  should vary continuously as long as no eigenvalues approach the contours  $\Gamma_{\pm}$  (cf. Figure [2\)](#page-15-0). Unfortunately we cannot prove the continuity of  $B \mapsto P_+(B) \in \mathscr{B}(L^2(\Sigma, E_{\Sigma}))$ if we equip  $\text{Diff}^1(\Sigma; E_{\Sigma})$  say with the norm  $\|\cdot\|_{1,0}$ ; we cannot prove it for any other norm either. We will come back to this problem below in Subsection [7.5,](#page-48-0) where we will give a criterion for the continuity of  $P_+$  in special cases.

PROOF. The Theorem follows from Proposition [5.10,](#page-28-3) Proposition [7.8,](#page-46-3) Propo-sition [7.12,](#page-47-1) and Theorem [7.9.](#page-46-2)

The discussion in Subsection [7.5](#page-48-0) will give at least the following result:

<span id="page-43-0"></span>**Corollary 7.4.** Denote by  $\mathcal{E}_{\text{UCP}}^{sa}(M;E,F)$  the subspace of  $\mathcal{E}_{\text{UCP}}(M;E,F)$  consisting of pairs  $(A, T)$  where the corresponding tangential operator  $B_0(A)$  has selfadjoint leading symbol. Then for  $s \in [-1/2, 1/2]$  the map

$$
(\mathscr{E}_{\mathrm{UCP}}^{\mathrm{sa}}(M;E,F),d_{str})\longrightarrow \mathscr{B}(L^2_s(\Sigma,E_{\Sigma}))
$$

sending  $(A, T)$  to the Calderón projection  $C_+(A, T)$  is continuous.

PROOF. We may adopt the language of a family  $(A(z), T(z))_{z\in Z}$  (with  $Z =$  $\mathscr{E}_{\text{UCP}}^{\text{sa}}(M;E,F)!$  of the previous Theorem [7.2.](#page-42-0)

The point is that locally (cf. Convention [3.7\)](#page-0-0) one can choose a continuous family  $z \mapsto P_+(B_0(z))$ . Then the result follows from Theorem [7.2b](#page-42-0).

To see this we recall from Remark [5.2](#page-25-3) that if the tangential operator has selfadjoint leading symbol the  $B_0 = B_0(A(z))$  in [\(5.11\)](#page-24-2) can be chosen to be selfadjoint. Hence let  $B_0(z)$  now denote this selfadjoint operator in the representation [\(5.11\)](#page-24-2). To show continuity at  $z_0$  pick a spectral cut c for  $B_0(z_0)$ . Then by Proposition [7.15](#page-49-0) below the family  $P_+(B_0(z)) := 1_{[c,\infty)}(B_0(z)) \in \mathscr{B}(L^2_s(\Sigma, E_{\Sigma}))$  depends continuously on z in a neighborhood of  $z_0$ . Hence Theorem [7.2b](#page-42-0) yields the claim.  $\square$ 

Remark 7.5. [[BBF98](#page-56-3), Section 3.3] gives a purely functional analytic proof of the continuous variation of the Cauchy data spaces as subspaces of the von-Neumann quotient  $\beta(A)$  of all natural boundary values, as defined above in Paragraph [6.2.4,](#page-34-0) in great generality: only the symmetry of  $A$ , weak inner UCP and the existence of a selfadjoint Fredholm extension are assumed. In particular, no product form near the boundary or symmetry of a tangential operator is assumed. However, loc. cit. is restricted to continuous variation by bounded perturbations, i.e., perturbations of lower order in the operator norm, whereas the preceding corollary admits arbitrary continuous variations, though in the strong metric.

Below in Proposition [7.13](#page-48-1) we shall explain why lower order perturbations of a fixed operator lead to continuous variation of the sectorial projection in our setting, and hence to continuous variation of the Calderón projection according to the preceding Theorem.

We now proceed to give criteria for the continuity of the maps  $(1)-(2)$  $(1)-(2)$ ,  $(1')$ . We start with some basic estimates.

<span id="page-44-0"></span>7.1. Some estimates. We fix a first order elliptic differential operator  $B \in$ Diff<sup>1</sup>( $\Sigma$ ;  $E_{\Sigma}$ ) such that  $B-\lambda$  is parameter dependent elliptic in a conic neighborhood of iR (cf. Subsection [3.1\)](#page-11-1). Choose contours  $\Gamma_{+}$  accordingly as in Figure [2.](#page-15-0)

Before we address the continuous dependence of the sectorial projections on the data, i.e., on B, we shall give some useful estimates.

We will frequently use that for  $V \in CL^1(\Sigma; E_{\Sigma})$  we have by duality  $||V||_{0,-1} =$  $||V^t||_{1,0}.$ 

Our first result is the following perturbation lemma.

<span id="page-44-1"></span>**Lemma 7.6.** Let  $V \in \text{Diff}^1(\Sigma; E_{\Sigma})$ . If  $||V||_{1,0} + ||V^t||_{1,0}$  is sufficiently small, then  $B + V - \lambda$  is parameter dependent elliptic in a conic neighborhood of iR containing  $\Gamma_+$ ,  $\Gamma_-$ .

Furthermore, for  $|s|, |s'|, |s - s'| \leq 1$  we have for  $\lambda \in \Gamma_- \cup \Gamma_+$ 

$$
\begin{aligned} ||(\lambda - (B + V))^{-1} - (\lambda - B)^{-1}||_{s,s'} \\ &\le C(s, s', B) \left( ||V||_{1,0} + ||V^t||_{1,0} \right) |\lambda|^{-1+s'-s} .\end{aligned}
$$

PROOF. The first claim is clear. The second follows from a straightforward application of the Neumann series for the resolvent of  $B + V$  and complex interpolation. For the convenience of the reader we present some details of the estimate.

For  $s \in \{0, 1\}$  Equation [\(3.20\)](#page-18-1) yields for  $\lambda \in \Gamma_+ \cup \Gamma_-$ 

(7.5) 
$$
\|(B - \lambda)^{-1}V\|_{s,s} \leq \|(B - \lambda)^{-1}\|_{s-1,s} (\|V\|_{1,0} + \|V^t\|_{1,0})
$$

$$
\leq C(s) (\|V\|_{1,0} + \|V^t\|_{1,0}),
$$

and similarly

(7.6) 
$$
||V(B - \lambda)^{-1}||_{s,s} \leq C(s) (||V||_{1,0} + ||V^t||_{1,0}).
$$

Furthermore, complex interpolation (or Hadamard's three line theorem) gives that there is a constant C such that

(7.7) 
$$
\sup_{0 \le s \le 1, \lambda \in \Gamma_+ \cup \Gamma_-} \|V(B - \lambda)^{-1}\|_{s,s} + \sup_{0 \le s \le 1, \lambda \in \Gamma_+ \cup \Gamma_-} \|(B - \lambda)^{-1}V\|_{s,s} \le C (\|V\|_{1,0} + \|V^t\|_{1,0}).
$$

Choose V such that  $C(||V||_{1,0} + ||V^t||_{1,0}) < 1/2$ . Consequently,  $B + V - \lambda$  is invertible for all  $\lambda \in \Gamma_+ \cup \Gamma_-$  and as operator  $L^2_s \to L^2_{s-1}$ ,  $0 \le s \le 1$ , its inverse is given by the Neumann series

$$
(B + V - \lambda)^{-1} = (\text{Id} + (B - \lambda)^{-1}V)^{-1}(B - \lambda)^{-1}
$$
  
= 
$$
\sum_{n \ge 0} (-1)^n ((B - \lambda)^{-1}V)^n (B - \lambda)^{-1}.
$$

Hence

$$
(7.8) \quad ||(B+V-\lambda)^{-1} - (B-\lambda)^{-1}||_{s,s'} \leq \sum_{n\geq 1} ||((B-\lambda)^{-1}V)^{n} (B-\lambda)^{-1}||_{s,s'}.
$$

Now one has to check case by case.

1. Let 
$$
s' \ge 0
$$
,  $s' - s \ge 0$ . Then  
\n
$$
\left\| \left( (B - \lambda)^{-1} V \right)^n (B - \lambda)^{-1} \right\|_{s,s'}
$$
\n
$$
\le \left\| (B - \lambda)^{-1} \right\|_{s,s'} \left\| (B - \lambda)^{-1} V \right\|_{s',s'}^{n}
$$
\n(7.9)  
\n(3.20)  
\n(3.20)  
\n(4.320)  
\n(5.21)  
\n(6.41)  
\n(7.9)  
\n(8.42)  
\n(9.41)  
\n(10.41)  
\n(11.0)  $|V^t|_{1,0}$ )\n(11.0 +  $||V^t||_{1,0}$ ).

Summing up gives the claim in this case. 2. Let  $-1 \leq s \leq s' \leq 0$ . Then

$$
\begin{aligned} \left\| \left( (B - \lambda)^{-1} V \right)^n (B - \lambda)^{-1} \right\|_{s,s'} \\ &\leq \| (B - \lambda)^{-1} V \|_{0,s'} \left\| \left( (B - \lambda)^{-1} V \right)^{n-1} \right\|_{0,0} \| (B - \lambda)^{-1} \|_{s,0} \\ &\leq C \, |\lambda|^{-1+s'+1} \left( \| V \|_{1,0} + \| V^t \|_{1,0} \right) (1/2)^{n-1} \, |\lambda|^{-1-s} . \end{aligned}
$$

Again, summing up gives the claim also in this case.

For estimating  $\left\| \left( (B - \lambda)^{-1} V \right)^n (B - \lambda)^{-1} \right\|_{s,s'}$ , the roles of s and s' are symmetric. Hence the cases  $s' \leq s$  follow analogously.

We shall investigate the stability of the sectorial projections under perturbation of the input data B by V and show that the operator norm of  $\varphi(Q_{\pm}(B+V) - Q_{\pm}(B))$ from  $L_s^2(\Sigma, E_\Sigma)$  to  $L_{s', comp}^2(\mathbb{R}_+ \times \Sigma, E_\Sigma)$  is bounded by a constant depending on  $s, s', B, \varphi \text{ times } (\|V\|_{1,0} + \|V^t\|_{1,0}).$ 

<span id="page-45-2"></span>**Proposition 7.7.** Let  $\varphi \in C_0^{\infty}(\mathbb{R}_+)$  and V as in Lemma [7.6](#page-44-1).  $Q_{\pm}(B)$  and  $Q_{\pm}(B +$ V) are the operator families of B respectively  $B + V$  introduced in Definition [3.8](#page-15-3).

(a) For  $-1/2 < s \leq s' < s + 1/2, s' \leq 1$  we have for  $\xi \in L_s^2(\Sigma, E_{\Sigma})$ 

$$
(7.10) \quad \left\|\varphi\big(Q_{\pm}(B+V)-Q_{\pm}(B)\big)\xi\right\|_{s'}\leq C(s,s',B,\varphi)\left(\|V\|_{1,0}+\|V^t\|_{1,0}\right)\|\xi\|_s.
$$

(b) For  $-1 \leq s \leq 0$  we have for  $\xi \in L^2_s(\Sigma, E_{\Sigma})$ 

$$
(7.11) \left\| \mathrm{id}_{\mathbb{R}_+} \varphi \big( Q_\pm(B+V) - Q_\pm(B) \big) \xi \right\|_{s+1} \le C(s, B, \varphi) \left( \|V\|_{1,0} + \|V^t\|_{1,0} \right) \|\xi\|_s.
$$

PROOF. We use Lemma [7.6](#page-44-1) and estimate for  $-1 \leq s \leq s' \leq s+1, s' \leq 1$ ,  $\xi \in L^2_s(\Sigma,E_{\Sigma})$  and  $m \in \{0,1\}$ 

<span id="page-45-0"></span>
$$
\|x^{m}\varphi(x)\big(Q_{\pm}(B+V)-Q_{\pm}(B)\big)(x)\,\xi\|_{s'}
$$
\n
$$
\leq \frac{x^{m}}{2\pi}\varphi(x)\int_{\Gamma_{\pm}}|e^{-x\lambda}|\big|\big((\lambda-(B+V))^{-1}-(\lambda-B)^{-1}\big)\xi\big\|_{s'}\,|d\lambda|
$$
\n
$$
\leq C(s,s',B)\frac{x^{m}\varphi(x)}{2\pi}\int_{\Gamma_{\pm}}|e^{-x\lambda}|\,|\lambda|^{-1+s'-s}\,|d\lambda|\,(\|V\|_{1,0}+\|V^t\|_{1,0})\,\|\xi\|_{s}
$$
\n
$$
\leq C(s,s',B)(\|V\|_{1,0}+\|V^t\|_{1,0})\,\|\xi\|_{s}\,x^{m-s'+s}|\log x|\varphi(x).
$$

Analogously and using the previous estimate [\(7.12\)](#page-45-0) we find for  $0 \le s \le s' = 1$ 

<span id="page-45-1"></span>
$$
(7.13) \quad \left\| \partial_x \left( x^m \varphi(x) \left( Q_\pm(B+V) - Q_\pm(B) \right) \xi(x) \right) \right\|_{L^2(\Sigma, E_\Sigma)} \leq C(s, B, \varphi) (\|V\|_{1,0} + \|V^t\|_{1,0}) \, \|\xi\|_s \times \left( x^s \partial_x (x^m \varphi(x)) + \varphi(x) x^{m-1+s} \right) |\log x|.
$$

The  $\log x$ –terms on the right of [\(7.12\)](#page-45-0), [\(7.13\)](#page-45-1) are necessary only in the case  $s = s'$ . They are obsolete if  $s < s'$ .

From these two estimates both claims will follow in a straightforward manner:

(a). It suffices to prove the claim for  $-1/2 < s < s' = 0$  and for  $1/2 < s \leq s' = 1$ . It is clear that it then holds for  $-1/2 < s \leq s' \leq 0$  and  $1/2 < s \leq s' \leq 1$ . The general case then follows from the complex interpolation method.

So let us start with the case  $-1/2 < s \leq s' = 0$ . Since  $s > -1/2$  we may integrate the square of [\(7.12\)](#page-45-0) and reach the conclusion.

If  $1/2 < s \leq s' = 1$  then apply [\(7.12\)](#page-45-0) and [\(7.13\)](#page-45-1) with  $m = 0$ . Squaring and integrating the inequality gives the claim in view of [\(3.21\)](#page-18-2).

(b). By interpolation theory it is enough to deal with the cases  $s = -1$  and  $s = 0$ . If  $s = -1$  apply [\(7.12\)](#page-45-0) with  $s' = s+1 = 0$  and if  $s = 0$  apply (7.12) with  $s' = s+1 = 1$ and [\(7.13\)](#page-45-1). Again referring to [\(3.21\)](#page-18-2) we are done.

<span id="page-46-0"></span>**7.2. Continuous dependence of**  $\varphi R_T(B_0)$ **.** Since  $R_T$  is the multiplication of  $R$  by a simple matrix containing Id and  $T$  it suffices to study the dependence of  $R = R(B_0)$  on  $B_0$ :

<span id="page-46-3"></span>**Proposition 7.8.** Let  $(A, T) \in \mathcal{E}(M; E, F)$ . Fix a real number  $c > 0$  such that spec  $B_0 \cap \{z \in \mathbb{C} \mid |z| = c\} = \emptyset$ , cf. Convention [3.7](#page-0-0). Let  $V \in \text{Diff}^1(\Sigma; E_{\Sigma})$  be a first order differential operator. According to Lemma [7.6](#page-44-1) assume that  $(||V||_{1,0} + ||V^t||_{1,0})$ is small enough so that  $B_0 + V - \lambda$  is parameter dependent elliptic in a conic neighborhood of i $\mathbb R$  containing  $\Gamma_+ \cup \Gamma_-$ .

Then for  $-1/2 < s \leq s' < s + 1/2, s' \leq 1$  we have

$$
(7.14) \qquad \|\varphi\big(R(B_0+V)-R(B_0)\big)\big\|_{s,s'}\leq C(s,s',B_0,\varphi)(\|V\|_{1,0}+\|V^t\|_{1,0}).
$$

PROOF. This follows immediately from Proposition [7.7](#page-45-2) and Equations [\(5.18\)](#page-25-0) and [\(5.19\)](#page-25-1).

<span id="page-46-1"></span>7.3. Continuous dependence of the invertible double. Dealing with the generalized inverse would make the discussion of the parameter dependence of the invertible double rather tedious. Therefore we assume in this Subsection [7.3](#page-46-1) that UCP holds. Although with some care the results of this section carry over to families of operators where the dimensions of the spaces of "ghost solutions"  $Z_{+,0}(A), Z_{-,0}(A),$  see [\(4.12\)](#page-21-2), remain fixed (cf. [**[HKL04](#page-57-6)**, Thm. 3.16]).

Recall the definition of spaces and norms of Definition [7.1.](#page-42-5) The goal of this subsection is to prove:

<span id="page-46-2"></span>**Theorem 7.9.** The map  $(\mathscr{E}_{\text{UCP}}, d_0) \longrightarrow \mathscr{B}(L^2(M, F \oplus E), L^2_1(M, E \oplus F)), (A, T) \mapsto$  $\widetilde{A}^{-1}_{P(T)}$  is continuous.

Note that this is much more than just graph continuity of  $(A,T) \mapsto \widetilde{A}^{-1}_{P(T)}$ . Namely, by construction of the metric  $d_0$  this means that also  $(A,T) \mapsto ((\widetilde{A}_{P(T)})^*)^{-1} =$  $(\widetilde{A}_{P(-T^{-1})}^t)^{-1}$  (cf. [\(4.7\)](#page-20-7)) is continuous as a map to  $\mathscr{B}(L^2(M, F \oplus E), L^2_1(M, E \oplus F)$ . Graph continuity of  $(A, T) \mapsto \tilde{A}_{P(T)}^{-1}$  means that  $(A, T) \mapsto A_{P(T)}^{-1} \in \mathcal{B}(L^2, L^2)$  and  $(A, T) \mapsto ((A_{P(T)})^*)^{-1} \in \mathscr{B}(L^2, L^2)$  are continuous.

The result should not come as a surprise. We should take the invertible double as a guideline. Under more restrictive assumptions on A one can construct from A an invertible operator  $\widetilde{A}$  on the double  $\widetilde{M}$  of M. If A varies continuously in the metric  $d_0$  then the geometric invertible double is a continuously varying family in  $\mathscr{B}(L_1^2(\tilde{M}, \tilde{E} \oplus \tilde{F}), L^2(\tilde{M}, \tilde{F} \oplus \tilde{E}))$  and hence<sup>[6](#page-46-4)</sup> its inverse varies continuously in  $\mathscr{B}(L^2(\overline{M}, \tilde{F} \oplus \tilde{E}), L_1^2(\overline{M}, \tilde{E} \oplus \tilde{F}))$ . In the case of a geometric invertible double the nice thing is that the domains of all first order elliptic differential operators coincide with  $L_1^2$ .

<span id="page-46-4"></span><sup>6</sup>Note that for any pair of Banach spaces X, Y the inversion map  $\mathscr{B}(X,Y) \longrightarrow \mathscr{B}(Y,X), S \mapsto$  $S^{-1}$  is continuous.

The difficulty we are facing here is that the domains  $L_{1,T}^2$  vary with T. So our first task will be to transform, at least locally, the whole situation to families of operators with constant domain.

<span id="page-47-3"></span>**Definition 7.10.** Let  $e: L_s^2(\Sigma, F_\Sigma) \to L_{s+1/2}^2(M, F), s > 0$ , be a linear rightinverse to the trace map (cf. e.g. Remark [2.5.](#page-10-1)1 and [[BBW93](#page-56-1), Definition 11.7e]). For  $T, T' \in \text{Diff}^0(\Sigma; E_{\Sigma}, F_{\Sigma})$  we put

(7.15) 
$$
\Phi_{T,T'}\begin{pmatrix} f_+ \\ f_- \end{pmatrix} := \begin{pmatrix} f_+ \\ f_- + e(T'-T)\varrho f_+ \end{pmatrix}.
$$

We record some properties of  $\Phi_{T,T'}$  which are straightforward to verify.

<span id="page-47-2"></span>**Lemma 7.11.**  $\Phi_{T,T'} \in \mathscr{B}(L_1^2(M,E\oplus F))$  and we have  $\Phi_{T',T''} \circ \Phi_{T,T'} = \Phi_{T,T''}$ . In particular  $\Phi_{T,T'}$  is invertible with inverse  $\Phi_{T',T}$ .

Furthermore,  $\Phi_{T,T'}$  maps  $L^2_{1,T}(M, E \oplus F)$  bijectively onto  $L^2_{1,T'}(M, E \oplus F)$ . Finally, we have

(7.16) 
$$
\|\Phi_{T,T'} - \text{Id}\|_{1,1} \leq C(e) \|T - T'\|_{1/2,1/2}.
$$

After these preparations, the proof of Theorem [7.9](#page-46-2) is straightforward:

PROOF OF THEOREM [7.9](#page-46-2). The map  $(A,T) \mapsto \widetilde{A}_{P(T)}^{-1}$  can be factorized as follows: fix a  $T_0 \in \text{Diff}^0(\Sigma; E_\Sigma, F_\Sigma)$ . Then the following maps are continuous:

(7.17) 
$$
\mathscr{E} \longrightarrow G\mathscr{B}(L_{1,T_0}^2, L^2) \stackrel{\text{inversion}}{\longrightarrow} \mathscr{B}(L^2, L_{1,T_0}^2)
$$

$$
(A, T) \longmapsto \widetilde{A}_T \circ \Phi_{T_0, T}.
$$

Here,  $G\mathscr{B}(L_{1,T_0}^2, L^2)$  denotes the *invertible* bounded linear maps between  $L_{1,T_0}^2$  and  $L^2$ .

The continuity is seen as follows:

$$
\|\tilde{A}_{P(T)} \circ \Phi_{T_0, T} - \tilde{A}'_{P(T')} \circ \Phi_{T_0, T'} \|_{L^2_{1, T_0} \to L^2}
$$
\n
$$
\leq \|\tilde{A}_{P(T)} \circ \Phi_{T_0, T} - \tilde{A}'_{P(T')} \circ \Phi_{T_0, T'} \|_{1,0}
$$
\n
$$
\leq \|\tilde{A}\|_{1,0} \|\Phi_{T_0, T} - \Phi_{T_0, T'} \|_{1,1}
$$
\n
$$
+ \|\tilde{A} - \tilde{A}' \|_{1,0} \|\Phi_{T_0, T'} \|_{1,1}.
$$

Furthermore, the map

(7.19) 
$$
\text{Diff}^{0}(\Sigma; E, F) \times \mathscr{B}(L^{2}, L_{1, T_{0}}^{2}) \longrightarrow \mathscr{B}(L^{2}, L_{1}^{2})
$$

$$
(T, T) \longmapsto \Phi_{T_{0}, T} \circ T
$$

is continuous in view of Lemma [7.11.](#page-47-2) Hence

(7.20) 
$$
(A,T) \mapsto \widetilde{A}_{P(T)}^{-1} = \Phi_{T_0,T} \circ (\widetilde{A} \circ \Phi_{T_0,T} | L_{1,T_0}^2)^{-1}
$$

is continuous as claimed.  $\Box$ 

<span id="page-47-0"></span>**7.4. Continuous dependence of**  $S(A, T)$ . Next we give a simple criterion for the continuous dependence of  $S(A, T)$  on the input data.

<span id="page-47-1"></span>**Proposition 7.12.** The map  $(A, T) \mapsto S(A, T) \in \mathscr{B}(L^2(\Sigma, E_{\Sigma}), L^2_{\text{comp}}(M, E \oplus F))$ is continuous with respect to the norm  $N_1$  introduced in Definition [7.1](#page-42-5).

PROOF. This follows immediately from  $(5.18)$  and  $(5.19)$ .

<span id="page-48-0"></span>7.5. Continuous dependence of  $P_{\pm}$  on input data. Finally we study the dependence of  $P_{\pm}$  on B, where  $B \in \text{Diff}^1(\Sigma; E_{\Sigma})$  satisfies the usual assumptions (cf. Subsection [7.1\)](#page-44-0).

The definition of  $P_+ = P_+(B)$  for  $B \in \text{Diff}^1(\Sigma; E_{\Sigma})$  requires a choice of a spectral cut, i.e., a  $c > 0$  such that

<span id="page-48-2"></span>(7.21) 
$$
\operatorname{spec} B \cap \{z \in \mathbb{C} \mid |z| = c\} = \emptyset.
$$

Obviously, for a choice of c the map  $B \mapsto P_+(B)$  has a discontinuity at B's where [\(7.21\)](#page-48-2) is violated.

Apart from that  $P_+(B)$  should depend continuously on B with respect to the norm  $V \mapsto ||V||_{1,0} + ||V^t||_{1,0}$ . Unfortunately, we cannot prove or disprove this conjecture. Instead we mention two simple continuity criteria. The first deals with lower order perturbations of a fixed operator and the second deals with selfadjoint operators where the Spectral Theorem yields continuity in a rather simple fashion.

<span id="page-48-1"></span>**Proposition 7.13.** Let  $B \in \text{Diff}^1(\Sigma; E_{\Sigma})$  be a first order elliptic differential operator such that  $B - \lambda$  is parameter dependent elliptic for  $\lambda$  in a conic neighborhood of iR. Furthermore, let  $V \in CL^{\alpha}(\Sigma; E_{\Sigma})$ ,  $\alpha < 1$ , with  $||V||_{\alpha,0}$  sufficiently small. Then  $B + V - \lambda$  is also parameter dependent elliptic in a conic neighborhood of i $\mathbb R$ and for  $c > 0$  such that  $\operatorname{spec}(B + tV) \cap \{z \in \mathbb{C} \mid |z| = c\} = \emptyset$ ,  $0 \le t \le 1$ , we have the estimate

$$
||P_+(B+V) - P_+(B)||_{0,0} \le C(B)||V||_{\alpha,0}.
$$

PROOF. By a Neumann series argument it is clear that such a c exists. Let  $\Gamma_{+}$  (see Figure [2\)](#page-15-0) be the usual contour. Analogously to Lemma [7.6](#page-44-1) one shows the estimate

$$
(7.22) \qquad ||(\lambda - (B + V))^{-1} - (\lambda - B)^{-1}||_0 \le C(B) ||V||_{\alpha,0} |\lambda|^{-2 + \alpha}, \quad \lambda \in \Gamma_+,
$$

from which the claim, thanks to  $-2 + \alpha < -1$ , follows by invoking the contour integral.

We now turn to (formally) selfadjoint  $B$ . We first give a slight improvement of [[Les05](#page-57-25), Prop. 2.2] on the continuity of the  $Riesz$ —map (cf. loc. cit. Equ.  $(2.2)$ ).

<span id="page-48-4"></span>**Proposition 7.14.** Let  $Ell^{1,sa}(\Sigma; E_{\Sigma}) \subset Diff^{1}(\Sigma; E_{\Sigma})$  denote the space of selfadjoint first order elliptic differential operators. Then the Riesz–map

$$
B \mapsto B(\mathrm{Id} + B^2)^{-1/2}
$$

is continuous  $(Ell^{1,sa}, \|\cdot\|_{1,0}) \to \mathscr{B}(L^2_s(\Sigma, E_{\Sigma}))$  for all  $s \in [-1/2, 1/2]$ .

PROOF. For  $s = 0$  this was proved in [[Les05](#page-57-25), Prop. 2.2]. The proof in loc. cit., however shows the claimed stronger statement (cf.  $[Les 05, Equ. (2.19)]$ ). For the convenience of the reader let us present the argument.

It suffices to prove the continuity of F for  $s = 1/2$ . Since  $F(B)$  is selfadjoint  $||F(B) - F(\tilde{B})||_{1/2,1/2} = ||F(B) - F(\tilde{B})||_{-1/2,-1/2}$  and hence by complex inter-polation (cf. [[Les05](#page-57-25), Appendix])  $||F(B) - F(\tilde{B})||_{s,s} \leq ||F(B) - F(\tilde{B})||_{1/2,1/2}$  for  $|s| \leq 1/2$ .

Fix a  $B \in \text{Ell}^{1,\text{sa}}(\Sigma, E_{\Sigma})$  and we have to prove the continuity of F at B. Let  $0 < q < \frac{1}{2}$  and consider  $\widetilde{B} \in \text{Ell}^{1,\text{sa}}(\Sigma, E_{\Sigma})$  with

<span id="page-48-3"></span>(7.23) 
$$
\| (B - \widetilde{B})(B \pm i)^{-1} \|_{0,0} + \| (B \pm i)^{-1} (B - \widetilde{B}) \|_{0,0} \le q.
$$

Note that by ellipticity the graph norm of  $B$  is equivalent to the Sobolev norm  $\|\cdot\|_1$ , hence the left hand side of [\(7.23\)](#page-48-3) induces a metric on Ell<sup>1,sa</sup>(Σ, E<sub>Σ</sub>) which is equivalent to the metric induced by  $\|\cdot\|_{1,0}$ . The Neumann series then immediately implies

(7.24) 
$$
\|(\widetilde{B}+i)^{-1}(B+i)\|_{0,0} \leq \frac{1}{1-q}.
$$

Thus, for  $f \in L^2(\Sigma, E_{\Sigma})$  we have

(7.25) 
$$
\|(\widetilde{B}+i)^{-1}f\|_0 \leq \frac{1}{1-q} \|(B+i)^{-1}f\|_0
$$

and

$$
(7.26)\ \|(B+i)^{-1}f\|_{0} \leq \|(B+i)^{-1}(\widetilde{B}+i)\|_{0,0}\|(\widetilde{B}+i)^{-1}f\|_{0} \leq (1+q)\|(\widetilde{B}+i)^{-1}f\|_{0}.
$$

This implies the operator inequalities

(7.27) 
$$
\frac{1}{(1+q)^2}|B+i|^{-2} \leq |\widetilde{B}+i|^{-2} \leq \frac{1}{(1-q)^2}|B+i|^{-2}.
$$

Since the square root is an operator–monotonic increasing function ([[KR97](#page-57-26), Prop. 4.2.8]) we may take the square root of these inequalities and after subtracting  $|B + i|^{-1}$  we arrive at

(7.28) 
$$
-\frac{q}{1+q}|B+i|^{-1} \leq |\tilde{B}+i|^{-1} - |B+i|^{-1} \leq \frac{q}{1-q}|B+i|^{-1}.
$$

This gives

(7.29) 
$$
\| |B + i|^{1/2} |\widetilde{B} + i|^{-1} |B + i|^{1/2} - \text{Id} \|_{0,0} \le \frac{q}{1-q}.
$$

After these preparations we find

$$
||F(B) - F(B)||_{1/2,1/2}
$$
  
\n
$$
= || |i + B|^{1/2} (F(B) - F(\tilde{B})) |i + B|^{-1/2} ||_{0,0}
$$
  
\n
$$
= || |i + B|^{-1/2} (F(B) - F(\tilde{B})) |i + B|^{1/2} ||_{0,0}
$$
  
\n
$$
\le || |i + B|^{-1/2} (B - \tilde{B}) |i + B|^{-1/2} ||_{0,0}
$$
  
\n(7.30)  
\n
$$
+ || |B + i|^{-1/2} (\tilde{B}(|i + B|^{-1} - |i + \tilde{B}|^{-1})) |i + B|^{1/2} ||_{0,0}
$$
  
\n
$$
\le || |i + B|^{-1} (B - \tilde{B}) ||_{0,0}
$$
  
\n
$$
+ || |i + B|^{-1/2} \tilde{B} |i + B|^{-1/2} || || \text{Id} - |i + B|^{1/2} |i + \tilde{B}|^{-1} |i + B|^{1/2} ||_{0,0}
$$
  
\n
$$
\le q + || |i + B|^{-1} \tilde{B} ||_{0,0} \frac{q}{1 - q}
$$
  
\n
$$
\le q(1 + \frac{1 + q}{1 - q}).
$$

Here we have used that for a first order operator  $V$  one has

(7.31) 
$$
\| |i + B|^{-1/2} V |i + B|^{-1/2} \|_{0,0} \le \| |i + B|^{-1} V \|_{0,0}.
$$

This inequality also follows from complex interpolation (see [[Les05](#page-57-25), Appendix]). This shows that if  $||B_n - B||_{1,0} \to 0$  then  $F(B_n) \to F(B)$  in  $L^2_{1/2}(\Sigma, E_{\Sigma})$  and we are done.

<span id="page-49-0"></span>**Proposition 7.15.** Let  $Ell_c^{1,sa}(\Sigma; E_{\Sigma}) \subset Ell_c^{1,sa}(\Sigma; E_{\Sigma}) \subset Diff^1(\Sigma; E_{\Sigma})$  denote the space of selfadjoint first order elliptic differential operators B with  $\pm c \notin \text{spec } B$ . Then for  $|s| \leq 1/2$  the map

(7.32) 
$$
\left(Ell_c^1, \|\cdot\|_{1,0}\right) \longrightarrow \mathscr{B}(L_s^2(\Sigma, E_{\Sigma}))
$$

$$
B \mapsto 1_{[c,\infty)}(B)
$$

is continuous.

PROOF. We first note that for  $B \in \text{Ell}_c^{1,\text{sa}}$  we have by the Spectral Theorem

$$
1_{[c,\infty)}(B) = 1_{[F(c),\infty)}(F(B)).
$$

Since, independently of  $s \in [-1/2, 1/2]$ , we have

(7.33)  $\sec(F(B)) \subset [-1,1],$  $1_{[c,\infty)}(B)$  is given by the contour integral

(7.34) 
$$
\frac{1}{2\pi i} \oint_{|z - (F(\lambda) + 2)| = 2} \left( z - F(B) \right)^{-1} dz.
$$

In view of the previous Proposition [7.14](#page-48-4) this proves the claim. cf. also [[Les05](#page-57-25), Lemma 3.3].

<span id="page-50-0"></span>7.6. Continuity of families of well–posed selfadjoint Fredholm extensions. In Theorems 3.8 and 3.9 in [[BBLP05](#page-56-2)] the continuous dependence of the invertible double, the Calderón and Poisson operators and the graph continuity of realizations of well–posed boundary value problems are discussed in a special case. More precisely it was assumed that  $J^2 = -Id$  and that the tangential operator had a selfadjoint leading symbol.

Unfortunately the proof of Theorem 3.9 in [[BBLP05](#page-56-2)] was incomplete. Now we can present correct statements with complete proofs. Our result is more general than loc. cit. in the sense that we do neither have to assume that  $J^2 = -Id$ , nor do we have to assume a selfadjoint leading symbol of the tangential operator in all cases. On the negative side we must admit that the topology we have to impose on the space of differential operators is stronger than we hoped at the time of writing of [[BBLP05](#page-56-2)]. The correct replacement for Theorem 3.9a in loc. cit. is Theorem [7.2a](#page-42-0) and the correct replacement for Theorem 3.9b in loc. cit. are Theorem [7.2b](#page-42-0) and Corollary [7.4.](#page-43-0)

Theorem [7.9](#page-46-2) above generalizes Theorem 3.8 in loc. cit. and [[Nic97](#page-57-0), Proposition B.1]

Next we deal with families of realizations of well–posed boundary conditions (cf. Theorem 3.9d in loc. cit.).

<span id="page-50-1"></span>**Theorem 7.16.** Consider the space of pairs  $(A, P)$  where  $A \in \text{Diff}^1(M; E)$  is elliptic and formally selfadjoint and  $P \in CL^{0}(\Sigma; E_{\Sigma})$  is an orthogonal projection which is well–posed with respect to  $A$  and such that  $A_P$  is selfadjoint. Equip this space with the metric  $d_0$ , i.e.,

(7.35) 
$$
d_0((A, P), (A', P')) = N_0(A - A', P - P').
$$

Then the map  $(A, P) \mapsto (A_P + i)^{-1} \in \mathcal{B}(L^2(M, E), L^2_1(M, E))$  is continuous with respect to the  $d_0$  metric on the space of such pairs  $(A, P)$ . In particular  $(A, P) \mapsto$  $(A_P + i)^{-1}$  is continuous or, equivalently,  $(A, P) \mapsto A_P$  is graph continuous.

PROOF. The proof is basically the same as the proof of Theorem [7.9](#page-46-2) once the analogue of the maps  $\Phi_{T,T'}$  is established. Note that if  $P,Q \in CL^{0}(\Sigma; E_{\Sigma})$  are orthogonal projections with  $||P - Q||_{1/2,1/2} < 1$  they form an *invertible pair*, i.e.,  $P : \text{im } Q \longrightarrow \text{im } P$  is invertible. For such P, Q we put analogously to Definition [7.10](#page-47-3)

$$
\Psi_{P,Q}(f) := f - e(P - Q)\varrho f.
$$

Then as in Lemma [7.11](#page-47-2) we have

(7.37) 
$$
\|\Psi_{P,Q} - \text{Id}\|_{1,1} \leq C(e) \|P - Q\|_{1/2,1/2},
$$

and thus  $\Psi_{P,Q}$  is invertible for  $||P - Q||_{1/2,1/2}$  small enough. Furthermore,  $\Psi_{P,Q}$ maps  $\mathscr{D}(A_Q) = \{f \in L_1^2(M, E) \mid Q \varrho f = 0\}$  bijectively onto  $\mathscr{D}(A_P) = \{f \in L_1^2(M, E) \mid Q \varrho f = 0\}$  $L_1^2(M, E) | Q \varrho f = 0.$ 

#### 52 BERNHELM BOOSS-BAVNBEK, MATTHIAS LESCH, AND CHAOFENG ZHU

Now one mimics the proof of Theorem [7.9](#page-46-2) with  $\Psi_{P,Q}$  instead of  $\Phi_{T,T'}$  and  $A_P + i$  instead of  $A_{P(T)}$ .

Finally we state a more precise version of [[BBLP05](#page-56-2)], Theorem 3.9c. Note that the following version applies to a much wider class of operators than loc. cit.

**Theorem 7.17.** Let  $Ell_{\mathrm{UCP}}^{1,\mathrm{sa}}(M;E) \subset Diff^1(M;E)$  denote the space of formally selfadjoint elliptic differential operators acting on sections of the Hermitian vector bundle E which satisfy UCP and whose tangential operator  $B_0$  has a selfadjoint leading symbol. We equip this space with the strong metric induced by the embedding  $Ell_{\mathrm{UCP}}^{1,\mathrm{sa}}(M;E) \rightarrow \mathscr{E}_{\mathrm{UCP}}(M;E), A \mapsto (A,(J_0^t(A))^{-1})$ 

Then the map

 $Ell^{1,sa}(M;E) \longrightarrow \mathscr{B}(L_1^2(M,E), L^2(M,E)), \quad A \longmapsto A_{C_+}$ 

sending  $A$  to the selfadjoint well-posed realization associated to the Calderón projection is continuous. Here  $C_+$  denotes the version of the Calderón projection constructed from  $(J_0^t(A))^{-1}$ , cf. Proposition [5.12](#page-29-0).

PROOF. Note that  $A_{C_+}$  is selfadjoint by Theorem [6.1](#page-30-2) (II), Prop. [5.12.](#page-29-0)  $A_{C_+}$  is indeed a selfadjoint realization of a well–posed boundary value problem.

By Corollary [7.4](#page-43-0)  $A \mapsto (A, C_+)$  is now a continuous map from  $Ell_{\text{UCP}}^{1,\text{sa}}(M, E)$  to the space of pairs described in Theorem [7.16](#page-50-1) and hence Theorem [7.16](#page-50-1) yields the claim.

## <span id="page-52-0"></span>Appendix A. Smooth symmetric elliptic continuations with constant coefficients in normal direction

In this appendix, we restrict ourselves to formally selfadjoint operators. To begin with, we write  $D = J(\frac{d}{dx} + B)$  like in [\(2.9\)](#page-7-3) with the relations  $J^* = -J$  and  $JB = J' - B<sup>t</sup>J$  of [\(2.20\)](#page-8-6) without loss of generality.

Sometimes one is interested in operators satisfying additional relations. We shall consider the following cases:

- (I) D arbitrary symmetric elliptic, e.g. no additional relations.
- (II)  $J^2(x) = -Id$ . We will see below that then, after a suitable coordinate transformation, we can even obtain that  $J_x = J$  is constant. This is the Dirac operator case.
- (III)  $B_0 B_0^t$  of order 0. In view of [\(2.20\)](#page-8-6) this implies that  $J_0B_0 + B_0J_0$  is of order 0, too.
- (IV)  $B B<sup>t</sup>$  of order 0. Analogously then  $JB + BJ$  is of order 0, too.
- (V)  $D = J\left(\frac{d}{dx} + B\right) + \frac{1}{2}J' + C$  with  $J^* = -J$ ,  $B = B^t$ ,  $JB + BJ = 0$ ,  $J' = \frac{dJ}{dx}$ , and C of order 0. Then automatically  $C = C^*$ .

One can think of even more cases. But the preceding cases suffice for the moment. From now on we shall write

(A.1) 
$$
D = J\left(\frac{d}{dx} + B\right) + \frac{1}{2}J' + C
$$

in all five cases although the notation is redundant in cases I-IV. Recall, that here  $C$  is of order 0 and  $B$  is of order 1.

**Proposition A.1.** Consider the case (II). Then there is a smooth unitary gauge transformation  $U \in C^{\infty}([0,\varepsilon), \Gamma^{\infty}(\Sigma; \mathscr{U}(E_{\Sigma})))$  such that

<span id="page-52-1"></span>
$$
J_x = U(x)J_0U(x)^*.
$$

With the unitary transformation

$$
\Phi_1 : L^2([0, \varepsilon) \times \Sigma, E_\Sigma) \longrightarrow L^2([0, \varepsilon) \times \Sigma, E_\Sigma),
$$
  

$$
f \longmapsto Uf
$$

we find

(A.2) 
$$
\Phi_1^{-1} D \Phi_1 = J_0 \left( \frac{d}{dx} + U^* B U \right) + \widetilde{C}.
$$

Here and in the following we use the abbreviation  $E_{\Sigma} := E \restriction \Sigma$  introduced in Subsection [2.1](#page-5-1) and, by slight abuse of notation,  $E_{\Sigma} := E \mid [0, \varepsilon) \times \Sigma$  as well. Note that selfadjointness of B or C and the relation  $JB + BJ = 0$  (of lower order) are preserved under  $\Phi_1$ .

PROOF. We give a brief sketch; it is basically a standard fact often used in K–theory [[Bla86](#page-56-8), Prop. 4.3.3], see also [[BL01](#page-57-12), Section 3].

We only show a bit less, namely that the claim is true after making  $\varepsilon$  a bit smaller; but this is not really a loss of generality.

After possibly making  $\varepsilon$  smaller we may assume that

(A.3) 
$$
\| \operatorname{Id} + J_x J_0 \| = \| \operatorname{Id} - J_x^{-1} J_0 \| < 2, \text{ for } 0 \le x \le \varepsilon.
$$

Then the operator  $Z_x := \text{Id} - \frac{1}{2}(J_xJ_0 + \text{Id}) = \frac{1}{2}(\text{Id} - J_xJ_0)$  is invertible. Moreover, since  $J_x^2 = - \text{Id}$  we find  $J_x Z_x = \frac{1}{2}(J_x + J_0) = Z_x J_0$  and thus  $J_x = Z_x J_0 Z_x^{-1}$ .

One now checks by direct calculation that  $Z_x$  is normal and that  $Z_x Z_x^*$  commutes with  $J_x$  and  $J_0$  [[BL01](#page-57-12), (3.7)]. Hence we may put  $U(x) := Z_x^{-1}\sqrt{Z_x Z_x^*}$  to reach the conclusion.

The remaining assertions are now clear.

<span id="page-53-0"></span>**Remark A.2.** (1) One may ask under what conditions we can obtain unitary  $J_0$ , i.e.,  $J_0^2 = -Id$ ? It is not clear whether this question has a definite answer. E.g., it seems impossible to find a coordinate transformation preserving the symmetry of the fixed given differential operator D and providing a unitary leading symbol  $\tilde{J}$  in the new normal direction, unless all eigenvalues of the original  $J_0$  are of the form  $\pm i$ .

(2) For the symplectic Hilbert space  $(L^2(\Sigma, E_{\Sigma}), \langle \cdot, \cdot \rangle, \omega(\cdot, \cdot)),$  however, there is a simple answer. Here we set

$$
\langle f, g \rangle := \int_{\Sigma} \langle f(p), g(p) \rangle_{E_p} \, d\text{vol}
$$
 and  $\omega(f, g) := \langle J_0 f, g \rangle$ .

As always in symplectic Hilbert spaces (see, e.g., [[BBZ04](#page-57-21), Lemma 1.5]), we can preserve the symplectic form  $\omega$  while deforming the inner product  $\langle \cdot, \cdot \rangle$  of  $L^2(\Sigma, E_{\Sigma})$ over  $\Sigma$  smoothly into

$$
\langle \cdot, \cdot \rangle^{\tilde{}} := \langle Sf, g \rangle
$$
 with selfadjoint  $S := \sqrt{J_0^* J_0}$ 

in such a way that  $\omega(f, g) = \langle \tilde{J}f, g \rangle^{\sim}$  with  $\tilde{J}^* = -\tilde{J}$  and  $\tilde{J}^2 = -Id$ .

The notation  $(A.1)$  has another advantage in case  $(IV)$ . Namely, replacing B by  $\frac{1}{2}(B+B^t)$  and C by  $C+\frac{1}{2}J(B-B^t)$  if necessary we see that we may assume that  $B = B<sup>t</sup>$ . Note that this does not work so easily in case (III).

Summing up, the cases  $(I)-(V)$  may now be described as follows (when D is written as in  $(A.1)$ :

- (I) D arbitrary symmetric elliptic, e.g. no additional relations.
- (II)  $J^2 = -Id$ ,  $J_x = J_0$  constant. Again, this is called the Dirac operator case.
- (III)  $B_0 B_0^t$  and  $J_0 B_0 + B_0 J_0$  of order 0.

(IV)  $B = B^t$ ,  $JB + BJ$  of order 0.

(V)  $B = B^t$ ,  $JB + BJ = 0$ ,  $C = C^*$ .

We consider  $D$  as before. The goal of this appendix is to prove the following Theorem.

**Theorem A.3.** Let D be as in [\(A.1\)](#page-52-1) with families  $J_x, B_x, C_x$  smoothly depending on x. Then there is a  $\delta > 0$  and a symmetric elliptic first order differential operator D (i.e., with smooth coefficients) on  $\Gamma^{\infty}(-\delta,\varepsilon)\times\Sigma; E_{\Sigma}$ ) with the following properties:

- (1)  $\widetilde{D}\upharpoonright [0,\varepsilon)\times\Sigma=D$ , *i.e.*,  $\widetilde{D}$  extends D.
- (2)  $\widetilde{D}\upharpoonright[-\delta, -2/3\delta)\times\Sigma$  =  $J_0\left(\frac{d}{dx}+B_0\right)+C_0$ . In particular,  $\widetilde{D}$  has constant coefficients near the new boundary  $\{-\delta\}\times\Sigma$ , and the constant coefficients are given just by smoothly "rewinding" to the coefficients of D at 0.

Note that in the formula for  $\widetilde{D}$  near the boundary  $\frac{1}{2}J'$  is left out deliberately to make the constant coefficient operator symmetric.

Note also that due to the concrete formula for  $\overline{D}$  near the boundary the additional relations in the cases (II)–(V) still hold for the extended operator. It is not claimed (and it is open in some cases) that the relations in  $(IV),(V)$  are preserved on the whole interval  $[-\delta, 0]$ .

PROOF. 1. Let

(A.4) 
$$
D_0 := J_0 \left( \frac{d}{dx} + B_0 \right) + C_0.
$$

Since  $D_0$  has constant coefficients we may think of  $D_0$  as acting on  $\Gamma_0^{\infty}(\mathbb{R} \times \Sigma; E_{\Sigma})$ . Note that since D is formally selfadjoint, so is  $D_0$ . To see that we recall that

 $D = D<sup>t</sup>$  is (thanks to our singling-out of  $\frac{1}{2}J'$  in Equation [\(A.1\)](#page-52-1)) equivalent to the relations

(A.5) 
$$
J^* = -J, \quad -B^t J + C^* = JB + C.
$$

Then

(A.6) 
$$
D_0^t = J_0 \frac{d}{dx} - B_0^t J_0 + C_0^* = J_0 \frac{d}{dx} + J_0 B_0 + C_0 = D_0.
$$

2. Now we apply the definition of smoothness of maps defined on a manifold with boundary to conclude that  $J, B, C$  have smooth extensions to the whole negative half-line. More precisely there exist

$$
\widetilde{J} \in \Gamma^{\infty}((-\infty,\varepsilon) \times \Sigma; E_{\Sigma}), \quad \widetilde{C} \in \Gamma^{\infty}((-\infty,\varepsilon) \times \Sigma; E_{\Sigma}) \text{ and}
$$
  
 $\widetilde{B} \in C^{\infty}((-\infty,\varepsilon), \text{Diff}^{1}(\Sigma; E_{\Sigma}))$ 

such that

$$
\widetilde{J}\,[0,\varepsilon)\times\Sigma=J,\ \widetilde{C}\,[0,\varepsilon)\times\Sigma=C,\ \text{and}\ \widetilde{B}\,[0,\varepsilon)\times\Sigma=B.
$$

Replacing  $\tilde{J}$  by  $\frac{1}{2}(\tilde{J}-\tilde{J}^*)$  if necessary, we can, additionally, obtain that  $\tilde{J}^* = -\tilde{J}$ .

The extensions of  $J$  and  $C$  are immediate. However, for  $B$  one might feel a bit uneasy because of the target space  $\text{Diff}^1(\Sigma; E_{\Sigma})$ . Well, first extend the leading symbol of  $B$ , which works like for  $J$  and  $C$ . Then choose a right inverse to the symbol map on  $\Sigma$  to obtain an operator map  $\widetilde{B}_1$ . On  $[0, \varepsilon) \times \Sigma$ ,  $\widetilde{B}_1$  coincides with B up to order 0. The difference  $\widetilde{B}_1 - B[0,\varepsilon) \times \Sigma$  is again smooth. Extend it and subtract it to make up for the 0 order defect. This yields the wanted  $\widetilde{B}$ .

So, we can form the differential operator

(A.7) 
$$
\widetilde{D}_1 := \widetilde{J}\left(\frac{d}{dx} + \widetilde{B}\right) + \frac{1}{2}\widetilde{J}' + \widetilde{C}
$$

which is now defined on  $(-\infty, \varepsilon) \times \Sigma$ , has smooth coefficients and  $\widetilde{D}_1 | [0, \varepsilon) \times \Sigma = D$ .

So far  $D_1$  is not necessarily formally selfadjoint. Put

(A.8) 
$$
\widetilde{D}_2 := \frac{1}{2} \left( \widetilde{D}_1 + \widetilde{D}_1^t \right) =: \widetilde{J} \left( \frac{d}{dx} + \widetilde{B}_2 \right) + \frac{1}{2} \widetilde{J}' + \widetilde{C}_2.
$$

Next, consider a cut–off function  $\varphi\in C^\infty(\mathbb{R})$  with

(A.9) 
$$
\varphi(x) = \begin{cases} 1, & x \leq -\frac{2}{3}\delta, \\ 0, & x \geq -\frac{1}{3}\delta. \end{cases}
$$

Then we consider the operator

(A.10) 
$$
\widetilde{D} := \varphi D_0 + (1 - \varphi)\widetilde{D}_2 + \frac{1}{2}\varphi'(J_0 - \widetilde{J}).
$$

The last summand was left out in [[BBLP05](#page-56-2), (3.14)]; the additional term is, however, necessary to make  $\widetilde{D}$  formally selfadjoint.

 $\widetilde{D}$  has the following properties:

- (1)  $\widetilde{D}^t = \widetilde{D}$ . That follows immediately from the formal selfadjointness of  $D_0$ ,  $\widetilde{D}_2$  and the relations  $[D_0, \varphi] = J_0 \varphi', \; [\widetilde{D}_2, \varphi] = J \varphi'.$
- (2)  $\widetilde{D} \restriction [0, \varepsilon) \times \Sigma = D.$
- (3)  $\widetilde{D}\lceil[-\delta, -2/3\delta] \times \Sigma = J_0\left(\frac{d}{dx} + B_0\right) + C_0.$

It remains to discuss the ellipticity of  $\widetilde{D}$ . So let

$$
x \in [-\delta, 0], p \in \Sigma, \xi \in T_p^*(\Sigma) \text{ and } \lambda dx + \xi \in S_{x,p}^*(\mathbb{R} \times \Sigma),
$$

S<sup>\*</sup> denoting the cosphere bundle. Then  $|\lambda|^2 + |\xi|^2 = 1$  and we find for the leading symbol of  $\widetilde{D}$ 

(A.11)  
\n
$$
\sigma_{\tilde{D}}(x, p)[\lambda dx + \xi] = \left(\varphi(x)\sigma_{D_0}(x, p) + (1 - \varphi(x))\sigma_{\tilde{D}_2}(x, p)\right)[\lambda dx + \xi]
$$
\n
$$
= \varphi(x)J_0(i\lambda + \sigma_{B(0)}(\xi)) + (1 - \varphi(x))J_x(i\lambda + \sigma_{B(x)}(\xi))
$$
\n
$$
= J_0(i\lambda + \sigma_{B(0)}(\xi)) + (1 - \varphi(x))\{J_x(i\lambda + \sigma_{B(x)}(\xi)) - J_0(i\lambda + \sigma_{B(0)}(\xi))\}.
$$

Hence, by the compactness of  $\Sigma$  and by the continuity of  $J_x$  and  $\sigma_{B(x)}$  we may choose  $\delta$  so small that  $\widetilde{D}$  is elliptic.

The theorem is proved.

**Remark A.4.** We briefly discuss for the various cases  $(I)$ – $(V)$  whether the construction of  $D$  can be modified such that the relations continue to hold.

- (I) Since there are no relations, there is nothing to worry about.
- (II) Since J is constant, we may extend it constantly.
- (III) At the new boundary  $\{-\delta\}\times\Sigma$  we have by construction  $B(-\delta)=B(0)$ hence (III) also works fine.
- (IV), (V) By construction of  $D_0$  it is clear that  $\widetilde{D}$  also satisfies (IV) or (V) in the collar  $[-\delta, -2/3\delta] \times \Sigma$  of the new boundary. However, we do not know so far whether  $D$  can be constructed in such a way that the relations hold on the whole interval  $[-\delta, 0]$ .

We leave it to the reader to prove that the latter is indeed possible in the case  $(II)+(IV)$  or  $(II)+(V)$ .

In Subsection [2.1](#page-5-1) we first applied a unitary transformation to our operator. It remains to clarify what happens if we transform the whole construction back to the original situation. The result reads as follows:

**Theorem A.5.** Let  $(M, g_1)$  be a compact Riemannian manifold with boundary,  $(E, h_1)$  a Hermitian vector bundle over M and  $A: \Gamma_0^{\infty}(M; E) \to \Gamma_0^{\infty}(M; E)$  a first order elliptic differential operator which is formally selfadjoint in the Hilbert space  $L^2(M,E;g_1,h_1).$ 

Choose metrics q on M, h on E which are product near the boundary  $\Sigma = \partial M$ and such that g  $\partial M = g_1 \partial M$ ,  $h \partial M = h_1 \partial M$ . Let  $\Phi$  be the isometry of [\(2.8\)](#page-7-4) and assume that we have

(A.12) 
$$
\Phi A \Phi^{-1} = J\Big(\frac{d}{dx} + B\Big) + \frac{1}{2}J' + C.
$$

For  $\delta > 0$  form  $M_{\delta} := \left( [-\delta, 0] \times \Sigma \right) \cup_{\Sigma} M$ .

Then for  $\delta$  sufficiently small there are a Hermitian vector bundle  $(E_{\delta}, h_{1,\delta})$ over  $M_{\delta}$ , a Riemannian metric  $g_{1,\delta}$  on  $M_{\delta}$  and a first order symmetric elliptic differential operator  $A_{\delta}$  on  $M_{\delta}$  such that:

- (1)  $E_{\delta}$ ,  $g_{1,\delta}$ ,  $h_{1,\delta}$ ,  $A_{\delta}$  are extensions of E,  $g_1$ ,  $h_1$ , A respectively.
- (2)  $g_{1,\delta}$  and  $h_{1,\delta}$  are product metrics near  $\partial M_{\delta}$ . More precisely, we have on  $[-\delta, -2/3\delta] \times \Sigma$

<span id="page-55-0"></span>(A.13) 
$$
g_{1,\delta} = dx^2 \oplus g(0),
$$

$$
h_{1,\delta}(x) = h(0).
$$

(3) With the natural extension of  $\Phi$  to  $M_{\delta}$  we have on  $[-\delta, -2/3\delta] \times \Sigma$ :

(A.14) 
$$
\Phi A \Phi^{-1} = J_0 \Big( \frac{d}{dx} + B_0 \Big) + C_0.
$$

PROOF. We apply the previous Theorem to  $\Phi A \Phi^{-1}$  in  $L^2(M,E;g,h)$ . Since the spaces of metrics on  $M$  and on  $E$  are positive cones, we may extend the metrics  $g_1, h_1$  to smooth metrics  $\tilde{g}_\delta, h_{1,\delta}$  on  $M_\delta$  in such a way that on the collar  $[-\delta, -2/3\delta] \times \Sigma$  they are of the form  $(A.13)$ . Now consider the unitary transformation  $\Psi: L^2(M_\delta, E; g_{1,\delta}, h_{1,\delta}) \to L^2(M_\delta, E; g, h)$  as in Lemma [2.1](#page-6-1)  $(g, h \text{ extend})$ trivially to  $M_{\delta}$  since they are already product metrics). By the construction ex-plained before Lemma [2.1](#page-6-1) we see that  $\Psi$  [ $-\delta$ ,  $-2/3\delta$ ]  $\times \Sigma =$  Id and therefore we reach the conclusion from the previous theorem.  $\Box$ 

Remark A.6. (1) The preceding theorem shows that it is always possible to extend a given symmetric elliptic differential operator (of first order)  $A$  to a symmetric elliptic differential operator  $\tilde{A}$  by attaching a collar with finite cylindrical end with new boundary  $\Sigma'$  such that the Riemannian and Hermitian structures become product close to  $\Sigma'$  and the operators  $J_x$ ,  $B_x$  and  $C_x$  become independent of the normal variable x close to  $\Sigma'$ ; i.e., we can always bring the operator in product form near a new boundary by suitable prolongation under preservation of the symmetry property.

(2) The previous discussion shows that  $J_0^2 = -Id$  is perfectly enough to obtain, by coordinate transformation,  $J_x = J_0$  and hence  $J_x^2 = -Id$  in a whole neighbor-hood of the new boundary. See also Remark [A.2](#page-53-0) for conditions for  $J_0^2 = -Id$ .

(3) The previous discussion can be made parameter dependent with the right notion of parameter dependency, cf. our Section [7.](#page-41-1)

(4) The question remains whether weak inner UCP can be preserved under the symmetric prolongation. The short answer is

- yes in the cases (III)-(V), i.e., when the leading symbol of the tangential operator is symmetric;
- yes in a very restricted sense, namely that the UCP-defect dimension

 $(A.15)$   $d(x)$ 

$$
d(x) := \dim\{u \mid Au = 0 \text{ and } u \in \Sigma(x) = 0\}
$$

is constant on the last part of the collar, i.e., for sufficiently negative tangential coordinate  $x$  when constant coefficients in normal direction are obtained. Here  $\Sigma(x) := \{x\} \times \partial M$  denotes the parallel surface in the collar.

### References

- <span id="page-56-0"></span>[Bär99] C. Bär, Zero sets of semilinear elliptic systems of first order, Invent. Math. 138 (1999), 183–202.
- <span id="page-56-6"></span>[BBC08] W. Ballmann, J. Brüning, and G. Carron, Regularity and index theory for Dirac-Schrödinger systems with Lipschitz coefficients, J. Math. Pures Appl. 89:5 (2008), 429–476, [math/0703185.](http://arxiv.org/abs/math/0703185)
- <span id="page-56-8"></span>[Bla86] B. Blackadar, K-theory for operator algebras, MSRI Publ., vol. 5, Springer–Verlag, Berlin–Heidelberg–New York, 1986.
- <span id="page-56-3"></span>[BBF98] B. Booß-Bavnbek and K. Furutani, The Maslov index: a functional analytical definition and the spectral flow formula, Tokyo J. Math. 21 (1998), 1–34.
- <span id="page-56-7"></span>[BBF99] , Symplectic functional analysis and spectral invariants, in: Geometric aspects of partial differential equations (B. Booß–Bavnbek and K.P. Wojciechowski, eds.), Contemp. Math., vol. 242, Amer. Math. Soc., 1999, pp. 53–83.
- <span id="page-56-5"></span>[BBFO01] B. Booß-Bavnbek, K. Furutani, and N. Otsuki, Criss-cross reduction of the Maslov index and a proof of the Yoshida-Nicolaescu theorem, Tokyo J. Math. 24 (2001), 113– 128.
- <span id="page-56-4"></span>[BBL09] B. Booß-Bavnbek and M. Lesch, The invertible double of elliptic operators, Lett. Math. Phys. 87 (2009), 19–46.
- <span id="page-56-2"></span>[BBLP05] B. Booß-Bavnbek, M. Lesch, and J. Phillips, Unbounded Fredholm operators and spectral flow, Canad. J. Math.  $57$  (2) (2005), 225-250, math. FA/0108014.
- <span id="page-56-1"></span>[BBW93] B. Booß-Bavnbek and K. P. Wojciechowski, Elliptic boundary problems for Dirac operators, Birkhäuser, Basel, 1993.
- <span id="page-57-21"></span>[BBZ04] B. Booß-Bavnbek and C. Zhu, Weak symplectic functional analysis and general spectral flow formula, [math.DG/0406139,](http://arxiv.org/abs/math/0406139) 2004.
- <span id="page-57-2"></span>[BBZ05] B. Booß-Bavnbek and C. Zhu, General spectral flow formula for fixed maximal domain, Centr. Eur. J. Math. 3 (3) (2005), 1–20, [math.DG/0504125.](http://arxiv.org/abs/math/0504125)
- <span id="page-57-23"></span>[BBZ09] , Weak symplectic functional analysis, Maslov index under symplectic reduction, and general spectral flow formula, in preparation, draft available online: http://milne.ruc.dk/∼[Booss/ell/SpectralFlow2Maslov.pdf](http://milne.ruc.dk/~Booss/ell/SpectralFlow2Maslov.pdf)
- <span id="page-57-20"></span>[Bra02] M. Braverman, New proof of the cobordism invariance of the index, Proc. Amer. Math. Soc. 130 (4) (2002), 1095–1101, [math.SP/0011237.](http://arxiv.org/abs/math/0011237)
- <span id="page-57-12"></span>[BL01] J. Brüning and M. Lesch, On boundary value problems for Dirac type operators: I. Regularity and self–adjointness, J. Funct. Anal. 185 (2001), 1–62, [math.FA/9905181.](http://arxiv.org/abs/math/9905181)
- <span id="page-57-8"></span>[Bur70] T. Burak, On spectral projections of elliptic operators, Ann. Scuola Norm. Sup. Pisa (3) 24 (1970), 209–230.
- <span id="page-57-19"></span>[Fre05] C. Frey, On non-local boundary value problems for elliptic operators, PhD Thesis, Köln, [http://www.sfbtr12.uni-koeln.de/dissertationfrey.pdf,](http://www.sfbtr12.uni-koeln.de/dissertationfrey.pdf) 2005.
- <span id="page-57-16"></span>[Gru96] G. Grubb, Functional calculus of pseudodifferential boundary problems, 2nd ed., Birkhäuser, Boston–Basel–Berlin, 1996.
- <span id="page-57-17"></span>[Gru99] , Trace expansions for pseudodifferential boundary problems for Dirac–type operators and more general systems, Ark. Mat. 37 (1999), 45–86.
- <span id="page-57-5"></span>[Gru09]  $\_\_\_\_\_$ , Distributions and operators, Graduate Texts in Mathematics, vol. 252. Springer–Verlag, Berlin–Heidelberg–New York, 2009.
- <span id="page-57-6"></span>[HKL04] B. Himpel, P. Kirk, and M. Lesch, Calderón projector for the Hessian of the perturbed Chern–Simons function on a 3–manifold with boundary, Proc. London Math. Soc. 89 (2004), 241–272, [math.GT/0302234.](http://arxiv.org/abs/math/0302234)
- <span id="page-57-3"></span>[Hör85] L. Hörmander, The analysis of linear partial differential operators III, Grundlehren der Mathematischen Wissenschaften, vol. 274, Springer–Verlag, Berlin–Heidelberg– New York, 1985.
- <span id="page-57-26"></span>[KR97] R. V. Kadison and J. R. Ringrose, Fundamentals of the theory of operator algebras I, Graduate Studies in Mathematics, vol. 15, Amer. Math. Soc., Providence, RI, 1997.
- <span id="page-57-22"></span>[KL04] P. Kirk and M. Lesch, The  $\eta$ -invariant, Maslov index, and spectral flow for Diractype operators on manifolds with boundary, Forum Math. 16 (2004), 553–629, [math.DG/0012123.](http://arxiv.org/abs/math/0012123)
- <span id="page-57-25"></span>[Les05] M. Lesch, The uniqueness of the spectral flow on spaces of unbounded self–adjoint Fredholm operators, in: Spectral geometry of manifolds with boundary and decomposition of manifolds (B. Booß-Bavnbek, G. Grubb, and K.P. Wojciechowski, eds.), Contemp. Math., vol. 366, Amer. Math. Soc., 2005, pp. 193–224, [math.FA/0401411.](http://arxiv.org/abs/math/0401411)
- <span id="page-57-11"></span>[LM72] J.L. Lions and E. Magenes, Non-homogeneous boundary value problems and applications I, Grundlehren der Mathematischen Wissenschaften, vol. 181, Springer–Verlag, Berlin–Heidelberg–New York, 1972.
- <span id="page-57-0"></span>[Nic97] L. Nicolaescu, Generalized symplectic geometries and the index of families of elliptic problems, Mem. Amer. Math. Soc. 128 (1997), no. 609, 1–80.
- <span id="page-57-1"></span>[Pli61] A. Plis, A smooth linear elliptic differential equation without any solution in a sphere, Comm. Pure Appl. Math. 14 (1961), 599–617.
- <span id="page-57-9"></span>[Pon06] R. Ponge, Spectral asymmetry, zeta functions, and the noncommutative residue, Internat. J. Math. 17 (2006), 1065–1090, [math.DG/0310102.](http://arxiv.org/abs/math/0310102)
- <span id="page-57-10"></span>[Ral70] J. V. Ralston, *Deficiency indices of symmetric operators with elliptic boundary condi*tions, Comm. Pure Appl. Math. 23 (1970), 221–232.
- <span id="page-57-18"></span>[Sco95] S. G. Scott, Determinants of Dirac boundary value problems over odd-dimensional manifolds, Comm. Math. Phys. 173 (1995), 43–76.
- <span id="page-57-24"></span>[See66] R. T Seeley, Singular integrals and boundary value problems, Amer. J. Math. 88 (1966), 781–809.
- <span id="page-57-13"></span>[See67] , Complex powers of an elliptic operator, Proc. Sympos. Pure Math. 10 (1967), 288–307.
- <span id="page-57-4"></span>[See69] , Topics in pseudo–differential operators, in: C.I.M.E., Conference on pseudo– differential operators 1968, Edizioni Cremonese, Roma, 1969, pp. 169–305.
- <span id="page-57-7"></span>[Shu80] M. A. Shubin, Pseudodifferential operators and spectral theory, Springer–Verlag, Berlin–Heidelberg–New York, 1980.
- <span id="page-57-15"></span>[Tay96] M. E. Taylor, Partial differential equations I, Applied Mathematical Sciences, vol. 115, Springer–Verlag, Berlin–Heidelberg–New York, 1996.
- <span id="page-57-14"></span>[Woj85] K. P. Wojciechowski, Spectral flow and the general linear conjugation problem, Simon Stevin 59 (1985), 59–91.

# Index

<span id="page-58-0"></span>adjoint formal, 8 functional analytic, 9

basic data, 8 boundary condition, 10 Šapiro-Lopatinskii, 12 dual, 11 local elliptic, 12 regular, 10 strongly regular, 10

Calderón projection, 29 Seeley's classical, 29 Cauchy data space, 22 Cobordism Theorem, 31 complex interpolation theory, 19, 24 conic set, 13

data, see also basic data domain, see also operator dual, see also boundary condition

elliptic, see also operator

Fredholm Lagrangian Grassmannian, 36 Fredholm pair, 36, 39, 40

generalized eigenspace, 14 Green's formula, 8, 12, 21, 23, 27, 31, 38

idempotent, 15 unbounded, 15 isotropic, 34

Lagrangian, 33

operator Dirac, 6 classical pseudodifferential, 13 domain, 9 elliptic, 10 parameter dependent elliptic, 10, 13, 18 pseudodifferential with parameter, 12, 13 symbol, 13 sectorial, 15, 20 tangential, 12, 18 unbounded, 9 idempotent, 15 weakly sectorial, 15 orthogonalization, 15, 29, 31

Poisson operator, 29 pseudo–inverse, 27 pseudodifferential, see also operator

Riesz–map, 49

sectorial, see also operator, spectral projection Sobolev space, 9 spectral projection sectorial, 17 strong metric, 43

symbol classical, 13 leading, 9 parametric leading symbol, 13 symplectic Hilbert space, 33, 36 strongly, 33 weakly, 33

trace map, 10  $L^2$ -dual, 25

UCP, see also unique continuation property unique continuation property weak inner, 23

```
60 Notation
```
## Notation

<span id="page-59-0"></span>

Department of Science, Systems, and Models, Roskilde University, DK-4000 Roskilde, Denmark

 $E$ -mail address: booss@ruc.dk

URL: http://milne.ruc.dk/∼booss

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, ENDENICHER ALLEE 60, D-53115 BONN, Germany

E-mail address: ml@matthiaslesch.de, lesch@math.uni-bonn.de URL: http://www.matthiaslesch.de, [http://www.math.uni-bonn.de/](http://www.math.uni-bonn.de/~lesch)∼lesch

Chern Institute of Mathematics, Nankai University, Tianjin 300071, P. R. China  $\it E\mbox{-}mail\;address:$ zhucf@nankai.edu.cn