

Creation of neutron spinless pairs in the superfluid liquid ^4He .

Minasyan V.N. and Touryan K.J.

American University of Armenia, Yerevan, Armenia

June 21, 2024

Abstract

First, the creation of a free neutron spinless pairs is predicted in an inelastic neutron-scattering experiments, connected with investigation of the superfluid liquid ^4He . The novelty of the result presented is that an application of a new model for the superfluid liquid helium, which represents as an extension of the Bogoliubov model of the Bose gas with strong repulsive interaction between atoms. The new result of the theory presented herein, shows that the strong repulsive interaction between the Bogoliubov modes and the density modes of the neutrons vanishes by inducing of the attractive interaction mediated via the neutrons. This fact leads to a bound state on a spinless neutron pair which can exist at temperatures bellow than the critical temperature T_c of the Bose gas.

PACS 67.40. – *w*

1. Introduction.

The motivation for our theoretical study of the very low-temperature properties of the dilute hard sphere Bose gas is an attempt at a microscopic understanding of superfluidity in helium ^4He . We proceed by discussing some experimental and theoretical investigations.

The connection between the ideal Bose gas and superfluidity in helium was first made by London [1] in 1938. The ideal Bose gas undergoes a phase transition at sufficiently low temperatures to a condition in which the zero-momentum quantum state is occupied by a finite fraction of the atoms. This momentum-condensed phase was postulated by London to represent the superfluid component of liquid ^4He . With this hypothesis, the beginnings of a two- fluid hydrodynamic model of superfluids was developed by Landau [2] where he predicted the notation of a collective excitations so- called phonons and rotons.

The purely microscopic theory with mostly utilizes-technique, was first described by Bogoliubov [3] within the model of weakly non-ideal Bose-gas, with the inter-particle S- wave scattering. Based on the application of the presence of a macroscopic number of condensate atoms $N_0 \approx N$ (where N is the total number of atoms), Bogoliubov has dropped a density operator term which describes the atoms above the momentum -condensed phase, and in turn the dispersion curve for single particle Bogoliubov excitations was obtained.

The dispersion curve of an excitations excited in superfluid helium has been accurately measured a function from momentum [4]. Within this experiment, the position of a sharp peak inelastic neutron scattering intensity defines by energy of the single particle excitations, and there is appearing a broad component in inelastic neutron scattering intensity, at higher momenta of atoms. For explanation of the appearance of a broad component in inelastic neutron scattering intensity, the authors of papers [5-7] proposed to consider the presence of collective modes in the superfluid liquid ^4He , which are represented a density excitations. In this respect, the authors of this letter predicted that the collective modes represent as the density quasiparticles [8]. In these works, these density excitations and density quasiparticles are appeared due to remained density operator term for describing atoms above the condensate, which was neglected by Bogoliubov [3].

In this letter, we present a new model of a nonideal Bose gas for describing of the superfluid liquid helium. The given model is based on the application of the Penrose-Onsager definition of the Bose condensation [9] (where there is a basic condition as $\frac{N_0}{N} = \text{const}$), which in turn explains a broken of Bose-symmetry law for atoms of the superfluid liquid helium in the condensate, and leads to the creation of the Bogoliubov excitations.

Further, we investigate a helium gas-neutron gas mixture where exists

the repulsive strong interaction between atoms of superfluid helium as well as the repulsive strong interaction between atoms and neutrons. Due to the Penrose-Onsager definition of the Bose condensation by using of the method calculation proposed by Frölich [10] the term of interaction between the Bogoliubov modes and the density modes of the neutrons vanishes by inducing an effective attractive interaction between neutrons which determines a bound state on neutron pair.

2. New model of a nonideal Bose gas.

For beginning, we present a new model of the nonideal Bose gas with strong repulsive interaction between Bose-particles. The starting point our investigation is a main part of Hamiltonian of a system consisting of N identical atoms with mass m in box of volume V which is expressed in the second quantization form as:

$$\hat{H} = \sum_{\vec{p} \neq 0} \frac{p^2}{2m} \hat{a}_{\vec{p}}^+ \hat{a}_{\vec{p}} + \frac{1}{2V} \sum_{\vec{p} \neq 0} U_{\vec{p}} \hat{\varrho}_{\vec{p}} \hat{\varrho}_{\vec{p}}^+ \quad (1)$$

Here $\hat{a}_{\vec{p}}^+$ and $\hat{a}_{\vec{p}}$ are, respectively, the "creation" and "annihilation" operators of a free atoms with momentum \vec{p} ; $U_{\vec{p}}$ is the Fourier transform of a S-wave pseudopotential in the momentum space:

$$U_{\vec{p}} = \frac{4\pi d \hbar^2}{m} \quad (2)$$

where d is the scattering amplitude; and the Fourier component of the density operator presents as

$$\hat{\varrho}_{\vec{p}} = \sum_{\vec{p}_1} \hat{a}_{\vec{p}_1 - \vec{p}}^+ \hat{a}_{\vec{p}_1} \quad (3)$$

According to the Bogoliubov's theory [3], it is a necessary to separate the atoms in the condensate from those atoms filling states above the condensate. In this respect, the operators \hat{a}_0 and \hat{a}_0^+ are replaced by c-numbers $\hat{a}_0 = \hat{a}_0^+ = \sqrt{N_0}$ within approximation the presence of a macroscopic number of condensate atoms $N_0 \gg 1$. This assumption leads to a broken of Bose-symmetry law for atoms occupying in the condensate state. To be refused a broken from Bose-symmetry law for bosons in the condensate, we apply the Penrose-Onsager's definition of the Bose condensation [9]:

$$\lim_{N_0, N \rightarrow \infty} \frac{N_0}{N} = \text{const} \quad (4)$$

This reasoning is a very important factor for microscopic investigation of the model non-ideal Bose gas because the presence of a macroscopic number of atoms in the condensate means $N_0 \gg N_{\vec{p} \neq 0}$ (where $N_{\vec{p} \neq 0}$ is the occupation

number of atoms with momentum $\vec{p} \neq 0$), which also can be presented by a following approximation for an occupation number of atoms with momentum \vec{p} :

$$\lim_{N_0 \rightarrow \infty} \frac{N_{\vec{p}}}{N_0} = \delta_{\vec{p},0} \quad (5)$$

Now, using of a new method of perturbation theory presented in textbook of the Huang [10], we try to write down the matrix elements of interacting part included in the Hamiltonian of system (1) with inserting of a value $\hat{\rho}_{\vec{p}}$ in (3):

$$\begin{aligned} \langle m | \frac{1}{2V} \sum_{\vec{p}_1} \sum_{\vec{p}_2} \sum_{\vec{p} \neq 0} U_{\vec{p}} \hat{a}_{\vec{p}_1+\vec{p}}^+ \hat{a}_{\vec{p}_2-\vec{p}}^+ \hat{a}_{\vec{p}_1} \hat{a}_{\vec{p}_2} | n \rangle &= \\ = \frac{N_0}{2V} \sum_{\vec{p}_1} \sum_{\vec{p}_2} \sum_{\vec{p} \neq 0} \frac{U_{\vec{p}} \sqrt{(N_{\vec{p}_1+\vec{p}}+1)(N_{\vec{p}_2-\vec{p}}+1)N_{\vec{p}_1}N_{\vec{p}_2}}}{N_0} & \end{aligned} \quad (6)$$

where $|n\rangle$ and $|m\rangle$ are, respectively, the unperturbed wave functions, which satisfy to the condition $|n\rangle \neq |m\rangle$. Consequently,

$$|n\rangle = |\dots, N_{\vec{p}_1}, \dots, N_{\vec{p}_2}, \dots, N_{\vec{p}_1+\vec{p}}, \dots, N_{\vec{p}_2-\vec{p}}, \dots\rangle$$

$$|m\rangle = |\dots, N_{\vec{p}_1} - 1, \dots, N_{\vec{p}_2} - 1, \dots, N_{\vec{p}_1+\vec{p}} + 1, \dots, N_{\vec{p}_2-\vec{p}} + 1, \dots\rangle$$

Thus, we emphasize here that the approximation (5) presents as a main point for obtaining of an extension Bogoliubov model of the Bose gas by applications $\frac{N_{\vec{p}_1}}{N_0} = \delta_{\vec{p}_1,0}$; $\frac{N_{\vec{p}_2}}{N_0} = \delta_{\vec{p}_2,0}$; $\frac{N_{\vec{p}_1+\vec{p}}}{N_0} = \delta_{\vec{p}_1,-\vec{p}}$ and $\frac{N_{\vec{p}_2-\vec{p}}}{N_0} = \delta_{\vec{p}_2,\vec{p}}$ which lead to transformation of a matrix elements of interaction in (6)

$$\begin{aligned} \langle n | \frac{1}{2V} \sum_{\vec{p}_1} \sum_{\vec{p}_2} \sum_{\vec{p} \neq 0} U_{\vec{p}} \hat{a}_{\vec{p}_1+\vec{p}}^+ \hat{a}_{\vec{p}_2-\vec{p}}^+ \hat{a}_{\vec{p}_1} \hat{a}_{\vec{p}_2} | m \rangle &= \\ = \sum_{\vec{p} \neq 0} \frac{N_0 U_{\vec{p}}}{V} \left(2\sqrt{(N_{\vec{p}}+1)N_{-\vec{p}}} + \sqrt{(N_{\vec{p}}+1)(N_{-\vec{p}}+1)} + \sqrt{N_{\vec{p}}N_{-\vec{p}}} \right) & \end{aligned} \quad (7)$$

In this respect, the Bose operators $\hat{a}_{\vec{p}}^+$ and $\hat{a}_{\vec{p}}$ satisfy to a following expressions:

$$\hat{a}_{\vec{p}}^+ |n\rangle = \sqrt{N_{\vec{p}}+1} |m\rangle$$

and

$$\hat{a}_{\vec{p}} |n\rangle = \sqrt{N_{\vec{p}}} |m\rangle$$

Consequently, the Hamiltonian of system presented in (1) by support of (7) reproduces an extension form of the Bogoliubov Hamiltonian, at approximation $\frac{N_0}{N} = \text{const}$:

$$\hat{H} = \sum_{\vec{p} \neq 0} \frac{p^2}{2m} \hat{a}_{\vec{p}}^+ \hat{a}_{\vec{p}} + \sum_{\vec{p} \neq 0} m v^2 \left(2 \hat{a}_{\vec{p}}^+ \hat{a}_{\vec{p}} + \hat{a}_{-\vec{p}}^+ \hat{a}_{\vec{p}}^+ + \hat{a}_{\vec{p}} \hat{a}_{-\vec{p}} \right) \quad (8)$$

where $v = \sqrt{\frac{4\pi d \hbar^2 N_0}{m^2 V}}$ is the velocity of sound in the Bose gas which depends on the density atoms in the condensate.

In this context, the density operator $\hat{\varrho}_{\vec{p}}$ presents a form of the Bogoliubov theory [3]:

$$\hat{\varrho}_{\vec{p}} = \sqrt{N_0} \left(\hat{a}_{-\vec{p}}^+ + \hat{a}_{\vec{p}} \right) \quad (9)$$

Further, we show that a form of the density operator presented by (9) is an important factor for describing the repulsive strong interaction between atoms of superfluid helium as well as the repulsive strong interaction between atoms and neutrons, which are participating in an inelastic neutron-scattering experiments.

Now, we can prove that the approximation (5), within application of a new method of perturbation theory destroys the theories connected with presence of the density quasiparticles [8] as well as the density excitations [5-7] because the Bose-operators take new properties:

$$\lim_{N_0 \rightarrow \infty} \frac{\hat{a}_{\vec{p}}^+}{\sqrt{N_0}} = \lim_{N_0 \rightarrow \infty} \frac{\hat{a}_{\vec{p}}}{\sqrt{N_0}} = \delta_{\vec{p},0} \quad (10)$$

and

$$\lim_{N_0 \rightarrow \infty} \frac{\hat{a}_{\vec{p}_1 - \vec{p}}^+}{\sqrt{N_0}} = \lim_{N \rightarrow \infty} \frac{\hat{a}_{\vec{p}_1 - \vec{p}}}{\sqrt{N_0}} = \delta_{\vec{p}_1, \vec{p}} \quad (11)$$

As it is known from [8], the density operator $\hat{\varrho}_{\vec{p}}$ in (3) contains a description of the density-quasiparticles presented by authors of this letter:

$$\hat{\varrho}_{\vec{p}} = \sqrt{N_0} \left(\hat{a}_{-\vec{p}}^+ + \sqrt{2} \hat{c}_{\vec{p}} \right) \quad (12)$$

where $\hat{c}_{\vec{p}}$ is the Bose-operator "annihilation" of a free density-quasiparticles with momentum \vec{p} .

The application of (11) transforms the operator $\hat{c}_{\vec{p}}$ to a term of Bose-operator "annihilation" of a free atom:

$$\hat{c}_{\vec{p}} = \frac{1}{\sqrt{2N_0}} \sum_{\vec{p}_1 \neq 0} \hat{a}_{\vec{p}_1 - \vec{p}}^+ \hat{a}_{\vec{p}_1} = \frac{1}{\sqrt{2}} \sum_{\vec{p}_1 \neq 0} \delta_{\vec{p}_1, \vec{p}} \hat{a}_{\vec{p}_1} = \frac{1}{\sqrt{2}} \hat{a}_{\vec{p}}$$

which displays that the density operator $\hat{\varrho}_{\vec{p}}$ (3) coincides by a form presented (9). This fact implies that the density quasiparticles are absent.

The identical picture is observed in the case of the density excitations predicted by the Glyde, Griffin and Stirling [5-7], because

$$\hat{\varrho}_{\vec{p}} = \sqrt{N_0} \left(\hat{a}_{-\vec{p}}^+ + \hat{a}_{\vec{p}} + \tilde{\varrho}_{\vec{p}} \right) \quad (13)$$

where terms involving $\vec{p}_1 \neq 0$ and $\vec{p}_1 \neq \vec{p}$ are written separately; and the operator $\tilde{\varrho}_{\vec{p}}$ describes the density-excitations:

$$\tilde{\varrho}_{\vec{p}} = \frac{1}{\sqrt{N_0}} \sum_{\vec{p}_1 \neq 0, \vec{p}_1 \neq \vec{p}} \hat{a}_{\vec{p}_1 - \vec{p}}^+ \hat{a}_{\vec{p}_1} \quad (14)$$

At inserting of (10,11) into (14), the term, representing as the density-excitations, vanishes because $\tilde{\varrho}_{\vec{p}} = 0$. Reduction of the Bose-operators $\hat{d}_{\vec{p}}^+$ and $\hat{d}_{\vec{p}}$ by using of the Bogoliubov linear transformation [3]:

$$\hat{a}_{\vec{p}} = \frac{\hat{d}_{\vec{p}} + L_{\vec{p}} \hat{d}_{-\vec{p}}^+}{\sqrt{1 - L_{\vec{p}}^2}} \quad (15)$$

Substitution of (16) into (8) leads to

$$\hat{H} = \sum_{\vec{p}} \varepsilon_{\vec{p}} \hat{d}_{\vec{p}}^+ \hat{d}_{\vec{p}} \quad (16)$$

hence we infer that $\hat{d}_{\vec{p}}^+$ and $\hat{d}_{\vec{p}}$ are the "creation" and "annihilation" operators of a Bogoliubov's quasiparticles with energy:

$$\varepsilon_{\vec{p}} = \left[\left(\frac{p^2}{2m} \right)^2 + p^2 v^2 \right]^{1/2} \quad (17)$$

with

$$L_{\vec{p}}^2 = \frac{\frac{p^2}{2m} + mv^2 - \varepsilon_{\vec{p}}}{\frac{p^2}{2m} + mv^2 + \varepsilon_{\vec{p}}} \quad (18)$$

3. The effective attractive potential interaction between neutrons.

We attempt to describe the interaction of scattering neutrons with atoms of the superfluid liquid helium by using of the above-mentioned model of a Bose gas for the superfluid liquid helium. In this context, we consider a helium gas of atoms -neutron gas mixture where N atoms of helium interact with n neutrons via the repulsive neutron-atom S-wave scattering. To describe a helium gas-neutron gas mixture we apply the well-known Frölich

approach [11]. At solving of the problem superconductivity, the Frölich described a electron gas-phonon gas mixture where exist an interaction between electrons (fermions) and phonons (bosons) of lattice. Due to an application of the Frölich method we arrive to the Hamiltonian of a helium gas of atoms -neutron gas mixture which is expressed in the second quantization form:

$$\hat{H}_n = \sum_{\vec{p}} \varepsilon_{\vec{p}} \hat{d}_{\vec{p}}^+ \hat{d}_{\vec{p}} + \frac{1}{2V} \sum_{\vec{p} \neq 0} U_{\vec{p},n} \hat{\varrho}_{\vec{p}} \hat{\varrho}_{-\vec{p},n} + \sum_{\vec{p},\sigma} \frac{p^2}{2m_n} \hat{b}_{\vec{p},\sigma}^+ \hat{b}_{\vec{p},\sigma} \quad (19)$$

where $U_{\vec{p},n}$ is the Fourier transform of the attractive S-wave scattering between neutrons and atoms

$$U_{\vec{p},n} = \frac{4\pi d_n \hbar^2}{\mu} = U_0 \quad (20)$$

where d_n is the scattering amplitude between atom and neutron; $\mu = \frac{m \cdot m_n}{m + m_n}$; $\hat{b}_{\vec{p},\sigma}^+$ and $\hat{b}_{\vec{p},\sigma}$ are, respectively, the operators of creation and annihilation for free neutron with momentum \vec{p} , by the value of its spin z-component $\sigma = \pm \frac{1}{2}$. These operators satisfy to the Fermi commutation relations $[\cdot \cdot \cdot]_+$ as:

$$[\hat{b}_{\vec{p},\sigma}, \hat{b}_{\vec{p}',\sigma'}^+]_+ = \delta_{\vec{p},\vec{p}'} \cdot \delta_{\sigma,\sigma'} \quad (21)$$

$$[\hat{b}_{\vec{p},\sigma}, \hat{b}_{\vec{p}',\sigma'}]_+ = 0 \quad (22)$$

$$[\hat{b}_{\vec{p},\sigma}^+, \hat{b}_{\vec{p}',\sigma'}^+]_+ = 0 \quad (23)$$

The density operator of neutrons with spin σ in momentum \vec{p} is defined as

$$\hat{\varrho}_{\vec{p},n} = \sum_{\vec{p}_1,\sigma} \hat{b}_{\vec{p}_1-\vec{p},\sigma}^+ \hat{b}_{\vec{p}_1,\sigma} \quad (24)$$

where $\hat{\varrho}_{\vec{p},n}^+ = \hat{\varrho}_{-\vec{p},n}$

The operator of total number of neutrons is

$$\sum_{\vec{p},\sigma} \hat{b}_{\vec{p},\sigma}^+ \hat{b}_{\vec{p},\sigma} = \hat{n} \quad (25)$$

Hence, we note the operators describing of the Bogoliubov quasiparticles $\hat{d}_{\vec{p}}$ and neutrons $\hat{b}_{\vec{p},\sigma}$ is an independent.

The Hamiltonian \hat{H}_n (19) may describe a helium gas of Bogoliubov quasiparticles -neutron gas mixture because the density operator in the term of the Bogoliubov quasiparticles is expressed by following form, to application (15) into (9):

$$\hat{\varrho}_{\vec{p}} = \sqrt{N_0} \sqrt{\frac{1+L_{\vec{p}}}{1-L_{\vec{p}}}} \left(\hat{d}_{-\vec{p}}^+ + \hat{d}_{\vec{p}} \right) \quad (26)$$

Inserting (26) into (19) leads to reducing the Hamiltonian of system \hat{H}_n :

$$\hat{H}_n = \sum_{\vec{p}} \varepsilon_{\vec{p}} \hat{d}_{\vec{p}}^+ \hat{d}_{\vec{p}} + \frac{U_0 \sqrt{N_0}}{2V} \sum_{\vec{p}} \sqrt{\frac{1+L_{\vec{p}}}{1-L_{\vec{p}}}} \left(\hat{d}_{-\vec{p}}^+ + \hat{d}_{\vec{p}} \right) \hat{\varrho}_{-\vec{p},n} + \sum_{\vec{p},\sigma} \frac{p^2}{2m_n} \hat{b}_{\vec{p},\sigma}^+ \hat{b}_{\vec{p},\sigma} \quad (27)$$

To allocate anomalous terms in the Hamiltonian \hat{H}_n (27), we apply the Frölich approach [11] which allows to do the canonical transformation for an operator \hat{H}_n within introducing a new operator \tilde{H} :

$$\tilde{H} = \exp(\hat{S}^+) \hat{H}_n \exp(\hat{S}) \quad (28)$$

which decays

$$\tilde{H} = \exp(\hat{S}^+) \hat{H}_n \exp(\hat{S}) = \hat{H}_n - [\hat{S}, \hat{H}_n] + \frac{1}{2} [\hat{S}, [\hat{S}, \hat{H}_n]] - \dots \quad (29)$$

where the operators represent as:

$$\hat{S}^+ = \sum_{\vec{p}} \hat{S}_{\vec{p}}^+$$

and

$$\hat{S} = \sum_{\vec{p}} \hat{S}_{\vec{p}}$$

and satisfy to a condition $\hat{S}^+ = -\hat{S}$

In this respect, we assume that

$$\hat{S}_{\vec{p}} = A_{\vec{p}} \left(\hat{\varrho}_{\vec{p},n} \hat{d}_{\vec{p}} - \hat{\varrho}_{\vec{p},n}^+ \hat{d}_{\vec{p}}^+ \right) \quad (30)$$

where $A_{\vec{p}}$ is the unknown real symmetrical function from a momentum \vec{p} . To find $A_{\vec{p}}$, we substitute (30) and (27) into (29), and then:

$$[\hat{S}, \hat{H}_n] = -\frac{1}{V} \sum_{\vec{p}} A_{\vec{p}} U_0 \sqrt{N_0} \sqrt{\frac{1+L_{\vec{p}}}{1-L_{\vec{p}}}} \hat{\varrho}_{\vec{p}} \hat{\varrho}_{-\vec{p}} - \sum_{\vec{p}} A_{\vec{p}} \varepsilon_{\vec{p}} \hat{\varrho}_{\vec{p}} \left(\hat{d}_{-\vec{p}}^+ + \hat{d}_{\vec{p}} \right) \quad (31)$$

$$\frac{1}{2} [\hat{S}, [\hat{S}, \hat{H}_n]] = - \sum_{\vec{p}} A_{\vec{p}}^2 \varepsilon_{\vec{p}} \hat{\varrho}_{\vec{p},n} \hat{\varrho}_{-\vec{p},n} \quad (32)$$

and $[\hat{S}, [\hat{S}, [\hat{S}, \hat{H}_n]]] = 0$ within a Bose commutation relation $[\hat{\varrho}_{\vec{p}_1,n}, \hat{\varrho}_{\vec{p}_2,n}] = 0$; $[\hat{b}_{\vec{p}_1,\sigma}^+, \hat{b}_{\vec{p}_1,\sigma}, \hat{\varrho}_{\vec{p}_2,n}] = 0$.

Thus, the form of new operator \tilde{H} in (29) is reduced to a following form:

$$\begin{aligned}
\tilde{H} = & \sum_{\vec{p}} \varepsilon_{\vec{p}} \hat{d}_{\vec{p}}^+ \hat{d}_{\vec{p}} + \frac{1}{2V} \sum_{\vec{p}} U_0 \sqrt{N_0} \sqrt{\frac{1+L_{\vec{p}}}{1-L_{\vec{p}}}} \left(\hat{d}_{-\vec{p}}^+ + \hat{d}_{\vec{p}} \right) \hat{\varrho}_{-\vec{p},n} + \\
& + \sum_{\vec{p},\sigma} \frac{p^2}{2m_n} \hat{b}_{\vec{p},\sigma}^+ \hat{b}_{\vec{p},\sigma} + \frac{1}{V} \sum_{\vec{p}} A_{\vec{p}} U_0 \sqrt{N_0} \sqrt{\frac{1+L_{\vec{p}}}{1-L_{\vec{p}}}} \hat{\varrho}_{\vec{p},n} \hat{\varrho}_{-\vec{p},n} + \\
& + \sum_{\vec{p}} A_{\vec{p}} \varepsilon_{\vec{p}} \hat{\varrho}_{\vec{p},n} \left(\hat{d}_{-\vec{p}}^+ + \hat{d}_{\vec{p}} \right) - \sum_{\vec{p}} A_{\vec{p}}^2 \varepsilon_{\vec{p}} \hat{\varrho}_{\vec{p},n} \hat{\varrho}_{-\vec{p},n}
\end{aligned} \tag{33}$$

Removing a second and fifth terms in right part of (33) implies a transformation of interacting part between atoms and neutrons. In this respect, we obtain a quantity for $A_{\vec{p}}$:

$$A_{\vec{p}} = -\frac{U_0 \sqrt{N_0}}{2\varepsilon_{\vec{p}} V} \cdot \sqrt{\frac{1+L_{\vec{p}}}{1-L_{\vec{p}}}} \tag{34}$$

Then, the introducing (34) into a fourth and sixth terms in right side of (33), and making a some calculations in turn yields the new form of Hamiltonian system:

$$\tilde{H} = \sum_{\vec{p}} \varepsilon_{\vec{p}} \hat{d}_{\vec{p}}^+ \hat{d}_{\vec{p}} + \sum_{\vec{p},\sigma} \frac{p^2}{2m_n} \hat{b}_{\vec{p},\sigma}^+ \hat{b}_{\vec{p},\sigma} + \frac{1}{2V} \sum_{\vec{p}} V_{\vec{p}} \hat{\varrho}_{\vec{p},n} \hat{\varrho}_{-\vec{p},n} \tag{35}$$

where $V_{\vec{p}}$ is the effective potential of interaction between two neutrons in momentum space:

$$V_{\vec{p}} = 2A_{\vec{p}} U_0 \sqrt{N_0} \sqrt{\frac{1+L_{\vec{p}}}{1-L_{\vec{p}}}} - 2A_{\vec{p}}^2 \varepsilon_{\vec{p}} V = -\frac{3U_0^2 N_0 (1+L_{\vec{p}})}{2V \varepsilon_{\vec{p}} (1-L_{\vec{p}})} \tag{36}$$

The effective potential interaction between neutrons, presented in (36) is always attractive in momentum space because $V_{\vec{p}} \leq 0$, at $\varepsilon_{\vec{p}} \geq 0$ which in turn provides $\frac{1+L_{\vec{p}}}{1-L_{\vec{p}}} > 0$ with taking into account Eq.(18).

This attractive potential interaction (36) is not depending on the kinetic energy of the incoming single neutrons in a scattering event but depends on the density atoms in the condensate and the momentum of atoms as well as masses of an atom and neutron.

On other hand, as we show, this potential between two neutrons may define a bound state on neutron pair.

In this letter, we consider following cases: 1. At low momenta of atoms $p \ll 2mv$, the Bogoliunovs quasiparticles represent as phonons with energy $\varepsilon_{\vec{p}} \approx pv$ in (15) which leads to $L_{\vec{p}}^2 \approx \frac{1-\frac{p}{mv}}{1+\frac{p}{mv}} \approx \left(1-\frac{p}{mv}\right)^2$ in (18) or $L_{\vec{p}} \approx 1-\frac{p}{mv}$.

In this context, the attractive potential between two neutrons in momentum space, presented in (36), takes a following form:

$$V_{\vec{p}} \approx -\frac{3mU_0^2N_0}{Vp^2} \quad (37)$$

which can be rewritten by the term of so-called an effective charge e_* :

$$V_{\vec{p}} \approx -\frac{4\pi\hbar^2e_*^2}{p^2} \quad (38)$$

where

$$e_* = \frac{U_0}{2\hbar} \sqrt{\frac{3mN_0}{V\pi}}$$

2. At high momenta of atoms $p \gg 2mv$, we obtain $\varepsilon_{\vec{p}} \approx \frac{p^2}{2m} + mv^2$ in (15) which in turn defines $L_{\vec{p}} \approx 0$ in (18). These results present the attractive potential between two neutrons (36) by a form presented in (37).

Obviously, in both considering cases, as at small momenta of atoms as well as at high momenta of atoms, the effective potential between two neutrons is transformed in a coordinate space by following form:

$$V(\vec{r}) = \frac{1}{V} \sum_{\vec{p}} V_{\vec{p}} \cdot e^{i\vec{p}\vec{r}} = -\frac{e_*^2}{r} \quad (39)$$

Thus, we reach to the interaction between two neutrons in a coordinate space which has a similar form as the attractive Coulomb interaction between two charged particles with opposite charges. In analogy of the problem Hydrogen atom, the two neutrons with opposite spins, as two identical charged particles with opposite charges and spins, is bound as the spinless neutron pair with binding energy:

$$E_n = -\frac{m_n e_*^4}{4\hbar^2 l^2} = -\frac{1.4 \cdot 10^3 \hbar^2 d_n^4 N_0^2}{m_n n^2 V^2} \quad (40)$$

where n is the natural number which determines a bound state on a neutron pair.

Consequently, there is a spinless neutron pair with mass $m_0 = 2m_n$ which is created in the helium gas by the neutrons scattering on the atoms of the Bose gas. The bound state on a neutron pair is determined by the binding energy (40), which depends on the density of atoms in the condensate, and therefore, defines the state of temperatures $0 \leq T < T_c$ for existing of neutron pair. The later is absent, at temperatures $T \geq T_c$, because the density atoms in the condensate is a zero $\frac{N_0}{V} = 0$ which implies that $E_n = 0$ in (40), and therefore, the spinless neutron pair is decayed on two free neutrons.

References

1. F. London , Nature, **141**, 643 (1938)
2. L. Landau , J. Phys.(USSR), **5**, 77 (1941); Phys.(USSR), **11**, 91 (1947).
3. N.N. Bogoliubov , Jour. of Phys.(USSR), **11**, 23 (1947)
4. N.M. Blagoveshchenskii et al.,Phys. Rev. B **50**, 16550 (1994)
5. H.R. Glyde and A. Griffin ., Phys.Rev.Lett. **65**, 1454 (1990).
6. W.G. Stirling ,H.R. Glyde , Phys.Rev.B. **41**, 4224 (1990)
7. H.R. Glyde , Phys.Rev.B. **45**, 7321 (1992)
8. V.N. Minasyan and K.J. Touryan , Phys.Rev.Lett. **90**, 235301 (2003)
9. O. Penrose and L. Onsager , Phys. Rev., **104** , 576 (1956)
10. K. Huang , Statistical Mechanics, (Wiley 1963)
11. H. Frölich , Phys.Rev., **79**, 845, (1950); Proc.Roy. Soc, **215**, 291 (1952).