

Two-Level Chromophore and Irreversibility

Erich N. Wolf erich@uoregon.edu

Department of Chemistry, University of Oregon, Eugene, OR 97403, USA

Abstract:

We re-examine the traditional model of a two-level chromophore in a radiation field at resonant frequencies while undergoing stochastic collisions with a buffer gas, making explicit the distinction between reversible and irreversible work at the quantum level. An expectation value for the steady-state excited-state probability is constructed based on the microscopic behavior of a chromophore undergoing an infinite number of time-ordered, irreversible, state-changing transitions between the two (“pure”) quantum levels in the chromophore. We distinguish among three independent mechanisms for absorption and emission of photons: collision-induced state-changes, spontaneous state-changes (for both absorption and emission), and field-induced state-changes. A comparison to experimental data reveals the existence and nature of a change in the phase factor of the chromophore wavefunction for collision-induced state-changing events; this is interpreted as a parameterization of the impulsive nature of collision-induced state-changing events in the chromophore. “Hidden” in this parameterization are the details of the irreversible work done during state-changing photon absorption and emission events. The relationship between irreversibility in the quantum and macroscopic descriptions of the natural world then readily follows using the usual arguments of thermodynamics and statistical mechanics.

7.0 Introduction

This chapter will re-examine the traditional model of a two-level chromophore, making explicit the distinction between reversible and irreversible work at the quantum level. An expectation value for the steady-state excited-state probability is constructed based on the microscopic behavior of a chromophore undergoing an infinite number of time-ordered, irreversible state-changing transitions between the two (“pure”) quantum states in the chromophore. These two quantum levels are “connected” at a resonant frequency ω by an electromagnetic radiation field. Three (independent) mechanisms for absorption and emission of photons will be described. One mechanism accounts for state-changing “collisions” with a buffer gas and is referred to as a “collision-induced” state-change; a second mechanism alters slightly the traditional interpretation of spontaneous (“natural”) state-changes to include the ability to both absorb and emit photons in the presence of a radiation field turned on at a resonant frequency of the chromophore; the third mechanism recognizes that sufficiently “strong” radiation fields have the ability to perturb the electronic distribution of the chromophore so as to directly induce photon absorption and emission and is referred to as a “field-induced” state-change.

Along with a preliminary comparison to pressure-dependent experimental data (pressure broadening and pressure shift coefficients) an analysis of the two-level model presented in this chapter reveals the existence and nature of an ensemble-average change in the phase factor $\Delta\alpha$ of the chromophore wavefunction for collision-induced state-changing events; this

is interpreted as an expectation value that parameterizes the impulsive nature of collision-induced state-changing events in the chromophore. “Hidden” in this parameterization are the details of the irreversible work done during state-changing photon absorption or emission events. Similarly, an asymptotic-limit (approximate) value of the change in the phase factor $\Delta\alpha$ of the wavefunction for spontaneous absorption and emission events (such as have been extensively modeled with the mechanism of vacuum-state fluctuations) can be readily deduced. The relationship between irreversibility in the quantum and macroscopic descriptions of the natural world then readily follows using the usual arguments of thermodynamics and statistical mechanics.

7.1 The Hamiltonian

The derivation of the steady-state (frequency domain) line-shape for a two-level chromophore at the microscopic level begins, as usual, with a description of the Hamiltonian. The time-dependent Hamiltonian $\mathcal{H} = H + V$ for a two-level chromophore is first transformed to a projection representation. The pure quantum states of the “isolated” (unperturbed) chromophore are characterized in the usual manner as:

$$H |g\rangle = E_g |g\rangle; \quad H |e\rangle = E_e |e\rangle \quad (7.1)$$

Equation 7.1 indicates that the Hamiltonian operator H acting on the wavefunction (or state-function corresponding to a particular quantum-state) will return the energy (eigenvalue) E_g and E_e for the ground and excited states, $|g\rangle$ and $|e\rangle$, respectively.

The “closure” relation [1, 2] (or “completeness” relation, as in a complete set of functions; choosing which of these two words to use appears to change among well known teachers [3, 4]) applicable to the two-level model is given by:

$$|g\rangle\langle g| + |e\rangle\langle e| = 1 \quad (7.2)$$

Multiplying both sides of each of equations 7.1 by equation 7.2 (and recalling that $\langle e|e\rangle = \langle g|g\rangle = 1$) gives the isolated chromophore Hamiltonian as two “unconnected” (and thus isolated) stationary-states (levels):

$$H = H_g + H_e = E_g |g\rangle\langle g| + E_e |e\rangle\langle e| \quad (7.3)$$

The Hamiltonian also has what is assumed to be a small (in “strength”; see also section 5.1.1) time-dependent perturbation (interaction) term to account for collision-induced electronic state-changes of the chromophore. These interactions can be expressed as matrix elements in the form $V_{eg} = \langle g|V_m|e\rangle$ and $V_{ge} = \langle e|V_m|g\rangle$. However, our goal here is to convert the Hamiltonian to an operator representation that has the irreversible transition between “pure” electronic states given by $|e\rangle\langle g|$ and $|g\rangle\langle e|$. Furthermore, there is an implicit assumption that

the impulsive interactions between the chromophore and buffer gas are an integral part of the collision-induced state-changing process.

It can then be surmised – and this “analysis” will be described further below – that the irreversible state-changes of the chromophore due to collision-induced photon absorption and emission are given by:

$$V(t) = -\hbar \sum_{m=0}^{\infty} P_m \mu \cdot E(t) (|e\rangle \langle g| + |g\rangle \langle e|) \quad (7.4)$$

The radiation source is assumed to be monochromatic with a center frequency ω and field amplitude E_0 ; that is to say, $E(t) = E_0 \cos(\omega t)$. For the results presented in this thesis the finite frequency width of the radiation source is neglected on the assumptions that it is relatively narrow (compared to the Doppler broadened chromophore line width) and constant in shape and intensity for all frequencies ω . The transition (dipole) moment μ is assumed to be independent of the intra-molecular nuclear coordinates, which is a declaration of neglecting changes in the transition moment as a function of inter-atomic separation. The non-vanishing interaction of the chromophore transition moment and the electric field of the radiation source is implicit in writing $\mu \cdot E(t)$; that is to say, an observed transition is an allowed transition.

Equation 7.4 separates the state-changes associated with collision-induced absorption and emission of photons into independent contributions with definite (deterministic) endpoints characterized by the “pure” states $|e\rangle$ and $|g\rangle$. The probability P_m of the m^{th} collision-induced absorption (or collision-induced emission) state-change in the time interval t is given by the Poisson distribution:

$$P_m = \left(\frac{t}{2\tau_0} \right)^m \frac{1}{m!} \exp\left(-\frac{t}{2\tau_0} \right) \quad (7.5)$$

As equation 7.5 indicates, the average time interval between successive collision-induced absorption events is $2\tau_0$, and likewise for the time interval between successive collision-induced emission events. This corresponds to an average time interval τ_0 between collision-induced state-changes. It is necessary to indulge your patience here, as we (tentatively) recognize that: when the radiation source is initially turned-on at a particular frequency ω the absorption of photons is predominantly driven by pressure-*in*dependent mechanisms (spontaneous and/or field-induced absorption). It then follows that this initial ensemble of excited state chromophore is on average $\tau_0/2$ away from the next collision-induced state-changing interaction with the buffer gas. However, the average time interval between collision-induced photon absorption or emission events in equation 7.5 ignores any ensembles that were initially created (in the “neighborhood” of $t = 0$), which is consistent with the chromophore “forgetting” its initial conditions when making the passage from one (thermodynamic) equilibrium configuration with the nearly monochromatic radiation source

turned off to a different steady-state configuration with the radiation source turned on at a particular frequency ω for an essentially infinite period of time [5].

The use of a Poisson distribution to characterize the ensemble-average probability of collision-induced state-changing events of the chromophore as it traverses a path to the steady-state condition between absorption and emission of photons corresponds to the situation of these relatively brief state-changing events being well separated from each other in both time and space. These conditions can generally be satisfied for pressures of buffer gas below about 760 torr (approximately 1 atmosphere).

7.2 The Microscopic Steady-State Line-Shape

The steady-state signal obtained from a linear absorption experiment as a function of radiation frequency ω can be predicted from a first-order time-dependent perturbation analysis for the case of weak radiation fields (linear response). We begin in the Schrödinger picture with the Born-Oppenheimer approximation, which allows the chromophore wavefunction to be expressed as the product of the nuclear wavefunction $|\psi_g\rangle$ and the electronic ground-state wavefunction $|g\rangle$.

We choose the initial values of the phase α and time t : $\alpha = 0$ at $t = 0$. The chromophore wavefunction then begins as an “isolated” quantum system as:

$$\begin{aligned}
 |\Psi(t)\rangle &= \exp(-i \hbar^{-1} H t) \exp(i \alpha) |g\rangle |\psi_g(t)\rangle \\
 \Rightarrow |\Psi(0)\rangle &= |g\rangle |\psi_g(0)\rangle
 \end{aligned}
 \tag{7.6}$$

It is natural to separate a molecular wavefunction into the product of two contributions operating on different time scales (Born-Oppenheimer approximation), so that the nuclear wavefunction $|\psi_g\rangle$ becomes a convenient marker during the course of this derivation.

However, this model is expected to be applicable to both atoms and molecules in the role of chromophore and so it is worth mentioning at this point that the focus of this derivation will quickly shift to the time evolution of the phase factor α (during a state-changing interaction with the buffer gas), which will be given in the ensemble-average impulsive limit as $\Delta\alpha$.

This derivation is facilitated by a transformation to an interaction picture that allows us to “observe” the chromophore as it undergoes collision-induced state-changes. It is basically a method of isolating the perturbation term (equation 7.4) from the total Hamiltonian. The “rotating frame” provides a perspective that can “see” on either side of the projection operators $|e\rangle\langle g|$ and $|g\rangle\langle e|$, thus revealing the natural (as in “causality preserving”) division of the action of the operators in the Hamiltonian on the wavefunction during these rather abrupt, time-ordered, irreversible, collision-induced state-changing events. It also isolates (in time) the action on the two levels of the chromophore into independent contributions, which is at

least consistent with our notions of causality. This transformation in turn allows us to reduce the calculation of the steady-state excited-state expectation value to integrals that have analytic solutions.

The Schrödinger picture is related to the interaction picture by:

$$|\tilde{\Psi}(t)\rangle = \exp(i \hbar^{-1} H_I t) |\Psi(t)\rangle \quad (7.7)$$

The interaction picture Hamiltonian H_I will be defined further below (equation 7.12); we will first review the construction of the basic form of the time-ordered first-order perturbation solution in the interaction picture. The choice of initial conditions in equation 7.6 gives:

$$|\tilde{\Psi}(0)\rangle = |g\rangle |\psi_g(0)\rangle \quad (7.8)$$

We are choosing an interaction picture for which the Schrodinger equation takes the form:

$$\frac{d}{dt} |\tilde{\Psi}(t)\rangle = i \hbar^{-1} \tilde{V}(t) |\tilde{\Psi}(t)\rangle \quad (7.9)$$

The interaction picture Hamiltonian can be written in a general form as:

$$\tilde{V}(t) = \exp(i \hbar^{-1} H_I t) V(t) \exp(-i \hbar^{-1} H_I t) \quad (7.10)$$

We are seeking a perturbation solution across the long averaging time intervals of a frequency domain observation, which is basically an assumption that the measured signal (with the radiation field at a particular, “single” frequency) corresponds to a well determined steady-state average, often referred to as steady-state conditions. In the limit of low intensity radiation fields we can neglect multi-photon absorption events and seek a perturbation solution of equation 7.9 to first order in the time-dependent perturbation term in the interaction picture [6], which takes the form:

$$\begin{aligned}
 |\tilde{\Psi}(\infty)\rangle &\cong \left(1 - i\hbar^{-1} \int_0^\infty \tilde{V}(t) dt\right) |\tilde{\Psi}(0)\rangle \\
 &= \left(1 - i\hbar^{-1} \int_0^\infty \tilde{V}(t) dt\right) |g\rangle |\psi_g(0)\rangle
 \end{aligned}
 \tag{7.11}$$

The upper limit of integration in equation 7.11 is based on the assumption that the time interval over which the detector voltage is averaged is much, much longer than the mean time between time-ordered, irreversible, collision-induced state-changes of the chromophore, essentially an infinite period of time, as far as the chromophore is concerned, so that we are sure to have “established” steady-state conditions.

The interaction picture Hamiltonian H_I of equation 7.10 is chosen so as to be centered on the energy of the chromophore as it interacts m times with the buffer gas in successive collision-induced absorption (or collision-induced emission) events. The following definitions are useful in clarifying the meaning of the operator H_I :

$$\begin{aligned} H_f &= H_e + m P_f \\ H_i &= H_g + m P_i \\ H_I &= H_f + H_i \end{aligned} \tag{7.12}$$

Equation 7.12 is a statement of conservation of energy for the impulsive effect of the m^{th} irreversible state-changing collision of the chromophore. P_f and P_i in equation 7.12 are, respectively, the time-dependent interactions between the excited and ground electronic states with the buffer gas, conveniently separated by the “instantaneous” action of the projection operators $|e\rangle\langle g|$ and $|g\rangle\langle e|$ in the interaction picture. What might (at first) appear to be an unusual structure in equations 7.12, can be interpreted as a separation of the state-changing process of the chromophore into reversible and irreversible portions as it undergoes the “infinite” cycle of collision-induced absorption and emission. (This method of separating the work done during a state-changing event into reversible and irreversible components will be discussed further; please be patient, as there are several threads to weave together in this chapter.)

We are anticipating that most of the interaction between the chromophore and buffer gas occurs in a time interval that is relatively short compared to the mean time between

irreversible (electronic) state-changes. On the other hand, the “suddenness” [7] of an irreversible state-change of the chromophore during the collision event should also be emphasized. The transition moment in the chromophore (driven by the electric field of the laser at a wave length near 675 nm) will be oscillating with a period of about 2.5 fs. This time interval is roughly a factor of a thousand smaller than the duration of a state-changing collision that takes place across a distance of about 10 Å at a relative speed of approximately 400 m/s. It is thus reasonable to expect that there is considerable opportunity for a net amount of irreversible work to be done; this will be shown below to be due to the differential interactions of the two levels of the chromophore with the buffer gas. It may even follow, as we shall see, that the possibility of doing irreversible work is necessary for these state-changing events to be observable in the first place.

Application of what is often referred to as the rotating-wave approximation amounts to writing the cosine term in the radiation field as a sum of counter-propagating waves and then dropping the term that rotates in the opposite “direction” to the rotation transformation into the interaction picture; this is often used as justification for neglecting (what are often described as) rapidly oscillating and thus ineffectual terms in the integrands of equation 7.11 for one direction of rotation of the radiation field [8]. The trigonometric function $\cos(\omega t)$, however, merely describes rotation, which can progress in two indistinguishable directions when being expressed as an exponential functions containing an imaginary argument.

It follows then that application of the “rotating wave” description is not an approximation, but is an exact relationship:

$$E(t) = E_0 \cos(\omega t) = \left(\frac{E_0}{2}\right)(e^{-i \omega t} + e^{i \omega t}) \Rightarrow E = E_0 e^{-i \omega t} \quad (7.13)$$

The solution for the opposite direction of rotation is obtained by rotating into the interaction picture in the opposite direction in equation 7.7. These two counter propagating electromagnetic waves contribute equally to the steady-state excited-state expectation value, and invoking the idea of counter-propagating waves in the solutions to equation 7.9 does not appear to be necessary. Instead, when the solution obtained from considering one direction of rotation of the electric field (as was done in equation 7.7) is multiplied by two (as has been done in equation 7.13) the result is an exact solution, not an approximation. Furthermore, the idea of a “negative” frequency that appears in this exponential representation is artificial; that is to say, the real world energy transfer experienced by the chromophore is not altered by this mathematical formalism. This particular factor of two is not terribly significant in the results presented in this thesis, but it is important when measuring and/or calculating the absolute value of the transition moment.

The interaction picture Hamiltonian expanded to first-order (equation 7.10) can then be written as:

$$\begin{aligned} \tilde{V}(t) = & -\hbar \sum_{m=0}^{\infty} P_m \mu E_0 \exp\left(i \hbar^{-1} (H_e + m P_f) t\right) \\ & \times |e\rangle \langle g| \exp\left(-i \hbar^{-1} (H_g + m P_i) t\right) \exp(-i \omega t) + H.c. \end{aligned} \quad (7.14)$$

The Hermitian conjugate (*H.c.*), as usual, refers to the previous explicitly stated sum of terms in equation 7.14. The structure of equation 7.14 can also be interpreted as conforming to our notions of causality with regard to a time-ordering of the interaction between the two levels of the chromophore with the buffer gas: the non-vanishing terms, both before and after the electronic state-change, are on either side of the “transition projection” operator $|e\rangle\langle g|$ in equation 7.14. (That is to say, we can perhaps dispense with the notion that superposition states are real objects and take the more rational point of view, as Einstein was arguing all along, that causality will not be violated and the rest is merely illusion, much of it the result of a hidden variable that will soon be discovered in this derivation; symbolically we will designate this variable as $\Delta\alpha$.)

It will hopefully make the (time-ordered) algebra more transparent by reducing the action of a transition between states in the chromophore to the instantaneous action of a transition-operator \mathcal{T} [7] mediated by the probability of successive binary interactions between the chromophore and buffer gas that lead to collision-induced absorption events:

$$\begin{aligned} \mathcal{T} &= -\hbar\mu E_0 \int_0^\infty \left(\sum_{m=0}^\infty P_m \exp(i\hbar^{-1}(H_e - H_g)t - i\omega t) \right. \\ &\quad \left. \times \exp(i\hbar^{-1}m(P_f - P_i)t) \right) dt \\ &= -\hbar\mu E_0 \int_0^\infty \left(\sum_{m=0}^\infty P_m \exp(i\omega_0 t - i\omega t) \exp(i m \Delta\alpha) \right) dt \end{aligned} \quad (7.15)$$

The usual conversion between energy and frequency, namely $E = \hbar\omega$, has been applied between the second and third lines of equation 7.15 after accounting for the action of the operators. The unperturbed (“isolated” chromophore) line-center frequency is given by $\omega_0 = \hbar^{-1}(E_e - E_g)$.

The last line of equation 7.15 makes explicit the conversion of the irreversible component of a state-changing interaction between the chromophore and the buffer gas to an “instantaneous” impulsive process. Furthermore, The irreversible work done on the chromophore is cumulative (while the reversible work is not) so that the change in phase factor experienced by the chromophore in traversing m times to the upper level with the chromophore initially in the lower level is $m\Delta\alpha$. Similarly, but with a reversal of operator action for the Hermitian

conjugate of equation 7.15, the collision-induced emission events begin in the upper level and accounts for the change in phase factor $m\Delta\alpha$ associated with making m transitions to the lower level. From this perspective we recognize that $\Delta\alpha$ is the ensemble-average change in the phase factor of the chromophore wavefunction for collision-induced state-changes that are separated by the average time interval τ_0 . Since absorption and emission are related by complex conjugation, the sign of this quantity for collision-induced absorption is opposite (in “direction”) to that of collision-induced emission. The parameter $\Delta\alpha$ appears then to be an expectation value (associated with an eigenvalue) that characterizes the irreversible work done in an impulsive state-changing event for all resonant frequencies of the radiation field and all pressures of the buffer gas at a given temperature. And yet the (ensemble-average) irreversible work done on the chromophore during these state-changing events is opposite in sign for absorption and emission. The notion that the chromophore has a “memory” for the irreversible work and not the reversible work and the manner in which irreversible work “appears” at the macroscopic level will be further explored later in this chapter.

The integrals of equation 7.11 then take the form:

$$\int_0^{\infty} \tilde{V}(t) dt = \mathcal{T} |e\rangle \langle g| + \mathcal{T}^\dagger |g\rangle \langle e| \quad (7.16)$$

The ‘ \dagger ’ symbol (or perhaps ‘ \ddagger ’, depending on type-setting software) in equation 7.16 indicates the Hermitian conjugate of the object immediately to its left. It should also be

recognized that when identifying \mathcal{T} with collision-induced absorption the Hermitian conjugate term corresponds to the collision-induced emission events.

The excited-state (probability) amplitude obtained through collision-induced absorption can now be “projected” out of equation 7.11 through multiplication on the left by $\langle e|$ to obtain:

$$\begin{aligned}
 \langle e | \tilde{\Psi}(\infty) \rangle &\cong \langle e | \left(1 - i \hbar^{-1} \int_0^\infty \tilde{V}(t) dt \right) |g\rangle | \psi_g(0) \rangle \\
 &= \langle e | \left(1 - i \hbar^{-1} \left(\mathcal{T} |e\rangle \langle g| + \mathcal{T}^\dagger |g\rangle \langle e| \right) |g\rangle | \psi_g(0) \rangle \right) \\
 &= -i \hbar^{-1} \mathcal{T} | \psi_g(0) \rangle
 \end{aligned} \tag{7.17}$$

The expectation value $\langle e(\omega, t) \rangle$ of observing a given chromophore in the excited-state at steady-state conditions as a function of radiation frequency ω is given by the (scalar) product of the projected amplitudes due to collision-induced emission and collision-induced absorption:

$$\langle e(\omega, \infty) \rangle \cong \langle e | \tilde{\Psi}(\infty) \rangle \langle \tilde{\Psi}(\infty) | e \rangle = \hbar^{-2} (\mathcal{T}^\dagger \times \mathcal{T}) \tag{7.18}$$

The expectation value given by equation 7.18, often referred to as “resolving the superposition-state wavefunction”, contains hidden information about the nature of the irreversible work done during a collision-induced state-change. Since this point appears to be

fundamental to measurement theory [9], it is worth mentioning that later in this derivation we will reach the conclusion that in order to observe a steady-state ensemble-average value (detector signal) of equation 7.18 at the macroscopic level (in linear absorption spectra), it must be possible for irreversible work to be done during these (irreversible) state-changes.

To find a solution to equation 7.18 we first express the change in phase factor as a real and an imaginary component with the following definitions:

$$\exp(i m \Delta\alpha) = (A + i B)^m; \quad A = \cos(\Delta\alpha); \quad B = \sin(\Delta\alpha) \quad (7.19)$$

(The notation used in equation 7.19 is consistent with that used by Foley [10]. Please do not confuse the symbols A and B with the “Einstein A and B coefficients”.)

Performing the summation over all possible state-changes m of the chromophore (due to interactions with the buffer gas) that appears in the integrals in equation 7.15 gives:

$$\begin{aligned} & \sum_{m=0}^{\infty} P_m \exp(i m \Delta\alpha) \\ &= \sum_{m=0}^{\infty} \left(\frac{t}{2 \tau_0} \right)^m \frac{1}{m!} \exp\left(-\frac{t}{2 \tau_0}\right) (A + i B)^m \\ &= \exp\left(\frac{(A - 1 + i B) t}{2 \tau_0}\right) \end{aligned} \quad (7.20)$$

The solution of the transition integral \mathcal{T} (equation 7.15) is then given by:

$$\begin{aligned} \mathcal{T} &= \hbar \mu E_0 \int_0^\infty \exp\left(\frac{(A-1+iB)t}{2\tau_0}\right) \\ &\quad \times \exp(-\nu_n t) \exp(i\omega_0 t - i\omega t) dt \\ &= -\frac{\hbar \mu E_0}{\left(-\nu_n + \frac{A-1}{2\tau_0}\right) + i\left(\omega_0 - \omega + \frac{B}{2\tau_0}\right)} \end{aligned} \quad (7.21)$$

The ‘natural’ decay rate has been added in the usual ad-hoc (or heuristic) manner to equation 7.21 as a real (first-order) decay rate ν_n of the wavefunction amplitude. It is worth noting that when the natural decay rate is set equal to zero the solution to equation 7.21 strictly requires that A is less than one ($A < 1$).

According to equation 7.18 the expectation value for the steady-state excited-state probability $\langle e(\omega, t) \rangle$ corresponds to $t = \infty$ and is given by:

$$\begin{aligned} \langle e(\omega, \infty) \rangle &\cong \hbar^{-2} (\mathcal{T}^\dagger \times \mathcal{T}) \\ &= \frac{(\mu E_0)^2}{\left(\nu_n + \frac{1-A}{2\tau_0}\right)^2 + \left(\omega_0 - \omega + \frac{B}{2\tau_0}\right)^2} \end{aligned} \quad (7.22)$$

Equation 7.22 is the well known Lorentz (Cauchy) distribution so that normalization gives the “ideal” ensemble-average expectation value (probability) for the excited-state of a two-level chromophore (to first-order in the perturbation “expansion”):

$$\langle e(\omega, \infty) \rangle \cong \frac{(\mu E_0)^2 \left(\frac{2}{2\pi} \right) \left(\nu_n + \frac{1-A}{2\tau_0} \right)}{\left(\nu_n + \frac{1-A}{2\tau_0} \right)^2 + \left(\omega_0 - \omega + \frac{B}{2\tau_0} \right)^2} \quad (7.23)$$

In the limit of infinitely weak radiation fields the approximation inherent in equation 7.23 becomes a perfect equality; however, we expect that this approximation can be treated as a perfect equality when care is taken to ensure the radiation fields are sufficiently “weak”. It is perhaps worth noting that the first term in the denominator of equation 7.23 (containing A) is in units of events per time (usually referred to as cycles per time); while the other term (containing B) is in units of angular frequency, which is equal to “events per time” multiplied by 2π .

7.3 The Macroscopic Steady-State Linear Absorption Signal

Electronic detectors return a time-averaged (steady-state) voltage (signal) that is proportional to the intensity of the radiation field incident on these (macroscopic) devices. It is generally most convenient to arrange linear absorption experiments so that the radiation fields are well collimated beams. The total path length L of the absorbing medium and the chromophore

number density in a typical steady-state linear absorption experiment are chosen so that the strongest absorbing feature of interest does not absorb all of the incident radiation: $I/I_0 > 0$, where I_0 is the transmitted radiation intensity in the absence of an absorbing medium and I is the transmitted radiation intensity at frequency ω in the presence of an absorbing medium of fixed path length and at constant temperature and pressure. Furthermore, a working definition of what it means to be in the linear response region (of a perturbation analysis) with regard to changes in the radiation intensity can be realized experimentally by the observation that incident radiation intensities that differ by an order of magnitude (or preferably more) will give the same value for I/I_0 at the same strongly resonant frequency ω and all other conditions (temperature, pressure, and path length) held fixed. An analysis of the line-shape associated with the predicted excited-state probability distribution of equation 7.23 requires that both of the above conditions be satisfied.

The time-averaged intensity of a radiation field is proportional to the square of the electric field amplitude E_0 at a given location in space. Defining the radiation propagation direction as z , we first recognize that E_0^2 in equation 7.23 corresponds to the radiation field that interacts with a chromophore located in a thin slice in the region $z \rightarrow z + dz$. This recognition suggests that we separate this experimental parameter from the steady-state excited-state probability of equation 7.23 according to $\langle e(\omega, \infty) \rangle \Rightarrow (\mu E_0)^2 \langle e(\omega, \infty) \rangle$, and similarly for $\langle g(\omega, \infty) \rangle$.

A macroscopic steady-state linear absorption signal can then be characterized (as a proportion) by the differential change in beam intensity $I(z)$ passing through an optically thin slice of the total path length z as:

$$\frac{dI}{dz} \sim (N_e - N_g) = N(2\langle e(\omega, \infty) \rangle - 1)I \quad (7.24)$$

In constructing equation 7.24 the proportional relationship between the (time-averaged) intensity I of the radiation source (at a given distance along the total path length z) to the square of its amplitude E_0 (at that point) has been invoked; the probability amplitudes $\langle e(\omega, \infty) \rangle$ and $\langle g(\omega, \infty) \rangle$ obtained from equation 7.23 are thus reduced by this same factor $(\mu E_0)^2$, and we are assuming that there are no unaccounted for factors that have a functional dependence on frequency. Also, in keeping with the two-level chromophore model, two conservation equations have been used. The first is that the total probability (before reduction by the factor $(\mu E_0)^2$) is normalized: $\langle e(\omega, \infty) \rangle + \langle g(\omega, \infty) \rangle = 1$. The second is that the chromophore number density available in the gas cell for resonant interaction with the radiation field at frequency ω is the sum of the ground and excited electronic state number densities: $N = N_e + N_g = N(\langle e(\omega, \infty) \rangle + \langle g(\omega, \infty) \rangle)$.

Rearranging equation 7.24 and integrating gives the steady-state (macroscopic) absorption signal:

$$\begin{aligned}
 I(\omega) &\sim I_0 \exp(N (2 \langle e(\omega, \infty) \rangle - 1) L) \\
 &\sim I_0 \exp(2 N \langle e(\omega, \infty) \rangle L)
 \end{aligned}
 \tag{7.25}$$

The beam intensity of the radiation source after passing through an evacuated gas cell is given by I_0 in equation 7.25, and the second line of this equation assumes that the chromophore number density N and optical path length L are constant for a particular experimental configuration. A line-shape analysis must obviously also take into account the exponential relationship between the excited-state probability and the radiation intensity measurements (Beer-Lambert Law; see also section 5.1.1.). In the limit that $I/I_0 \approx 1$ this relationship can be taken as the linear limit of linear absorption (for sufficiently small x , $e^x \approx 1 + x$), which was done for the results presented throughout this thesis, since the direct (un-modulated) absorption signal at each frequency ω for an absorption feature had $I/I_0 \approx 0.97$.

The derivation in this section is merely a sketch. It is possible that other factors which are relatively slowly varying functions of frequency have been forgotten or neglected [11]. More important, though, is the question of connecting a macroscopic measurement (“observation”) to microscopic behavior. After all it can not be the same chromophore in a given thin slice of the absorbing medium that undergoes more than one let alone an infinite number of transitions between the two quantum states (levels) in the microscopic model leading to equation 7.23. A statistical mechanics formulation of this problem will (no doubt) invoke the

fact that from our perspective the particles in a macroscopic-size sample are indistinguishable. The indistinguishable nature of the quantum systems in macroscopic-size experiments and the randomness of the state-changing process suggest the need for a mechanism that allows the sample to “forget” its original thermodynamic state as it makes a transition to another one. As discussed near the end of section 7.1, this condition of “forgetting” has been satisfied in the derivation of the line-shape given in equation 7.23.

There is another perspective worth briefly exploring before moving on; one that has to do with the “memory” that macroscopic systems display. The “canonical distributions” of statistical mechanics [12] that are based on the Boltzmann (energy) factor $k_B T$ (and the underlying assumption that each chromophore in the macroscopic ensemble has an equal probability of being in either of the two levels consistent with a fixed “temperature”, where T is the temperature and k_B is the Boltzmann constant) do not allow for inversion of the steady-state populations; an equilibrium configuration that began without inversion of the populations in the macroscopic sample (i.e. $\langle g(\omega, \infty) \rangle > \langle e(\omega, \infty) \rangle$) can not end in another (properly time-averaged) macroscopic steady-state with the populations inverted, at least not without some additional assumptions (mechanisms). However, focusing only on the two-level chromophore model for the moment, it is the difference in steady-state populations of the two chromophore levels that provides the link between irreversible work at the microscopic (quantum) level and the appearance of a flow of heat between (finite-sized) macroscopic systems; the reason that heat must flow is because the macroscopic steady-state populations of the two levels (within the finite region of space defined by the overlap of the radiation field and the absorbing medium) can not be made equal. The “memory” that is

present in the two-level chromophore derivation, which accounted for all of the equal and opposite irreversible work done in absorption and emission (as expressed in equation 7.12) is thus in the “ideal” limit of no heat exchange with the outside world, which can only happen for a macroscopic system that is infinite in size or in some other manner could be made to be perfectly isolated from the rest of the universe.

And so it follows that the observation of deviations from this “ideal” Lorentz line shape (in a finite size experiment) may, at least in part, be an indication of heat flow between finite-sized macroscopic systems, one of which is “defined” by the overlap of the radiation fields and the absorbing medium. Of course, it is quite non-trivial to account for all of the experimental factors that contribute to the observed line shape. Furthermore, at longer wavelengths, there is the possibility of macroscopic effects of a quasi-static (slow response) nature that do not appear to have been accounted for in the traditional absorption-dispersion formulation [13]; this may be a source of asymmetric line shapes that have long been observed in pure rotation spectra.

7.4 Pressure Broadening and Pressure Shift Coefficients

De-convolution of the constant (inhomogeneous) Doppler width from an observed line-shape in a Doppler-limited high resolution absorption spectrum provides a measure of the Lorentz full width at half maximum height (FWHM). The change in Lorentz width as a function of pressure is representative of the changes in the collision-induced state-changing rate; all other

contributions to the homogeneous state-changing rate are assumed to be constant for a given molecular line. As well, the internally referenced spectrometers used for data collection in this project provided high quality measurements of the rate of change in the shift of the line-center from a nearly “isolated” line-center frequency ω_0 as a function of buffer gas pressure.

The relevant pieces of the “hard sphere” collision model (in the framework of the “ideal” gas model) necessary to relate the observed line-width and line-center shift to the buffer gas pressure and the state-changing cross-section are given by:

$$p = n k_B T ; \quad v = \sqrt{\frac{8 k_B T}{\pi \mu}} ; \quad \mu = \frac{m_1 m_2}{m_1 + m_2}$$

$$z = \frac{1}{\tau_0} = n \sigma v = \frac{p \sigma v}{k_B T} ; \quad \sigma = \pi d^2 \tag{7.26}$$

The parameters in equation 7.26 have the following definitions: p is the pressure; n is the number density of the buffer gas (which is assumed to be at least more than an order of magnitude larger than that of the total chromophore number density N); k_B is the Boltzmann constant; T is the temperature; z is the collision frequency; τ_0 is the mean time interval between state-changing interactions of the chromophore with the buffer gas; σ is the integrated (collision-induced) state-changing cross-section between the chromophore and buffer gas; d is the interaction distance associated with the integrated cross-section; v is the average relative speed between the chromophore and the buffer gas for thermal equilibrium conditions in a gas cell in the absence of the radiation source; μ is the reduced mass (not to be

confused with the use of this symbol as the transition moment beginning in equation 7.4), m_1 is the mass of a chromophore particle; and m_2 is the mass of a buffer gas particle.

In terms of the “hard sphere” collision model we can readily identify the full width at half maximum height ($\Delta\omega_L = 2\pi\Delta\nu_L$) and shift ($\Delta\omega_0 = 2\pi\Delta\nu_0$) in the Lorentz distribution of equation 7.23:

$$\begin{aligned}\Delta\omega_L &= 2\pi(\Delta\nu_L) = 2\left(\nu_n + \frac{(1-A)}{2\tau_0}\right) \\ &= 2\nu_n + \frac{2(1-A)}{2} \frac{p\sigma v}{k_B T}\end{aligned}\tag{7.27}$$

$$\Delta\omega_0 = 2\pi(\Delta\nu_0) = \frac{B}{2\tau_0} = \frac{B}{2} \frac{p\sigma v}{k_B T}\tag{7.28}$$

The pressure-dependent portion of the above two equations gives rise to the pressure broadening (B_p) and pressure shift (S_p) coefficients; these are, respectively, the rate at which the line broadens and the line-center shifts per unit change in pressure:

$$B_p \equiv \frac{\partial \Delta\nu_L}{\partial p} = \frac{(1 - A)}{2\pi} \frac{\sigma v}{k_B T} \quad (7.29)$$

$$S_p \equiv \frac{\partial \Delta\nu_0}{\partial p} = \frac{B}{4\pi} \frac{\sigma v}{k_B T} \quad (7.30)$$

The pressure broadening and pressure shift coefficients of equation 7.29 and 7.30 are obtained through an analysis of the de-convolved line-shape data. These two experimentally determined parameters can be rearranged to obtain values of A and B and thus σ and $\Delta\alpha$.

The ratio of the pressure broadening to pressure shift coefficients is given by:

$$R_{bs} \equiv \frac{B_p}{S_p} = \frac{2(1 - A)}{B} = \frac{2(1 - \cos(\Delta\alpha))}{\sin(\Delta\alpha)} \quad (7.31)$$

Equation 7.31 is plotted in figure 7.1.




Figure 7.1: Plot of the ratio of pressure broadening coefficient to pressure shift coefficient, R_{bs} , as a function of the ensemble-average change in phase factor, $\Delta\alpha$.

Solving equation 7.31 for the ensemble-average change in phase factor $\Delta\alpha$ gives:

$$\Delta\alpha = \cos^{-1}\left(\frac{4 \pm R_{bs}^2}{4 + R_{bs}^2}\right) \quad (7.32)$$

One solution of equation 7.31 is always $\Delta\alpha = 0$, which corresponds to $A = 1$ and $B = 0$, so that there is no line-width and no line-center shift associated with this solution; it is not observable in steady-state linear absorption spectroscopy. (It is not yet clear if this is an appropriate solution; technically speaking, it is often recognized that the phase of a complex number is undefined when the imaginary portion is zero ($B = 0$). Or it may be hiding a deeper, as yet unresolved meaning about the relationship of reversible and irreversible processes at the microscopic level and “observations” at the macroscopic level.) Consideration of equation

7.32 and figure 7.1 reveal that for all values of $\Delta\alpha$ the sign of R_{bs} follows that of $\sin(\Delta\alpha)$. All buffer gases used in this project produced a negative pressure shift coefficient, which results in $R_{bs} < 0$. This implies that $-180^\circ \leq \Delta\alpha < 0$ (see figure 7.1). The ensemble-average changes in the phase factor $\Delta\alpha$ of the wavefunction for diatomic iodine in the presence of noble gases (calculated from equation 7.32) are listed in table 7.1.

Table 7.1: Average phase factor change $\Delta\alpha$ of diatomic iodine in the presence of noble gases (at a temperature of 292 K and laser wavelength near 675 nm) based on the pressure broadening and pressure shift coefficients presented in chapter 6 (of this thesis) and equations 7.27 and 7.31. The statistical uncertainties of the values are given in parenthesis in units of the last significant figure of the quoted value.

Buffer Gas	B_p (MHz/torr)	S_p (MHz/torr)	R_{bs}	$\Delta\alpha$ (degrees)
He	8.34(31)	-0.201(50)	-41.5(104)	-174(1)
Ne	6.17(39)	-0.713(29)	-8.65(65)	-154(2)
Ar	7.70(41)	-1.32(11)	-5.83(58)	-142(4)
Kr	6.86(66)	-1.40(6)	-4.90(52)	-136(3)
Xe	6.47(44)	-1.71(10)	-3.78(34)	-124(4)

The ensemble-average expectation value of the change in phase factor $\Delta\alpha$ parameterizes the state-changing process at all frequencies (of a given resonance transition) and all pressures of buffer gas consistent with the use of a Poisson distribution in the impulsive limit of binary-collisions. However, this expectation value is changing sign between absorption and emission so that this can not be the source of the line-center shift. The origin of the observed line-center shift as a “mixing” of distinct line centers, much like mixing paint pigments to create a “perceived” (average) color, will be outlined in the next section.

It is also possible to make a qualitative comparison between the relative amounts of irreversible work done in the state-changing collisions with the buffer gases listed in table 7.1. The irreversible work can be proportionally approximated by dividing the change in phase $\Delta\alpha$ by an interaction time interval. Since, at constant temperature T , the relative speed between the chromophore and buffer gas increases with decreasing buffer gas mass, and thus the interaction time decreases, it follows (see table 7.1) that the ensemble-average irreversible work done during a collision-induced state-change increases as the mass of the buffer gas decreases.

The observed width (FWHM) of a line in steady-state frequency domain spectrum has traditionally been “shown” to have the form [14]:

$$\begin{aligned}\Delta\omega_L &= 2\pi(\Delta\nu_L) = 2\left(\nu_n + \frac{1}{\tau_0}\right) \\ &= 2\left(\nu_n + \frac{p\sigma v}{k_B T}\right)\end{aligned}\tag{7.33}$$

$\Delta\nu_L$ in equation 7.33 is the observed Lorentz component of the line-width (FWHM obtained from a line-shape analysis) in units of Hz. Except for a portion of the last figure of chapter 6, all of the results reported in previous chapters (of this thesis) are based on equation 7.33. In writing equation 7.33 we have used the property that Lorentz widths are additive in forming (through convolution) a composite line-shape. The traditional form of the Lorentz width corresponds to setting $A = -1$ and $B = 0$ in equation 7.23, so that $\Delta\alpha = \pm\pi$; it is not possible to

distinguish between the two possible signs of $\Delta\alpha$ in this case. Comparison of equations 7.27 and 7.33 reveals that the cross sections reported in previous chapters differ by a factor of $(1 - A) \div 2$ from the cross-section σ predicted by the two-level chromophore model presented in this chapter.

7.5 Analysis and Interpretation

Application of the traditional formulation of the postulates of (modern) quantum mechanics to equation 7.15 implies the existence of a Hermitian operator that “accounts for” (parameterizes) the irreversible work done between the chromophore and a buffer gas during an irreversible, collision-induced, impulsive state-changing interaction. This irreversible work arises due to the differential interaction between the two-levels of the chromophore with the buffer gas. That is to say, the impulsive action of the collision and the establishment of conditions necessary for resonant excitation of the chromophore by the radiation field are concerted events. The ensemble-average expectation value associated with this Hermitian operator is the change in the phase factor $\Delta\alpha$ of the chromophore wavefunction for an irreversible, collision-induced state-changing event between “pure” quantum states, the value of which can be extracted from the changes in the entire line shape (FWHM) as a function of pressure (equation 7.32).

According to the perturbation term (equation 7.4) the collision-induced absorption and collision-induced emission events are considered separately. Also, this equation explicitly

“says” that the chromophore is passing from one “pure” state to another; the formation of a superposition-state does not occur in this model. The perturbation analysis then investigates the probabilities associated with the change between these two “pure” states in the limit of an “infinite” averaging time interval (slow measurement). The choice of initial values for the parameters t and α (as was done in equation 7.6) leads to a steady-state limit in which it is not possible to distinguish (in a deterministic sense) which event came first – collision-induced absorption or collision-induced emission – or even when it occurred; in this sense the initial conditions have been “forgotten”. It is also evident from equations 7.16 and 7.18 that the action of the operators for these two events is in opposite “directions”. In the process of clearly specifying the direction of time (through the time-ordering property of the Poisson distribution) we arrive at an interpretation in which the sign of the change in phase factor $\Delta\alpha$ reverses between collision-induced absorption and collision-induced emission events, and from this “action-reversal” symmetry a steady-state excited-state probability distribution (as a function of radiation frequency ω) can be constructed that leads to an explicit expression for the ensemble-average change in the phase factor $\Delta\alpha$ for irreversible collision-induced state-changes.

The ensemble-average change in the phase factor $\Delta\alpha$ of the chromophore wavefunction for each interaction with a buffer gas (that leads to an irreversible state-change event) alters in a rather significant manner how we interpret pressure broadening and pressure shift data acquired from the analysis of pressure-dependent steady-state frequency domain spectra. The cross-sections reported in previous chapters do not take into account the impulsive nature of the interaction of a chromophore with a buffer gas during the time-dependent “collision” that

leads to a state-change in the chromophore. The monotonic increase of the change in phase shift $\Delta\alpha$ as state-changing collisions becomes “harder” (more elastic, which is to say there is less deformation of the charge distribution of the buffer gas during the collision) is readily seen in the progression of these values in table 7.1. This is the starting point in forming the following interpretation: collision-induced absorption and collision-induced emission of photons (which is equivalent to electronic state-changes of the chromophore in the two-level model) occur when the phase of the chromophore wavefunction is “suddenly” disrupted to some critical extent (impulsive limit), the nature of which is encoded in the (ensemble) average value of $\Delta\alpha$ for the collision-induced state-changing process. Collisions between these complex objects can be expected to include deformations of the charge distributions (polarization) and inelastic energy transfer. Further developments of the “ideal” two-level model presented in this chapter that make use of the experimental results presented in this thesis (pressure broadening and pressure shift coefficients of iodine), should no doubt also make use of the inelastic collision data for iodine that, in the words of the authors, “should be adequate in both quantity and accuracy” [15]. (Reference 15 is the first in a series of articles on (diatomic) iodine by J. Steinfeld that began during his graduate studies in The Klemperer Group.)

These experiments have also revealed that the line-center shift decreases monotonically as the state-changing collisions become “harder”. It is thus evident that in the limit of a non-polarizing state-changing collision, for which there is no deformation (polarization) of the charge distribution, the line-center shift will vanish, for which $A = -1$, $B = 0$, and $\Delta\alpha = \pm \pi$. It thus becomes reasonable to also “hypothesize” the existence of a pressure-*in*dependent

interaction, such as “vacuum-state fluctuations” [16], that gives rise to spontaneous absorption *and* spontaneous emission of photons by the chromophore and approximate the change in the phase factor of the wavefunction as $\pm \pi$, which is consistent with the manner in which equation 7.23 was derived. Similarly, the traditionally assumed form of equation 7.33 corresponds to this notion of a perfectly non-polarized state-changing collision between the chromophore and the buffer gas. However, in the limit that the “states” produced by these fluctuations are not identical to the “pure” states of the two-level chromophore model, it can be anticipated (perhaps as a hypothesis) that the ensemble-average change in phase factor for these events can only approach $|\pi|$ asymptotically from smaller absolute values. That is to say, perfectly non-polarized state-changes do not occur in nature.

The line-center shift of equation 7.23 can then be interpreted as a competition between the spontaneous and collision-induced rates of state-changing events corresponding to two idealized limits for the energy of the line-center. The two-level model presented in this chapter and the results presented in this thesis only explore the collision-induced line-center shift in the impulsive impact region of collision dynamics near the “isolated” chromophore line-center. According to gas-kinetic collision theory of “hard spheres” and equation 7.23 the impulsive region of collision dynamics is expected to result in a linear line-center shift [14]. Observation of the other line-center limit takes place in molecular beam experiments that study molecular (Van der Waals) complexes of the chromophore and buffer gas, which corresponds to the idealized case of a perfectly inelastic binary collision. As an example, it is well documented [17] that the transition energies of the argon-iodine (Ar-I₂) complex are shifted to smaller wave number values by several wave numbers (cm⁻¹) relative to the

“isolated” chromophore. (Keep in mind that this shift in energy is only a measure of the difference in energy between the X and B electronic states of the molecular complex and the isolated chromophore; it does not reveal unambiguously the nature of the shift of each of the levels in the chromophore.) What we “observe” on the macroscopic scale is then quite similar in fashion to mixing paint pigments; as the pressure of the buffer gas is increased the time-averaged “color” of the macroscopic sample shifts slightly further towards that of the bound state complex.

The results presented so far (in this chapter) bring us to a place where we may wish to consider more carefully the implications for the EPR paradox [18]. For example, is it true that a probability based theory, such as quantum mechanics, is incompatible with concepts of causality, locality, and determinism? The derivation leading to the expression for the expectation value of the steady-state excited-state probability (equation 7.23) did not require the use of a superposition-state wavefunction, but it does reveal the existence of the expectation value for the change in phase of the wave function $\Delta\alpha$ that parameterizes the irreversible component of work for these state-changing interactions at a given “temperature”. This parameter $\Delta\alpha$ essentially says that determining the “state” of a system can *only* be done by “disturbing the system”. This point of view can perhaps be extended to claim that superposition-states are not real objects, but are simply convenient constructs present in the theoretical foundations of quantum mechanics by way of the mathematical property of “linearity”.

This property of “linearity” appears to be at the heart of dynamical theories. In quantum mechanics it is used quite often to construct an ensemble-average total-wavefunction that is the “weighted” amplitude-sum of the wavefunctions, in this case the two “pure” states of the isolated chromophore. In the two-level model this is typically expressed as $|\Psi\rangle = a|g\rangle + b|e\rangle$ and the wavefunction probability-amplitudes “a” and “b” are in general complex numbers that satisfy the relation $|a|^2 + |b|^2 \equiv a^* \times a + b^* \times b = 1$; where “a*” and “b*” are the complex conjugates of “a” and “b”, and these amplitudes are in general functions of time. That is to say, the effects of interference that appear in many solutions to quantum mechanics problems are not a denial of “the law of causality”; rather they are a manifestation of the averaging processes necessary to construct a “description” of the microscopic world as viewed from the macroscopic level, perhaps better thought of as a probabilistic-determinism. The equations of quantum mechanics are, however, derived from the same conceptual framework that gave rise to Hamilton’s Principle [19], and so we might instead expect that there are no “hidden variables”, that quantum mechanics is a complete theory (at least as far as Gödel’s “Incompleteness Theorem” is concerned), and that the inability to violate causality should perhaps be considered a law of nature. It follows then that the difficulty in interpreting quantum mechanics may simply have been a matter of discovering all of the possible measurable (“observable”) parameters consistent with the “completeness” of quantum mechanics.

Part of the difficulty in discovering the “observable” change in phase factor $\Delta\alpha$ may be due to some considerable confusion in the spectroscopy community about how to properly use and interpret time-dependent perturbation theory (for which I am no exception, as this is

obviously a difficult problem that has for decades rendered this observable parameter as a “hidden variable”). As an example of the source of this confusion, it is often mistakenly assumed [20] that the solution to the steady-state excited-state expectation value for the two-level chromophore model obtained from using what are explicitly first-order perturbation terms [21] is valid for all radiation field amplitudes. This assumption is obviously inconsistent with what it means to construct a (Taylor) series expansion that accounts for the smoothly varying nature of the perturbation “strength” from the region of weak perturbations (linear) to strong perturbations (non-linear) [22]. And so it follows that such experiments are likely to be reaching the strong-field limit long before reaching the first maximum in the Rabi frequency. Accurate modeling of a line-shape in those types of spectroscopy experiments that make use of “strong” radiation fields can thus be expected to require the use of higher-order (non-linear) expansion terms in obtaining solutions to the two-level chromophore model.

Furthermore, in as much as we can expect molecular beam experiments to provide a source of “isolated”, collision-free chromophore, the state-changing mechanisms in studies like those performed of reference 20 will be dominated by spontaneous *and* field-induced photon absorption and emission. The “observed” spectrum of figure 4a in the molecular beam studies of reference 20 suggest that there are (at least) two independent mechanisms for photon absorption and emission that are not perfectly synchronized with each other. Also, in that same spectrum in reference 20, the slow and non-linear convergence to the asymptotic ensemble-average expectation value of equalized level-populations at the infinite-strength limit of the radiation field (i.e. $\langle g(\omega, t_R) \rangle = \langle e(\omega, t_R) \rangle = 1/2$; where t_R is the time spent in the resonance region in a molecular beam experiment) is certainly not consistent with describing

that spectrum by linear perturbation theory alone. It will require careful considerations, though, to construct a transition moment μ for linear and higher-order perturbation solution for the two-level chromophore model that accounts for field-induced absorption due to “strong” radiation fields. This is a project for the future and so I’ll only mention in passing that the form of the transition moment μ might more appropriately be decomposed into a term that is independent of the radiation field (accounting for spontaneous and collision-induced state-changes) and another term that does depend on the radiation field amplitude (accounting for field-induced absorption). And this field-induced dependence is such that for sufficiently small values of the radiation field amplitude the contribution due to field-induced absorption can be neglected, as was done in deriving equation 7.23. (And, of course, this is not to imply that there is no relationship between spontaneous and field-induced state-changes of the chromophore; we’ll simply leave these sorts of questions unanswered for now.)

Another useful example highlighting the “ideal” limits of the model presented in this chapter can be found in the polarization spectroscopy experiments performed in the steady-state saturation limit that “show” the magnitude of the interaction of the B electronic state of diatomic iodine with oxygen (as the buffer gas) to be a bit larger than that of the X electronic state [23]. In the “ideal” limit of the two-level chromophore model presented in this chapter (leading to equation 7.23) the interactions that give rise to these cross-sections are equal in magnitude (and opposite in sign). The B electronic state, however, offers additional “decay” paths, such as magnetically induced dissociation [24], which implies, at least theoretically, that the opportunity exists to create a steady-state inverted population; i.e. a population involving “pure” states (different from the original states) in which $\langle g(\omega, \infty) \rangle < \langle e(\omega, \infty) \rangle$.

It is well known that a population inversion can then be used to “create” a laser. In the context of this chapter, a laser will be active when the radiation field amplitude inside the laser cavity is sufficiently large to initiate field-induced absorption and emission of photons; and this process will be independent of the random processes that lead to decay. Thus, the actual ability to “detect” (observe) the emitted laser beam that leaks out of an “ideal” (lossless) laser cavity is the only limitation in this “ideal” two-level model; all that is necessary to produce laser radiation (aside from constructing the “ideal” laser cavity) is to begin with an initial steady-state inverted population. The mechanical precision of the laser cavity is the framework upon which the coherent state of the laser field is created; otherwise, it is quite difficult to fathom how the random absorption and emission events that are the result of collision-induced and/or spontaneous processes could result in a source of coherent radiation fields. Furthermore, the coherent steady-state created in a laser cavity can be associated with being an “attractor” of the initial inverted steady-state population; that is to say, coherence can act as an attractor for equilibrium conditions [5].

7.6 State-changing Rates and Collision Cross-Sections

In time-resolved experiments such as fluorescent decay and optical photon echo a short pulse of light is used to create a non-equilibrium state. The mechanism driving the absorption (and emission) of photons in a time interval much shorter than the time between collisions with a buffer gas is expected (based on the hypothesis discussed in the previous section) to be

dominated by field-induced and spontaneous absorption (and emission). The ensemble of excited-state atoms or molecules prepared in this manner is on average $\tau_0/2$ away from the next collision-induced state-changing collision with the buffer gas. The observed (ensemble-average) pressure-dependent decay rate is thus $2/\tau_0$ for the excited-state ensemble prepared in these experiments. The relationship of the observed spontaneous (“natural”) decay rate will depend on the intensity and duration of the pulse of radiation used to create the non-equilibrium state; it will be observed in these types of experiments as being between ν_n and $\nu_n/2$, where ν_n is the spontaneous (“natural”) decay rate.

The frequency domain experiments described in this thesis used “weak” and nearly monochromatic (and nearly collimated) radiation fields, thus making it possible to explore steady-state dynamics for which τ_0 is the average time between collision-induced state-changing events (typically present at a pressure such that $1/\tau_0$ is much greater than ν_n). The measurement (“observation”) process at each frequency ω of the radiation source is configured to average the signal for a time interval much longer than τ_0 , so that the initial non-equilibrium state (described above for time-resolved experiments) is completely “forgotten” and thus absent from the recorded signal. The independent mechanisms driving the cycle of absorbing and emitting photons for these experimental conditions (of relatively low intensity radiation fields) are spontaneous *and* collision-induced absorption and emission of photons. The observed transition rate (obtained from an analysis of the line-shape) is thus given by $1/\tau_0 + \nu_n$ for these experiments. (However, it is well known that the value of ν_n obtained from such experiments is in general not reliable.)

There is a long tradition of defining the non-equilibrium decay rate in time-resolved experiments as $1/T_1$ (the so called “transverse relaxation time”), which has perhaps obscured the relationship between cross-sections obtained from non-equilibrium decay experiments and steady-state absorption experiments. A careful comparison between the reported cross-sections for these two classes of experiments for the sodium D-lines [25, 26, 27] and (independently) for transitions between the X and B electronic states of diatomic iodine [28, 29] in the presence of noble gases reveals that the collision cross-sections obtained from time-resolved experiments are often “seen” to be roughly a factor of two larger than those from frequency domain experiments. Reports on the time-resolved experiments, however, typically consider the time to the next collision as τ_0 (i.e. T_1), instead of $\tau_0/2$, so that there is nothing inconsistent with the results between the two classes of experiments. Rather it is a matter of recognizing the difference in the definition of the ensemble-average time interval to the next collision-induced state-change for experiments that measure a non-equilibrium decay rate and those that measure a steady-state transition rate (that has been properly time-ordered by the Poisson distribution). The collision cross-sections presented in this thesis are based on the steady-state time interval τ_0 between the irreversible collision-induced state-changing events of the chromophore in the presence of a radiation source “turned-on” at frequency ω for an essentially infinite period of time.

7.7 Epilogue: Final Thoughts

An obvious path of exploration is to study the collision dynamics as a function of $\Delta\alpha$, which can be achieved, at least in part, by obtaining the pressure broadening and pressure shift coefficients over a wide range of temperatures [30]. However, it is hoped that the results presented in this chapter will provide insights for other researchers in their pursuit of comprehending and describing the natural world. It appears that whenever we clearly distinguish between illusion and reality in our “perceptions” of the real world, it often becomes possible to harness that knowledge in the pursuit of “advancing” science and technology.

As a child of about six years old it was the “understanding” presented by scholars in the physical sciences on the structure of the cosmos that grabbed my attention. At about eleven years old, I came to recognize that the fate of this era of human civilization depends a good deal on discovering “safe” methods of transforming energy from natural sources that are nearly infinite in supply (such as the sun) into forms suitable for use by humans. As the years passed and my psychological and political awareness grew, it became apparent that this was not enough:

“Many people have believed that science and technology could solve all our problems. Lately, however, we have witnessed a change in attitude. It has become clear that external progress alone cannot bring mental peace. People have begun to pay greater attention to inner science, the path of mental investigation and development. Through

our own experience we have arrived at a point where there is a new awareness of the importance and value of inner mental qualities. Therefore, the explanation of the mind and its working presented by the ancient scholars...are becoming increasingly valuable in our time. The strength of these traditions is related to developing mental peace. Science and technology are related to material progress. But a combination of these two can provide the complete conditions for obtaining real human happiness.” [31]

References:

1. C. Cohen-Tannoudji, B. Diu, F. Laloë; *Quantum Mechanics*, page 122, volume 1, John Wiley & Sons, Singapore (2005); ISBN 0-471-16433-X.
2. R. Loudon, *The Quantum Theory of Light*, 2nd Edition, p 153, Oxford University Press, New York, New York (1991); ISBN 0-19-851155-8.
3. R. Shankar; *Principles of Quantum Mechanics*, 2nd Edition, page 23, Plenum Press, New York, New York (1994); ISBN 0-306-44790-8.
4. A. Messiah; *Quantum Mechanics*, page 174, Dover Publications Inc., Mineola, New York (1999); ISBN 0-486-40924-4.
5. I. Prigogine; *From Being To Becoming: Time and Complexity in the Physical Sciences*, page 8, W. H. Freeman and Company, New York, New York (1980); ISBN 0-7167-1108-7.
6. S. Mukamel; *Nonlinear Optical Spectroscopy*, chapter 1, Oxford University Press, New York, New York (1995); ISBN 0-19-513291-2.
7. J. A. Cina; *Journal of Chemical Physics*, 113, 9488-9496 (2000).
8. L. Allen and J. H. Eberly; *Optical Resonance and Two-Level Atoms*, chapter 2, Dover Publications, Inc., Mineola, New York (1987); ISBN 0-486-65533-4.
9. Same as reference 5, but not limited to page 8. More specifically, I. Prigogine points to the work of B. d'Espagnat; *Conceptual Foundations of Quantum Mechanics*, 2nd Edition, Addison-Wesley Publishing Company, Redwood City, California (1989); ISBN 0-201-09388-X.
10. H. M. Foley; *Physical Review*, 69, 616-628 (1946).
11. J. D. Jackson; *Classical Electrodynamics*, 3rd Edition, chapter 7, John-Wiley & Sons, Inc., Hoboken, New Jersey (1999) ISBN 0-471-30932-X.
12. F. Reif, *Fundamentals of Statistical and Thermal Physics*, chapter 6, McGraw-Hill Book Company, New York, New York (1965); ISBN 07-051800-9.
13. C. H. Townes and A. L. Schawlow; *Microwave Spectroscopy*, chapter 13, Dover

Publications Inc., New York, New York (1975); ISBN 0-486-61798-X.

14. P.F. Bernath; *Spectra of Atoms and Molecules*, chapter 1, Oxford University Press, New York, New York (1995); ISBN 0-19-507598-6.
15. J. I. Steinfeld and W. Klemperer; *Journal of Chemical Physics*, 42, 3475-3497 (1965).
16. Page 619 of reference 1.
17. A. Buchachenko, N. Halberstadt, B. Lepetit, O. Roncero; *International Reviews in Physical Chemistry*, 22, 153-202 (2003).
18. A. Einstein, B. Podolsky, N. Rosen; *Physical Review*, 47, 777-780 (1935).
19. R. P. Feynman; *Physical Review*, 76, 769-789 (1949).
20. T. R. Dyke, G. R. Tomasevich, and W. Klemperer; *Journal of Chemical Physics*, 57, 2277-2284 (1972).
21. N. F. Ramsey; *Molecular Beams*, chapter V, Oxford University Press, London (1956).
22. E. Butkov; *Mathematical Physics*, chapter 15 Addison-Wesley Publishing Company, Reading, Massachusetts (1968); ISBN 0-201-00727-4.
23. J. C. D. Brand and R. J. Hayward; *Chemical Physics Letters*, 68, 369-373 (1979).
24. J. C. Lehmann; *Contemporary Physics*, 19, 449-468 (1978).
25. R. Kachru, T. W. Mossberg, and S. R. Hartmann; *Journal of Physics B; Atomic and Molecular Physics*, 13, L363-L368 (1980).
26. J. F. Kielkopf; *Journal of Physics B; Atomic and Molecular Physics*, 13, 3813-3821 (1980).
27. R. Walkup, A. Spielfiedel, D. Ely, and W. D. Phillips; *Journal of Physics B; Atomic and Molecular Physics*, 14, 1953-1961 (1981).
28. M. Comstock, V. V. Lozovoy, M. Dantus; *Journal of Chemical Physics*, 119, 6546-6553 (2003).
29. J. A. Eng, J. L. Hardwick, J. A. Raasch, E. N. Wolf; *Spectrochimica Acta Part A*, 60, 3413-3419 (2004).
30. For example, data obtained by J. L. Hardwick on pressure broadening and pressure shift of acetylene as a function of temperature (which is most likely still in press).
31. H. H. The Fourteenth Dalai Lama; *Sleeping, Dreaming, and Dying*, page 1, Wisdom Publications, Boston, Massachusetts (1997); ISBN 0-86171-123-8.