GENERALIZED NEWMAN PHENOMENA AND DIGIT CONJECTURES ON PRIMES

VLADIMIR SHEVELEV

ABSTRACT. We prove that the ratio of the Newman sum over numbers multiple of a fixed integer which is not multiple of 3 and the Newman sum over numbers multiple of a fixed integer divisible by 3 is o(1) when the upper limit of summing tends to infinity. We also discuss a connection of our results with a digit conjecture on primes.

1. INTRODUCTION

Denote for $x, m \in \mathbb{N}$

(1)
$$S_m(x) = \sum_{0 \le n < x, n \equiv 0 \pmod{m}} (-1)^{s(n)}$$

where s(n) is the number of 1's in the binary expansion of n. Sum (1) is a Newman digit sum.

From the fundamental paper of A.O.Gelfond [4] it follows that

(2)
$$S_m(x) = O(x^{\lambda}), \quad \lambda = \frac{\ln 3}{\ln 4}$$

The case m = 3 was studied in detail [5], [2], [7]. So, from the Coquet's theorem [2], [1] it follows that

(3)
$$-\frac{1}{3} + \frac{2}{\sqrt{3}}x^{\lambda} \le S_3(3x) \le \frac{1}{3} + \frac{55}{3}\left(\frac{3}{65}\right)^{\lambda}x^{\lambda}$$

with a microscopic improvement [7]

(4)
$$\frac{2}{\sqrt{3}}x^{\lambda} \le S_3(3x) \le \frac{55}{3} \left(\frac{3}{65}\right)^{\lambda} x^{\lambda}, \quad x \ge 2,$$

and moreover,

¹⁹⁹¹ Mathematics Subject Classification. 11A63; Key words and phrases: Gelfond digit theorem, Coquet's theorem, Newman phenomena, Newman sums over multiples of

(5)
$$\left\lfloor 2\left(\frac{x}{6}\right)^{\lambda} \right\rfloor \le S_3(x) \le \left\lceil \frac{55}{3}\left(\frac{x}{65}\right)^{\lambda} \right\rceil$$

These estimates give the most exact modern limits of the so-called Newman phenomena. Note that M.Drmota and M.Skalba [3] using a close function $(S_m^{(m)}(x))$ proved that if m is a multiple of 3 then for sufficiently large x,

(6)
$$S_m(x) > 0, \quad x \ge x_0(m).$$

In this paper we study a general case for $m \ge 5$ (in the cases of m = 2and m = 4 we have $|S_m(n)| \le 1$).

To formulate our results put for $m \geq 5$

(7)
$$\lambda_m = 1 + \log_2 b_m,$$

(8)
$$\mu_m = \frac{2b_m + 1}{2b_m - 1},$$

where

(9)
$$b_m^2 = \begin{cases} \sin(\frac{\pi}{3}(1+\frac{3}{m}))(\sqrt{3} - \sin(\frac{\pi}{3}(1+\frac{3}{m}))), & if \ m \equiv 0 \pmod{3} \\ \sin(\frac{\pi}{3}(1-\frac{1}{m}))(\sqrt{3} - \sin(\frac{\pi}{3}(1-\frac{1}{m}))), & if \ m \equiv 1 \pmod{3} \\ \sin(\frac{\pi}{3}(1+\frac{1}{m}))(\sqrt{3} - \sin(\frac{\pi}{3}(1+\frac{1}{m}))), & if \ m \equiv 2 \pmod{3}. \end{cases}$$

Directly one can see that

(10)
$$\frac{\sqrt{3}}{2} > b_m \ge \begin{cases} 0.86184088\dots, & if \ (m,3) = 1, \\ 0.85559967\dots, & if \ (m,3) = 3, \end{cases}$$

and thus,

(11)
$$\lambda_m < \lambda$$

and

(12)
$$3.73205080\ldots < \mu_m \le \begin{cases} 3.76364572\ldots, & if (m,3) = 1, \\ 3.81215109\ldots, & if (m,3) = 3. \end{cases}$$

Below we prove the following results.

Theorem 1. If (m, 3) = 1 then

$$|S_m(x)| \le \mu_m x^{\lambda_m}.$$

Theorem 2. (Generalized Newman phenomena). If m > 3 is a multiple of 3 then

(14)
$$\left|S_m(x) - \frac{3}{m}S_3(x)\right| \le \mu_m x^{\lambda_m}.$$

Using Theorem 2 and (5) one can estimate $x_0(m)$ in (6). E.g., one can prove that $x_0(21) < e^{985}$.

2. Explicit formula for $S_m(N)$

We have

(15)
$$S_m(N) = \sum_{n=0,m|n}^{N-1} (-1)^{s(n)} = \frac{1}{m} \sum_{t=0}^{m-1} \sum_{n=0}^{N-1} (-1)^{s(n)} e^{2\pi i (\frac{nt}{m})} =$$
$$= \frac{1}{m} \sum_{t=0}^{m-1} \sum_{n=0}^{N-1} e^{2\pi i (\frac{t}{m}n + \frac{1}{2}s(n))}.$$

Note that the interior sum has the form

(16)
$$F_{\alpha}(N) = \sum_{n=0}^{N-1} e^{2\pi i (\alpha n + \frac{1}{2}s(n))} \quad 0 \le \alpha < 1.$$

Lemma 1. If $N = 2^{\nu_0} + 2^{\nu_1} + \ldots + 2^{\nu_r}$, $\nu_0 > \nu_1 > \ldots > \nu_r \ge 0$, then

(17)
$$F_{\alpha}(N) = \sum_{h=0}^{r} e^{2\pi i (\alpha \sum_{j=0}^{h-1} 2^{\nu_j} + \frac{h}{2})} \prod_{k=0}^{\nu_h - 1} (1 + e^{2\pi i (\alpha 2^k + \frac{1}{2})}),$$

where as usual $\sum_{j=0}^{-1} = 0$, $\prod_{k=0}^{-1} = 1$.

Proof. Let r = 0. Then by (16)

(18)
$$F_{\alpha}(N) = \sum_{n=0}^{N-1} (-1)^{s(n)} e^{2\pi i \alpha n} = 1 - \sum_{j=0}^{\nu_0 - 1} e^{2\pi i \alpha 2^j} + \sum_{0 \le j_1 < j_2 \le \nu_0 - 1} e^{2\pi i \alpha (2^{j_1} + 2^{j_2})} - \dots = \prod_{k=0}^{\nu_0 - 1} (1 - e^{2\pi i \alpha 2^k}),$$

which corresponds to (17) for r = 0.

Assuming that (17) is valid for every N with s(N) = r + 1 let us consider $N_1 = 2^{\nu_r}a + 2^{\nu_{r+1}}$ where a is odd, s(a) = r + 1 and $\nu_{r+1} < \nu_r$. Let

$$N = 2^{\nu_r} a = 2^{\nu_0} + \ldots + 2^{\nu_r}; \quad N_1 = 2^{\nu_0} + \ldots + 2^{\nu_r} + 2^{\nu_{r+1}}.$$

Notice that for $n \in [0, 2^{\nu_{r+1}})$ we have

$$s(N+n) = s(N) + s(n).$$

Therefore,

$$F_{\alpha}(N_{1}) = F_{\alpha}(N) + \sum_{n=N}^{N_{1}-1} e^{2\pi i (\alpha n + \frac{1}{2}s(n))} =$$
$$= F_{\alpha}(N) + \sum_{n=0}^{2^{\nu_{r+1}}-1} e^{2\pi i (\alpha n + \alpha N + \frac{1}{2}(s(N) + s(n)))} =$$
$$= F_{\alpha}(N) + e^{2\pi i (\alpha N + \frac{1}{2}s(N))} \sum_{n=0}^{2^{\nu_{r+1}}-1} e^{2\pi i (\alpha n + \frac{1}{2}s(n))}.$$

Thus, by (17) and (18),

$$F_{\alpha}(N_{1}) = \sum_{h=0}^{r} e^{2\pi i (\alpha \sum_{j=0}^{h-1} 2^{\nu_{j}} + \frac{h}{2})} \prod_{k=0}^{\nu_{h}-1} (1 + e^{2\pi i (\alpha 2^{k} + \frac{1}{2})} + e^{2\pi i (\alpha \sum_{j=0}^{r} 2^{\nu_{j}} + \frac{r+1}{2})} \prod_{k=0}^{\nu_{r+1}-1} \left(1 + e^{2\pi i (\alpha 2^{k} + \frac{1}{2})}\right) = \sum_{h=0}^{r+1} e^{2\pi i (\alpha \sum_{j=0}^{h-1} 2^{\nu_{j}} + \frac{h}{2})} \prod_{k=0}^{\nu_{h}-1} \left(1 + e^{2\pi i (\alpha 2^{k} + \frac{1}{2})}\right) .$$

Formulas (15)-(17) give an explicit expression for $S_m(N)$ as a linear combination of the products of the form

(19)
$$\prod_{k=0}^{\nu_h - 1} \left(1 + e^{2\pi i (\alpha 2^k + \frac{1}{2})} \right), \quad \alpha = \frac{t}{m}, \quad 0 \le t \le m - 1.$$

Remark 1. On can extract (17) from a very complicated general Gelfond formula [4], however, we prefer to give an independent proof.

3. Proof of Theorem 1

Note that in (17)

(20)
$$r \le \nu_0 = \left\lfloor \frac{\ln N}{\ln 2} \right\rfloor.$$

By Lemma 1 we have

(21)
$$|F_{\alpha}(N)| \leq \sum_{\nu_{h}=\nu_{0},\nu_{1},...,\nu_{r}} \left| \prod_{k=1}^{\nu_{h}} \left(1 + e^{2\pi i (\alpha 2^{k-1} + \frac{1}{2})} \right) \right| \leq \sum_{h=0}^{\nu_{0}} \left| \prod_{k=1}^{h} \left(1 + e^{2\pi i (\alpha 2^{k-1} + \frac{1}{2})} \right) \right|.$$

Furthermore,

$$1 + e^{2\pi i (2^{k-1}\alpha + \frac{1}{2})} = 2\sin(2^{k-1}\alpha\pi)(\sin(2^{k-1}\alpha\pi) - i\cos(2^{k-1}\alpha\pi))$$

and, therefore,

(22)
$$\left|1 + e^{2\pi i (2^{k-1}\alpha + \frac{1}{2})}\right| \le 2 \left|\sin(2^{k-1}\alpha \pi)\right|.$$

According to (21) let us estimate the product

(23)
$$\prod_{k=1}^{h} (2|\sin(2^{k-1}\alpha\pi)|) = 2^{h} \prod_{k=1}^{h} |\sin(2^{k-1}\alpha\pi)|,$$

where by (15)

(24)
$$\alpha = \frac{t}{m}, \quad 0 \le t \le m - 1.$$

Repeating arguments of [4], put

(25)
$$\left|\sin(2^{k-1}\alpha\pi)\right| = t_k.$$

Considering the function

(26)
$$\rho(x) = 2x\sqrt{1-x^2}, \ 0 \le x \le 1,$$

we have

(27)
$$t_k = 2t_{k-1}\sqrt{1 - t_{k-1}^2} = \rho(t_{k-1}).$$

Note that

(28)
$$\rho'(x) = 2(\sqrt{1-x^2} - \frac{x^2}{\sqrt{1-x^2}}) \le -1$$

for $x_0 \leq x \leq 1$, where

$$(29) x_0 = \frac{\sqrt{3}}{2}$$

is the only positive root of the equation $\rho(x) = x$. Show that either

(30)
$$t_k \le \sin\left(\frac{\pi}{m} \left\lfloor \frac{m}{3} \right\rfloor\right) = \sin\left(\frac{\pi}{m} \left\lceil \frac{2m}{3} \right\rceil\right) = g_m < \frac{\sqrt{3}}{2}$$

or simultaneously $t_k > g_m$ and

$$t_k t_{k+1} \le \max_{0 \le l \le m-1} \left(\left| \sin \frac{l\pi}{m} \right| \left(\sqrt{3} - \left| \sin \frac{l\pi}{m} \right| \right) \right) =$$

$$(31) = \begin{cases} \left(\sin\left(\frac{\pi}{m} \lfloor \frac{m}{3} \rfloor\right)\right) \left(\sqrt{3} - \sin\left(\frac{\pi}{m} \lfloor \frac{m}{3} \rfloor\right)\right), & if \ m \equiv 1 \pmod{3} \\ \left(\sin\left(\frac{\pi}{m} \lceil \frac{m}{3} \rceil\right)\right) \left(\sqrt{3} - \sin\left(\frac{\pi}{m} \lceil \frac{m}{3} \rceil\right)\right), & if \ m \equiv 2 \pmod{3} \end{cases} = h_m < \frac{3}{4}.$$

Indeed, let for a fixed values of $t \in [0, m-1]$ and $k \in [1, n]$

(32)
$$t2^{k-1} \equiv l \pmod{m}, \ 0 \le l \le m-1.$$

Then

(33)
$$t_k = \left| \sin \frac{l\pi}{m} \right|.$$

Now distinguish two cases: 1) $t_k \leq \frac{\sqrt{3}}{2}$ 2) $t_k > \frac{\sqrt{3}}{2}$. In case 1)

$$t_k = \frac{\sqrt{3}}{2} \leftrightarrows \frac{l\pi}{m} = \frac{r\pi}{3}, \ (r,3) = 1$$

and since $0 \le l \le m - 1$ then

$$m = \frac{3l}{r}, \quad r = 1, 2.$$

Because of the condition (m, 3) = 1, we have $t_k < \frac{\sqrt{3}}{2}$. Thus, in (33)

$$l \in \left[0, \left\lfloor \frac{m}{3} \right\rfloor\right] \cup \left[\left\lceil \frac{2m}{3} \right\rceil, m\right]$$

and (30) follows.

In case 2) let $t_k > \frac{\sqrt{3}}{2} = x_0$. For $\varepsilon > 0$ put

(34)
$$1 + \varepsilon = \frac{t_k}{x_0} = \frac{2}{\sqrt{3}} \left| \sin(\pi 2^{k-1} \alpha) \right|,$$

such that

$$1 - \varepsilon = 2 - \frac{2}{\sqrt{3}} \left| \sin(\pi 2^{k-1} \alpha) \right|$$

and

(35)
$$1 - \varepsilon^2 = \frac{4}{3} \left| \sin(\pi 2^{k-1} \alpha) \right| \left(\sqrt{3} - \left| \sin(\pi 2^{k-1} \alpha) \right| \right).$$

By (27) and (34) we have

$$t_{k+1} = \rho(t_k) = \rho((1+\varepsilon)x_0) = \rho(x_0) + \varepsilon x_0 \rho'(c),$$

where $c \in (x_0, (1 + \varepsilon)x_0)$.

Thus, according to (28) and taking into account that $\rho(x_0) = x_0$, we find

$$t_{k+1} \le x_0(1+\varepsilon)$$

while by (34)

$$t_k = x_0(1+\varepsilon).$$

Now in view of (35) and (29)

$$t_k t_{k+1} \le \left| \sin \pi 2^{k-1} \alpha \right| \left(\sqrt{3} - \left| \sin(\pi 2^{k-1} \alpha) \right| \right)$$

and according to (32),(33) we obtain that

$$t_k t_{k+1} \le h_m,$$

where h_m is defined by (31).

Notice that from simple arguments and according to (9)

$$g_m \le \sqrt{h_m} = b_m$$

Therefore,

$$\prod_{k=1}^{h} \left| \sin(\pi 2^{k-1} \alpha) \right| \le (b_m^{\lfloor \frac{h}{2} \rfloor})^2 \le b_m^{h-1}.$$

 $\prod_{k=1} \left| \sin(\pi 2^{k-1} \alpha) \right| \le (b_m^{\lfloor \frac{h}{2} \rfloor})^2 \le b_m^{h-1}.$ Now, by (21)- (22), for $\alpha = \frac{t}{m}, \ t = 0, 1, ..., m-1$, we have

$$\left| F_{\frac{t}{m}}(N) \right| \le \sum_{h=0}^{\nu_0} \left| \prod_{k=1}^h (1 + e^{2\pi i (\alpha 2^{k-1} + \frac{1}{2})}) \right| \le \sum_{h=0}^{\nu_0} 2^h \prod_{k=1}^h \left| \sin(2^{k-1} \alpha \pi) \right| \le 1 + 2 \sum_{h=1}^{\nu_0} (2b_m)^{h-1} \le 1 + 2 \frac{(2b_m)^{\nu_0}}{2b_m - 1}.$$

Note that, according to (7) and (20)

$$(2b_m)^{\nu_0} = 2^{\lambda_m \nu_0} \le 2^{\lambda_m \log_2 N} = N^{\lambda_m}.$$

Thus,

$$|F_{\frac{t}{m}}(N)| \le 1 + \frac{2}{2b_m - 1} N^{\lambda_m} \le \frac{2}{2b_m - 1 - \gamma_m} N^{\lambda_m},$$

where γ_m is defined by the equality

$$\frac{1}{2b_m - 1 - \gamma_m} - \frac{1}{2b_m - 1} = \frac{1}{2}.$$

Hence, we find

$$\gamma_m = \frac{(2b_m - 1)^2}{2b_m + 1}$$

and, consequently, by (8),

$$|F_{\frac{t}{m}}(N)| \le \frac{2b_m + 1}{2b_m - 1}N^{\lambda_m} = \mu_m N^{\lambda_m}.$$

Thus, the theorem follows from (15).

4. Proof of Theorem 2.

Select in (15) the summands which correspond to $t = 0, \frac{m}{3}, \frac{2m}{3}$. We have

$$mS_m(N) = \sum_{n=0}^{N-1} \left(e^{\pi i s(n)} + e^{2\pi i (\frac{n}{3} + \frac{1}{2}s(n))} + e^{2\pi i (\frac{2n}{3} + \frac{1}{2}s(n))} \right) + \frac{m-1}{2\pi i (\frac{n}{3} + \frac{1}{2}s(n))}$$

(36)
$$+\sum_{t=1,t\neq\frac{m}{3},\frac{2m}{3}}^{m-1}\sum_{n=0}^{N-1}e^{2\pi i(\frac{t}{m}n+\frac{1}{2}s(n))}.$$

Since the chosen summands do not depend on m and for m = 3 the latter sum is empty then we find

(37)
$$mS_m(N) = 3S_3(N) + \sum_{t=1, t \neq \frac{m}{3}, \frac{2m}{3}}^{m-1} \sum_{n=0}^{N-1} e^{2\pi i (\frac{t}{m}n + \frac{1}{2}s(n))}.$$

Further, the last double sum is estimated by the same way as in Section 3 such that

(38)
$$\left|S_m(N) - \frac{3}{m}S_3(N)\right| \le \mu_m N^{\lambda_m} \blacksquare.$$

Remark 2. Notice that from elementary arguments it follows that if $m \ge 5$ is a multiple of 3 then

$$\left(\sin\frac{\pi}{m}\left\lfloor\frac{m-1}{3}\right\rfloor\right)\left(\sqrt{3}-\sin\frac{\pi}{m}\left\lfloor\frac{m-1}{3}\right\rfloor\right) \le \le \left(\sin\frac{\pi}{m}\left\lceil\frac{m+1}{3}\right\rceil\right)\left(\sqrt{3}-\sin\frac{\pi}{m}\left\lceil\frac{m+1}{3}\right\rceil\right).$$

The latter expression is the value of b_m^2 in this case (see (9)).

Example. Let us find some x_0 such that $S_{21}(x) > 0$ for $x \ge x_0$. Supposing that x is multiple of 3 and using (4) we obtain that

$$S_3(x) \ge \frac{2}{3^{\lambda + \frac{1}{2}}} x^{\lambda}.$$

Therefore, putting m = 21 in Theorem 2, we have

$$S_{21}(x) \ge \frac{1}{7} S_3(x) - \mu_{21} x^{\lambda_{21}} \ge \frac{2}{7 \cdot 3^{\lambda + \frac{1}{2}}} x^{\lambda} - \mu_{21} x^{\lambda_{21}}.$$

Now, calculating λ and λ_m by (2) and (8), we find a required x_0 :

$$x_0 = (3.5 \cdot 3^{\lambda + \frac{1}{2}} \mu_{21})^{\frac{1}{\lambda - \lambda_{21}}} = e^{984.839\dots}.$$

Corollary. For m which is not a multiple of 3, denote $U_m(x)$ the set of the positive integers not exceeding x which are multiples of m and not multiples of 3. Then

$$\sum_{e \in U_m(x)} (-1)^{s(n)} = -\frac{1}{m} S_3(x) + O(x^{\lambda_m}).$$

In particular, for sufficiently large x we have

$$\sum_{n \in U_m(x)} (-1)^{s(n)} < 0$$

Proof. Since

$$|U_m(x)| = S_m(x) - S_{3m}(x)$$

then the corollary immediately follows from Theorems 1, 2.

5. On Newman sum over primes

In [6] we put the following binary digit conjectures on primes. Conjecture 1. For all $n \in \mathbb{N}$, $n \neq 5, 6$

$$\sum_{p \le n} (-1)^{s(p)} \le 0,$$

where the summing is over all primes not exceeding n. Moreover, by the observations, $\sum_{p \leq n} (-1)^{s(p)} < 0$ beginning with n = 31. Conjecture 2.

$$\lim_{n \to \infty} \frac{\ln\left(-\sum_{p \le n} (-1)^{s(p)}\right)}{\ln n} = \frac{\ln 3}{\ln 4}.$$

A heuristic proof of Conjecture 2 was given in [8]. For a prime p, denote $V_p(x)$ the set of positive integers not exceeding x for which p is the least prime divisor. Show that the correctness of Conjectures 1 (for $n \ge n_0$) follows from the following very plausible statement, especially in view of the above estimates.

Conjecture 3. For sufficiently large n we have

(39)
$$\left|\sum_{5 \le p \le \sqrt{n}} \sum_{j \in V_p(n), j > p} (-1)^{s(j)} \right| < \sum_{j \in V_3(n)} (-1)^{s(j)} = S_3(n) - S_6(n).$$

Indeed, in the "worst case" (really is not satisfied) in which for all $n \ge p^2$

(40)
$$\sum_{j \in V_p(n), j > p} (-1)^{s(j)} < 0, \quad p \ge 5.$$

we have a decreasing but *positive* sequence of sums

$$\sum_{j \in V_3(n), j > 3} (-1)^{s(j)}, \sum_{j \in V_3(n), j > 3} (-1)^{s(j)} + \sum_{j \in V_5(n), j > 5} (-1)^{s(j)},$$
$$\dots, \sum_{j \in V_3(n), j > 3} (-1)^{s(j)} + \sum_{5 \le p < \sqrt{n}} \sum_{j \in V_p(n), j > p} (-1)^{s(j)} > 0.$$

Hence, the "balance condition" for odd numbers [8]

(41)
$$\left|\sum_{j \le n, j \text{ is odd}} (-1)^{s(j)}\right| \le 1$$

must be ensured permanently by the excess of the odious primes. This explains Conjecture 1.

It is very interesting that for some primes p most likely indeed (40) is satisfied for all $n \ge p^2$. Such primes we call "resonance primes". Our numerous observations show that all resonance primes not exceeding 1000 are:

$11, 19, 41, 67, 107, 173, 179, 181, 307, 313, 421, 431, 433, 587, \\601, 631, 641, 647, 727, 787.$

In conclusion, note that for $p \geq 3$ we have

(42)
$$\lim_{n \to \infty} \frac{|V_p(n)|}{n} = \frac{1}{p} \prod_{2 \le q < p} \left(1 - \frac{1}{q} \right)$$

such that

(43)
$$\lim_{n \to \infty} \left(\sum_{p \ge 3} \frac{|V_p(n)|}{n} \right) = \frac{1}{2}.$$

Thus, using Theorems 1, 2 in the form

(44)
$$S_m(n) = \begin{cases} o(S_3(n)), & (m,3) = 1\\ \frac{3}{m}S_3(n)(1+o(1)), & 3|m \end{cases}$$

and inclusion-exclusion for $p \ge 5$, we find

(45)
$$\sum_{j \in V_p(n)} (-1)^{\sigma(j)} = -\frac{3}{3p} \prod_{2 \le q < p, q \ne 3} (1 - \frac{1}{q}) S_3(n) (1 + o(1)) = -\frac{3}{2p} \prod_{2 \le q < p} (1 - \frac{1}{q}) S_3(n) (1 + o(1)).$$

Now in view of (5) we obtain the following absolute result as an approximation of Conjectures 1, 2.

Theorem 3. For every prime number $p \ge 5$ and sufficiently large $n \ge n_p$ we have

$$\sum_{j \in V_p(n)} (-1)^{s(j)} < 0$$

and, moreover,

$$\lim_{n \to \infty} \frac{\ln(-\sum_{j \in V_p(n)} (-1)^{s(j)})}{\ln n} = \frac{\ln 3}{\ln 4}.$$

References

- J.-P. Allouche and J.Shallit. Automatic Sequences: Theory, Applications, Generalizations, Cambridge University Press, 2003.
- [2] J.Coquet, A summation formula related to the binary digits, Invent, Math.73 (1983),107-115.
- [3] M.Drmota and M.Skalba, Rarified sums of the Thue-Morse sequence, Tranc. of the AMS 352, no.2 (1999), 609-642.
- [4] A.O.Gelfond, Sur les nombres qui ont des proprietes additives et multuplicatives donnees, Acta ArithmeticaXIII (1968),259-265.
- [5] D.I.Newman, On the number of binery digit in a multiple of three, Proc. AMS 21 (1969), 719-721.
- [6] V.Shevelev, A conjecture on primes and a step towards justification, arXiv (math. NT math. CO), 0706.0786
- [7] . V.Shevelev, Two algorithms for exact evalution of the Newman digit sum and the sharp estimates, arXiv (math. NT), 0709.0885 v1.
- [8] . V.Shevelev, On excess of the odious primes, arXiv (math. NT), 0707.1761.

DEPARTMENTS OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA 84105, ISRAEL. E-MAIL:SHEVELEV@BGU.AC.IL