Bent Rectangles[∗]

Sergey Agievich

National Research Center for Applied Problems of Mathematics and Informatics Belarusian State University Nezavisimosti av. 4, 220030 Minsk, Belarus agievich@bsu.by

Abstract

We study generalized regular bent functions using a representation by bent rectangles, that is, special matrices with restrictions on rows and columns. We describe affine transformations of bent rectangles, propose new biaffine and bilinear constructions, study partitions of a vector space into affine planes of the same dimension and use such partitions to build bent rectangles. We illustrate the concept of bent rectangles by examples for the Boolean case.

Introduction

Boolean bent functions were introduced by Rothaus [\[14\]](#page-18-0) and Dillon [\[7\]](#page-18-1), and since then have been widely studied owing to their interesting algebraic and combinatorial properties and because of their applications in signal processing, coding theory, cryptography. The current results on the construction, classification, enumeration of bent functions can be found in the surveys [\[5,](#page-17-0) [9\]](#page-18-2).

In the present paper we study bent functions using a representation by bent rectangles. Rows and normalized columns of such rectangles consist of the Walsh–Hadamard coefficients of Boolean functions and the Boolean functions of rows are simply restrictions of a target bent function to a subset of variables.

The notion of bent rectangles was introduced in [\[2\]](#page-17-1) and has analogues in other papers. For example, a special kind of rectangles, all elements of which are equal in magnitude, was proposed in [\[1\]](#page-17-2). Authors of [\[3\]](#page-17-3) also intensively dealt with restrictions of bent functions and actually used 2- and 4-row bent rectangles. In [\[10,](#page-18-3) preliminary version], \mathbb{Z} -bent squares that are defined for integer-valued functions are used to construct Boolean bent functions recursively.

We extend the results of [\[2\]](#page-17-1) in the following directions. In Sections [1,](#page-1-0) [2](#page-2-0) we transfer the concept of bent rectangles to the generalized regular bent functions over an arbitrary quotient ring of integers (see [\[11\]](#page-18-4)). In Section [3](#page-4-0) we describe affine transformations of bent rectangles. In Section [5](#page-10-0) we propose new constructions of bent functions which are based on biaffine and bilinear mappings. In Section [6](#page-12-0) we study partitions of a vector space into affine planes of the

[∗]Submitted to: Proceedings of the NATO Advanced Study Institute on Boolean Functions in Cryptology and Information Security (Moscow, September 8–18, 2007).

same dimension and use such partitions to construct bent rectangles. Additionally, in Section [4](#page-6-0) we illustrate some known properties of Boolean bent functions by using bent rectangles.

The author plans to continue this paper and discuss further dual bent functions, bent rectangles with a small number of rows, and cubic Boolean bent functions.

1 Functions

Let \mathbb{Z}_q be the ring of integers modulo q which we identify with the set $\{0, 1, \ldots, q-1\}$. Denote by \mathbb{Z}_q the set of all qth roots of unity in $\mathbb C$ and introduce the additive character $\chi: \mathbb{Z}_q \to \mathbb{Z}_q$, $\chi(a) = \exp(2\pi i a/q)$, where $i =$ √ $\overline{-1}$. The ring of integers modulo a prime q is a field. We emphasize this fact by writing \mathbb{F}_q instead of \mathbb{Z}_q .

The set $V_n = \mathbb{Z}_q^n$ consists of all vectors $\mathbf{a} = (a_1, \ldots, a_n)$, $a_i \in \mathbb{Z}_q$, and forms a group under vector addition. If q is prime, then V_n is the n-dimensional vector space over the field \mathbb{F}_q .

Let \mathcal{F}_n be the set of all functions $V_n \to \mathbb{Z}_q$. A function $f \in \mathcal{F}_n$ depends on n variables coordinates of the argument $\mathbf{x} = (x_1, \ldots, x_n)$. Given $J = \{j_1, \ldots, j_m\} \subseteq \{1, \ldots, n\}$, the restriction of $f(\mathbf{x})$ to $(x_{j_1},...,x_{j_m})$ is a function obtained from f by keeping variables x_j , $j \notin J$, constant.

Starting from f , construct the function $^{\circ}$ $f: V_n \to$ $\overset{\circ}{\mathbb{Z}}_{q},\overset{\circ}{f}$ $f(\mathbf{x}) = \chi(f(\mathbf{x}))$, and the function $\hat{f}: V_n \to \mathbb{C},$

$$
\hat{f}(\mathbf{u}) = \sum_{\mathbf{x} \in V_n} \hat{f}(\mathbf{x}) \overline{\chi(\mathbf{u} \cdot \mathbf{x})}, \quad \mathbf{u} \in V_n.
$$

Here $\mathbf{u} \cdot \mathbf{x} = u_1 x_1 + \ldots + u_n x_n$ and the bar indicates complex conjugation. It is convenient to assume that \mathcal{F}_0 consists of constant functions $f \equiv c, c \in \mathbb{Z}_q$, for which \hat{f} $\hat{f} = \hat{f} \equiv \chi(c)$. Denote ◦ $\mathfrak{F}_n = \{$ ◦ $\hat{f} : f \in \mathcal{F}_n$, $\hat{\mathcal{F}}_n = \{ \hat{f} : f \in \mathcal{F}_n \}.$

The conversion $\hat{f} \mapsto \hat{f}$ (or $f \mapsto \hat{f}$) is called the *Walsh–Hadamard transform*. Since

$$
\sum_{\mathbf{u}\in V_n} \chi(\mathbf{a}\cdot\mathbf{u}) = \begin{cases} q^n, & \mathbf{a} = \mathbf{0}, \\ 0 & \text{otherwise} \end{cases}
$$

for each $\mathbf{a} \in V_n$, the inverse transform $\hat{f} \mapsto$ $^{\circ}$ f can be defined as follows:

$$
\hat{f}(\mathbf{x}) = q^{-n} \sum_{\mathbf{y} \in V_n} \hat{f}(\mathbf{y}) \sum_{\mathbf{u} \in V_n} \chi(\mathbf{u} \cdot (\mathbf{x} - \mathbf{y}))
$$

= $q^{-n} \sum_{\mathbf{u} \in V_n} \sum_{\mathbf{y} \in V_n} \hat{f}(\mathbf{y}) \overline{\chi(\mathbf{u} \cdot \mathbf{y})} \chi(\mathbf{u} \cdot \mathbf{x})$
= $q^{-n} \sum_{\mathbf{u} \in V_n} \hat{f}(\mathbf{u}) \chi(\mathbf{u} \cdot \mathbf{x}).$

The Walsh–Hadamard transform is often used in cryptography, coding theory, signal processing to derive some properties of f . In many cases it is important to use functions f with the characteristic max_u $\hat{f}(\mathbf{u})$ as small as possible. Due to the Parseval's identity

$$
\sum_{\mathbf{u}\in V_n} |\hat{f}(\mathbf{u})|^2 = q^{2n},
$$

we have $\max_{\mathbf{u}} |\hat{f}(\mathbf{u})| \geq q^{n/2}$, where the equality holds if and only if

$$
|\hat{f}(\mathbf{u})| = q^{n/2}, \quad \mathbf{u} \in V_n.
$$
 (1)

If f satisfies [\(1\)](#page-2-1), then it is called a *bent* function. If, additionally to (1), $\hat{f}(\mathbf{u}) \in q^{n/2} \mathbb{Z}_q$ for all **u**, then f is called a regular bent function. Let \mathcal{B}_n be the set of all regular bent functions of n variables.

Further all subscripts, superscripts and other notations defined for \mathcal{F}_n are automatically transferred to all subsets of \mathcal{F}_n . For example, $^{\circ}$ $\mathfrak{B}_n = \{$ $^{\circ}$ \hat{f} : $f \in \mathcal{B}_n$, $\hat{\mathcal{B}}_n = \{\hat{f} : f \in \mathcal{B}_n\}$. Note that $\mathring{\mathcal{B}}_n = q^{n/2} \mathring{\mathcal{E}}$ \mathfrak{B}_n .

2 Rectangles

Let m, k be nonnegative integers and $f \in \mathcal{F}_{m+k}$. Define the function $\overrightarrow{f} : V_{m+k} \to \mathbb{C}$,

$$
\overline{f}(\mathbf{u}, \mathbf{v}) = \sum_{\mathbf{y} \in V_k} \hat{f}(\mathbf{u}, \mathbf{y}) \overline{\chi(\mathbf{v} \cdot \mathbf{y})}, \quad \mathbf{u} \in V_m, \quad \mathbf{v} \in V_k,
$$
\n(2)

and call it a *rectangle* of f. Denote by $\overline{\mathcal{F}}_{m,k}$ the set of all such rectangles. For the case $m = k$ we also call \bar{f} a square of f.

To each function f there correspond different rectangles of the sets $\mathcal{F}_{0,m+k} = \hat{\mathcal{F}}_{m+k}$, ${\mathfrak F}_{1,m+k-1}, \ldots, {\mathfrak F}_{m+k-1,1}, {\mathfrak F}_{m+k,0} =$ $^{\circ}$ \mathfrak{F}_{m+k} . These rectangles are connected with each other. For example, if $f \in \mathcal{F}_{m,k}$ and $f^* \in \mathcal{F}_{m+1,k-1}$, then

$$
\overrightarrow{f}(\mathbf{u}, v, \mathbf{v}') = \sum_{y \in \mathbb{Z}_q} \sum_{\mathbf{y}' \in V_{k-1}} \overrightarrow{f}(\mathbf{u}, y, \mathbf{y}') \overrightarrow{\chi(vy + \mathbf{v}' \cdot \mathbf{y}')}
$$

$$
= \sum_{y \in \mathbb{Z}_q} \overrightarrow{f}^*(\mathbf{u}, y, \mathbf{v}') \overrightarrow{\chi(vy)}, \quad \mathbf{v}' \in V_{k-1},
$$

and conversely,

$$
\overline{f}^*(\mathbf{u}, v, \mathbf{v}') = \frac{1}{q} \sum_{y \in \mathbb{Z}_q} \overline{f}(\mathbf{u}, y, \mathbf{v}') \chi(vy).
$$

For a fixed **u** call the mapping $\mathbf{v} \mapsto f(\mathbf{u}, \mathbf{v})$ a *column* of f. Analogously, for a fixed **v** call the mapping $\mathbf{u} \mapsto f(\mathbf{u}, \mathbf{v})$ a row of \overline{f} . By definition, each row of \overline{f} is an element of $\hat{\overline{f}}_k$. If furthermore each column of \int_a^{π} multiplied by $q^{(m-k)/2}$ is an element of $\hat{\mathcal{F}}_m$, then call the rectangle \overline{f} bent.

Our results are based on the following proposition, first stated in [\[2\]](#page-17-1) for the case $q = 2$. Note that under $m = 0$ this proposition can be considered as a definition of regular bent functions.

Proposition 1. A function $f \in \mathcal{F}_{m+k}$ is regular bent if and only if the rectangle $\overline{f} \in \overline{\mathcal{F}}_{m,k}$ is bent.

Proof. Let $f \in \mathcal{B}_n$, $n = m + k$. Define the function $g \in \mathcal{F}_n$ by the rule

$$
\overset{\circ}{g}(\mathbf{v},\mathbf{u})=q^{-n/2}\hat{f}(-\mathbf{u},\mathbf{v}),\quad \mathbf{u}\in V_m,\quad \mathbf{v}\in V_k,
$$

and determine the corresponding rectangle $\overline{g} \in \overline{\mathcal{F}}_{k,m}$:

 \Box

$$
\overline{g}(\mathbf{v}, \mathbf{u}) = q^{-n/2} \sum_{\mathbf{x} \in V_m} \hat{f}(-\mathbf{x}, \mathbf{v}) \overline{\chi(\mathbf{u} \cdot \mathbf{x})}
$$
\n
$$
= q^{-n/2} \sum_{\mathbf{x} \in V_m} \sum_{\mathbf{w} \in V_m} \sum_{\mathbf{y} \in V_k} \hat{f}(\mathbf{w}, \mathbf{y}) \overline{\chi(-\mathbf{x} \cdot \mathbf{w} + \mathbf{v} \cdot \mathbf{y} + \mathbf{u} \cdot \mathbf{x})}
$$
\n
$$
= q^{-n/2} \sum_{\mathbf{y} \in V_k} \sum_{\mathbf{w} \in V_m} \hat{f}(\mathbf{w}, \mathbf{y}) \overline{\chi(\mathbf{v} \cdot \mathbf{y})} \sum_{\mathbf{x} \in V_m} \chi((\mathbf{w} - \mathbf{u}) \cdot \mathbf{x})
$$
\n
$$
= q^{m-n/2} \sum_{\mathbf{y} \in V_k} \hat{f}(\mathbf{u}, \mathbf{y}) \overline{\chi(\mathbf{v} \cdot \mathbf{y})}
$$
\n
$$
= q^{(m-k)/2} \overline{\hat{f}}(\mathbf{u}, \mathbf{v}).
$$

Therefore, each column of \int_{0}^{π} multiplied by $q^{(m-k)/2}$ is an element of \hat{f}_m and \int_{0}^{π} is bent.

Conversely, if f is bent, then $g(\mathbf{v}, \mathbf{u}) = q^{(m-k)/2} f(\mathbf{u}, \mathbf{v})$ is well defined rectangle that $g(\mathbf{v}, \mathbf{u}) = q^{-n/2} \hat{f}(\mathbf{u}, \mathbf{v})$. Hence $\hat{f}(\mathbf{u}, \mathbf{v}) \in q^{n/2} \mathbb{Z}_q$ for all \mathbf{u}, \mathbf{v} $^{\circ}$ corresponds to the function and f is regular bent. $\overline{}$

It is convenient to identify $f \in \mathcal{F}_{m,k}$ with the $q^m \times q^k$ matrix F whose rows and columns are marked by lexicographically ordered vectors of V_m and V_k respectively and whose elements are the values $\bar{f}(\mathbf{u}, \mathbf{v})$.

The definition of \overline{F} puts restrictions on its rows. The bentness of \overline{F} puts additional restrictions on columns. We can draw an analogy with latin rectangles and this analogy justify the use of the term "bent *rectangle*".

If F corresponds to a bent rectangle $f \in \mathcal{B}_{m,k}$ and satisfies restrictions on rows and columns, then $G = q^{(m-k)/2} F^{T}$ also satisfies such restrictions and corresponds to a bent rectangle $g \in \mathcal{B}_{k,m}$. Call the transformation $f \mapsto g$ a transposition of f. During the proof of Proposition [1](#page-2-2) we actually showed that, under the transposition,

$$
\overset{\circ}{g}(\mathbf{v},\mathbf{u})=q^{-n/2}\hat{f}(-\mathbf{u},\mathbf{v}),\quad \mathbf{u}\in V_m,\quad \mathbf{v}\in V_k.
$$

As "material" for constructing bent rectangles we will often use affine functions $l(\mathbf{x}) =$ $\mathbf{b} \cdot \mathbf{x} + c$, where $\mathbf{x}, \mathbf{b} \in V_n$ and $c \in \mathbb{Z}_q$. Denote by \mathcal{A}_n the set of all affine functions of n variables.

The function \hat{l} that corresponds to l has the quite simple form:

$$
\hat{l}(\mathbf{u}) = \begin{cases} q^n \chi(c), & \text{if } \mathbf{u} = \mathbf{b}, \\ 0 & \text{otherwise.} \end{cases}
$$

This fact immediately gives us the following construction.

Example 1 (Maiorana–McFarland's construction). Consider a bent square $\overrightarrow{f} \in \mathcal{B}_{n,n}$ such that all its rows and columns belong to \hat{A}_n . The matrix \overline{F} associated with \overline{f} has the following form: each its row and each column contains exactly one element of the set $q^n \mathbb{Z}_q$, all other elements are zero. It means that there exists a permutation $\pi: V_n \to V_n$ and a function $\varphi \in \mathcal{F}_n$ such that

$$
\mathop{\mathcal{F}}\limits^{\square}(\mathbf{u}, \mathbf{v}) = \left\{ \begin{array}{l} \mathop{\phi}\limits^{\circ}(\mathbf{u}), & \text{if } \mathbf{v} = \pi(\mathbf{u}), \\ 0 & \text{otherwise.} \end{array} \right.
$$

Consequently, $f(\mathbf{x}, \mathbf{y}) = \pi(\mathbf{x}) \cdot \mathbf{y} + \varphi(\mathbf{x})$ and we obtain the well-known Maiorana-McFarland's bent function. \Box

3 Affine transformations

Let \mathbf{GL}_n (\mathbf{AGL}_n) be the general linear (affine) group of transformations of $V_n = \mathbb{Z}_q^n$. We identify \mathbf{GL}_n with the set of all invertible $n \times n$ matrices over \mathbb{Z}_q and denote by I_n the identity matrix of \mathbf{GL}_n . An element $\sigma \in \mathbf{AGL}_n$ is specified by a pair $(A, \mathbf{a}), A \in \mathbf{GL}_n$, $\mathbf{a} \in V_n$, and acts as follows: $\sigma(\mathbf{x}) = \mathbf{x}A + \mathbf{a}$. Extend the action of \mathbf{AGL}_n to functions f with domain V_n in a natural way:

$$
\sigma(f)(\mathbf{x}) = f(\mathbf{x}A + \mathbf{a}).
$$

Call functions $f, g \in \mathcal{F}_n$ affine equivalent if there exist $\sigma = (A, \mathbf{a}) \in \mathbf{AGL}_n$ and $l \in \mathcal{A}_n$, $l(\mathbf{x}) = \mathbf{b} \cdot \mathbf{x} + c$, such that

$$
g(\mathbf{x}) = \sigma(f)(\mathbf{x}) + l(\mathbf{x}) = f(\mathbf{x}A + \mathbf{a}) + \mathbf{b} \cdot \mathbf{x} + c.
$$
 (3)

If f and g are affine equivalent, then

$$
\hat{g}(\mathbf{u}) = \sum_{\mathbf{x} \in V_n} \chi(f(\mathbf{x}A + \mathbf{a}) + \mathbf{x} \cdot (\mathbf{b} - \mathbf{u}) + c)
$$

=
$$
\sum_{\mathbf{y} \in V_n} \chi(f(\mathbf{y}) + (\mathbf{y} - \mathbf{a})A^{-1} \cdot (\mathbf{b} - \mathbf{u}) + c)
$$

=
$$
\sum_{\mathbf{y} \in V_n} \chi(f(\mathbf{y}) - (\mathbf{y} - \mathbf{a}) \cdot (\mathbf{u} - \mathbf{b})(A^{-1})^{\mathrm{T}} + c)
$$

and

$$
\hat{g}(\mathbf{u}) = \chi(\mathbf{a} \cdot \mathbf{v} + c) \hat{f}(\mathbf{v}), \quad \mathbf{v} = (\mathbf{u} - \mathbf{b})(A^{-1})^{\mathrm{T}}.
$$
\n(4)

Therefore, the functions \hat{f} and \hat{g} are, in some sense, also affine equivalent: there exist $\sigma^* \in$ \mathbf{AGL}_n and $l^* \in \mathcal{A}_n$ such that

$$
\hat{g}(\mathbf{u}) = \sigma^*(\hat{f})(\mathbf{u})\overset{\circ}{l}^*(\mathbf{u}).
$$

It is useful to describe a connection between rectangles \overline{f} , $\overline{g} \in \overline{\mathcal{F}}_{m,k}$ of affine equivalent functions. For example, $g \in \mathcal{B}_n$ if and only if $f \in \mathcal{B}_n$ and a correspondence between f and g can be used to perform the affine classification of regular bent functions.

Let $n = m + k$, where $m, k \ge 1$. Divide the vector $\mathbf{x} \in V_n$ into two parts: $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2)$, $\mathbf{x}_1 \in V_m$, $\mathbf{x}_2 \in V_k$, and introduce the following *elementary transformations* $f \mapsto g$:

- A1) $g(\mathbf{x}) = f(\mathbf{x}_1 A_1 + \mathbf{a}_1, \mathbf{x}_2)$, where $A_1 \in \mathbf{GL}_m$, $\mathbf{a}_1 \in V_m$;
- A2) $g(\mathbf{x}) = f(\mathbf{x}_1, \mathbf{x}_2 A_2) + \mathbf{a}_2 \cdot \mathbf{x}_2$, where $A_2 \in \mathbf{GL}_k$, $\mathbf{a}_2 \in V_k$;
- B1) $q(\mathbf{x}) = f(\mathbf{x}) + \mathbf{b}_1 \cdot \mathbf{x}_1 + c$, where $\mathbf{b}_1 \in V_m$, $c \in \mathbb{Z}_q$;
- B2) $q(\mathbf{x}) = f(\mathbf{x}_1, \mathbf{x}_2 + \mathbf{b}_2)$, where $\mathbf{b}_2 \in V_k$;
- C1) $g(\mathbf{x}) = f(x_1, \ldots, x_{m-1}, x_m x_{m+1}, x_{m+1}, x_{m+2}, \ldots, x_n);$
- C2) $q(\mathbf{x}) = f(x_1, \ldots, x_{m-1}, x_m, x_{m+1} x_m, x_{m+2}, \ldots, x_n).$

Proposition 2. Every affine transformation $f \mapsto g$ of the form [\(3\)](#page-4-1) can be realized using only elementary transformations A1 – C2. Under these transformations, the rectangles \overrightarrow{f} , $\overrightarrow{g} \in \overrightarrow{T}_{m,k}$ are connected in the following manner:

A1)
$$
\overset{\Box}{g}(\mathbf{u}, \mathbf{v}) = \overset{\Box}{f}(\mathbf{u}A_1 + \mathbf{a}_1, \mathbf{v}),
$$

\nA2) $\overset{\Box}{g}(\mathbf{u}, \mathbf{v}) = \overset{\Box}{f}(\mathbf{u}, (\mathbf{v} - \mathbf{a}_2)B_2), \text{ where } B_2 = (A_2^{-1})^{\mathrm{T}};$
\nB1) $\overset{\Box}{g}(\mathbf{u}, \mathbf{v}) = \chi(\mathbf{b}_1 \cdot \mathbf{u} + c) \overset{\Box}{f}(\mathbf{u}, \mathbf{v});$
\nB2) $\overset{\Box}{g}(\mathbf{u}, \mathbf{v}) = \chi(\mathbf{b}_2 \cdot \mathbf{v}) \overset{\Box}{f}(\mathbf{u}, \mathbf{v});$
\nC1) $\overset{\Box}{g}(\mathbf{u}', u, v, \mathbf{v}') = \frac{1}{q} \sum_{x, y \in \mathbb{Z}_q} \overset{\Box}{f}(\mathbf{u}', x, y, \mathbf{v}') \overline{\chi((u - x)(v - y))}};$
\nC2) $\overset{\Box}{g}(\mathbf{u}', u, v, \mathbf{v}') = \overset{\Box}{f}(\mathbf{u}', u, v, \mathbf{v}') \overline{\chi(uv)}.$
\nHere $\mathbf{u} \in V_m$, $\mathbf{v} \in V_k$, $\mathbf{u}' \in V_{m-1}$, $\mathbf{v}' \in V_{k-1}$, $u, v \in \mathbb{Z}_q$.

Proof. To prove the first part of the proposition, we will establish that GL_n is generated by the matrices of its subgroups

$$
G_1 = \left\{ \begin{pmatrix} M_1 & 0 \\ 0 & I_k \end{pmatrix} : M_1 \in \mathbf{GL}_m \right\}, \quad G_2 = \left\{ \begin{pmatrix} I_m & 0 \\ 0 & M_2 \end{pmatrix} : M_2 \in \mathbf{GL}_k \right\}
$$

and additional matrices R and S of the following linear transformations:

$$
\mathbf{x} \mapsto \mathbf{x}R = (x_1, \dots, x_{m-1}, x_m - x_{m+1}, x_{m+1}, x_{m+2}, \dots, x_n),
$$

$$
\mathbf{x} \mapsto \mathbf{x}S = (x_1, \dots, x_{m-1}, x_m, x_{m+1} - x_m, x_{m+2}, \dots, x_n).
$$

It is known (see, for example, [\[15\]](#page-18-5)) that the group GL_n over the Euclidian ring \mathbb{Z}_m is generated by matrices of the following transformations of a vector x:

- 1) multiplication of the coordinates x_i by invertible elements of \mathbb{Z}_m , $1 \leq i \leq n$;
- 2) subtraction of x_j from x_i , $1 \leq i, j \leq n$, $i \neq j$.

The groups G_1 and G_2 contain matrices which realize all transformations of the first type and transformations of the second type for $1 \le i, j \le m$ and $m+1 \le i, j \le n$. We can subtract any different coordinates of x using additional matrices R and S . For example, the subtraction of x_j , $m+1 \le j \le n$, from x_i , $1 \le i \le m$, can be realized by the following steps: a) interchange x_i and x_m using some matrix of G_1 , b) interchange x_{m+1} and x_j using some matrix of G_2 , c) subtract x_{m+1} from x_m using R, d) interchange x_i and x_m , x_{m+1} and x_j again. Thus, the group generated by G_1, G_2, R, S contains all matrices of the first and the second types and this group coincides with GL_n .

The second part of the proposition is checked by direct calculations. Consider, for example, the transformation C1. We have

$$
\begin{split}\n\overset{\Box}{g}(\mathbf{u}', u, v, \mathbf{v}') &= \sum_{x \in \mathbb{Z}_q} \sum_{\mathbf{y}' \in V_{k-1}} \overset{\circ}{g}(\mathbf{u}', u, x, \mathbf{y}') \overline{\chi(vx + \mathbf{v}' \cdot \mathbf{y}')} \\
&= \sum_{x \in \mathbb{Z}_q} \sum_{\mathbf{y}' \in V_{k-1}} \overset{\circ}{f}(\mathbf{u}', u - x, x, \mathbf{y}') \overline{\chi(vx + \mathbf{v}' \cdot \mathbf{y}')} \\
&= \sum_{x \in \mathbb{Z}_q} \sum_{\mathbf{y}' \in V_{k-1}} \overset{\circ}{f}(\mathbf{u}', x, u - x, \mathbf{y}') \overline{\chi(v(u - x) + \mathbf{v}' \cdot \mathbf{y}')} \\
&= \frac{1}{q} \sum_{x, y, z \in \mathbb{Z}_q} \sum_{\mathbf{y}' \in V_{k-1}} \overset{\circ}{f}(\mathbf{u}', x, z, \mathbf{y}') \overline{\chi((u - x)(v - y) + yz + \mathbf{v}' \cdot \mathbf{y}')} \\
&= \frac{1}{q} \sum_{x, y \in \mathbb{Z}_q} \overset{\Box}{f}(\mathbf{u}', x, y, \mathbf{v}') \overline{\chi((u - x)(v - y))}\n\end{split}
$$

and the required identity holds.

Therefore, we can realize any affine transformation of a rectangle \overrightarrow{F} by affine permutations of its rows $(A1)$ and columns $(A2)$ and by multiplying its elements by qth roots of unity $(B1,$ B2, C2). The remaining transformation C1 is a single method of changing the elements of \overline{F} in magnitude.

 \Box

Example 2 (cells). For the case $q = 2$ it is convenient to illustrate the transformations C1, C2 in the following way. Divide \overline{F} into the *cells*, that is, the submatrices

$$
\begin{pmatrix}\n\int_{a}^{B} (\mathbf{u}', 0, 0, \mathbf{v}') & \int_{a}^{B} (\mathbf{u}', 0, 1, \mathbf{v}') \\
\int_{a}^{B} (\mathbf{u}', 1, 0, \mathbf{v}') & \int_{a}^{B} (\mathbf{u}', 1, 1, \mathbf{v}')\n\end{pmatrix}, \quad \mathbf{u}' \in V_{m-1}, \quad \mathbf{v}' \in V_{k-1},
$$

and during the transformations modify all of the cells simultaneously by the rules

C1:
$$
\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \mapsto \frac{1}{2} \begin{pmatrix} \alpha + \beta + \gamma - \delta & \alpha + \beta - \gamma + \delta \\ \alpha - \beta + \gamma + \delta & -\alpha + \beta + \gamma + \delta \end{pmatrix}
$$
, C2: $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \mapsto \begin{pmatrix} \alpha & \beta \\ \gamma & -\delta \end{pmatrix}$.

4 Illustrations

In this section we consider the case $q = 2$ and give examples to illustrate the usage of bent rectangles for the analysis of some known properties of bent functions. Note that in the Boolean case each bent function is regular and $\mathcal{B}_n \neq \emptyset$ only for even n.

Start with some useful definitions and facts. Firstly, identify a Boolean function $f \in \mathcal{F}_n$ of $\mathbf{x} = (x_1, \ldots, x_n)$ with its algebraic normal form, that is, a polynomial of the ring $\mathbb{F}_2[x_1, \ldots, x_n]$ reduced modulo the ideal $(x_1^2-x_1,\ldots,x_n^2-x_n)$. Denote by deg f the degree of such a polynomial.

Following [\[16\]](#page-18-6), introduce the set $\mathcal{P}_{n,r} \subseteq \mathcal{F}_n$ of plateaued functions of order $r: f \in \mathcal{P}_{n,r}$ if $|\hat{f}(\mathbf{u})| \in \{0, 2^{n-r/2}\}\$ for all $\mathbf{u} \in V_n$ (more precisely, \hat{f} has exactly 2^r nonzero values $\pm 2^{n-r/2}$). It is clear that $A_n = \mathcal{P}_{n,0}, \mathcal{B}_n = \mathcal{P}_{n,n}$.

Finally, recall the following result of [\[2\]](#page-17-1).

Lemma 1. Let $q = 2$, $f_1, f_2, f_3, f_4 \in \mathcal{F}_n$ and

$$
\hat{g}(\mathbf{u}) = \frac{1}{2}(\hat{f}_1(\mathbf{u}) + \hat{f}_2(\mathbf{u}) + \hat{f}_3(\mathbf{u}) + \hat{f}_4(\mathbf{u})), \quad \mathbf{u} \in V_n.
$$

The function $\hat{g} \in \hat{\mathcal{F}}_n$ if and only if

$$
f_1(\mathbf{x}) + f_2(\mathbf{x}) + f_3(\mathbf{x}) + f_4(\mathbf{x}) = 1, \quad \mathbf{x} \in V_n.
$$

Under this condition, $g(\mathbf{x}) = f_1(\mathbf{x})f_2(\mathbf{x}) + f_1(\mathbf{x})f_3(\mathbf{x}) + f_2(\mathbf{x})f_3(\mathbf{x}).$

Turn to examples.

Example 3 (sums of bent functions). It is well-known (see [\[14\]](#page-18-0)) that if $f_1 \in \mathcal{B}_m$ and $f_2 \in \mathcal{B}_k$, then $f(\mathbf{x}, \mathbf{y}) = f_1(\mathbf{x}) + f_2(\mathbf{y})$ is also bent. Indeed, the rectangle $\overline{f} \in \overline{\mathcal{F}}_{m,k}$ has the form ◦ \int_{1}^{∞} \int_{1}^{∞} \int_{1}^{∞} and obviously satisfies the restrictions on columns. $f\left(\mathbf{u}, \mathbf{v}\right) =$ \Box

Example 4 (degrees of bent functions). Let $f \in \mathcal{B}_{2n}$, $n \geq 2$. In [\[14\]](#page-18-0), Rothaus proved that $\deg f \leq n$. Give an alternative proof of this fact.

Suppose to the contrary that deg $f = k > n$, i.e. the polynomial $f(x_1, \ldots, x_{2n})$ contains a monomial of degree k . Without loss of generality, assume that f contains the monomial $x_{m+1}x_{m+2}...x_{2n}$, $m=2n-k$. Then there are an odd number of 1's among the values $g(\mathbf{y})=$ $f(\mathbf{0}, \mathbf{y}), \mathbf{y} \in V_k$, and $\hat{g}(\mathbf{0}) = 2r$, where r is odd. It is impossible when $k = 2n$, since in this case $(g(\mathbf{0}) = \hat{f}(\mathbf{0}) \in {\{\pm 2^n\}}$. Therefore $k < 2n$ and consequently $m > 0$. Consider $\overline{f} \in \mathring{B}_{m,k}$. We have $\int_a^{\pi} (0,0) = \int_a^{\pi} (0,0) dx = 2r$ and $2^{(m-k)/2} \int_a^{\pi} (0,0)$ is not even integer. Since functions of $\hat{\mathcal{F}}_m$ take only even values, the restrictions on columns of \overline{f} are not satisfied, a contradiction. \Box

Example 5 (2-row bent rectangles). Let $\overline{f} \in \mathcal{B}_{1,n-1}$. By definition, the columns of \overline{f} belong to the set $2^{n/2-1}\mathcal{F}_1$. Since $\mathcal{F}_1 = \mathcal{A}_1$, each column takes exactly one nonzero value $\pm 2^{n/2}$. It means that all rows of \overrightarrow{f} are in $\hat{\mathcal{P}}_{n-1,n-2}$. This fact was pointed out in [\[16\]](#page-18-6).

In Proposition [5](#page-12-1) we will describe how to construct $(2^{n}-2)|\mathcal{B}_{n-2}|^2$ bent rectangles of $\mathcal{B}_{1,n-1}$. For example, such a construction allows to obtain all 896 elements of $\overline{B}_{1,3}$. \Box

Example 6 (4-row bent rectangles). Consider a rectangle $\overrightarrow{f} \in \mathcal{B}_{2n-2}$. Since every function of \mathcal{F}_2 is either affine or bent, the possible values of f are exhausted by 0, $\pm 2^{n/2-1}$, $\pm 2^{n/2}$.

If we restrict to the numbers $\pm 2^{n/2-1}$, then each column of \int_{a}^{π} belongs to $2^{n/2-1} \mathcal{L}$ B_2 , takes an odd number of negative values, and a product of these values is exactly -2^{2n-4} . This result was proved in [\[13\]](#page-18-7).

If we restrict to the numbers $0, \pm 2^{n/2}$, then we obtain a bent rectangle all rows of which are in $\hat{\mathcal{P}}_{n-2,n-4}$. Proposition [5](#page-12-1) will give $8(2^{n-2}-1)(2^{n-3}-1)(7 \cdot 2^{n-3}-13)|\mathcal{B}_{n-4}|^4$ such rectangles for $n > 4$.

Example 7 (Rothaus' construction). Let $f_1, f_2, f_3, f_4 \in \mathcal{B}_n$ satisfy $f_1(\mathbf{y}) + f_2(\mathbf{y}) + f_3(\mathbf{y}) + f_4(\mathbf{y})$ $f_4(\mathbf{y}) = 0$ for all $\mathbf{y} \in V_n$. In [\[14\]](#page-18-0), Rothaus showed that the function

$$
f(u_1, u_2, \mathbf{y}) = f_1(\mathbf{y})f_2(\mathbf{y}) + f_1(\mathbf{y})f_3(\mathbf{y}) + f_2(\mathbf{y})f_3(\mathbf{y}) + u_1(f_1(\mathbf{y}) + f_2(\mathbf{y})) + u_2(f_1(\mathbf{y}) + f_3(\mathbf{y})) + u_1u_2
$$

is bent. Give another proof of this fact using the rectangle $\overline{f} \in \overline{\mathcal{F}}_{2,n}$.

Lemma [1](#page-7-0) implies that \overline{f} has the following form:

$$
\overrightarrow{f}(u_1, u_2, \mathbf{v}) = \frac{1}{2} \begin{cases} \hat{f}_1(\mathbf{v}) + \hat{f}_2(\mathbf{v}) + \hat{f}_3(\mathbf{v}) - \hat{f}_4(\mathbf{v}), & u_1 = u_2 = 0, \\ \hat{f}_1(\mathbf{v}) - \hat{f}_2(\mathbf{v}) + \hat{f}_3(\mathbf{v}) + \hat{f}_4(\mathbf{v}), & u_1 = 0, u_2 = 1, \\ \hat{f}_1(\mathbf{v}) + \hat{f}_2(\mathbf{v}) - \hat{f}_3(\mathbf{v}) + \hat{f}_4(\mathbf{v}), & u_1 = 1, u_2 = 0, \\ \hat{f}_1(\mathbf{v}) - \hat{f}_2(\mathbf{v}) - \hat{f}_3(\mathbf{v}) - \hat{f}_4(\mathbf{v}), & u_1 = u_2 = 1. \end{cases}
$$
(5)

Choose an arbitrary **v** and construct the function $h \in \mathcal{F}_2$: $\stackrel{\circ}{h}(0,0) = 2^{-n/2} \hat{f}_1(\mathbf{v}),$ $^{\circ}$ $h(0,1) =$ $2^{-n/2} \hat{f}_2(\mathbf{v}),$ $\stackrel{\circ}{h}(1,0) = 2^{-n/2} \stackrel{\wedge}{f}_3(\mathbf{v}),$ ◦ $\hat{h}(1,1) = -2^{-n/2} \hat{f}_4(\mathbf{v})$. By [\(5\)](#page-8-0), the vth column of \hat{f} coincides with $2^{n/2-1} \hat{h}$. Hence f satisfies restrictions on columns and f is bent.

Observe that the function $f(u_1, u_2, y)$ can take all the values $0, \pm 2^{n/2-1}, \pm 2^{n/2}$ and the Rothaus' construction provides more subtle 4-row rectangles than ones considered in the previous example. \Box

Example 8 (Carlet's transformation). Let $f \in \mathcal{B}_{2n}$, $E \subset V_{2n}$ be an affine plane of dimension $k \geq n$, and $\phi_E \in \mathcal{F}_{2n}$ be the support of E, that is, $\phi_E(\mathbf{x}) = 1$ if only if $\mathbf{x} \in E$. In [\[6,](#page-18-8) p. 94], Carlet obtained conditions which provide the bentness of the function $g(\mathbf{x}) = f(\mathbf{x}) + \phi_E(\mathbf{x})$. Let us prove yet another condition: $g \in \mathcal{B}_{2n}$ if and only if $f_E \in \mathcal{P}_{k,2(k-n)}$, where f_E is a restriction of f to E, that is, $f_E(\mathbf{y}) = f(\varphi^{-1}(\mathbf{y}))$ for some affine bijection $\varphi: E \to V_k$.

Without loss of generality, assume that $E = \{(\mathbf{0}, \mathbf{a}) : \mathbf{a} \in V_k\}$ and $\varphi: (\mathbf{0}, \mathbf{a}) \mapsto \mathbf{a}$. Consider the rectangles $\overline{f} \in \mathring{B}_{2n-k,k}$ and $\overline{g} \in \mathring{F}_{2n-k,k}$. They differ only in the first row:

$$
\overset{\square}{f}(\mathbf{0},\mathbf{v})=\hat{f}_{E}(\mathbf{v}),\quad \overset{\square}{g}(\mathbf{0},\mathbf{v})=-\hat{f}_{E}(\mathbf{v}).
$$

Let $\hat{f}_E(\mathbf{v}) \neq 0$ for some **v**. Consider the normalized columns $\hat{h}(\mathbf{u}) = 2^{n-k} \hat{f}(\mathbf{u}, \mathbf{v})$ and $\hat{h}'(\mathbf{u}) =$ $2^{n-k} \overline{g}(\mathbf{u}, \mathbf{v})$. The functions $\hat{h} \in \mathcal{F}_{2n-k}$ and \hat{h}' differ only in signs of their (nonzero) values at $\mathbf{u} = \mathbf{0}$. Therefore, $\hat{h}' \in \hat{\mathcal{F}}_{2n-k}$ if and only if $\hat{h}(\mathbf{0}) = \pm 2^{2n-k}$. Under this condition,

$$
\hat{f}_E(\mathbf{v}) = \hat{f}(\mathbf{0}, \mathbf{v}) = 2^{k-n} \hat{h}(\mathbf{0}) = \pm 2^n.
$$

Consequently, \overline{g} is bent if and only if $\overline{\hat{f}}_E$ takes only the values $0, \pm 2^n$, that is, $f_E \in \mathcal{P}_{k,2(k-n)}$. \Box **Example 9** (normal bent functions). Following [\[8\]](#page-18-9), call a function $f \in \mathcal{B}_{2n}$ (weakly) normal if its restriction to some *n*-dimesional affine plane is affine. In other words, f is normal if there exists an affine equivalent function q such that

$$
M(g) = \max_{\mathbf{u},\mathbf{v}\in V_n} |\overset{\square}{g}(\mathbf{u},\mathbf{v})| = 2^n, \quad \overset{\square}{g} \in \overset{\square}{\mathcal{B}}_{n,n}.
$$

Using the affine classification of bent functions of 6 variables (see [\[14\]](#page-18-0)), one can check that every $f \in \mathcal{B}_6$ is normal. We give a direct proof of this fact by applying the results of Section [3.](#page-4-0)

Write $\varphi \sim \alpha_1^{d_1} \alpha_2^{d_2} \dots$ for a function φ that takes d_1 values α_1 , d_2 values α_2 , and so on. If $\hat{h} \in \hat{\mathcal{F}}_3$, then $|\hat{h}| \sim 8^10^7$ or $|\hat{h}| \sim 6^12^7$ or $|\hat{h}| \sim 4^40^4$. Since all rows and columns of $\hat{f} \in \mathring{B}_{3,3}$ are in $\hat{\mathcal{F}}_3$, we have $M(f) \in \{4, 6, 8\}.$

Let $M(f) = 4$. Applying the elementary transformations A1, A2 (i.e. permuting rows and columns) and B1, B2 (i.e. changing signs of rows and columns), we can arrange elements of \ddot{f} into one of the following cells:

$$
\begin{pmatrix} 4 & 4 \ 4 & 0 \end{pmatrix}, \quad \begin{pmatrix} 4 & 4 \ 4 & -4 \end{pmatrix}, \quad \begin{pmatrix} 4 & 4 \ 4 & 4 \end{pmatrix}.
$$

Then apply C1, C2:

$$
\begin{pmatrix} 4 & 4 \ 4 & 0 \end{pmatrix} \xrightarrow{C_1} \begin{pmatrix} 6 & 2 \ 2 & 2 \end{pmatrix}, \quad \begin{pmatrix} 4 & 4 \ 4 & -4 \end{pmatrix} \xrightarrow{C_1} \begin{pmatrix} 8 & 0 \ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 4 & 4 \ 4 & 4 \end{pmatrix} \xrightarrow{C_2} \begin{pmatrix} 4 & 4 \ 4 & -4 \end{pmatrix} \xrightarrow{C_1} \begin{pmatrix} 8 & 0 \ 0 & 0 \end{pmatrix},
$$

and obtain a rectangle that takes the value ≥ 6 . So it is sufficient to treat the case $M(f) = 6$ only. In this case we proceed in a similar manner:

$$
\xrightarrow{\text{A1},\text{A2},\text{B1},\text{B2},\text{C2}} \begin{pmatrix} 6 & 2 \\ 2 & -6 \end{pmatrix} \xrightarrow{\text{C1}} \begin{pmatrix} 8 & 0 \\ 0 & -4 \end{pmatrix},
$$

 \Box

and obtain a rectangle that takes the value 8, which is what we wanted.

Example 10 (the number of bent functions). If $g \in \mathcal{P}_{n,2}$, that is, $\hat{g}(\mathbf{a}_i) = 2^{n-1}\chi(b_i)$ for some $b_i \in \mathbb{F}_2$ and distinct $\mathbf{a}_i \in V_n$, $i = 1, 2, 3, 4$ $i = 1, 2, 3, 4$ $i = 1, 2, 3, 4$, then by Lemma 1

- (i) $a_1 + a_2 + a_3 + a_4 = 0$,
- (ii) $b_1 + b_2 + b_3 + b_4 = 1$.

In [\[2\]](#page-17-1), we proposed an algorithm to construct bent squares $\overline{f} \in \mathcal{B}_{n,n}$ all rows and columns of which are elements of $\hat{A}_n \cup \hat{P}_{n,2}$. The algorithm arranges nonzero elements in the matrix \overline{P} subject to the condition (i) and then places signs of elements subject to (ii).

Counting different outputs of the algorithm, we obtain constructive lower bounds for $|\mathcal{B}_{n,n}|$. In particular, the algorithm provides $1559994535674013286400 > 2^{70.4}$ distinct bent squares of $\mathcal{B}_{4,4}$ and, consequently, $|\mathcal{B}_8| > 2^{70.4}$. \Box

5 Biaffine and bilinear bent squares

Consider a mapping $\pi: V_n \times V_n \to V_n$. As previously, for a fixed **v** call the mapping $\mathbf{u} \mapsto \pi(\mathbf{u}, \mathbf{v})$ a restriction of π to **u**, and for a fixed **u** call the mapping $\mathbf{v} \mapsto \pi(\mathbf{u}, \mathbf{v})$ a restriction to **v**. Say that π is *biaffine* (*bilinear*) if all its restrictions to **u** and **v** are affine (linear) transformations of V_n . A biaffine mapping is *nonsingular* if all its restrictions are invertible. A bilinear mapping is nonsingular if all its restrictions to **u** for $v \neq 0$ and to **v** for $u \neq 0$ are invertible.

Choose an arbitrary function $g \in \mathcal{F}_n$ and construct a bent square $\overline{f} \in \overline{\mathcal{B}}_{n,n}$ such that the corresponding matrix \overline{F} consists almost only of the values of \hat{g} permuted in some order and multiplied by elements of \mathbb{Z}_q . Informally speaking, we "scatter" values of \hat{g} over \overline{F} . The connection [\(4\)](#page-4-2) between the Walsh–Hadamard coefficients of affine equivalent functions allows us to make such a scattering using a nonsingular biaffine or bilinear mapping. The obtained constructions are given by the following two easily verified propositions.

Proposition 3 (biaffine bent squares). Let π be a nonsingular biaffine mapping $V_n \times V_n \to V_n$, $g \in \mathcal{F}_n$, $\varphi \in \mathcal{F}_{2n}$, and all restrictions of $\varphi(\mathbf{u}, \mathbf{v})$ to $\mathbf{u} \in V_n$ and $\mathbf{v} \in V_n$ are affine functions. Then the square

$$
\overset{\square}{f}(\mathbf{u},\mathbf{v}) = \overset{\circ}{\varphi}(\mathbf{u},\mathbf{v}) \hat{g}(\pi(\mathbf{u},\mathbf{v}))
$$

is bent.

Proposition 4 (bilinear bent squares). Let π be a nonsingular bilinear mapping $V_n \times V_n \to V_n$, $g \in \mathcal{F}_n$, φ be defined as in the previous proposition, and $h, h' \in \mathcal{F}_n$ be such that

$$
\stackrel{\wedge}{h}(\mathbf{0})=\stackrel{\wedge}{h}'(\mathbf{0}), \quad \stackrel{\wedge}{h}(\mathbf{u})=\stackrel{\circ}{\varphi}(\mathbf{u},\mathbf{0})\stackrel{\wedge}{g}(\mathbf{0}), \quad \stackrel{\wedge}{h}'(\mathbf{v})=\stackrel{\circ}{\varphi}(\mathbf{0},\mathbf{v})\stackrel{\wedge}{g}(\mathbf{0}), \quad \mathbf{u},\mathbf{v}\in V_n\setminus\{\mathbf{0}\}.
$$

Then the square

$$
\overrightarrow{f}(\mathbf{u}, \mathbf{v}) = \begin{cases} \hat{h}(\mathbf{0}), & \mathbf{u} = \mathbf{v} = \mathbf{0}, \\ \varphi(\mathbf{u}, \mathbf{v}) \hat{g}(\pi(\mathbf{u}, \mathbf{v})) & otherwise, \end{cases}
$$

is bent.

Note that any function of the form

$$
\varphi(u_1,\ldots,u_n,v_1,\ldots,v_n)=\sum_{i,j=1}^n\alpha_{ij}u_iv_j+\sum_{i=1}^n\beta_iu_i+\sum_{i=1}^n\gamma_iv_i+\delta,\ \alpha_{ij},\beta_i,\gamma_j,\delta\in\mathbb{Z}_q,
$$

satisfies the condition of the above propositions. Note also that in Proposition [4](#page-10-1) we can easily construct h and h' if $\hat{g}(\mathbf{0}) = 0$. Indeed, in this case it is sufficiently to choose $h = h' \equiv c$, $c \in \mathbb{Z}_q$.

In Proposition [3](#page-10-2) we can use the following nonsingular biaffine mapping

$$
\pi(\mathbf{u}, \mathbf{v}) = \mathbf{u}A + \mathbf{v}B + (\mathbf{u}C_1\mathbf{v}^T, \dots, \mathbf{u}C_n\mathbf{v}^T) + \mathbf{d},
$$

where $\mathbf{d} \in V_n$ and A, B, C_1, \ldots, C_n are $n \times n$ matrices over \mathbb{Z}_q such that $A + (C_1 \mathbf{v}^T, \ldots, C_n \mathbf{v}^T)$ and $B + (C_1^T \mathbf{u}^T, \dots, C_n^T \mathbf{u}^T)$ are invertible for every **v** and **u**.

Further, each nonsingular bilinear mapping π has the form

$$
\pi(\mathbf{u}, \mathbf{v}) = \mathbf{u} A_{\mathbf{v}},\tag{6}
$$

where $A_{\mathbf{v}}$ are $n \times n$ matrices over \mathbb{Z}_q such that

- (i) $A_0 = 0$ and $A_v \in GL_n$ for all nonzero $v \in V_n$;
- (ii) $A_{\mathbf{v}+\mathbf{v}'} = A_{\mathbf{v}} + A_{\mathbf{v}'}$ for all $\mathbf{v}, \mathbf{v}' \in V_n$.

Indeed, considering restrictions of π to **u**, we get [\(6\)](#page-11-0). Since each such restriction for $v \neq 0$ must be invertible, $A_v \in GL_n$ for all nonzero v. For each fixed **u** the mapping $v \mapsto uA_v$ must be linear. Therefore, $\mathbf{u}A_{\mathbf{v}+\mathbf{v}'} = \mathbf{u}(A_{\mathbf{v}} + A_{\mathbf{v}'})$ for all v and v', which yields (ii) and the first part of (i).

Assume further that q is prime. The arising structure $R = \{A_v : v \in V_n\}$ is connected with some concepts of projective geometry. Consider the following linear subspaces of V_{2n} :

$$
E_{\infty} = \{ (\mathbf{0}, \mathbf{v}) : \mathbf{v} \in V_n \}, \quad E_{\mathbf{v}} = \{ (\mathbf{u}, \mathbf{u}A_{\mathbf{v}}) : \mathbf{u} \in V_n \}, \quad \mathbf{v} \in V_n. \tag{7}
$$

Each such subspace is of dimension n , two different subspaces intersect only in the zero vector and, consequently, the union of subspaces is V_{2n} . A set of subspaces with these properties is called a *spread* of V_{2n} .

Observe that the condition (ii) is too strong for the set [\(7\)](#page-11-1) to be a spread. We need only that $A - A' \in GL_n$ for all distinct $A, A' \in R$ (with the additional property $I_n \in R$ such a set R is called a quasifield).

Spreads are used in the following well-known construction of bent functions.

Example 11 (Dillon's construction). Consider a spread [\(7\)](#page-11-1) determined by a set $R = \{A_v : v \in$ V_n . Choose $c \in \mathbb{F}_q$ and $g \in \mathcal{F}_n$ such that $\hat{g}(0) = 0$. In [\[7\]](#page-18-1), Dillon actually proved that the function $f \in \mathcal{F}_{2n}$,

$$
f(\mathbf{x}) = \begin{cases} c, & \mathbf{x} \in E_{\infty}, \\ g(\mathbf{v}), & \mathbf{x} \in E_{\mathbf{v}} \setminus \{\mathbf{0}\}, \end{cases}
$$

is regular bent.

The corresponding bent square $\overrightarrow{f} \in \mathcal{B}_{n,n}$ has the form

$$
\overline{f}(\mathbf{u}, \mathbf{v}) = \begin{cases} q^n \chi(c), & \mathbf{u} = \mathbf{v} = \mathbf{0}, \\ \sum_{\mathbf{y} \in V_n} g(\mathbf{y}) \overline{\chi(\mathbf{u} A_{\mathbf{y}} \cdot \mathbf{v})} & \text{otherwise.} \end{cases}
$$

We see that the Dillon's bent squares are similar to the bilinear ones. In particular, if elements of R satisfy the conditions (i) and (ii), then the Dillon's construction is covered by Proposition [3](#page-10-2) under the choice $h = h' \equiv c, \varphi \equiv 0, \pi(\mathbf{u}, \mathbf{v}) = \mathbf{v} B_{\mathbf{u}}^{\mathrm{T}}$, where the matrices $B_{\mathbf{u}}$ are such that $uA_y = yB_u$ for all $u, y \in V_n$. \Box

Remark that the Maiorana–McFarland's bent squares can be represented in the form of Proposition [4](#page-10-1) with only the refinement that g is necessarily affine. The simple form of g in this case allows to relax the conditions of the proposition: we can use an arbitrary φ and choose $\pi(\mathbf{u}, \mathbf{v})$ such that all its restrictions to **u** and **v** are bijections (possibly not affine).

6 Partitions into affine planes

Let q be prime and L be a linear subspace of $V_n = \mathbb{F}_q^n$ of dimension r. Write the latter as $L < V_n$, dim $L = r$. Recall that the number of distinct r-dimensional subspaces of V_n is given by the Gaussian coefficient

$$
\binom{n}{r}_q = \frac{(q^n - 1)(q^{n-1} - 1)\dots(q^{n-r+1} - 1)}{(q^r - 1)(q^{r-1} - 1)\dots(q - 1)}, \quad 0 \le r \le n.
$$

Let $E = L + \mathbf{b}$ be an affine plane obtained by a shift of L by a vector $\mathbf{b} \in V_n$. The plane E is the image of the affine mapping $\pi: V_r \to V_n$, $\mathbf{w} \mapsto \mathbf{w}A + \mathbf{b}$, under a suitable choice of the $r \times n$ matrix A over \mathbb{F}_q of rank r.

Given $g \in \mathcal{F}_r$, construct the function $h \in \mathcal{F}_n$,

$$
h(\mathbf{x}) = g(\mathbf{x}A^{\mathrm{T}}) + \mathbf{b} \cdot \mathbf{x}, \quad \mathbf{x} \in V_n.
$$
 (8)

Call the transformation $g \mapsto h$ a *stretching* of g to the plane E. Under the stretching

$$
\hat{h}(\mathbf{v}) = \sum_{\mathbf{x} \in V_n} \hat{g}(\mathbf{x}A^{\mathrm{T}}) \chi((\mathbf{b} - \mathbf{v}) \cdot \mathbf{x})
$$

\n
$$
= q^{-r} \sum_{\mathbf{x} \in V_n} \sum_{\mathbf{w} \in V_r} \hat{g}(\mathbf{w}) \chi(\mathbf{x}A^{\mathrm{T}} \cdot \mathbf{w} + (\mathbf{b} - \mathbf{v}) \cdot \mathbf{x})
$$

\n
$$
= q^{-r} \sum_{\mathbf{w} \in V_r} \hat{g}(\mathbf{w}) \sum_{\mathbf{x} \in V_n} \chi((\pi(\mathbf{w}) - \mathbf{v}) \cdot \mathbf{x})
$$

and, consequently,

$$
\hat{h}(\mathbf{v}) = \begin{cases}\n q^{n-r} \hat{g}(\pi^{-1}(\mathbf{v})), & \mathbf{v} \in E, \\
 0 & \text{otherwise.} \n\end{cases}
$$
\n(9)

Describe bent rectangles $\overline{f} \in \overline{\mathcal{B}}_{m,n}$, $m = n - r$, which have all rows of the form [\(9\)](#page-12-2) and all columns in the set $q^{r/2} \hat{A}_m$. Such bent rectangles were first introduced by Carlet [\[4\]](#page-17-4) for the case $q = 2$. Note that after the transposition of f we obtain the rectangle $f^* \in \mathcal{B}_{n,m}$ that corresponds to the so called (see [\[12\]](#page-18-10)) partial affine bent function $f^* \in \mathcal{B}_{n+m}$: each restriction of $f^*(\mathbf{x}, \mathbf{y})$ to $\mathbf{x} \in V_n$ is affine.

Proposition 5 (partitions into affine planes). Let m, r be nonnegative integers, $n = m + r$, $\mathbf{u} \in V_m$, $\mathbf{v} \in V_n$, and

$$
\mathop{f}^{n}(\mathbf{u}, \mathbf{v}) = \begin{cases} q^{m} \mathop{g}^{n} \mathop{u}(\pi_{\mathbf{u}}^{-1}(\mathbf{v})), & \mathbf{v} \in E_{\mathbf{u}}, \\ 0 & otherwise, \end{cases}
$$

where $g_{\mathbf{u}} \in \mathcal{B}_r$ and $\pi_{\mathbf{u}}$ are mappings $V_r \to V_n$ such that $E_{\mathbf{u}} = \pi_{\mathbf{u}}(V_r)$ are affine planes of dimension r. If the planes ${E_{\bf u}}$ are disjoint and therefore determine a partition of V_n , then $\bar{f} \in$ $\bar{\mathcal{B}}_{m,n}$.

Example 12. Let $q = 2$. Choose the following partition of V_4 into the planes of dimension 2:

$$
E_{(0,0)} = \{(0,0,0,0), (0,0,0,1), (0,0,1,0), (0,0,1,1)\},
$$

\n
$$
E_{(0,1)} = \{(0,1,0,0), (0,1,0,1), (1,1,0,0), (1,1,0,1)\},
$$

\n
$$
E_{(1,0)} = \{(0,1,1,0), (0,1,1,1), (1,0,0,0), (1,0,0,1)\},
$$

\n
$$
E_{(1,1)} = \{(1,0,1,0), (1,0,1,1), (1,1,1,0), (1,1,1,1)\}.
$$

Using this partition, construct a rectangle $\overline{f} \in \overline{B}_{2,4}$ with the matrix

$$
\vec{F} = \left(\begin{array}{cccccccccccc} \pm 8 & \pm 8 & \pm 8 & \pm 8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \pm 8 & \pm 8 & 0 & 0 & 0 & 0 & 0 & \pm 8 & \pm 8 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \pm 8 & \pm 8 & \pm 8 & \pm 8 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \pm 8 & \pm 8 & 0 & 0 & \pm 8 & \pm 8 \end{array} \right),
$$

where signs of the elements are determined by unspecified bent functions $g_{\mathbf{u}} \in \mathcal{B}_2$.

It is convenient to assume that every one-element subset of a vector space is an affine plane of dimension 0 and that $\mathcal{B}_0 = \mathcal{F}_0$. Then under $r = 0$ the above proposition gives us all Maiorana–McFarland's bent squares.

In general, Proposition [5](#page-12-1) allows to construct

$$
(q^m)!c_q(n,m)|\mathcal{B}_{n-m}|^{q^m}
$$

distinct regular bent functions of $n + m$ variables. Here $c_q(n,m)$ is the number of distinct partitions of $V_n = \mathbb{F}_q^n$ into q^m affine planes of dimension $n - m$.

To obtain estimates for $c_q(n,m)$ consider some partition $\{E_1, E_2, \ldots, E_{q^m}\}\$ counted by $c_q(n,m)$. Let $E_i = L_i + \mathbf{b}_i$, where $L_i < V_n$, $\mathbf{b}_i \in V_n$, and

$$
W=L_1\cap L_2\cap\ldots\cap L_{q^m}.
$$

Call the partition ${E_i}$ primitive if $W = \{0\}$. Denote by $c_q^*(n,m)$ the number of distinct primitive partitions of V_n into planes of dimension $n - m$.

Suppose that ${E_i}$ is not primitive, that is, $d = \dim W \ge 1$. Then V_n can be represented as the direct sum $U \oplus W$, $U < V_n$, dim $U = n - d$, and each plane E_i takes the form $\{u + w : u \in$ $E'_i, \mathbf{w} \in W$, where $E'_i = E_i \cap U$ is an affine plane of U of dimension $n - m - d$. The planes $E'_1, E'_2, \ldots, E'_{q^m}$ determine a primitive partition of U. There are $c_q^*(n-d,m)$ ways to choose such a partition, $\binom{n}{d}$ $\binom{n}{d}$ ways to choose W and, consequently,

$$
c_q(n,m) = \sum_{d=0}^{n-m} \binom{n}{d}_q c_q^*(n-d,m).
$$
 (10)

 \Box

Denote by $L_i + L_j$ the subspace of V_n consisting of the sums $\mathbf{v} + \mathbf{v}'$, where \mathbf{v} runs over L_i and **v'** runs over L_j . For each distinct $i, j \in \{1, 2, ..., q^m\}$ we have

$$
\dim(L_i \cap L_j) = \dim L_i + \dim L_j - \dim(L_i + L_j) \ge 2(n - m) - n = n - 2m.
$$

If $\dim(L_i \cap L_j) = n - 2m$, then $L_i + L_j = V_n$ and $\mathbf{b}_j - \mathbf{b}_i = \mathbf{v} + \mathbf{v}'$ for some $\mathbf{v} \in L_i$, $\mathbf{v}' \in L_j$. It means that

$$
|E_i \cap E_j| = |L_i \cap (L_j + \mathbf{v} + \mathbf{v}')| = |(L_i - \mathbf{v}) \cap (L_j + \mathbf{v}')| = |L_i \cap L_j| \neq 0,
$$

a contradiction. Hence

$$
\dim(L_i \cap L_j) \ge n - 2m + 1. \tag{11}
$$

Under $m = 1$ this inequality yields that $L_1 = \ldots = L_q = W$, where dim $W = n - 1$. Therefore, $c_q^*(1,1) = 1$, $c_1^*(n,1) = 0$ for $n \ge 2$, and

$$
c_q(n,1) = {n \choose n-1}_q c_q^*(1,1) = \frac{q^n-1}{q-1}.
$$

For the case $q = 2$ we also obtain the estimate

$$
c_2(n,2) = {n \choose n-2}_2 + 98 {n \choose n-3}_2 = \frac{1}{3}(2^n - 1)(2^{n-1} - 1)(7 \cdot 2^{n-1} - 13)
$$

(cf. Example [6\)](#page-7-1) using the following result.

Lemma 2. $c_2^*(2,2) = 1$, $c_2^*(3,2) = 98$, and $c_2^*(n,2) = 0$ for $n \ge 4$.

Proof. The first equality is trivial. Further, if $q = 2$ then each partition of V_3 into two-element subsets is also the partition into affine planes. There are $\frac{8!}{4!2^4} = 105$ such distinct partitions and

$$
c_2^*(3,2) = 105 - \binom{3}{1}_2 c_2^*(2,2) = 98
$$

of them are primitive.

We prove the third equality by a contradiction. Suppose that $n \geq 4$ and $\{E_i = L_i + \mathbf{b}_i : i =$ 1, 2, 3, 4} is a primitive partition of V_n into planes of dimension $n-2$. Let us analyze the properties of such hypothetical partition which help to reveal a contradiction.

A. Let $U = L_1 + L_2 + L_3 + L_4$. Prove that $U = V_n$.

Note that dim $U \ge \dim L_1 = n - 2$. If $\dim U = n - 2$, then $L_1 = L_2 = L_3 = L_4$ and the partition ${E_i}$ is not primitive for $n > 3$.

Suppose that dim $U = n - 1$ and let, without loss of generality, $U = \{(\mathbf{x}, 0): \mathbf{x} \in V_{n-1}\}.$ Denote $U_i = E_i \cap U$. Since $\{E_i\}$ is a partition, $E_i = U_i + (0, \ldots, 0, b_i)$ for some $b_i \in \mathbb{F}_2$, where there are two 0's and two 1's among ${b_i}$. Assume for simplicity that $b_1 = b_2$ and therefore $b_3 = b_4$. Then $U_1 \cup U_2 = V_{n-1}$ and $U_3 \cup U_4 = V_{n-1}$. As we show later, it yields $L_1 = L_2$ and $L_3 = L_4$. Now using [\(11\)](#page-14-0) we obtain

$$
\dim(L_1 \cap L_2 \cap L_3 \cap L_4) = \dim(L_1 \cap L_3) \ge n - 3
$$

and the partition $\{E_i\}$ is not primitive for $n \geq 4$.

B. Prove that $d_{ij} = \dim(L_i \cap L_j) = n - 3$ for all $1 \leq i < j \leq 4$.

By [\(11\)](#page-14-0), the numbers $d_{ij} \in \{n-3, n-2\}$. Suppose that $d_{ij} = n-2$ for some i and j, say for $i = 3$ and $j = 4$. Then $L_3 = L_4$,

$$
\dim(L_1 \cap L_2 \cap L_3 \cap L_4) = \dim(L_1 \cap L_2 \cap L_3)
$$

=
$$
\dim(L_1 + L_2 + L_3) - \dim L_1 - \dim L_2 - \dim L_3
$$

+
$$
d_{12} + d_{13} + d_{23}
$$

$$
\geq n - 3(n - 2) + 3(n - 3) = n - 3,
$$

and the partition $\{E_i\}$ is not primitive for $n \geq 4$.

C. Prove that $d_{ijk} = \dim(L_i \cap L_j \cap L_k) \in \{n-4, n-3\}$ for all $1 \leq i < j < k \leq 4$ and moreover $d_{ijk} \neq n - 3$ for $n > 4$.

Indeed, $d_{ijk} \leq d_{ij} = n-3$ and

$$
d_{ijk} = \dim L_i + \dim(L_j \cap L_k) - \dim(L_i + (L_j \cap L_k))
$$

$$
\geq (n-2) + (n-3) - \dim(L_i + L_j) = n - 4.
$$

If some coefficient d_{ijk} is equal to $n-3$, say $d_{123} = n-3$, then

$$
\dim((L_1 \cap L_2 \cap L_3) + L_4) = d_{123} + \dim L_4 - \dim(L_1 \cap L_2 \cap L_3 \cap L_4)
$$

= $(n-3) + (n-2) + 0 = 2n - 5.$

On the other hand,

$$
\dim((L_1 \cap L_2 \cap L_3) + L_4) \le \dim(L_1 + L_4) = n - 1
$$

and we obtain the inequality $2n - 5 \leq n - 1$ that doesn't hold for $n > 4$.

Gathering A, B and C, we get

$$
\dim(L_1 \cap L_2 \cap L_3 \cap L_4) = \sum \dim L_i - \sum d_{ij} + \sum d_{ijk} - \dim \left(\sum L_i\right)
$$

= 4(n-2) - 6(n-3) + $\sum d_{ijk} - n$
 $\geq 4(n-2) - 6(n-3) + 4(n-4) - n = n - 6.$

Consequently, the partition $\{E_i\}$ is primitive only if $n \leq 6$ and $\sum d_{ijk} = 3n - 10$. Check these conditions.

If $n = 4$, then two coefficients d_{ijk} are equal to $n - 3 = 1$, say $d_{123} = d_{124} = 1$. Then $\dim(L_1 \cap L_2 \cap L_3 \cap L_4) = 1$, a contradiction. If $n = 5$, then there exists $d_{ijk} = n - 3 = 2$, a contradiction to the second part of C. Finally, if $n = 6$, then

$$
\dim(L_1 \cap L_3) = \dim L_1 + \dim L_3 - \dim(L_1 + L_3)
$$

\n
$$
\leq 8 - \dim((L_1 \cap L_2) + (L_3 \cap L_4))
$$

\n
$$
= 8 - (\dim(L_1 \cap L_2) + \dim(L_3 \cap L_4) - \dim(L_1 \cap L_2 \cap L_3 \cap L_4))
$$

\n
$$
= 8 - (3 + 3 - 0) = 2
$$

that contradicts B.

 \Box

The above lemma allows to account all partitions of V_{r+2} into planes of dimension r during the proof of the following result.

Proposition 6. Let $q = 2, r \geq 2, \tilde{f} \in \mathcal{B}_{2,r+2}$ be constructed by Proposition [5](#page-12-1) and $f(\mathbf{u}, \mathbf{x})$, $u \in V_2$, $x \in V_{r+2}$, be a corresponding bent function. By separate affine permutations of the variables \bf{u} and \bf{x} , \bf{f} can be transformed into one of the following functions:

$$
g(\mathbf{u}, \mathbf{x}) = u_1 u_2(g_1(x_3, \mathbf{y}) + g_2(x_3, \mathbf{y}) + g_3(x_3, \mathbf{y}) + g_4(x_3, \mathbf{y}))
$$
(12)
+ $u_1(g_1(x_3, \mathbf{y}) + g_3(x_3, \mathbf{y}) + x_1)$
+ $u_2(g_1(x_3, \mathbf{y}) + g_2(x_3, \mathbf{y}) + x_2)$
+ $g_1(x_3, \mathbf{y}),$

$$
g(\mathbf{u}, \mathbf{x}) = u_1 u_2(g_1(x_3, \mathbf{y}) + g_2(x_3, \mathbf{y}) + g_3(x_2, \mathbf{y}) + g_4(x_2, \mathbf{y}) + x_2 + x_3)
$$
(13)
+ $u_1(g_1(x_3, \mathbf{y}) + g_3(x_2, \mathbf{y}) + x_1)$
+ $u_2(g_1(x_3, \mathbf{y}) + g_2(x_3, \mathbf{y}) + x_2)$
+ $g_1(x_3, \mathbf{y}),$

$$
g(\mathbf{u}, \mathbf{x}) = u_1 u_2(g_1(x_3, \mathbf{y}) + g_2(x_1, \mathbf{y}) + g_3(x_1 + x_2 + x_3, \mathbf{y}) + g_4(x_2, \mathbf{y}) + x_1)
$$
(14)
+ $u_1(g_1(x_3, \mathbf{y}) + g_3(x_1 + x_2 + x_3, \mathbf{y}) + x_2 + x_3)$
+ $u_2(g_1(x_3, \mathbf{y}) + g_2(x_1, \mathbf{y}) + x_2)$
+ $g_1(x_3, \mathbf{y}),$

where $\mathbf{x} = (x_1, x_2, x_3, \mathbf{y}), \mathbf{y} \in V_{r-1}, q_i \in \mathcal{B}_r.$

Proof. Consider partitions of V_3 into planes of dimension 1. Denote such a plane $\{b, b + e\}$ by $[\mathbf{b}, \mathbf{e}]$. Let \mathbf{e}_i be the vector of V_3 having only the *i*th coordinate nonzero. It is easy to check that every partition of V_3 can be transformed into one of the following

$$
\begin{aligned} & \{ [\mathbf{0},\mathbf{e}_3], [\mathbf{e}_2,\mathbf{e}_3], [\mathbf{e}_1,\mathbf{e}_3], [\mathbf{e}_1+\mathbf{e}_2,\mathbf{e}_3] \}, \\ & \{ [\mathbf{0},\mathbf{e}_3], [\mathbf{e}_2,\mathbf{e}_3], [\mathbf{e}_1,\mathbf{e}_2], [\mathbf{e}_1+\mathbf{e}_3,\mathbf{e}_2] \}, \\ & \{ [\mathbf{0},\mathbf{e}_3], [\mathbf{e}_2,\mathbf{e}_1], [\mathbf{e}_2+\mathbf{e}_3,\mathbf{e}_1+\mathbf{e}_2+\mathbf{e}_3], [\mathbf{e}_1+\mathbf{e}_3,\mathbf{e}_2] \} \end{aligned}
$$

by an affine permutation of coordinates.

Choose some of these partitions, replace each its plane $[\mathbf{b}, \mathbf{e}] = {\mathbf{b} + \alpha \mathbf{e} : \alpha \in \mathbb{F}_2}$ by ${(b, 0) + \alpha(e, y): \alpha \in \mathbb{F}_2, y \in V_{r-1}}$ and obtain the partition of V_{r+2} into the planes of dimension r. Call such a partition canonical (for example, a canonical partition was used in Example [12\)](#page-13-0). Lemma [2](#page-14-1) implies that each partition of V_{r+2} can be transformed into some canonical by an affine permutation of coordinates. Also introduce a canonical ordering of planes of partitions by which the lexicographically minimal vectors of consecutive planes are increased (cf. Example [12\)](#page-13-0).

Let a bent rectangle \overrightarrow{f} be constructed by Proposition [5](#page-12-1) using a partition $\{E_{\mathbf{u}}\}$. By an affine permutation of columns of \overrightarrow{f} , the partition $\{E_{\mathbf{u}}\}$ can be transformed into some canonical and by an affine permutation of rows, the canonical ordering of planes can be achieved. Thus, it is sufficient to consider bent rectangles \overline{g} built by the canonical partitions ${E_{\bf{u}}}$ under the canonical ordering and determine the corresponding bent functions $q(\mathbf{u}, \mathbf{x})$. To do this, we can use the stretching equations [\(8\)](#page-12-3), [\(9\)](#page-12-2) to calculate the restrictions of q to **x** and then utilize the representation

$$
g(u_1, u_2, \mathbf{x}) = \sum_{\alpha_1, \alpha_2 \in \mathbb{F}_2} (u_1 + \alpha_1 + 1)(u_2 + \alpha_2 + 1)g(\alpha_1, \alpha_2, \mathbf{x}).
$$

For example, if g is built by the third canonical partition, then its restrictions to x look as follows:

$$
g(0, 0, \mathbf{x}) = g_1(x_3, \mathbf{y}),
$$

\n
$$
g(0, 1, \mathbf{x}) = g_2(x_1, \mathbf{y}) + x_2,
$$

\n
$$
g(1, 0, \mathbf{x}) = g_3(x_1 + x_2 + x_3, \mathbf{y}) + x_2 + x_3,
$$

\n
$$
g(1, 1, \mathbf{x}) = g_4(x_2, \mathbf{y}) + x_1 + x_3
$$

and we obtain [\(14\)](#page-16-0).

It is interesting that if $\overline{f} \in \mathring{B}_{2,r+2}$ is constructed by Proposition [5](#page-12-1) and $f_1(\mathbf{x})$, $f_2(\mathbf{x})$, $f_3(\mathbf{x})$, $f_4(\mathbf{x})$ are the restrictions of $f(\mathbf{u}, \mathbf{x})$ to x, then all the functions $f_i + f_j$, $1 \leq i < j \leq 4$, are balanced, that is, they take the values 0 and 1 equally often.

 \Box

Indeed, if $g(\mathbf{u}, \mathbf{x})$ has one of the forms $(12) - (14)$ $(12) - (14)$ $(12) - (14)$, then each sum of two its distinct restrictions to x is balanced. It is easily follows from the form of g and the fact that $h(\mathbf{x}) = h^*(x_1, \ldots, x_{k-1}, x_{k+1}, \ldots, x_{r+2}) + x_k$ and all derived functions $\sigma(h)(\mathbf{x}), \sigma \in \mathbf{AGL}_{r+2}$, are balanced. By Proposition [6,](#page-16-2) f can be obtained from q by separate affine permutations of \bf{u} and x . But a permutation of **u** only rearrange the restrictions, and a permutation of x doesn't change the balance of their sums.

References

- [1] C. A. Adams, S. E. Tavares, Generating and counting binary bent sequences, IEEE Trans. on Inform. Theory **IT-36** (1990), 1170-1173.
- [2] S. Agievich, On the representation of bent functions by bent rectangles, Probabilistic Methods in Discrete Mathematics: Fifth International Conference (Petrozavodsk, Russia, June 1–6, 2000), Utrecht, Boston: VSP, 2002, 121–135.
- [3] A. Canteaut, P. Charpin, Decomposing bent functions, IEEE Trans. on Inform. Theory IT-49 (2003), 2004–2019.
- [4] C. Carlet, On the confusion and diffusion properties of Maiorana–McFarland's and extended Maiorana–McFarland's functions, Special Issue "Complexity Issues in Coding and Cryptography", dedicated to Prof. H. Niederreiter on the occasion of his 60th birthday, J. of Complexity 20 (2004), p. 182–204.
- [5] C. Carlet, Recent results on binary bent functions, J. of Combinatorics, Information and System Sciences 24 (1999), 275–291.
- [6] C. Carlet, Two new classes of bent functions, Advances in Cryptology: Proceedings of EUROCRYPT'93 (Lofthus, Norway, May 23–27, 1993), Lecture Notes in Computer Science 765 (1994), Springer, 77–101.
- [7] J. Dillon, A survey of bent functions. NSA Technical Journal Special Issue (1972), 191–215.
- [8] H. Dobbertin, Construction of bent functions and balanced Boolean functions with high nonlinearity, Fast Software Encryption: Second International Workshop (Leuven, Belgium, December 14–16, 1994), Lecture Notes in Computer Science 1008 (1995), Springer, 61–74.
- [9] H. Dobbertin, G. Leander, A survey of some recent results on bent functions. Sequences and Their Applications: Third International Conference (SETA'2004, Seoul, Korea, October 24–28, 2004), Lecture Notes in Computer Science 3486 (2005), Springer, 1–29.
- [10] H. Dobbertin, G. Leander, Cryptographer's Toolkit for Construction of 8-Bit Bent Functions, Cryptology ePrint Archive, Report 2005/089, 2005, <http://eprint.iacr.org>.
- [11] P.V. Kumar, R.A. Scholtz, and L.R. Welch, Generalized bent functions and their properties, *J. of Comb. Theory* **A 40** (1985), 90–107.
- [12] O.A. Logachev, A.A. Salnikov, and V.V. Yashchenko, Bent functions and partitions of a Boolean cube, Formal Power Series and Algebraic Combinatorics: 12th International Conference (FPSAC'00, Moscow, Russia, June 26–30, 2000), 2000, 43–48 (In Russian).
- [13] B. Preneel, W. Van Leekwijck, L. Van Linden, R. Goevarts, and J. Vanderwalle, Propagation characteristics of Boolean functions, Advances in Cryptology: Proceedings of EU-ROCRYPT'90 (Aarhus, Denmark, May 21–24, 1990), Lecture Notes in Computer Science 437 (1991), Springer, 161–173.
- [14] O.S. Rothaus, On "bent" functions, *J. Comb. Theory* **A 20** (1976), 300–305.
- [15] B.L. van der Waerden, Algebra II, Springer-Verlag, Berlin-Heidelberg-New York, 1967.
- [16] Y. Zheng, X.M. Zhang, Plateaued functions, Information and Communication Security: Second International Conference (ICICS'99 Sydney, Australia, November 9–11, 1999), Lecture Notes in Computer Science 1726, Springer, 284–300.