

Disk single Hurwitz numbers.

S.M. Natanzon

1. INTRODUCTION.

The classical Hurwitz numbers are weighted numbers of coverings over a compact surface with prescribed types of critical values [5]. For present the full definition let us consider coverings with ramifications $\varphi : \Omega \rightarrow S$ over compact surface S of genus g by surfaces Ω . We consider that two coverings φ and $\varphi' : \Omega' \rightarrow S$ are equivalent if there exists a homeomorphism $f : \Omega \rightarrow \Omega'$ such that $\varphi'f = \varphi$. Denote by $\text{Aut}(\varphi)$ the group of autoequivalence of φ and by $|\text{Aut}(\varphi)|$ its order.

The covering φ is equivalent to a covering $z \mapsto z^{n(y)}$ on a neighborhood of any point $y \in \Omega$. Consider formal variables $\{a_i\}$. The monomial $a_{n(y_1)} \dots a_{n(y_k)}$, where $\varphi^{-1}(x) = \{y_1, \dots, y_k\}$, is called *type of the value* $x \in S$. A value of a type $a_2 a_1^k$ is called *simple*.

Fix now points $x_1, \dots, x_v \in S$. Fix also monomials a^1, \dots, a^v . *The (classical) Hurwitz number* is the sum $\langle a^1, \dots, a^v \rangle_g = \sum \frac{1}{|\text{Aut}(\varphi)|}$, by all equivalent classes of coverings, that have critical values of types a^1, \dots, a^v in the points x_1, \dots, x_v and have not other critical values. A Hurwitz number $\langle a \rangle^m = \langle a, a_2 a_1^k, \dots, a_2 a_1^k \rangle_0$, where $m+1$ is the number of critical value is called *classical single Hurwitz number*. The classical single Hurwitz number connect with the intersection theory on the moduli space of complex algebraic curves [4].

Correspond variables p_i to variables a_i and correspond monomials $p_a = p_{i_1} \dots p_{i_r}$ to monomials $a = a_{i_1} \dots a_{i_r}$. Generating function for classical single Hurwitz number is $\Phi(\lambda, p_1, p_2, \dots) = \sum_{m \geq 0} \frac{\lambda^m}{m!} \sum_a \langle a \rangle^m p_a$, where second sum is the sum by all monomials. According to [3], it satisfy to the "cut-and-join" differential equation

$$\frac{\partial \Phi}{\partial \lambda} = L_\lambda \Phi, \quad L_\lambda = \frac{1}{2} \sum_{ij} (i+j) p_i p_j \frac{\partial}{\partial p_{i+j}} + \sum_{ij} i j p_{i+j} \frac{\partial^2}{\partial p_i \partial p_j}.$$

This is a corollary from the fact that classical Hurwitz number generate a closed topological field theory [2]. The "cut-and-join" equation has some important properties. It is in particular a differential equation for generating function of Hodge integrals [6].

A definition of Hurwitz numbers for surfaces with boundary and non-orientable surfaces where suggested in [1]. In preset paper I define and investigate *disc single Hurwitz numbers*. They correspond to covering of disk with single non-simple boundary critical value. It is found, in particular, recursive equations defining all disk single Hurwitz numbers.

Generating function H for disc single Hurwitz numbers depend from complex parameters α, β (analog of λ from classical situation) and 4 infinite series of variables $\acute{p}_i, \grave{p}_i, \bar{p}_i, \dot{p}_i$, describing topological type of non-simple critical value (analog of p_i from classical situation). An analog of "cut-and-join" differential equation $\frac{\partial \Phi}{\partial \lambda} = L_\lambda \Phi$ is 2 differential equation, corresponding to interior and boundary simple critical values.

These differential equation follow, from the fact, that Hurwitz numbers for surfaces with boundary form a open-closed (non-commutative) topological field theory [1].

2. HURWITZ NUMBERS.

Let D be a closed disk $\{z \in \mathbb{C} \mid |z| \leq 1\}$ with oriented boundary ∂D . We will be consider coverings with ramification $\varphi : \Omega \rightarrow D$ of degree k , where Ω is a compact surface with a boundary $\partial\Omega$.

Preimage $\varphi^{-1}(x)$ of an interior point $x \in D^\circ = D \setminus \partial D$ consists of $n = n(x) \leq k$ points. Consider a small simple contour $r \in D^\circ$, around x . Its preimage $\varphi^{-1}(r)$ form simple contours $C_1, \dots, C_n \in \Omega^\circ$. The set of degrees of restrictions $(\deg(\varphi|_{C_1}), \dots, \deg(\varphi|_{C_n}))$ is called *(topological) type of interior value* $x \in D^\circ$. Correspond the monomial $a_1^{t_1} \cdots a_k^{t_k}$ to this topological type. Here a_i are commutated formal variables and t_i is the number of indexes j such that $\deg(\varphi|_{C_j}) = i$ and $a_i^0 = 1$.

Values of the type a_1^k are called non-critical. All other interior values are called *interior critical values*. Any covering has only finite number of interior critical values. Interior critical values of the type $a_1^{k-1}a_2$ are called *simple*.

A preimage $\varphi^{-1}(y)$ of boundary point $y \in \partial D$ also consists of $n = n(y) \leq k$ points. Consider a simple small interval $l \subset D$ around y with ends ∂D . The preimage $\varphi^{-1}(l) \subset \Omega$ form a graph with k edges. The vertexes of the graph form two groups that correspond to two ends of l . For convenience, we will call one of the ends and the groups "left" and other "right", considering that the moving on ∂D from left to right via y correspond to the orientation of ∂D .

The bipartite graph $\varphi^{-1}(l)$ is called *(topological) type of boundary critical value* $y \in \partial D$ [1]. The valency of any vertex of it is not more than 2. Thus connected components of $\varphi^{-1}(l)$ belong to one of the next type.

- \acute{b}_i is a graph with i left and $i + 1$ right vertexes;
- \grave{b}_i is a graph with $i + 1$ left and i right vertexes;
- \bar{b}_i is a graph with i left and i right vertexes;
- \grave{b}_i is a closed graph with i left and i right vertexes.

Correspond to the graphs $\acute{b}_i, \bar{b}_i, \grave{b}_i, \dot{b}_i$ commutative formal variables $\acute{b}_i, \bar{b}_i, \grave{b}_i, \dot{b}_i$. Correspond the monomial to join of graphs. Thus a topological of any boundary value is a monomial $b = \acute{b}_1^{s_1} \cdots \acute{b}_n^{s_n} \bar{b}_1^{s_1} \cdots \bar{b}_n^{s_n} \bar{b}_1^{\bar{s}_1} \cdots \bar{b}_n^{\bar{s}_n} \dot{b}_1^{s_1} \cdots \dot{b}_n^{s_n}$, where $k = \sum_{i=1}^n 2s_i + \sum_{i=1}^n (2\bar{s}_i - 1) + \sum_{i=1}^n 2\dot{s}_i + \sum_{i=1}^n 2\dot{s}_i$. Denote by $\text{Aut}(b)$ the group of automorphisms of the graph that correspond to the monomial b . Let $|\text{Aut}(b)|$ be the order of the group.

The changing of the order of the vertexes generates the involution $b \mapsto b^*$ on the set of monomials. In particular $\acute{b}_i^* = \grave{b}_i, \bar{b}_i^* = \bar{b}_i, \dot{b}_i^* = \dot{b}_i, \dot{b}_i^* = \dot{b}_i$.

The values of types $\bar{b}_i^{\bar{s}_i} \dot{b}_i^{s_i}$ are called non-critical. All other boundary values are called *boundary critical values*. Any covering has only finite number of boundary critical values. Boundary critical values of the type $\bar{b}_1 \bar{b}_1^{\bar{s}_1} \dot{b}_1^{s_1}$ and $\dot{b}_1 \bar{b}_1^{\bar{s}_1} \bar{b}_1^{s_1}$ are called *simple boundary critical values*. We call critical values of types $\bar{b}_1 \bar{b}_1^{\bar{s}_1} \dot{b}_1^{s_1}$ and $\dot{b}_1 \bar{b}_1^{\bar{s}_1} \bar{b}_1^{s_1}$ as *acute -points* and *grave-points* respectively.

Covering $\varphi_1 : \Omega_1 \rightarrow D$ and $\varphi_2 : \Omega_2 \rightarrow D$ are called equivalent if there exists a homeomorphism $\phi : \Omega_1 \rightarrow \Omega_2$ such that $\varphi_1 = \varphi_2 \phi$. The automorphisms groups $\text{Aut}(\varphi_i)$ of equivalent coverings are isomorphic. Denote by $|\text{Aut}(\varphi_i)|$ the order of its.

Fix interior points $x_1, \dots, x_v \in D \setminus \partial D$ and boundary points $y_1, \dots, y_w \in \partial D$ on a disk D . We consider that the numeration of the boundary points is convenient with the orientation of ∂D . Fix monomials a^1, \dots, a^v from the variable a_i and monomials

b^1, \dots, b^w from the variable $\acute{b}_i, \bar{b}_i, \grave{b}_i, \dot{b}_i$. The number $\langle a^1, \dots, a^v, (b^1, \dots, b^w) \rangle = \sum \frac{1}{|\text{Aut}(\varphi)|}$ is called *Hurwitz number*. Here the sum is taken by all equivalent classes of coverings, having critical values in points $x_1, \dots, x_v, y_1, \dots, y_w$ of types $a^1, \dots, a^v, b^1, \dots, b^w$ and have not other critical values. The Hurwitz number don't depend from position of critical points. It is kept by any permutation of a^i cyclic permutation of b^i . Denote by $\{F^{pq}\}$ the the inverse matrix to $\langle (\beta_p, \beta_q) \rangle$, where $\{\beta_p\}$ is the set of all connected graphs.

Lemma 2.1. *If $\langle (c, b) \rangle \neq 0$, then $c = b^*$ and $\langle (c, b) \rangle = \frac{1}{|\text{Aut}(b)|}$.*

If $\langle (c, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, b) \rangle \neq 0$, then this is one from the next cases

- $\langle (\acute{b}_i d^*, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \grave{b}_i d) \rangle = \frac{1}{2 |\text{Aut}(d)|}$;
- $\langle (\bar{b}_{i+j} d^*, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \bar{b}_i \grave{b}_j d) \rangle = \frac{1}{|\text{Aut}(d)|}$;
- $\langle (\acute{b}_{i+j} d^*, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \grave{b}_i \grave{b}_j d) \rangle = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\text{Aut}(d)|}$;
- $\langle (\grave{b}_{i+j-1} d^*, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \bar{b}_i \bar{b}_j d) \rangle = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\text{Aut}(d)|}$.

If $\langle (c, \grave{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, b) \rangle \neq 0$, then this is one from the next cases

- $\langle (\acute{b}_i d, \grave{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \acute{b}_i d^*) \rangle = \frac{1}{2 |\text{Aut}(d)|}$;
- $\langle (\bar{b}_i \acute{b}_j d, \grave{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \bar{b}_{i+j} d^*) \rangle = \frac{1}{|\text{Aut}(d)|}$;
- $\langle (\acute{b}_i \acute{b}_j d, \grave{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \acute{b}_{i+j} d^*) \rangle = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\text{Aut}(d)|}$;
- $\langle (\bar{b}_i \bar{b}_j d, \grave{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \acute{b}_{i+j-1} d^*) \rangle = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\text{Aut}(d)|}$.

Proof. The first statement is evident. Let $\langle (c, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, b) \rangle \neq 0$. Then there exists a covering $\varphi : \Omega \rightarrow D$ with boundary critical values $y_1, y_2, y_3 \in \partial D$ of types $c, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, b$ (respectively) and without other critical values. Let us consider points $z_1 \in (y_1, y_2)$ and $z_2 \in (y_2, y_3)$. The preimage of z_1 consists of \bar{m} simple points, where φ is local homeomorphism, and $\dot{m} + 1$ double points where φ is local two-sheeted. The preimage of z_2 consists of $\bar{m} + 2$ simple points and \dot{m} double points.

Let $p_1, p_2 \in \varphi^{-1}(z_2)$ be simple points, corresponding to \acute{b}_1 . They come to simple points q_1, q_2 of b . Then we have one from the next cases

- q_1 and q_2 belong to one connected component of type \acute{b}_i ;
- q_1 and q_2 belong to connected components of types \bar{b}_i and \dot{b}_j ;
- q_1 and q_2 belong to different connected components of types \bar{b}_i and \bar{b}_j ;
- q_1 and q_2 belong to different connected components of types \dot{b}_i and \dot{b}_j .

In the first case $b = \acute{b}_i d$. Let us consider the restriction $\varphi' : \Omega' D$ of φ on the connected component, containing the points q_1, q_2 . Consider the restriction $\varphi'' : \Omega'' D$ of φ on the complement $\Omega'' = \Omega \setminus \Omega'$. The covering φ'' has the critical values $y_1, y_3 \in \partial D$ of the types d^*, d . Therefore φ' has critical values $y_1, y_2, y_3 \in \partial D$ of types $\acute{b}_i, \acute{b}_1 e, \acute{b}_i$. Thus φ has critical values $y_1, y_2, y_3 \in \partial D$ of types $\acute{b}_i d^*, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \acute{b}_i d$.

The equivalent class of φ' (respectively φ'') contain of all coverings that have the same types of critical values as φ' (respectively φ''). Moreover, $\text{Aut}(\varphi) = \text{Aut}(\varphi') \times \text{Aut}(\varphi'')$. Thus $\langle (\acute{b}_i d^*, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \acute{b}_i d) \rangle = \frac{1}{|\text{Aut}(\varphi)|} = \frac{1}{|\text{Aut}(\varphi')| |\text{Aut}(\varphi'')|} = \frac{1}{|\text{Aut}(\varphi')| |\text{Aut}(d)|}$. The group $\text{Aut}(\varphi')$ is generated by involution that transpose the points p_1 and p_2 . Thus $\langle (\acute{b}_i d^*, \acute{b}_1 \bar{b}_1^{\bar{m}} \dot{b}_1^{\dot{m}}, \acute{b}_i d) \rangle = \frac{1}{2 |\text{Aut}(d)|}$.

Analogical arguments prove other cases. The main difference consists of properties of $\text{Aut}(\varphi')$. This group is non trivial only for $i = j$ in the last two cases.

Changing of orientation of the boundary of the disk we found that $\langle (a, b, c) \rangle = \langle (c^*, b^*, a^*) \rangle$. Thus the second statement of the lemma follow from the first.

□

We denote graphs \acute{b}_i and \grave{b}_i by \hat{b}_i in situations when differences between them are not important. Denote by $|b|$ and call length the number of edges in connected graph b . In particular, $|\bar{b}_i| = 2i - 1$ $|\hat{b}_i| = |\acute{b}_i| = |\grave{b}_i| = 2i$. Denote by $|b|$ the minimum of lengths of connected components of arbitrary graph b .

Lemma 2.2. *If $\langle a_1^m a_2, (c, b) \rangle \neq 0$, then this is one from the next cases*

- $\langle a_1^m a_2, (\acute{b}_i \acute{b}_j d^*, \acute{b}_{i+j} d) \rangle = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\text{Aut}(d)|}$;
- $\langle a_1^m a_2, (\acute{b}_i \bar{b}_j d^*, \bar{b}_{i+j} d) \rangle = \frac{|\bar{b}_j|}{|\text{Aut}(d)|}$;
- $\langle a_1^m a_2, (\grave{b}_i \acute{b}_j d^*, \acute{b}_{i+j} d) \rangle = \frac{|\acute{b}_j|}{|\text{Aut}(d)|}$;
- $\langle a_1^m a_2, (\acute{b}_i \acute{b}_j d^*, \grave{b}_{i+j} d) \rangle = \frac{|\acute{b}_j|}{|\text{Aut}(d)|}$;
- $\langle a_1^m a_2, (\bar{b}_i \acute{b}_j d^*, \bar{b}_k \acute{b}_l d) \rangle = \delta_{(i+j)(k+l)} \frac{|\bar{b}_i \acute{b}_j \bar{b}_k \acute{b}_l|}{|\text{Aut}(d)|}$;
- $\langle a_1^m a_2, (\acute{b}_i \bar{b}_j d^*, \bar{b}_k \bar{b}_l d) \rangle = \delta_{(i+j+1)(k+l)} \frac{|\acute{b}_i \bar{b}_j \bar{b}_k \bar{b}_l|}{|\text{Aut}(d)|}$;
- $\langle a_1^m a_2, (\acute{b}_i \acute{b}_j d^*, \acute{b}_k \acute{b}_l d) \rangle = (1 - \frac{\delta_{ij} \delta_{kl}}{2}) \delta_{(i+j)(k+l)} \frac{|\acute{b}_i \acute{b}_j \acute{b}_k \acute{b}_l|}{|\text{Aut}(d)|}$;
- $\langle a_1^m a_2, (\bar{b}_i \bar{b}_j d^*, \bar{b}_k \bar{b}_l d) \rangle = (1 - \frac{\delta_{ij} \delta_{kl}}{2}) \delta_{(i+j)(k+l)} \frac{|\bar{b}_i \bar{b}_j \bar{b}_k \bar{b}_l|}{|\text{Aut}(d)|}$;

Proof. Let $\langle a_1^m a_2, (c, b) \rangle \neq 0$. Then there exists a covering $\varphi : \Omega \rightarrow D$ with boundary critical values $x, y_1, y_2 \in \partial D$ of types $a_1^m a_2, c, b$ (respectively) and without other critical values. Let us consider a segment $l \subset \partial D$, connecting y_1 and y_2 by x . Consider also a simple small interval $l_i \subset D$ near y_i with ends on $\partial D \setminus (\partial D \cap U)$. Then the graph $b_i = \varphi^{-1}(l_i)$ is the topological type of y_i .

Consider a homotopy $\varphi : [1, 2] \rightarrow D$ between l_1 and l_2 such that $\varphi(t') \cap \varphi(t'') = \emptyset$ for $t' \neq t''$. Preimage $\varphi^{-1}(x_t)$ of $x_t = \varphi(t) \cap l$ consists of $\deg \varphi$ points if $x_t \neq x$ and consists of $\deg \varphi - 1$ points if $x_t = x$. This describe the reconstruction of the graph $b_t = \varphi^{-1}(l_t)$ in the moment, when $x_t \neq x$. The same reconstruction map the graph b_1 to the graph b_2^* . This consists of a cutting of 2 edges of b_1 and the gluing of its with the changing. For realization of this reconstruction by a covering $\varphi : \Omega \rightarrow D$ it is necessary that the orientation of the edges, that appear from order of vertexes, are generated by an orientation of ∂D . All pairs of graphs b_1, b_2 of such type and its reconstructions are presented in lemma 2.2. This gives also and corresponding Hurwitz numbers.

□

3. DISK SINGLE HURWITZ NUMBERS.

We will be say that a boundary critical values p' of a covering φ precedes a boundary critical values p'' of φ if the orientation of ∂D generate the orientation of (p', p'') from p' to p'' and (p', p'') does not contain any critical values.

Denote by $\mathcal{H}(m, \hat{m}, b)$ the sen of equivalent classe of covering with m interior simple critical values, with \hat{m} boundary simple critical values and with single non obviously simple critical value of type b (that is called *special*). The set $\mathcal{H}(m, \hat{m}, b)$ decompose into subsets $\mathcal{H}(m, \acute{m}, \grave{m}, b)$, that consists of coverings with m interior critical values, \acute{m} acute-points, \grave{m} grave-points. The set $\mathcal{H}(m, \acute{m}, \grave{m}, b)$ decompose into 2 subsets $\acute{\mathcal{H}}(m, \acute{m}, \grave{m}, b)$

and $\mathcal{H}(m, \acute{m}, \grave{m}, b)$. The first (respectively second) consists of covering where a acute-point (respectively a grave-point) precedes the special critical value.

Denote by $\mathsf{H}(m, \acute{m}, b)$ (respectively $\mathsf{H}(m, \acute{m}, \grave{m}, b)$, $\acute{\mathsf{H}}(m, \acute{m}, \grave{m}, b)$, $\grave{\mathsf{H}}(m, \acute{m}, \grave{m}, b)$) the sum of Hurwitz numbers from $\mathcal{H}(m, \acute{m}, b)$ (respectively $\mathcal{H}(m, \acute{m}, \grave{m}, b)$, $\acute{\mathcal{H}}(m, \acute{m}, \grave{m}, b)$, $\grave{\mathcal{H}}(m, \acute{m}, \grave{m}, b)$). Extend the definitions of the numbers $\mathsf{H}(m, \acute{m}, b)$, $\mathsf{H}(m, \acute{m}, \grave{m}, b)$, $\acute{\mathsf{H}}(m, \acute{m}, \grave{m}, b)$ and $\grave{\mathsf{H}}(m, \acute{m}, \grave{m}, b)$ on quotients of monomials $b = \frac{b^1}{b^2}$. Here we consider that these numbers are 0, if b is not monomial. The number $\mathsf{H}(m, \acute{m}, b)$ is called a *disk single Hurwitz number*.

Example 3.1. If $\mathsf{H}(0, 0, b) > 0$, then $b = \bar{b}_1^{\acute{s}_1} \dots \bar{b}_n^{\acute{s}_n} \bar{b}_1^{\grave{s}_1} \dots \bar{b}_n^{\grave{s}_n}$ and $\mathsf{H}(0, 0, b) = \prod_{i=1}^n \frac{1}{\acute{s}_i!} \prod_{i=1}^n \frac{1}{\grave{s}_i!}$.

Lemma 3.1. Let $b = \acute{b}_1^{\acute{s}_1} \dots \acute{b}_n^{\acute{s}_n} \acute{b}_1^{\grave{s}_1} \dots \acute{b}_n^{\grave{s}_n} \bar{b}_1^{\acute{s}_1} \dots \bar{b}_n^{\acute{s}_n} \bar{b}_1^{\grave{s}_1} \dots \bar{b}_n^{\grave{s}_n}$. Then $\acute{\mathsf{H}}(m, \acute{m} + 1, \grave{m}, b) = \sum_i \frac{i}{2} (\acute{s}_i + 1) \mathsf{H}(m, \acute{m}, \grave{m}, b \frac{\acute{b}_i}{b_i}) + \sum_{ij} ((\bar{s}_{i+j} + 1) \mathsf{H}(m, \acute{m}, \grave{m}, b \frac{\acute{b}_{i+j}}{b_i b_j}) + (\acute{s}_{i+j} + 1) \mathsf{H}(m, \acute{m}, \grave{m}, b \frac{\acute{b}_{i+j}}{b_i b_j}) + (\acute{s}_{i+j-1} + 1) \mathsf{H}(m, \acute{m}, \grave{m}, b \frac{\acute{b}_{i+j-1}}{b_i b_j}))$.

Proof. Consider a covering $\varphi : \Omega \rightarrow D$ from $\acute{\mathcal{H}}(m, \acute{m} + 1, \grave{m}, b)$. Denote by y and y' the special point and the precedes of it critical boundary point respectively. Collapse to a point y_l a segment $l \in D$ with ends on ∂D , that separate the points y, y' from other critical values. Then we have two disks D' and D'' and two coverings $\varphi' : \Omega' \rightarrow D'$, $\varphi'' : \Omega'' \rightarrow D''$. The critical values of φ' are y_l, y' and y . Comparing the disks D and D'' we see, that $\varphi'' \in \mathcal{H}(m, \acute{m}, \grave{m}, c)$ for some monomial c . Thus, according to [1], $\acute{\mathsf{H}}(m, \acute{m} + 1, \grave{m}, b) = \sum_{pq} H(m, \acute{m}, \grave{m}, \beta_p) > F^{pq} < (\beta_q, \acute{b}_1 \bar{b}_1^{\acute{n}} \acute{b}_1^{\acute{n}}, b) >$, where $\{\beta_p\}$ is the set of all bipartite graphs, $\{F^{pq}\}$ is the invert matrix to $< (\beta_p, \beta_q) >$ and the sum is given by all pairs of bipartite graphs. It is follow from lemma 2.1, that $F^{pq} = \delta_{\beta_p, \beta_q^*} = |\mathbf{Aut}(\beta_p)|$. Moreover it is follow from lemma 2.1, that the sum in the right parte of decompose to 4 subsumes. These subsumes depend from forms of b and $\beta_q = \beta_p^*$.

Let $b = \acute{b}_i d$ and $\beta_q = \acute{b}_i d^*$. Then $|\mathbf{Aut}(\beta_p)| = |\mathbf{Aut}(\acute{b}_i d)| = \frac{|\mathbf{Aut}(d)|}{i^{\acute{s}_i} \acute{s}_i!} i^{\acute{s}_i+1} (\acute{s}_i + 1)! = |\mathbf{Aut}(d)| i (\acute{s}_i + 1)$. Moreover $< (\beta_q, \acute{b}_1 \bar{b}_1^{\acute{n}} \acute{b}_1^{\acute{n}}, b) > = < (\acute{b}_i d^*, \acute{b}_1 \bar{b}_1^{\acute{n}} \acute{b}_1^{\acute{n}}, \acute{b}_i d) > = \frac{1}{2 |\mathbf{Aut}(d)|}$ according to lemma 2.1. Thus this subsume is $\sum_i \frac{i}{2} (\acute{s}_i + 1) \mathsf{H}(m, \acute{m}, \grave{m}, b \frac{\acute{b}_i}{b_i})$. Let $b = \bar{b}_i \acute{b}_j d$ and $\beta_q = \bar{b}_{i+j} d^*$. Then $|\mathbf{Aut}(\beta_p)| = |\mathbf{Aut}(\bar{b}_{i+j} d)| = \frac{|\mathbf{Aut}(d)|}{\bar{s}_{i+j}!} (\bar{s}_{i+j} + 1)! = |\mathbf{Aut}(d)| (\bar{s}_{i+j} + 1)$. Moreover $< (\beta_q, \acute{b}_1 \bar{b}_1^{\acute{n}} \acute{b}_1^{\acute{n}}, b) > = < (\bar{b}_{i+j} d^*, \acute{b}_1 \bar{b}_1^{\acute{n}} \acute{b}_1^{\acute{n}}, \bar{b}_i \acute{b}_j d) > = \frac{1}{|\mathbf{Aut}(d)|}$. Thus this subsume is $\sum_{ij} (\bar{s}_{i+j} + 1) \mathsf{H}(m, \acute{m}, \grave{m}, b \frac{\acute{b}_{i+j}}{b_i b_j})$. Let $b = \acute{b}_i \acute{b}_j d$ and $\beta_q = \acute{b}_{i+j} d^*$. Then $|\mathbf{Aut}(\beta_p)| = |\mathbf{Aut}(\acute{b}_{i+j} d)| = \frac{|\mathbf{Aut}(d)|}{2^{\acute{s}_{i+j}} \acute{s}_{i+j}!} 2^{(\acute{s}_{i+j}+1)} (\acute{s}_{i+j} + 1)! = 2 |\mathbf{Aut}(d)| (\acute{s}_{i+j} + 1)$. Moreover $< (\beta_q, \acute{b}_1 \bar{b}_1^{\acute{n}} \acute{b}_1^{\acute{n}}, b) > = < (\acute{b}_{i+j} d^*, \acute{b}_1 \bar{b}_1^{\acute{n}} \acute{b}_1^{\acute{n}}, \acute{b}_i \acute{b}_j d) > = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\mathbf{Aut}(d)|}$. Thus this subsume is $\sum_{ij} (\acute{s}_{i+j} + 1) \mathsf{H}(m, \acute{m}, \grave{m}, b \frac{\acute{b}_{i+j}}{b_i b_j})$. Let $b = \bar{b}_i \bar{b}_j d$ and $\beta_q = \bar{b}_{i+j-1} d^*$. Then $|\mathbf{Aut}(\beta_p)| = |\mathbf{Aut}(\bar{b}_{i+j-1} d)| = \frac{|\mathbf{Aut}(d)|}{2^{(\acute{s}_{i+j-1})} \acute{s}_{i+j-1}!} 2^{(\acute{s}_{i+j-1}+1)} (\acute{s}_{i+j-1} + 1)! = 2 |\mathbf{Aut}(d)| (\acute{s}_{i+j-1} + 1)$.

Moreover $\langle (\beta_q, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, b) \rangle = \langle (\acute{b}_{i+j-1} d^*, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \bar{b}_i \bar{b}_j d) \rangle = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\text{Aut}(d)|}$. Thus this subsume is $\sum_{ij} (\acute{s}_{i+j-1} + 1) \text{H}(m, \acute{m}, \acute{m}, b \frac{\acute{b}_{i+j-1}}{b_i b_j})$.

□

Lemma 3.2. Let $b = \acute{b}_1^{\acute{s}_1} \dots \acute{b}_n^{\acute{s}_n} \grave{b}_1^{\grave{s}_1} \dots \grave{b}_n^{\grave{s}_n} \bar{b}_1^{\bar{s}_1} \dots \bar{b}_n^{\bar{s}_1} \dots \bar{b}_1^{\bar{s}_n} \dots \bar{b}_n^{\bar{s}_n}$. Then $\text{H}(m, \acute{m}, \acute{m} + 1, b) = \sum_i (\acute{s}_i + 1) \text{H}(m, \acute{m}, \acute{m}, b \frac{\acute{b}_i}{b_i}) + \sum_{ij} (2(\bar{s}_i + 1)(\acute{s}_j + 1) \text{H}(m, \acute{m}, \acute{m}, b \frac{\bar{b}_i \acute{b}_j}{b_i b_j}) + 2((\acute{s}_i + 1)(\acute{s}_j + 1) + \delta_{ij}(\acute{s}_i + 1)) \text{H}(m, \acute{m}, \acute{m}, b \frac{\bar{b}_i \acute{b}_j}{b_{i+j}}) + \frac{1}{2}((\bar{s}_i + 1)(\bar{s}_j + 1) + \delta_{ij}(\bar{s}_i + 1)) \text{H}(m, \acute{m}, \acute{m}, b \frac{\bar{b}_i \bar{b}_j}{b_{i+j-1}}))$.

Proof. The proof of this lemma is the same that the previous, using the second part of lemma 2.1. If $(\beta_q, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, b) = (\acute{b}_i d^*, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \acute{b}_i d)$, Then $|\text{Aut}(\beta_p)| = |\text{Aut}(\acute{b}_i d)| = 2|\text{Aut}(d)|(\acute{s}_i + 1)$ and $\langle (\beta_q, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, b) \rangle = \langle (\acute{b}_i d^*, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \acute{b}_i d) \rangle = \frac{1}{2|\text{Aut}(d)|}$. Thus this subsume is $\sum_i (\acute{s}_i + 1) \text{H}(m, \acute{m}, \acute{m}, b \frac{\acute{b}_i}{b_i})$.

If $(\beta_q, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, b) = (\bar{b}_i \acute{b}_j d^*, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \bar{b}_{i+j} d)$, then $|\text{Aut}(\beta_p)| = |\text{Aut}(\bar{b}_i \acute{b}_j d)| = 2|\text{Aut}(d)|(\bar{s}_i + 1)(\acute{s}_j + 1)$ and $\langle (\beta_q, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, b) \rangle = \langle (\bar{b}_i \acute{b}_j d^*, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \bar{b}_{i+j} d) \rangle = \frac{1}{|\text{Aut}(d)|}$.

Thus this subsume is $2 \sum_{ij} (\bar{s}_i + 1)(\acute{s}_j + 1) \text{H}(m, \acute{m}, \acute{m}, b \frac{\bar{b}_i \acute{b}_j}{b_{i+j}})$. If

$(\beta_q, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, b) = (\acute{b}_i \acute{b}_j d^*, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \acute{b}_{i+j} d)$, then $|\text{Aut}(\beta_p)| = |\text{Aut}(\acute{b}_i \acute{b}_j d)| = 4|\text{Aut}(d)|((\acute{s}_i + 1)(\acute{s}_j + 1) + \delta_{ij}(\acute{s}_i + 1))$ and $\langle (\acute{b}_i \acute{b}_j d^*, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \acute{b}_{i+j} d) \rangle = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\text{Aut}(d)|}$

Thus this subsume is $2 \sum_{ij} ((\acute{s}_i + 1)(\acute{s}_j + 1) + \delta_{ij}(\acute{s}_i + 1)) \text{H}(m, \acute{m}, \acute{m}, b \frac{\acute{b}_i \acute{b}_j}{b_{i+j}})$.

If $(\beta_q, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, b) = (\bar{b}_i \bar{b}_j d^*, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \acute{b}_{i+j-1} d)$, then $|\text{Aut}(\beta_p)| = |\text{Aut}(\bar{b}_i \bar{b}_j d)| = |\text{Aut}(d)|((\bar{s}_i + 1)(\bar{s}_j + 1) + \delta_{ij}(\bar{s}_i + 1))$ and $\langle (\beta_q, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, b) \rangle = \langle (\bar{b}_i \bar{b}_j d, \acute{b}_1 \bar{b}_1^n \acute{b}_1^n, \acute{b}_{i+j-1} d^*) \rangle = (1 - \frac{\delta_{ij}}{2}) \frac{1}{|\text{Aut}(d)|}$. Thus this subsume is $\frac{1}{2} \sum_{ij} ((\bar{s}_i + 1)(\bar{s}_j + 1) + \delta_{ij}(\bar{s}_i + 1)) \text{H}(m, \acute{m}, \acute{m}, b \frac{\bar{b}_i \bar{b}_j}{b_{i+j-1}})$.

□

The previous lemmas give

Theorem 3.1. Let $b = \acute{b}_1^{\acute{s}_1} \dots \acute{b}_n^{\acute{s}_n} \grave{b}_1^{\grave{s}_1} \dots \grave{b}_n^{\grave{s}_n} \bar{b}_1^{\bar{s}_1} \dots \bar{b}_n^{\bar{s}_1} \dots \bar{b}_1^{\bar{s}_n} \dots \bar{b}_n^{\bar{s}_n}$. Then $\text{H}(m, \acute{m} + 1, b) = \sum_i \frac{i}{2} (\acute{s}_i + 1) \text{H}(m, \acute{m}, b \frac{\acute{b}_i}{b_i}) + \sum_{ij} ((\bar{s}_{i+j} + 1) \text{H}(m, \acute{m}, b \frac{\bar{b}_{i+j}}{b_i b_j}) + (\acute{s}_{i+j} + 1) \text{H}(m, \acute{m}, b \frac{\acute{b}_{i+j}}{b_i b_j}) + (\acute{s}_{i+j-1} + 1) \text{H}(m, \acute{m}, b \frac{\acute{b}_{i+j-1}}{b_i b_j})) + \sum_i (\acute{s}_i + 1) \text{H}(m, \acute{m}, b \frac{\acute{b}_i}{b_i}) + \sum_{ij} (2(\bar{s}_i + 1)(\acute{s}_j + 1) \text{H}(m, \acute{m}, b \frac{\bar{b}_i \acute{b}_j}{b_{i+j}}) + 2((\acute{s}_i + 1)(\acute{s}_j + 1) + \delta_{ij}(\acute{s}_i + 1)) \text{H}(m, \acute{m}, b \frac{\bar{b}_i \acute{b}_j}{b_{i+j}}) + \frac{1}{2}((\bar{s}_i + 1)(\bar{s}_j + 1) + \delta_{ij}(\bar{s}_i + 1)) \text{H}(m, \acute{m}, b \frac{\bar{b}_i \bar{b}_j}{b_{i+j-1}}))$.

Analogical theorem for interior points is

Theorem 3.2. Let $b = \acute{b}_1^{\acute{s}_1} \dots \acute{b}_n^{\acute{s}_n} \grave{b}_1^{\grave{s}_1} \dots \grave{b}_n^{\grave{s}_n} \bar{b}_1^{\bar{s}_1} \dots \bar{b}_n^{\bar{s}_1} \dots \bar{b}_1^{\bar{s}_n} \dots \bar{b}_n^{\bar{s}_n}$.

Then $\text{H}(m + 1, \acute{m}, b) =$

$$\sum_{ij} (\frac{i+j}{2} (\acute{s}_{i+j} + 1) \text{H}(m, \acute{m}, b \frac{\acute{b}_{i+j}}{b_i b_j}) + \frac{ij}{2} ((\acute{s}_i + 1)(\acute{s}_j + 1) + \delta_{ij}(\acute{s}_i + 1)) \text{H}(m, \acute{m}, b \frac{\acute{b}_i \acute{b}_j}{b_{i+j}}) + |\bar{b}_j| (\bar{s}_{i+j} + 1) \text{H}(m, \acute{m}, b \frac{\bar{b}_{i+j}}{b_i b_j}) + i |\bar{b}_j| (\acute{s}_i + 1) (\bar{s}_j + 1) \text{H}(m, \acute{m}, b \frac{\bar{b}_i \bar{b}_j}{b_{i+j}}) +$$

$$\begin{aligned}
& 2|\hat{b}_j|(\hat{s}_{i+j} + 1)\mathbb{H}(m, \hat{m}, \hat{m}, b_{\frac{\hat{b}_i \hat{b}_j}{\hat{b}_i \hat{b}_j}}) + 2i|\hat{b}_j|(\hat{s}_i + 1)(\hat{s}_j + 1)\mathbb{H}(m, \hat{m}, \hat{m}, b_{\frac{\hat{b}_i \hat{b}_j}{\hat{b}_{i+j}}}) + \\
& 2|\hat{b}_j|(\hat{s}_{i+j} + 1)\mathbb{H}(m, \hat{m}, b_{\frac{\hat{b}_i \hat{b}_j}{\hat{b}_i \hat{b}_j}}) + 2i|\hat{b}_j|(\hat{s}_i + 1)(\hat{s}_j + 1)\mathbb{H}(m, \hat{m}, b_{\frac{\hat{b}_i \hat{b}_j}{\hat{b}_{i+j}}}) + \\
& \sum_{i+j=k+l} (2|\bar{b}_i \bar{b}_j \bar{b}_k \bar{b}_l|(\bar{s}_i + 1)(\hat{s}_j + 1)\mathbb{H}(m, \hat{m}, b_{\frac{\bar{b}_i \bar{b}_j}{\bar{b}_k \bar{b}_l}}) + 2|\bar{b}_i \bar{b}_j \bar{b}_k \bar{b}_l|(\bar{s}_i + 1)(\hat{s}_j + 1)\mathbb{H}(m, \hat{m}, b_{\frac{\bar{b}_i \bar{b}_j}{\bar{b}_k \bar{b}_l}})) + \\
& \sum_{i+j+1=k+l} 2(1 + \delta_{kl})|\bar{b}_i \bar{b}_j \bar{b}_k \bar{b}_l|(\hat{s}_i + 1)(\hat{s}_j + 1)\mathbb{H}(m, \hat{m}, b_{\frac{\bar{b}_i \bar{b}_j}{\bar{b}_k \bar{b}_l}}) + \\
& \sum_{i+j=k+l+1} \frac{1}{2}(1 + \delta_{ij})|\bar{b}_i \bar{b}_j \bar{b}_k \bar{b}_l|((\bar{s}_i + 1)(\bar{s}_j + 1) + \delta_{ij}(\bar{s}_i + 1))\mathbb{H}(m, \hat{m}, b_{\frac{\bar{b}_i \bar{b}_j}{\bar{b}_k \bar{b}_l}}) + \\
& \sum_{i+j=k+l} (1 + \delta_{ij})(1 + \delta_{kl}) (\frac{1}{4}|\bar{b}_i \bar{b}_j \bar{b}_k \bar{b}_l|((\bar{s}_i + 1)(\bar{s}_j + 1) + \delta_{ij}(\bar{s}_i + 1))\mathbb{H}(m, \hat{m}, b_{\frac{\bar{b}_i \bar{b}_j}{\bar{b}_k \bar{b}_l}}) + \\
& |\hat{b}_i \hat{b}_j \hat{b}_k \hat{b}_l|((\hat{s}_i + 1)(\hat{s}_j + 1) + \delta_{ij}(\hat{s}_i + 1))\mathbb{H}(m, \hat{m}, b_{\frac{\hat{b}_i \hat{b}_j}{\hat{b}_k \hat{b}_l}}) + \\
& |\hat{b}_i \hat{b}_j \hat{b}_k \hat{b}_l|((\hat{s}_i + 1)(\hat{s}_j + 1) + \delta_{ij}(\hat{s}_i + 1))\mathbb{H}(m, \hat{m}, b_{\frac{\hat{b}_i \hat{b}_j}{\hat{b}_k \hat{b}_l}})),
\end{aligned}$$

Proof. Consider a covering $\varphi : \Omega \rightarrow D$ from $\mathcal{H}(m + 1, \hat{m}, b)$. Denote by y and x the special critical point and an interior critical point respectively. Collapse to a point y_l a segment $l \in D$ with ends on ∂D , that separate the points y, x from other critical values. Then we have two disks D' and D'' and two coverings $\varphi' : \Omega' \rightarrow D'$, $\varphi'' : \Omega'' \rightarrow D''$. The critical values of φ' are y_l, y and x . Comparing the disks D and D'' we see, that $\varphi'' \in \mathcal{H}(m, \hat{m}, c)$ for some monomial c . Thus, according to [1], $\hat{H}(m + 1, \hat{m}, b) = \sum_{pq} H(m, \hat{m}, \beta_p) > F^{pq} < (\beta_q, \hat{b}_1 \bar{b}_1 \hat{b}_1^{\hat{n}}, b >$.

It is follow from lemma 2.2 that the sum from the right parte decompose on 15 subsumes. Any subsume is calculated similar as in lemmas 3.1 and 3.2.

□

Theorems 3.1, 3.2 give an algorithm for calculation of all Hurwitz numbers $\mathbb{H}(m, \hat{m}, b)$, starting from $\mathbb{H}(0, 0, b)$.

4. DIFFERENTIAL EQUATIONS ON GENERATING THE FUNCTION.

Consider the algebra of formal power series, generated by commutative variables $\dot{p}_i, \dot{\bar{p}}_i, \bar{p}_i, \dot{\bar{p}}_i$. Correspond the monomial $p_b = \dot{p}_1^{\hat{s}_1} \dots \dot{p}_n^{\hat{s}_n} \dot{\bar{p}}_1^{\hat{s}_1} \dots \dot{\bar{p}}_n^{\hat{s}_n} \bar{p}_1^{\bar{s}_1} \dots \bar{p}_n^{\bar{s}_n} \bar{p}_1^{\bar{s}_1} \dots \bar{p}_n^{\bar{s}_n}$ to monomial $b = \hat{b}_1^{\hat{s}_1} \dots \hat{b}_n^{\hat{s}_n} \hat{b}_1^{\hat{s}_1} \dots \hat{b}_n^{\hat{s}_n} \bar{b}_1^{\bar{s}_1} \dots \bar{b}_n^{\bar{s}_n} \bar{b}_1^{\bar{s}_1} \dots \bar{b}_n^{\bar{s}_n}$.

Consider the generating function

$$H(\alpha, \beta | \hat{b}_1, \hat{b}_1, \bar{b}_1, \bar{b}_1, \hat{b}_2, \dots) = \sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \mathbb{H}(m, \hat{m}, b) p_b,$$

where the second sum is the sum by all monomial b .

In particulary, $H(0, 0 | \dots) = \Sigma \rho(p) p$, where the sum is given by all monomial in the form $p = \bar{p}_1^{\bar{s}_1} \dots \bar{p}_n^{\bar{s}_n} \dot{\bar{p}}_1^{\hat{s}_1} \dots \dot{\bar{p}}_n^{\hat{s}_n}$ and $\rho(p) = \prod_{i=1}^n \frac{1}{\bar{s}_i!} \prod_{i=1}^n \frac{1}{\hat{s}_i!}$. Moreover the invariance of Hurwitz numbers by changing of orientation of ∂D gives the invariance of H by changing $\hat{b}_i \leftrightarrow \bar{b}_i$ for all i .

Theorem 4.1.

$$\frac{\partial H}{\partial \beta} = L_{\hat{\beta}} H + L_{\bar{\beta}} H,$$

where

$$L_{\beta} = \sum_i \frac{i}{2} \dot{p}_i \frac{\partial}{\partial \dot{p}_i} + \sum_{ij} (\bar{p}_i \dot{p}_j \frac{\partial}{\partial \bar{p}_{i+j}} + \dot{p}_i \dot{p}_j \frac{\partial}{\partial \dot{p}_{i+j}} + \bar{p}_i \bar{p}_j \frac{\partial}{\partial \dot{p}_{i+j-1}})$$

and

$$L_{\beta} = \sum_{i=1}^{\infty} \dot{p}_i \frac{\partial}{\partial \dot{p}_i} + \sum_{ij} (2\bar{p}_{i+j+1} \frac{\partial^2}{\partial \bar{p}_i \partial \dot{p}_j} + 2\dot{p}_{i+j} \frac{\partial^2}{\partial \dot{p}_i \partial \dot{p}_j} + \frac{1}{2} \dot{p}_{i+j-1} \frac{\partial^2}{\partial \bar{p}_i \partial \bar{p}_j}).$$

Proof. Let $b = \dot{b}_1^{\dot{s}_1} \dots \dot{b}_{\dot{n}}^{\dot{s}_{\dot{n}}} \bar{b}_1^{\bar{s}_1} \dots \bar{b}_{\bar{n}}^{\bar{s}_{\bar{n}}} \dot{b}_1^{\dot{s}_1} \dots \dot{b}_{\dot{n}}^{\dot{s}_{\dot{n}}}$, where $\dot{n} = \dot{n}(b)$, $\dot{s}_i = \dot{s}_i(b)$, $\bar{n} = \bar{n}(b)$, $\bar{s}_i = \bar{s}_i(b)$, $\dot{n} = \dot{n}(b)$, $\dot{s}_i = \dot{s}_i(b)$.

It is follow from theorem 3.1 that

$$\begin{aligned} \frac{\partial H}{\partial \beta} &= \sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \mathcal{H}(m, \hat{m} + 1, b) p_b = \\ &\sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \sum_{i=1}^{\infty} \frac{i}{2} (\dot{s}_i + 1) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i}^{\dot{b}_i}) p_b + \\ &\sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} (\bar{s}_{i+j} + 1) \mathcal{H}(m, \hat{m}, b_{\bar{b}_i \bar{b}_j}^{\bar{b}_i \bar{b}_j}) p_b + \\ &\sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} (\dot{s}_{i+j} + 1) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i \dot{b}_j}^{\dot{b}_i \dot{b}_j}) p_b + \\ &\sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} (\dot{s}_{i+j-1} + 1) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i \dot{b}_j}^{\dot{b}_i \dot{b}_j - 1}) p_b + \\ &\sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \sum_{i=1}^{\infty} (\dot{s}_i + 1) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i}^{\dot{b}_i}) p_b + \\ &2 \sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} (\bar{s}_i + 1) (\dot{s}_j + 1) \mathcal{H}(m, \hat{m}, b_{\bar{b}_i \dot{b}_j}^{\bar{b}_i \dot{b}_j}) p_b + \\ &2 \sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \sum_{i=1}^{\infty} ((\dot{s}_i + 1) (\dot{s}_j + 1) + \delta_{ij} (\dot{s}_i + 1)) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i \dot{b}_j}^{\dot{b}_i \dot{b}_j}) p_b + \\ &\frac{1}{2} \sum_{m, \hat{m} \geq 0} \frac{\alpha^m \beta^{\hat{m}}}{m! \hat{m}!} \sum_b \sum_{i=1}^{\infty} ((\bar{s}_i + 1) (\bar{s}_j + 1) + \delta_{ij} (\bar{s}_i + 1)) \mathcal{H}(m, \hat{m}, b_{\bar{b}_i \bar{b}_j}^{\bar{b}_i \bar{b}_j - 1}) p_b. \end{aligned}$$

Make the changing $c = b_{\dot{b}_i}^{\dot{b}_i}$ in the first summand. Then

$$\begin{aligned} p_c &= p_b \frac{\dot{p}_i}{\dot{p}_i} \text{ and } p_b = p_c \frac{\dot{p}_i}{\dot{p}_i} = \dot{p}_i \frac{\partial p_c}{\partial \dot{p}_i} (\dot{s}_i + 1)^{-1}. \text{ Thus} \\ \sum_b \sum_{i=1}^{\infty} \frac{i}{2} (\dot{s}_i + 1) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i}^{\dot{b}_i}) p_b &= \sum_c \sum_{i=1}^{\infty} \frac{i}{2} \mathcal{H}(m, \hat{m}, a) \dot{p}_i \frac{\partial p_c}{\partial \dot{p}_i}, \text{ where the sum is} \\ &\text{given by all monomial } c. \end{aligned}$$

Make the changing $c = b_{\bar{b}_i \bar{b}_j}^{\bar{b}_i \bar{b}_j}$ in the second summand. Then

$$\begin{aligned} p_c &= p_b \frac{\bar{p}_{i+j}}{\bar{p}_i \bar{p}_j} \text{ and } p_b = p_c \frac{\bar{p}_i \bar{p}_j}{\bar{p}_{i+j}} = \bar{p}_i \bar{p}_j \frac{\partial p_c}{\partial \bar{p}_{i+j}} (\bar{s}_{i+j} + 1)^{-1}. \text{ Thus} \\ \sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} (\bar{s}_{i+j} + 1) \mathcal{H}(m, \hat{m}, b_{\bar{b}_i \bar{b}_j}^{\bar{b}_i \bar{b}_j}) p_b &= \sum_c \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \mathcal{H}(m, \hat{m}, c) \bar{p}_i \bar{p}_j \frac{\partial p_c}{\partial \bar{p}_{i+j}}. \end{aligned}$$

Analogically, $\sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} (\dot{s}_{i+j} + 1) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i \dot{b}_j}^{\dot{b}_i \dot{b}_j}) p_b =$

$$\begin{aligned} &\sum_c \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \mathcal{H}(m, \hat{m}, c) \dot{p}_i \dot{p}_j \frac{\partial p_c}{\partial \dot{p}_{i+j}} \text{ and } \sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} (\dot{s}_{i+j-1} + 1) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i \dot{b}_j}^{\dot{b}_i \dot{b}_j - 1}) p_b = \\ &\sum_c \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \mathcal{H}(m, \hat{m}, c) \bar{p}_i \bar{p}_j \frac{\partial p_c}{\partial \dot{p}_{i+j-1}}. \text{ Thus the sum of the first four summands is } L_{\beta} H. \end{aligned}$$

Make the changing $c = b_{\dot{b}_i}^{\dot{b}_i}$ in the fifth summand. Then $p_c = p_b \frac{\dot{p}_i}{\dot{p}_i}$

$$\begin{aligned} \text{and } p_b &= p_c \frac{\dot{p}_i}{\dot{p}_i} = \dot{p}_i \frac{\partial p_c}{\partial \dot{p}_i} (\dot{s}_i + 1)^{-1}. \text{ Thus } \sum_b \sum_{i=1}^{\infty} (\dot{s}_i + 1) \mathcal{H}(m, \hat{m}, b_{\dot{b}_i}^{\dot{b}_i}) p_b = \\ &\sum_c \sum_{i=1}^{\infty} \mathcal{H}(m, \hat{m}, c) \dot{p}_i \frac{\partial p_c}{\partial \dot{p}_i}, \text{ where the sum is given by all monomial} \\ &c. \end{aligned}$$

$$\begin{aligned} \text{Analogically, } \sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} (\bar{s}_i + 1) (\dot{s}_j + 1) \mathcal{H}(m, \hat{m}, b_{\bar{b}_i \dot{b}_j}^{\bar{b}_i \dot{b}_j}) p_b &= \\ \sum_c \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \mathcal{H}(m, \hat{m}, c) \bar{p}_{i+j+1} \frac{\partial^2 p_c}{\partial \bar{p}_i \partial \dot{p}_j}. \end{aligned}$$

Make the changing $c = b_{\bar{b}_i \dot{b}_j}^{\bar{b}_i \dot{b}_j}$ in the seventh summand. Then

$$p_c = p_b \frac{\dot{p}_i \dot{p}_j}{\dot{p}_{i+j-1}} \text{ and } p_b = p_c \frac{\dot{p}_{i+j-1}}{\dot{p}_i \dot{p}_j} = \dot{p}_{i+j-1} \frac{\partial^2 p_c}{\partial \bar{p}_i \partial \dot{p}_j} ((\dot{s}_i + 1) (\dot{s}_j + 1) + \delta_{ij} (\dot{s}_i + 1))^{-1}.$$

Thus $\sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} ((\dot{s}_i + 1)(\dot{s}_j + 1) + \delta_{ij}(\dot{s}_i + 1)) \mathbb{H}(m, \dot{m}, \dot{m}, b_{\frac{\dot{b}_i \dot{b}_j}{\dot{b}_{i+j-1}}}) p_b = \sum_c \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \mathbb{H}(m, \dot{m}, \dot{m}, c) \dot{p}_{i+j-1} \frac{\partial^2 p_c}{\partial \dot{p}_i \partial \dot{p}_j}$. Analogically, $\sum_b \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} ((\bar{s}_i + 1)(\bar{s}_j + 1) + \delta_{ij}(\bar{s}_i + 1)) \mathbb{H}(m, \dot{m}, \dot{m}, b_{\frac{\bar{b}_i \bar{b}_j}{\bar{b}_{i+j-1}}}) p_b = \sum_c \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \mathbb{H}(m, \dot{m}, \dot{m}, c) \dot{p}_{i+j-1} \frac{\partial^2 p_c}{\partial \bar{p}_i \partial \bar{p}_j}$. Thus the sum of the last four summands is $L_{\beta} H$.

□

Theorem 4.2.

$$\frac{\partial H}{\partial \alpha} = L_{\alpha} H,$$

where

$$\begin{aligned} L_{\alpha} = & \sum_{ij} \left(\frac{i+j}{2} \dot{p}_i \dot{p}_j \frac{\partial}{\partial \dot{p}_{i+j}} + \frac{ij}{2} \dot{p}_{i+j} \frac{\partial^2}{\partial \dot{p}_i \partial \dot{p}_j} + |\bar{b}_j| \dot{p}_i \bar{p}_j \frac{\partial}{\partial \bar{p}_{i+j}} + i |\bar{b}_j| \bar{p}_{i+j} \frac{\partial^2}{\partial \dot{p}_i \partial \bar{p}_j} + \right. \\ & 2 |\hat{b}_j| \dot{p}_i \dot{p}_j \frac{\partial}{\partial \dot{p}_{i+j}} + 2i |\hat{b}_j| \dot{p}_{i+j} \frac{\partial^2}{\partial \dot{p}_i \partial \dot{p}_j} + 2 |\hat{b}_j| \dot{p}_i \dot{p}_j \frac{\partial}{\partial \dot{p}_{i+j}} + 2i |\hat{b}_j| \dot{p}_{i+j} \frac{\partial^2}{\partial \dot{p}_i \partial \dot{p}_j} \Big) + \\ & \sum_{i+j=k+l} 2 (|\bar{b}_i \hat{b}_j \bar{b}_k \hat{b}_l| \bar{p}_k \dot{p}_l \frac{\partial^2}{\partial \bar{p}_i \partial \dot{p}_j} + |\bar{b}_i \hat{b}_j \bar{b}_k \hat{b}_l| \bar{p}_k \dot{p}_l \frac{\partial^2}{\partial \bar{p}_i \partial \dot{p}_j}) + \\ & \sum_{i+j+1=k+l} 2(1 + \delta_{kl}) |\hat{b}_i \hat{b}_j \bar{b}_k \bar{b}_l| \bar{p}_k \bar{p}_l \frac{\partial^2}{\partial \dot{p}_i \partial \bar{p}_j} + \sum_{i+j=k+l+1} \frac{1}{2} (1 + \delta_{ij}) |\bar{b}_i \bar{b}_j \hat{b}_k \hat{b}_l| \dot{p}_k \dot{p}_l \frac{\partial^2}{\partial \bar{p}_i \partial \bar{p}_j} + \\ & \sum_{i+j=k+l} (1 + \delta_{ij})(1 + \delta_{kl}) \left(\frac{1}{4} |\bar{b}_i \bar{b}_j \bar{b}_k \bar{b}_l| \bar{p}_k \bar{p}_l \frac{\partial^2}{\partial \bar{p}_i \partial \bar{p}_j} + |\hat{b}_i \hat{b}_j \hat{b}_k \hat{b}_l| \dot{p}_k \dot{p}_l \frac{\partial^2}{\partial \dot{p}_i \partial \dot{p}_j} + |\hat{b}_i \hat{b}_j \hat{b}_k \hat{b}_l| \dot{p}_k \dot{p}_l \frac{\partial^2}{\partial \dot{p}_i \partial \dot{p}_j} \right) \end{aligned}$$

The proof use theorem 3.2 and is given by the same method that the proof of theorem 4.1.

REFERENCES

- [1] Alexeevski A., Natanzon S., Noncommutative two-dimensional topological field theories and Hurwitz numbers for real algebraic curves. *Selecta Math.*, New ser. v.12,n.3, 2006, p. 307-377 (arXiv: math.GT/0202164).
- [2] Dijkgraaf R., Mirror symmetry and elliptic curves, *The moduli spaces of curves*, Progress in Math., 129 (1995), 149-163, Birkhäuser.
- [3] Goulden D., Jackson D.M., Vainshtein A., The number of ramified coverings of the sphere by torus and surfaces of higher genera, *Ann. of Comb.* 4(2000), 27-46, Birkhäuser.
- [4] Ekhidal T., Lando S., Shapiro M., Vainshtein A., Hurwitz numbers and intersections on moduli spaces of curves, *Invent.Math.* 146(2001), 297-327
- [5] Hurwitz A., Über Riemann'sche Flächen mit gegeben Verzweigungspunkten, *Math., Ann.*, Bn.39 (1891), 1-61.
- [6] Zhou J., Hodge integrals Hurwitz numbers and symmetric groups. (arXiv: math.AG/0308024).