## Quantum MERA Channels

V. Giovannetti,<sup>1</sup> S. Montangero,<sup>1</sup> and Rosario Fazio<sup>1,2</sup>

<sup>1</sup>NEST CNR-INFM & Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy

<sup>2</sup>International School for Advanced Studies (SISSA), Via Beirut 2-4, I-34014 Trieste, Italy

(Dated: November 7, 2018)

Tensor networks representations of many-body quantum systems can be described in terms of quantum channels. We focus on channels associated with the Multi-scale Entanglement Renormalization Ansatz (MERA) tensor network that has been recently introduced to efficiently describe critical systems. Our approach allows us to compute the MERA correspondent to the thermodynamic limit of a critical system introducing a transfer matrix formalism, and to relate the system critical exponents to the convergence rates of the associated channels.

PACS numbers: 03.67.-a,05.30.-d,89.70.-a

Understanding the properties of strongly interacting many-body quantum system is central in many areas of physics. Whenever it is hard to device reliable analytical approaches, as in many situations of experimental relevance, ingenious numerical methods are necessary to grasp the essential properties of these systems. Our ability of simulating them is based on the possibility to find an efficient description of their ground state. This is the case, for example, of White's Density Matrix Renormalization Group [1] which can be recasted in terms of Matrix Product States (MPS) [2, 3, 4, 5, 6]. Such representations are characterized by a simple tensor decomposition of the many-body wave-function which allows one i) to efficiently compute all the relevant observables of the system (e.g. energy, local observables, and correlation functions), and *ii*) to reduce the effective number of parameters over which the numerical optimization needs to be performed. MPS fulfill these requirements and can be used to describe faithfully the ground states of not critical, short range one-dimensional many-body Hamiltonians at zero-temperature. However MPS typically fail to provide an accurate description in other relevant situations, i.e. when the system is critical, in higher physical dimensions or if the model possesses longrange couplings. Several proposals have been put forward to overcome this problem. Projected Entangled Pair States (PEPS) [7] generalize MPS in dimensions higher than one. Weighted graph states [8] can deal with long-range correlations. In this Letter we focus on a solution recently proposed by Vidal [9] who introduced a tensor structure based on the so called Multiscale Entanglement Renormalization Ansatz (MERA). The MERA tensor network satisfies both the constraints i) and ii) and accommodates the scale invariance typical of critical systems [10, 11]. The relevance of this approach might represent a major breakthrough in our simulation capabilities [12] and motivates an intensive study of the MERA [13, 14].

Here we point out a previously unnoticed connection between the MERA and the theory of completely positive quantum maps [15] establishing a link between two important areas of quantum information science. This allows us to introduce a transfer matrix formalism in the same spirit as it has been done for MPS [3, 5], providing new tools to compute physical observables using MERA. The main outcomes of our work are i) a method for determining the properties of critical many-body systems in the thermodynamic limit and ii) a connection between the critical exponents governing the decay of correlation functions and the eigenvalues of the MERA transfer matrix. As a consequence this yields a full characterization of the asymptotic properties of one-dimensional critical systems. The paper is organized as follows: after a brief review of the MERA, we show how the local expectation values and correlations functions can be casted in terms of concatenated quantum channels. Using general properties of *mixing* quantum channels [16, 17, 18, 19] we then provide a way for expressing the thermodynamic limit of these quantities.

The MERA tensor network:- Consider a many-body quantum system composed by  $N = 2^n$  sites of dimension d (qudits). Any pure state can be expressed as  $|\psi\rangle = \sum \mathcal{T}_{\ell_1,\ell_2,\cdots,\ell_N} |\xi_{\ell_1},\xi_{\ell_2},\cdots,\xi_{\ell_N}\rangle$ , where for  $j \in$  $\{1, \cdots, N\}$  and  $\ell \in \{1, \cdots, d\}$  the vectors  $|\xi_{\ell}\rangle_i \in \mathcal{H}_d$ form the basis of the j-th d-dimensional system component. The MERA representation [9] of  $|\psi\rangle$  assumes a specific tensor decomposition of  $\mathcal{T}$  described in Fig. 1. Here the links emerging from the lowest part of the graph represent the N physical indices of  $\mathcal{T}$ . The nodes of the graph instead represent tensors. They are divided in three groups: the type- $\begin{pmatrix} 2\\2 \end{pmatrix}$  disentangler tensors  $\chi$  of elements  $\chi_{\ell_1,\ell_2}^{u_1,u_2}$  represented by the red Xs; the type- $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$  tensors  $\lambda$  of elements  $\lambda_{\ell_1,\ell_2}^{u_1}$  represented by the blue inverted Ys; and the type  $\begin{pmatrix} 0\\4 \end{pmatrix}$  tensor  $\mathcal{C}$  of elements  $\mathcal{C}_{\ell_1,\ell_2,\ell_3,\ell_4}$ , represented by the green semi-circle. As shown in Fig. 1 the  $\chi$ 's, the  $\lambda$ 's are coupled together to form a triangular structure with C as the closing element of the top: any two joined legs from any two distinct nodes indicate saturation of the associated indices. Consequently, the tensor  $\mathcal{T}$  associated with the N qudit state  $|\psi\rangle$  is written as a network of O(N) tensors organized in  $O(\log_2 N)$  different levels composed by one layer of  $\chi$  ten-




FIG. 1: Representation of a MERA decomposition of the tensor  $\mathcal{T}$  associated with a many-body state  $|\psi\rangle$  for N = 16 qudits (a dotted link emerging from the left side of the graph re-enter the figure as the corresponding dotted link on the right). The light blue region represents the causal cone [9] associated with the local operator  $\hat{\Theta}_j$  (black rectangle). It is composed by contraction of tensors  $\mathcal{M}_5$  (shown in the inset).

sors connected with one layer of  $\lambda$  tensors. In a generic MERA the  $\chi$ 's and the  $\lambda$ 's differ from node to node but the dimensions of their indices are upper-bounded by a fixed constant (for easy of notation we omit the labels expressing the position of the tensors in the network).

What makes the MERA decomposition a convenient one is the assumption that the  $\chi$ 's and  $\lambda$ 's satisfy special contraction rules. Specifically, for each  $\chi$  and  $\lambda$  let us define its adjoint  $\bar{\chi}$  and  $\bar{\lambda}$  as the tensors of elements  $\bar{\chi}_{u_1,u_2}^{\ell_1,\ell_2} = (\chi_{\ell_1,\ell_2}^{u_1,u_2})^*$  and  $\bar{\lambda}_{\ell_1}^{u_1,u_2} = (\lambda_{u_1,u_2}^{\ell_1})^*$ . With this definition the MERA contraction rules are  $\bar{\chi}_{\ell_1,\ell_2}^{\bullet,\diamond} \chi_{\bullet,\diamond}^{u_1,u_2} =$  $\chi_{u_1,u_2}^{\bullet,\diamond} = \bar{\chi}_{\ell_1,u_1} \delta_{\ell_2,u_2}$  and  $\bar{\lambda}_{\ell_1}^{\bullet,\diamond} \lambda_{\bullet,\diamond}^{u_1} = \delta_{\ell_1,u_1}$ , with  $\delta$  being the Kronecker delta and where the typographic symbols  $\bullet$  and  $\diamond$  indicate summation over the corresponding index. Under these conditions the expectation values of local observables on  $|\psi\rangle$  requires only to evaluate  $O(\log_2 N)$  non trivial tensor contractions [9].

Local observables and quantum channels:- We first show how the average of local observables can be related to the study of concatenated quantum channels. Given an operator  $\hat{\Theta}_j$  which acts not trivially on no more than three consecutive qudits (say the (j - 1)th, *j*th and (j + 1)th), the quantity  $\langle \hat{\Theta}_j \rangle \equiv \langle \psi | \hat{\Theta}_j | \psi \rangle$  requires to perform contractions only over the  $\chi$ 's and  $\lambda$ 's belonging to the causal cone [9] of the triple j - 1, j and j + 1. A compact expression is obtained by grouping these tensors in compounds composed by 2  $\chi$ 's and by 3  $\lambda$ 's (see inset of Fig. 1). This forms  $m = \log_2(N/4)$  non necessarily identical type- $\begin{pmatrix} 3\\6 \end{pmatrix}$  tensors  $\mathcal{M}_5 \equiv \lambda \chi \lambda \chi \lambda$ where the products  $\lambda \chi$  and  $\chi \lambda$  are defined by  $[\lambda \chi]_{\ell_1,\ell_2,\ell_3}^{u_1,u_2} \equiv \lambda_{\ell_1,\bullet}^{u_1} \chi_{\ell_2,\ell_3}^{\bullet,u_2}$ , and  $[\chi \lambda]_{\ell_1,\ell_2,\ell_3}^{u_1,u_2} \equiv \chi_{\ell_1,\ell_2}^{u_1,\bullet} \lambda_{\bullet,\ell_3}^{\bullet,u_2}$ . For each one of the m tensors  $\mathcal{M}_5$  we can then introduce two families of operators  $\{\hat{L}_r\}_r$  and  $\{\hat{R}_r\}_r$ acting on the Hilbert space  $\mathcal{H}_d^{\otimes 3}$  and labeled through

the composed index  $r \equiv (r_1, r_2, r_3)$  with  $r_{1,2,3}$  being d-dimensional [20]. In the computational basis they are defined by the matrices  $\langle \xi_{u_1}, \xi_{u_2}, \xi_{u_3} | \tilde{L}_r | \xi_{\ell_1}, \xi_{\ell_2}, \xi_{\ell_3} \rangle$ and  $\langle \xi_{u_1}, \xi_{u_2}, \xi_{u_3} | \hat{R}_r | \xi_{\ell_1}, \xi_{\ell_2}, \xi_{\ell_3} \rangle$  of elements  $[\mathcal{M}_5]^{u_1, u_2, u_3}_{r_1, \ell_1, \ell_2, \ell_3, r_2, r_3}$  and  $[\mathcal{M}_5]^{u_1, u_2, u_3}_{r_1, r_2, \ell_1, \ell_2, \ell_3, r_3}$  respectively. They are related through a reshuffling of the input and output qudits, i.e.  $\hat{R}_r = \Pi(\hat{L}_r) \equiv \hat{P}\hat{L}_r\hat{P}^{\dagger}$ , where  $\hat{P} = \hat{P}^{\dagger}$ is the unitary transformation which exchanges the first and the third qudit. Most importantly, according to the contraction rules defined previously, they satisfy the normalization conditions  $\sum_r \hat{L}_r \hat{L}_r^{\dagger} = \hat{I}^{\otimes 3} = \sum_r \hat{R}_r \hat{R}_r^{\dagger}$ with  $\hat{I}$  being the identity operator of  $\mathcal{H}_d$ . This implies that  $\{\hat{L}_r\}_r$  can be used to define a completely positive, unital, not necessarily trace preserving super-operators  $\Phi_{H}^{(L)}$  [15], which transforms the linear operator  $\hat{\Theta}$  of  $H_d^{\otimes 3}$  into  $\Phi_H^{(L)}(\hat{\Theta}) = \sum_r \hat{L}_r \hat{\Theta} \hat{L}_r^{\dagger}$ . Analogously  $\{\hat{R}_r\}_r$  defines the map  $\Phi_H^{(R)}$  which is related with  $\Phi_H^{(L)}$  through the identity  $\Phi_H^{(R)} = \Pi \circ \Phi_H^{(L)} \circ \Pi$ , where " $\circ$ " indicates the composition of super-operators. We also introduce the vector of  $\mathcal{H}_d^{\otimes 4}$ ,  $|hat\rangle \equiv \sum C_{\ell_1,\ell_2,\ell_3,\ell_4} |\xi_{\ell_1},\xi_{\ell_2},\xi_{\ell_3},\xi_{\ell_4}\rangle$ , which without loss of generality is assumed to be normalized, and define  $\hat{\rho}_C$  the three sites reduced density matrix obtained by tracing  $|hat\rangle\langle hat|$  over one of the 4 qudits. With these definitions one can finally write the expectation value of  $\hat{\Theta}_j$  as  $\langle \hat{\Theta}_j \rangle = \text{Tr}[\hat{\rho}_C \ \hat{B}_j^{(m)}]$ , with  $\hat{B}_{j}^{(m)} \equiv \Phi_{H}^{(m)} \circ \cdots \circ \Phi_{H}^{(1)}(\hat{\Theta}_{j}), \text{ and where (enumerating})$ from the lower MERA level of Fig. 1)  $\Phi_H^{(k)}$  is either the map  $\Phi_H^{(L)}$  or  $\Phi_H^{(R)}$  associated with the *k*-th tensor  $\mathcal{M}_5$  of the causal cone (which one depends upon N and j). The operator  $\hat{B}_{j}^{(m)}$  is thus obtained by applying to the observable  $\hat{\Theta}_j$  a sequence of *m* super-operators associated to the MERA causal cone. We can then write

$$\langle \hat{\Theta}_j \rangle = \text{Tr}[\Phi^{(1 \leftarrow m)}(\hat{\rho}_C) \ \hat{\Theta}_j],$$
 (1)

where  $\Phi^{(1 \leftarrow m)} \equiv \Phi^{(1)} \circ \cdots \circ \Phi^{(m)}$ , with  $\Phi^{(k)}$  being the super-operator  $\Phi_H^{(k)}$  in Schrödinger picture. By construction the  $\Phi^{(k)}$  (and hence  $\Phi^{(1 \leftarrow m)}$ ) are Completely Positive, Trace Preserving (CPT) maps, i.e. quantum channels with Kraus operators [15] defined by either the set  $\{\hat{L}_r^{\dagger}\}_r$  or  $\{\hat{R}_r^{\dagger}\}_r$ . Equation (1) establishes a formal equivalence between the MERA tensor network and the successive application of a family of CPT maps (the QuMERA family). Since it holds for *all* the local observable  $\hat{\Theta}_j$  this implies that  $\Phi^{(1 \leftarrow m)}(\hat{\rho}_C)$  coincides with the reduced density matrix  $\hat{\rho}_j$  of the input state  $|\psi\rangle$  associated with the qudits j - 1, j and j + 1, i.e.

$$\Phi^{(1 \leftarrow m)}(\hat{\rho}_C) = \hat{\rho}_j . \tag{2}$$

Critical systems in the thermodynamic limit: – Consider now a family of MERA states  $|\psi\rangle$  which describes the ground state of a many-body Hamiltonian  $\hat{H}$  at criticality. We are specifically interested in the thermodynamic limit of infinitely many sites (i.e.  $N \rightarrow$   $\infty$ ). According to the above derivation, in the limit of large m Eq. (2) converges toward the reduced density matrix  $\hat{\rho}_T$  of three consecutive qubits of the systems  $\lim_{m\to\infty} \Phi^{(1\leftarrow m)}(\hat{\rho}_C) = \hat{\rho}_T$ . Of course the above limit should not depend upon the particular causal cone "trajectory" one chooses to follow (the system is translational invariant). Without loss of generality we can thus pick the one associated with the central sites of the MERA, i.e. the one associated with the causal cone of N/2-th qudit. This allows us to identify all the  $\Phi^{(k)}$  of  $\Phi^{(1 \leftarrow m)}$ with maps of the form  $\Phi^{(R)}$ . A further simplification arises by enforcing the scale invariance property of the system. This can be done for instance by assuming that all the tensors  $\chi$ 's and  $\lambda$ 's of the MERA to be identical [9] and by requiring  $\Phi^{(L)} = \Phi^{(R)} = \Phi$  [21]. With this assumption all the sequence  $\Phi^{(1 \leftarrow m)}$  can now be written as a composition of m identical quantum channels, i.e.

$$\Phi^{(1 \leftarrow m)} = \Phi \circ \dots \circ \Phi = [\Phi]^m . \tag{3}$$

By general results on quantum channels the vast majority of CPT maps are known to be *mixing* (or *relaxing*) [16, 17, 18, 19]. This means that for a generic choice of  $\Phi$ , in the limit  $m \to \infty$  the transformation (3) will send all input states into a unique fix point identified as the unique eigen-operator of  $\Phi$  associated with its largest eigenvalue. This property allows us to identify the thermodynamic limit  $\hat{\rho}_T$  of the reduced density matrix  $\hat{\rho}_j$  with such an eigenstate. As in the case of MPS [2, 22], we can now provide a simplified expression for the thermodynamic limit of any local observable  $\hat{\Theta}$  of scale invariant MERAs. A convenient way to express this is obtained by moving in Liouville space [15, 23]. By doing so we can write

$$\langle \hat{\Theta} \rangle_T = \lim_{m \to \infty} \langle \langle \hat{\Theta} | (\hat{E}_{\Phi})^m | \hat{\rho}_C \rangle \rangle = \langle \langle \hat{\Theta} | \hat{\rho}_T \rangle \rangle , \qquad (4)$$

where the vectors  $|\hat{A}\rangle\rangle$  are the Liouville representations of the operators  $\hat{A}$ , while  $\hat{E}_{\Phi}$  is the Liouville operator associated to the map  $\Phi$ , i.e.

$$\hat{E}_{\Phi} \equiv \sum_{r} \hat{R}_{r} \otimes \hat{R}_{r}^{*} , \qquad (5)$$

(here  $\hat{R}_r^*$  is the complex conjugate of  $\hat{R}_r$  evaluated with respect to the canonical basis). Within this formalism the vector  $|\hat{\rho}_T\rangle\rangle$  which describes the fix point  $\hat{\rho}_T$  of the mixing map  $\Phi$  is also found as the the (unique) eigenvector correspondent to the unitary eigenvalue of  $\hat{E}_{\Phi}$ . Because of the close similarities between  $\hat{E}_{\Phi}$  and the MPS transfer matrix [2, 3, 5, 22] we dubbed the former the *transfer operator* of the MERA. The existence and uniqueness of the fix point is given by the physical assumption that the thermodynamical limit of a physical system exists and it is unique.

Correlation functions: – The computation of the long range correlation functions of  $|\psi\rangle$  has also a clear interpretation in terms of CPT maps and the thermodynamic




FIG. 2: Causal cone associated with the two point correlation function  $\langle \hat{\Theta}_i \hat{\Theta}_j \rangle$ . The light blue regions generate the tensor product channel  $\Phi^{(1 \leftarrow \bar{m})} \otimes \Phi^{(1 \leftarrow \bar{m})}$ . The magenta region instead corresponds to the CPT map  $\Psi^{(\bar{m} \leftarrow m)}$  which, acting on  $|hat\rangle$  produces the 6 qudits state  $\hat{\rho}_{ij}^{(\bar{m})}$ .

limit can be computed along the same lines presented for local observables. Here we specialize in the two point correlation functions as the generalization is straightforward. Consider then the expectation value  $\langle \hat{\Theta}_i \hat{\Theta}_j \rangle$  with  $\hat{\Theta}_i$  and  $\hat{\Theta}_j$  being two local operators acting on (say) the *i*-th and *j*-th qudit respectively. In this case the causal cone is formed by two single sites causal cones which intercept at the MERA level  $\bar{m} = \inf[\log_2(i-j)] - 1$ , (see Fig. 2). The resulting expectation values can then be written as

$$\langle \hat{\Theta}_i \hat{\Theta}_j \rangle = \operatorname{Tr}[(\Phi_i^{(1 \leftarrow \bar{m})} \otimes \Phi_j^{(1 \leftarrow \bar{m})})(\hat{\rho}_{ij}^{(\bar{m})}) (\hat{\Theta}_i \otimes \hat{\Theta}_j)], (6)$$

where  $\Phi_{i,j}^{(1\leftarrow\bar{m})}$  are the two CPT maps of the two singlesite causal cones associated with the sites *i* and *j* respectively (light blue regions of Fig. 2). The operator  $\hat{\rho}_{ij}^{(\bar{m})}$  instead is a 6 qudits state associated with the last  $m - \bar{m}$  levels of the MERA. It is obtained from the 4 qubit state  $|hat\rangle$  through the application of a quantum channel  $\Psi^{(\bar{m}\leftarrow\bar{m})}$  which, similarly to  $\Phi_{i,j}^{(1\leftarrow\bar{m})}$ , originates from a proper concatenation of CPT maps associated with  $\mathcal{M}_5$  or with the type- $\begin{pmatrix} 5\\10 \end{pmatrix}$  and type- $\begin{pmatrix} 4\\8 \end{pmatrix}$  tensors  $\mathcal{M}_9 \equiv \lambda \chi \lambda \chi \lambda \chi \lambda \chi \lambda \chi \lambda$  and  $\mathcal{M}_7 \equiv \lambda \chi \lambda \chi \lambda \chi \lambda \chi \lambda$ . Since this applies to all the two sites observable, we can then conclude that  $(\Phi_i^{(1\leftarrow\bar{m})} \otimes \Phi_j^{(1\leftarrow\bar{m})})(\hat{\rho}_{ij}^{(\bar{m})})$  must coincide with the reduced density matrix  $\hat{\rho}_{ij}$  of  $|hat\rangle$  associated with the sites *i* and *j*.

Let us focus then on the thermodynamic limit of the correlation function  $\Delta_{ij} \equiv \langle \hat{\Theta}_i \hat{\Theta}_j \rangle - \langle \hat{\Theta}_i \rangle \langle \hat{\Theta}_j \rangle$  which for Hamiltonian systems at criticality decays as  $|i - j|^{-\nu}$ . Under the same assumptions used to derive Eqs. (3) and (4) we can assume j to be the central site of the MERA (i.e. j = N/2). Suppose then that the associated map  $\Phi$  is mixing with fix point  $\hat{\rho}_T$ . For any input state  $\hat{\rho}_{ij}$  of the

sites *i* and *j* we then have  $\lim_{\bar{m}\to\infty} (\mathcal{I}_i \otimes \Phi_j^{(1\leftarrow\bar{m})})(\hat{\rho}_{ij}) = \hat{\rho}_i \otimes \hat{\rho}_T$ , with  $\hat{\rho}_i \equiv \operatorname{Tr}_j[\hat{\rho}_{ij}]$  and  $\mathcal{I}_i$  being the identity super-operator of the site *i*. The speed of convergence, evaluated through the trace distance, is exponentially fast [18, 19] in  $\bar{m}$  and, a part from some constant prefactor, can be upper-bounded by the quantity  $\bar{m}^{d^3} \kappa^{\bar{m}}$ with  $\kappa < 1$  being the modulus of the largest eigenvalue of  $\Phi$  whose associated eigenvector contribute in the expansion of Eq. (6). This is sufficient for claiming that the distance between  $(\Phi_i^{(1\leftarrow\bar{m})} \otimes \Phi_j^{(1\leftarrow\bar{m})})(\hat{\rho}_{ij}^{(\bar{m})})$  and  $\Phi_i^{(1\leftarrow\bar{m})}(\hat{\rho}_i) \otimes \Phi_j^{(1\leftarrow\bar{m})}(\hat{\rho}_j)$  is bounded by  $\propto \kappa^{2\bar{m}}$ . Thus we can write  $\log_2(|\Delta_{ij}|) \leq 2\bar{m}\log_2\kappa + \mathcal{O}(\log_2\bar{m})$ , which through the definition of  $\bar{m}$  provides a bound for the critical exponent  $\nu$  associated to the observable  $\hat{\Theta}$  in terms of the properties of the map  $\Phi$ , i.e.  $\nu \geq -2\log_2\kappa$ . In effect one can show that such bound is tight, i.e.

$$\nu = -2\log_2 \kappa . \tag{7}$$

This can be seen for instance by expressing Eq. (6) in the Liouville space formalism as in Eq. (4), and expanding the transfer matrices  $\hat{E}_{\Phi_i} \otimes \hat{E}_{\Phi_j}$  in Jordan blocks (the calculation is similar to the MPS analysis of Ref. [22]). A numerical test of Eq. (7) [24] on a MERA state approximating the ground energy of an Ising chain up to a  $10^{-4}$  accuracy yielded  $\kappa \simeq 0.915, 0.49, 0.52$  to be compared with the exact values  $\kappa_{th} \simeq 0.917, 0.46, 0.50$  associated with the x, y and z two-point correlation functions. Similar results have been obtained for the XXZ model.

Concluding remarks:- Equations. (5) and (7) constitute the main results of our analysis. As already discussed by Vidal [9], MERA networks are able to describe algebraic decaying correlations. In this work we put on firm grounds this observation giving an explicit expression of the critical exponents in terms of properties of the associated QuMERA channels. The combination of this approach with conformal field theory methods may provide a powerful tool to achieve a complete description of one-dimensional critical quantum systems. Similarly our findings yield a natural connection between the tensor network description of the thermodynamic limit of critical systems and the master equation formalism. Combining these results with the algorithms presented in [10, 13] one can exploit the introduction of the transfer operator (5) studying directly the infinite size system improving simulation efficiency.

The results presented here can be easily extended in several ways. For instance since a binary tree can be seen as a MERA with the disentanglers  $\chi$  set to the identity, all the arguments presented previously can be easily adapted to this case. Similarly also the thermodynamical limit for MPS [5] can be described in terms of repeated application of CPT maps (here the operator (5) reduces to the MPS transfer matrix). More generally our approach can be adapted to any tensor network by associating it with a family of CPT transformations which, properly concatenated, allows one to compute the local observables of the system. In this perspective the quantum circuit [9, 14] associated with the tensor network can be seen as a unitary dilation or, Stinespring representation [15], of the corresponding CPT family. Finally a generalization to higher spatial dimensions seems straightforward.

We thank M. Rizzi for useful discussions. This work was in part founded by the Quantum Information research program of Centro "Ennio De Giorgi" of SNS.

- S. R. White, Phys. Rev. Lett. **69**, 2863 (1992); Phys. Rev. B **48**, 10345 (1993).
- [2] M. Fannes, B. Nachtergaele, and R. F. Werner, Lett. Math. Phys. 25, 249 (1992).
- [3] S. Ostlund and S. Rommer, Phys. Rev. Lett. 75, 3537 (1995).
- [4] G. Vidal, Phys. Rev. Lett. **91**, 147902 (2003).
- [5] F. Verstraete, D. Porras, and J. I. Cirac, Phys. Rev. Lett. 93, 227205 (2004).
- [6] U. Schollwöck, Rev. Mod. Phys. 77 259 (2005); K. Hallberg, Adv. Phys. 55 477 (2006).
- [7] F. Verstraete, J. I. Cirac, Eprint arXiv:cond-mat/0407066; V. Murg, F. Verstraete, and J. I. Cirac, Phys. Rev. A 75, 033605 (2007).
- [8] W. Dür *et al.*, Phys. Rev. Lett. **94**, 097203 (2005); S. Anders *et al.*, Phys. Rev. Lett. **97**, 107206 (2006).
- [9] G. Vidal, Phys. Rev. Lett. 99, 220405 (2007); *ibid.* 101, 110501 (2008).
- [10] M. Rizzi, S. Montangero, and G. Vidal, Phys. Rev. A 77, 052328 (2008).
- [11] M. Aguado and G. Vidal, Phys. Rev. Lett. **100**, 070404 (2008); G. Evenbly and G. Vidal, Eprint arXiv:quantph/0801.2449.
- [12] G. Evenbly and G. Vidal, Eprint arXiv:quantph/0710.0692; L. Cincio, J. Dziarmaga, and M. M. Rams, Phys. Rev. Lett. **100**, 240603 (2008).
- [13] G. Vidal, Eprint: arXiv:0707.1454.
- [14] C. M. Dawson, J. Eisert, and T. J. Osborne, Phys. Rev. Lett. 100, 130501 (2008).
- [15] I. Bengstsson and K. Życzkowski, Geometry of Quantum States (Cambridge Univ. Press, Cambridge, 2006).
- [16] M. Raginsky, Phys. Rev. A 65, 032306 (2002).
- [17] R. Gohm, Noncommutative Stationary Processes (Springer, New York, 2004).
- [18] D. Burgarth and V. Giovannetti, New J. Phys. 9, 150 (2007).
- [19] B. M. Terhal and D. P. DiVincenzo, Phys. Rev. A 61 022301 (2000).
- [20] For easy of notation we consider the dimension D of the tensors  $\chi$  and  $\lambda$  to be equal to the physical dimension d. However, the subsequent equations hold for any D.
- [21] A necessary and sufficient condition [15] for  $\Phi^{(\tilde{L})} = \Phi^{(R)}$ is the existence of a unitary matrix  $U_{r,s}$ , such that  $\hat{L}_r = \sum_s U_{r,s} \hat{R}_s$ . This can then be easily casted into a necessary and sufficient condition for the tensor  $\mathcal{M}_5$ .
- [22] M. M. Wolf, et al. Phys. Rev. Lett. 97, 110403 (2006).
- [23] A. Royer, Phys. Rev. A 43, 44 (1991).
- [24] M. Rizzi, et al., in progress.