

# Extensions of the Quantum Fano Inequality

Naresh Sharma  
 Tata Institute of Fundamental Research  
 Mumbai 400 005, India  
 nsharma@tifr.res.in

February 6, 2020

## Abstract

We give extensions and an alternate derivation of the Quantum Fano inequality (QFI).

## 1 Introduction

Let  $R$  and  $Q$  be two quantum systems described by a Hilbert space  $\mathcal{H}_Q$  of finite dimension  $d$ , where  $d \geq 2$ . The joint system  $RQ$  is initially prepared in a pure entangled state

$$|\psi^{RQ}\rangle = \sum_{k=1}^d \sqrt{\lambda_k} |k^R\rangle |k^Q\rangle \quad (1)$$

where  $\boldsymbol{\lambda} = [\lambda_1 \cdots \lambda_d]$  is a probability vector, i.e.,  $\lambda_k \geq 0$ ,  $\sum_{k=1}^d \lambda_k = 1$ ,  $\{|k^R\rangle\}$  and  $\{|k^Q\rangle\}$ ,  $k = 1, \dots, d$ , are two orthonormal bases for  $\mathcal{H}_Q$ .  $|\psi^{RQ}\rangle$  is a purification of  $\rho$ , the state of system  $Q$ , and

$$\rho = \text{Tr}_R(|\psi^{RQ}\rangle\langle\psi^{RQ}|) = \sum_{k=1}^d \lambda_k |k^Q\rangle\langle k^Q| \quad (2)$$

The system  $Q$  undergoes a completely positive trace-preserving transformation or quantum operation  $\mathcal{E}$  and  $R$  is assumed to be isolated and its state remains the same. This quantum operation is also represented by  $\mathcal{I}_R \otimes \mathcal{E}$ , where  $\mathcal{I}_R$  is the identity superoperator on  $R$ .

We add subscript ‘1’ to denote the state of the system (joint or otherwise) after this quantum operation. So the state of the joint system is denoted by  $\rho^{R_1Q_1}$ . Note that  $\rho^{Q_1} = \mathcal{E}(\rho)$  and  $\rho^{R_1} = \rho^R$ .

The entanglement fidelity is defined by Schumacher [1] as

$$F(\rho, \mathcal{E}) = \langle \psi^{RQ} | \rho^{R_1Q_1} | \psi^{RQ} \rangle \quad (3)$$

and the entropy exchange as

$$S(\rho, \mathcal{E}) = S(\rho^{R_1Q_1}) \quad (4)$$

where  $S(\rho^{R_1Q_1})$  is the von-Neumann entropy of  $\rho^{R_1Q_1}$ . The QFI upper bounds  $S(\rho, \mathcal{E})$  by a function of the entanglement fidelity as [1]

$$S(\rho, \mathcal{E}) \leq H(F(\rho, \mathcal{E})) + (1 - F(\rho, \mathcal{E})) \log(d^2 - 1) \quad (5)$$

where we shall assume throughout the paper that  $\log$  denotes the natural logarithm. More details on the QFI can be found in [1, 2, 3].

Generalization of the Fano's inequality for the classical case was proposed by Han and Verdú [4], where various lower bounds to the mutual information are provided.

In this paper, we give extensions of the QFI. The QFI is shown to be a special case of the proposed inequalities and this also gives an alternate derivation of the QFI.

## 2 Extensions of the Quantum Fano inequality

Let  $R_2, Q_2$  be two ancilla quantum systems, possibly entangled, described by  $\mathcal{H}_Q$ . The joint system  $R_2Q_2$  is described by  $\mathcal{H}_{RQ} = \mathcal{H}_Q \otimes \mathcal{H}_Q$ , and let  $\{|k^{RQ}\rangle\}$  be an orthonormal basis for  $\mathcal{H}_{RQ}$ , and we define a set of projectors as

$$P_k = |k^{RQ}\rangle\langle k^{RQ}|, \quad \sum_{k=1}^{d^2} P_k = I^{RQ} \quad (6)$$

where we have chosen

$$|1^{RQ}\rangle = |\psi^{RQ}\rangle \quad (7)$$

and  $I^{RQ}$  is the  $d^2 \times d^2$  identity matrix. Then

$$S(\rho, \mathcal{E}) = S(\rho^{R_1Q_1}) \quad (8)$$

$$= -S(\rho^{R_1Q_1} || \rho^{R_2Q_2}) - \text{Tr}(\rho^{R_1Q_1} \log(\rho^{R_2Q_2})) \quad (9)$$

$$\leq -S\left(\sum_{k=1}^{d^2} P_k \rho^{R_1Q_1} P_k \middle| \middle| \sum_{k=1}^{d^2} P_k \rho^{R_2Q_2} P_k\right) - \text{Tr}(\rho^{R_1Q_1} \log(\rho^{R_2Q_2})) \quad (10)$$

$$= -D(\mathbf{p} || \mathbf{q}) - \text{Tr}(\rho^{R_1Q_1} \log(\rho^{R_2Q_2})) \quad (11)$$

$$(12)$$

where

$$S(\rho || \sigma) = \text{Tr}(\rho \log(\rho)) - \text{Tr}(\rho \log(\sigma)) \quad (13)$$

is the quantum relative entropy, in Eq. (10) we have used the fact that a trace-preserving completely positive transformation reduces the quantum relative entropy [5, 6],

$$\mathbf{p} = [p_1 \cdots p_{d^2}] \quad (14)$$

$$\mathbf{q} = [q_1 \cdots q_{d^2}] \quad (15)$$

$$p_k = \langle k^{RQ} | \rho^{R_1Q_1} | k^{RQ} \rangle \quad (16)$$

$$q_k = \langle k^{RQ} | \rho^{R_2Q_2} | k^{RQ} \rangle \quad (17)$$

and  $D(\cdot || \cdot)$  is the classical relative entropy given by

$$D(\mathbf{p} || \mathbf{q}) = \sum_{k=1}^{d^2} p_k \log\left(\frac{p_k}{q_k}\right) \quad (18)$$

Let

$$g(p, q) = D([p(1-p)] || [q(1-q)]) \quad (19)$$

Then

$$D(\mathbf{p}||\mathbf{q}) - g(p_1, q_1) = \sum_{k=2}^{d^2} p_k \log\left(\frac{p_k}{q_k}\right) - (1 - p_k) \log\left(\frac{1 - p_k}{1 - q_k}\right) \quad (20)$$

$$= \sum_{k=2}^{d^2} p_k \log\left(\frac{p_k(1 - q_1)}{q_k(1 - p_1)}\right) \quad (21)$$

$$\geq \sum_{k=2}^{d^2} p_k \left(1 - \frac{q_k(1 - p_1)}{p_k(1 - q_1)}\right) \quad (22)$$

$$= 0 \quad (23)$$

where in Eq. (22), we have used the fact that for  $x > 0$ ,  $\log(x) \geq 1 - 1/x$ , with equality iff  $x = 1$ . Hence, the equality condition for Eq. (23) is

$$\frac{q_k}{p_k} = \frac{1 - q_1}{1 - p_1}, \quad k = 2, \dots, d \quad (24)$$

More general lower bounds to the classical relative entropy are given by Blahut in [7]. Substituting Eq. (23) into Eq. (11), we get

$$S(\rho, \mathcal{E}) \leq -g(F(\rho, \mathcal{E}), q_1) - \text{Tr}(\rho^{R_1 Q_1} \log(\rho^{R_2 Q_2})) \quad (25)$$

where we have used the fact that  $p_1 = F(\rho, \mathcal{E})$ . There are different choices of the  $\rho^{R_2 Q_2}$  possible to give different upper bounds on  $S(\rho, \mathcal{E})$ . We consider a few such choices below.

### 3 Special Cases

Let

$$\rho^{R_2 Q_2} = \sum_{k=1}^d \gamma_k |k^R\rangle\langle k^R| \otimes \rho^{Q_2} \quad (26)$$

where  $\boldsymbol{\gamma} = [\gamma_1 \dots \gamma_d]$  is a probability vector, and we have not yet specified the state  $\rho^{Q_2}$ . This choice yields

$$q_1 = \sum_{i,j,k=1}^d \sqrt{\lambda_i \lambda_j} \gamma_k \langle i^R | \langle i^Q | \left( |k^R\rangle\langle k^R| \otimes \rho^{Q_2} \right) |j^R\rangle |j^Q\rangle \quad (27)$$

$$= \sum_{i,j,k=1}^d \sqrt{\lambda_i \lambda_j} \gamma_k \delta_{i,k} \delta_{k,j} \langle i^Q | \rho^{Q_2} |j^Q\rangle \quad (28)$$

$$= \sum_{k=1}^d \gamma_k \lambda_k \langle k^Q | \rho^{Q_2} |k^Q\rangle \quad (29)$$

where  $\delta_{i,k} = 1$  if  $i = k$  and is zero otherwise. Using Eq. (25), we get

$$S(\rho, \mathcal{E}) \leq -g(F(\rho, \mathcal{E}), q_1) - \sum_{k=1}^d \lambda_k \log(\gamma_k) - \text{Tr}(\mathcal{E}(\rho) \log(\rho^{Q_2})) \quad (30)$$

where we have used  $\rho^{Q_1} = \mathcal{E}(\rho)$ . Again, different choices of  $\rho^{Q_2}$  are possible. Let us consider

$$\rho^{Q_2} = \sum_{k=1}^d \xi_k |k^Q\rangle \langle k^Q| \quad (31)$$

where  $\boldsymbol{\xi} = [\xi_1 \cdots \xi_d]$  is a probability vector. With this choice and noting that

$$-\text{Tr} \left( \mathcal{E}(\rho) \log(\rho^{Q_2}) \right) = - \sum_k \log(\xi_k) \langle k_Q | \mathcal{E}(\rho) | k_Q \rangle \quad (32)$$

$$\leq - \log(\min_i \{\xi_i\}) \quad (33)$$

Eq. (30) reduces to

$$S(\rho, \mathcal{E}) \leq -g \left( F(\rho, \mathcal{E}), \sum_{k=1}^d \lambda_k \gamma_k \xi_k \right) - \sum_{k=1}^d \lambda_k \log(\gamma_k) - \log(\min_i \{\xi_i\}) \quad (34)$$

$$\begin{aligned} &= H(F(\rho, \mathcal{E})) + (1 - F(\rho, \mathcal{E})) \log \left( \frac{1}{\sum_{i=1}^d \lambda_i \gamma_i \xi_i} - 1 \right) + \log \left( \frac{\sum_{i=1}^d \lambda_i \gamma_i \xi_i}{\min_i \{\xi_i\}} \right) \\ &\quad - \sum_{k=1}^d \lambda_k \log(\gamma_k) \end{aligned} \quad (35)$$

where

$$H(x) = -x \log(x) - (1-x) \log(1-x) \quad (36)$$

is the binary entropy function.

The QFI follows as a special case by substituting  $\gamma_k = \xi_k = 1/d$ ,  $k = 1, \dots, d$ . Note that the above inequality holds for any probability vectors  $\boldsymbol{\gamma}$  and  $\boldsymbol{\xi}$ . We get the following simpler bound than Eq. (35) by choosing  $\xi_k = 1/d$ ,  $k = 1, \dots, d$ ,

$$S(\rho, \mathcal{E}) \leq H(F(\rho, \mathcal{E})) + (1 - F(\rho, \mathcal{E})) \log \left( \frac{d}{\sum_{i=1}^d \lambda_i \gamma_i} - 1 \right) + \log \left( \sum_{i=1}^d \lambda_i \gamma_i \right) - \sum_{k=1}^d \lambda_k \log(\gamma_k) \quad (37)$$

Eqs. (25), (30), (35), and (37) are various like Fano-like bounds that can be made tighter by appropriately choosing  $\rho^{R_2 Q_2}$ ,  $\{\boldsymbol{\gamma}, \rho^{Q_2}\}$ ,  $\{\boldsymbol{\gamma}, \boldsymbol{\xi}\}$ , and  $\boldsymbol{\gamma}$  respectively.

It might seem that one could get away from the dependence of the bounds on  $\boldsymbol{\lambda}$  by making the following choice of  $\rho^{R_2 Q_2}$ , which is different from Eq. (26). Let  $\beta_k$ ,  $k = 1, \dots, d^2$ , be the eigenvalues of  $\rho^{R_2 Q_2}$  and  $|\psi^{RQ}\rangle$  be one of the eigenvectors of  $\rho^{R_2 Q_2}$ . Let  $\beta_{\max} = \max_k \beta_k$ ,  $\beta_{\min} = \min_k \beta_k$ . Since the maximum of  $g(F, x)$ ,  $x \in [\beta_{\min}, \beta_{\max}]$ , occurs at the end-points, hence to make the bound tighter, one could choose the eigenvalue corresponding to the eigenvector  $|\psi^{RQ}\rangle$  as either  $\beta_{\min}$  or  $\beta_{\max}$ . The bound in Eq. (25) can be simplified to

$$S(\rho, \mathcal{E}) \leq -g(F(\rho, \mathcal{E}), q_1) - \log(\beta_{\min}) \quad (38)$$

where  $q_1 = \beta_{\max}$  or  $q_1 = \beta_{\min}$ . Suppose  $q_1 = \beta_{\max}$ , then to tighten the bound, one could choose  $\beta_{\min}$  as large as possible, or

$$\beta_{\min} = \frac{1 - \beta_{\max}}{d^2 - 1} \quad (39)$$

Substituting in Eq. (38), we get

$$S(\rho, \mathcal{E}) \leq H(F(\rho, \mathcal{E})) - F(\rho, \mathcal{E}) \log \left( \frac{1}{\beta_{\max}} - 1 \right) + \log(d^2 - 1) \quad (40)$$

We get the tightest bound by choosing minimum value of  $\beta_{\max}$  given by  $\beta_{\max} = 1/d^2$ , which reduces Eq. (40) to the QFI.

If  $q_1 = \beta_{\min}$ , then Eq. (38) reduces to

$$S(\rho, \mathcal{E}) \leq H(F(\rho, \mathcal{E})) + (1 - F(\rho, \mathcal{E})) \log \left( \frac{1}{\beta_{\min}} - 1 \right) \quad (41)$$

We get the tightest bound by choosing maximum value of  $\beta_{\min}$  given by  $\beta_{\min} = 1/d^2$ , which reduces Eq. (41) to the QFI. Hence, this choice of  $\rho^{R_2 Q_2}$  offers no improvement over the QFI.

## 4 An Example

We compute the QFI and the proposed inequality for the depolarizing channel for a single qubit ( $d = 2$ ) given by

$$\mathcal{E}(\rho) = \left(1 - \frac{3p}{4}\right) \rho + \frac{p}{4}(X\rho X + Y\rho Y + Z\rho Z) \quad (42)$$

where  $X, Y, Z$  are Pauli matrices. Let

$$\rho = U \begin{pmatrix} \lambda & 0 \\ 0 & 1 - \lambda \end{pmatrix} U^\dagger \quad (43)$$

where  $U$  is a randomly chosen  $2 \times 2$  Unitary matrix. It is easy to show that for any choice of  $U$

$$F(\rho, \mathcal{E}) = 1 + p \left( \lambda^2 - \lambda - \frac{1}{2} \right) \quad (44)$$

$$S(\rho, \mathcal{E}) = H_S \left( \left[ \frac{p\lambda}{2}, \frac{(1-\lambda)p}{2}, -\frac{p}{4} + \frac{1}{2} + \frac{\theta}{4}, -\frac{p}{4} + \frac{1}{2} - \frac{\theta}{4} \right] \right) \quad (45)$$

where  $H_S(\cdot)$  is the Shannon entropy, and

$$\theta = \sqrt{p^2 + 12p^2\lambda(1-\lambda) + 4(1-p) - 16p\lambda(1-\lambda)} \quad (46)$$

In Fig. 1, we compare  $S(\rho, \mathcal{E})$  with the QFI and the inequality in Eq. (37) numerically optimized over  $\gamma$  to give the tightest bound for  $\lambda = 0.1$ . The figure shows that the latter bound is tighter than the QFI. In Fig. 2, we plot the numerically computed value of  $\gamma_1$  that gives the tightest bound in Eq. (37). The QFI corresponds to  $\gamma_1 = 1/d = 0.5$ .

## References

- [1] B. Schumacher. Sending entanglement through noisy quantum channels. *Phys. Rev. A*, 54:2614–2628, Oct. 1996.

- [2] M. A. Nielsen and I. L. Chuang. *Quantum Computation and Quantum Information*. Cambridge University Press, Cambridge, 2000.
- [3] M. Hayashi. *Quantum Information: An Introduction*. Springer, 2006.
- [4] T. S. Han and S. Verdú. Generalizing the Fano inequality. *IEEE Trans. Inf. Theory*, 40:1247–1251, July 1994.
- [5] G. Lindblad. Completely positive maps and entropy inequalities. *Commun. Math. Phys.*, 40:147–151, June 1975.
- [6] A. Uhlmann. Relative entropy and the Wigner-Yanase-Dyson-Lieb concavity in an interpolation theory. *Commun. Math. Phys.*, 54:2–32, Feb. 1977.
- [7] R. E. Blahut. Information bounds of the Fano-Kullback type. *IEEE Trans. Inf. Theory*, 22:410–421, July 1976.




Figure 1: Plots of  $S(\rho, \mathcal{E})$ , the tightest bound from Eq. (37), and the QFI.




Figure 2:  $\gamma_1$  that gives the tightest bound in Eq. (37).