# Contribution of the second Landau level to the exchange energy of the three-dimensional electron gas in a high magnetic field

J. M. Morbec<sup>1</sup> and K. Capelle<sup>1,2,\*</sup>

<sup>1</sup>Departamento de Física e Informática, Instituto de Física de São Carlos

Universidade de São Paulo, Caixa Postal 369, São Carlos, 13560-970 SP, Brazil

<sup>2</sup>Theoretische Physik, Freie Universität Berlin, D-14195 Berlin, Germany

(Dated: October 30, 2018)

We derive a closed analytical expression for the exchange energy of the three-dimensional interacting electron gas in strong magnetic fields, which goes beyond the quantum limit (L = 0) by explicitly including the effect of the second, L = 1, Landau level and arbitrary spin polarization. The inclusion of the L = 1 level brings the fields to which the formula applies closer to the laboratory range, as compared to previous expressions, valid only for L = 0 and complete spin polarization. We identify, and explain, two distinct regimes, separated by a critical density  $n_c$ . Below  $n_c$ , the perparticle exchange energy is lowered by the contribution of L = 1, whereas above  $n_c$  it is increased. As special cases of our general equation we recover various known, more limited, results for higher fields, and identify and correct a few inconsistencies in some of these earlier expressions.

PACS numbers: 71.10.Ca,71.15.Mb,71.70.Di,97.60.Jd

#### I. INTRODUCTION

The calculation of the exchange energy of an interacting Fermi gas in high magnetic fields is a fundamental problem of many-body physics, with applications in fields as diverse as semiconductor physics,<sup>1,2,3</sup> astrophysics,<sup>1,4,5</sup> atomic and molecular physics<sup>1,4,5</sup> and density-functional theory.<sup>6,7,8,9</sup>

In a seminal 1971 paper<sup>10</sup> Danz and Glasser (hereafter DG) calculated the exchange energy,  $e_x$ , of a threedimensional electron gas in high magnetic fields, by means of Green's function techniques. A key result are analytical expressions for the dependence of  $e_x$  on density n and magnetic field B, valid if the electrons are fully spin polarized and occupy only the spin-down sublevel of the lowest Landau level. In an equally important 1974 paper Banerjee, Constantinescu and Rehak<sup>11</sup> (hereafter BCR) also calculated this exchange energy, and obtained a result that looks very similar to that of DG. The calculations of DG and BCR provided the background for a large body of later work on the exchange and correlation energy of the electron gas in high magnetic fields  $^{1,12,13,14,15,16,17,18}$  and are also frequently quoted as input for the local-density approximation to current-density-functional theory in strong magnetic fields.  $^{18,19,20,21,22}$ 

A major limitation of these early calculations is their restriction to complete spin polarization and to the lowest Landau level, which in three dimensions requires either magnetic fields that are beyond what is currently achievable in the laboratory, or restriction to low-density low-effective-mass systems. Here we extend the DG manybody calculations to the case of arbitrary spin polarization, and include the contribution of the second, L = 1, Landau level. As a consequence, the range of magnetic fields to which the resulting expression applies is extended towards weaker fields, as compared to earlier ex-

pressions.

Our formula reveals rather complex behaviour of the exchange energy, once higher Landau-levels are included: As a function of the density n,  $e_x$  first drops with a discontinuous derivative at density  $n_d$ , corresponding to the onset of occupation of L = 1, and then passes through two regimes, separated by a critical density  $n_c$ . Below  $n_c$ , the per-particle exchange energy is lowered (in modulus) by occupation of the L = 1 level, whereas above  $n_c$  it is increased (in modulus). The crossover between both regimes corresponds to a local minimum of  $e_x(n)$  at  $n_c$ . The physics of the drop and of both regimes can be understood in terms of the Landau-level structure. For currently achievable fields,  $n_c$  falls into the metallic density regime, and thus should be observable.

Next, we consider various special cases of our general expression, in order to make contact with more restricted results previously available in the literature, in particular those of DG and BCR. Scrutiny of these earlier expressions reveals a number of small inconsistencies and mistakes, which we correct on the basis of our more general expression.

### II. EXCHANGE ENERGY INCLUDING THE SECOND LANDAU LEVEL

The extension of the DG calculation to include higher Landau levels had, up to now, not been achieved in closed from. DG, and many other workers, <sup>1,11,12,13,14,15,16,17,18</sup> go beyond the quantum limit by rewriting the exchange integrals via expansion in infinite series, which cannot be resummed, or calculate them numerically. Neither approach yields analytical expressions that can be used, *e.g.* in the construction of density functionals for currentdensity-functional theory.<sup>6,7</sup> Motivated by the need for analytical expressions for lower fields, including L = 1, and by the observation of small inconsistencies in avail-

Three types of terms contribute: one,  $e_x^{(0)}$ , arises exclusively from L = 0, another,  $e_x^{(1)}$ , arises from L = 1, and a third,  $e_x^{(0,1)}$ , from inter-level exchange, involving contributions from L = 0 and L = 1. As a function of the occupation numbers  $n_L^{\sigma}$  of the spin up and down sublevels of the L = 0 and L = 1 Landau levels, the final result for the per-volume exchange energy can be written

$$e_x(n_0^{\uparrow}, n_0^{\downarrow}, n_1^{\uparrow}, n_1^{\downarrow}, B) = \frac{e^2}{8\pi^3} \left(\frac{m\omega_c}{\hbar}\right)^2 \sum_{\sigma} \left[e_x^{(0)}(n_0^{\sigma}, B) + e_x^{(0,1)}(n_0^{\sigma}, n_1^{\sigma}, B) + e_x^{(1)}(n_1^{\sigma}, B)\right]$$
(1)

where

$$e_x^{(0)}(n_0^{\sigma}, B) = C + \ln\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_0^{\sigma^2}\right) - \exp\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_0^{\sigma^2}\right)Ei\left(-\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_0^{\sigma^2}\right) \\ - \left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_0^{\sigma^2}\right)^{1/4}G_{23}^{22}\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_0^{\sigma^2}\right)\frac{3/4, 5/4}{3/4, 3/4, 1/4}\right),$$
(2)

$$e_x^{(0,1)}(n_0^{\sigma}, n_1^{\sigma}, B) = 4 \ln \left( \frac{n_0^{\sigma} + n_1^{\sigma}}{n_0^{\sigma} - n_1^{\sigma}} \right) + 2 \exp \left( \frac{2\pi^4 \hbar^3}{m^3 \omega_c^3} \left( n_0^{\sigma} - n_1^{\sigma} \right)^2 \right) Ei \left( -\frac{2\pi^4 \hbar^3}{m^3 \omega_c^3} \left( n_0^{\sigma} - n_1^{\sigma} \right)^2 \right) \\ -2 \exp \left( \frac{2\pi^4 \hbar^3}{m^3 \omega_c^3} \left( n_0^{\sigma} + n_1^{\sigma} \right)^2 \right) Ei \left( -\frac{2\pi^4 \hbar^3}{m^3 \omega_c^3} \left( n_0^{\sigma} + n_1^{\sigma} \right)^2 \right) \\ + \left( \frac{2\pi^4 \hbar^3}{m^3 \omega_c^3} \left( n_0^{\sigma} - n_1^{\sigma} \right)^2 \right)^{1/4} G_{23}^{22} \left( \frac{2\pi^4 \hbar^3}{m^3 \omega_c^3} \left( n_0^{\sigma} - n_1^{\sigma} \right)^2 \right)^{3/4} G_{23}^{4} \left( \frac{3/4}{m^3 \omega_c^3} \left( n_0^{\sigma} + n_1^{\sigma} \right)^2 \right) \\ - \left( \frac{2\pi^4 \hbar^3}{m^3 \omega_c^3} \left( n_0^{\sigma} + n_1^{\sigma} \right)^2 \right)^{1/4} G_{23}^{22} \left( \frac{2\pi^4 \hbar^3}{m^3 \omega_c^3} \left( n_0^{\sigma} + n_1^{\sigma} \right)^2 \right)^{3/4} G_{23}^{4} \left( \frac{3/4}{m^3 \omega_c^3} \left( n_0^{\sigma} + n_1^{\sigma} \right)^2 \right) \right)$$
(3)

and

$$e_x^{(1)}(n_1^{\sigma}, B) = C + \ln\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_1^{\sigma^2}\right) - \exp\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_1^{\sigma^2}\right)Ei\left(-\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_1^{\sigma^2}\right) \\ -\frac{3}{4}\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_1^{\sigma^2}\right)^{1/4}G_{23}^{22}\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_1^{\sigma^2}\right)\frac{3/4, 5/4}{3/4, 3/4, 1/4}\right) \\ +\frac{1}{2}\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_1^{\sigma^2}\right)\exp\left(\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_1^{\sigma^2}\right)Ei\left(-\frac{8\pi^4\hbar^3}{m^3\omega_c^3}n_1^{\sigma^2}\right).$$
(4)

Here  $\omega_c(B) = eB/mc$  is the cyclotron frequency, C = 0.57722 is Euler's constant,  $G_{23}^{22}$  is the Meijer Gfunction<sup>23</sup> and Ei is the exponential integral. In principle, this expression hold for arbitrary values of the gfactor, as long as  $L \leq 1$ , but we note that if the freeelectron value g = 2 is employed, an accidental degeneracy between the spin-up subband of L = 1 level and the spin-down subband of L = 2 occurs. Restriction to  $L \leq 1$ is thus only rigorously possible if either g < 2 (a common situation in semiconductors) or if the L = 1 level is fully

polarized, so that its spin-up subband is empty.

The condition  $L \leq 1$  implies a restriction on the allowable values of density and magnetic field. In terms of the magnetic length  $l(B) = \sqrt{\hbar c/(eB)}$  and the density parameter  $r_s(n)a_0 = [3/(4\pi n)]^{1/3}$ , where  $a_0 = \hbar^2/(me^2)$  is the Bohr radius, this restriction is conveniently written as

$$\frac{l}{r_s a_0} < \left[\frac{4}{3\pi} \left(1 + \sqrt{2}\right)\right]^{1/3} \tag{5}$$

for g = 2, and

$$\frac{l}{r_s a_0} < \left[\frac{4}{3\pi} \left(2 + \sqrt{2}\right)\right]^{1/3} \tag{6}$$

for g = 0. These conditions are derived in Appendix B.

Regarding the spin dependence, we note that spin up and spin down contributions do not mix, *i.e.*, their contribution to  $e_x$  can be evaluated separately. Equations (1) to (4) permit one to do this for arbitrary values of the occupation numbers  $n_L^{\sigma}$ , and thus also for arbitrary spin polarizations  $m \propto (n_{\uparrow} - n_{\downarrow})$ , where  $n_{\sigma} = \sum_{L}^{occ} n_L^{\sigma}$ .

As illustration of Eq. (1), Fig. 1 displays the exchange energy for a combination of densities and magnetic fields for which both L = 0 and L = 1 levels contribute, and compares it with the (erroneous) use of the L = 0 expression alone in the same regime.

Figure 1 reveals interesting behaviour of  $e_x$  that appears only once L > 0 is allowed for: Upon adding more particles (increasing the density), the per-particle exchange energy suddenly drops (in modulus) once the L = 1 level starts to be occupied at density  $n_d$ , and then passes through two regimes. Initially, it continuous to decrease, while for larger densities it increases (in modulus) up to values larger than those obtained by allocating all particles in L = 0. The first regime is entered with a discontinuous derivative in the  $e_x(n)$  curve, indicating a zero-temperature phase transition. The second regime is entered via a gradual crossover. Both regimes are separated by a critical density  $n_c$ , at which  $e_x(n)$  goes through a local minimum.

This intricate behaviour finds its explanation in the Landau-level structure: Due to the differences in the spatial part of the Landau-level wave functions, intra-level exchange integrals are larger than inter-level integrals. As long as only L = 0 contributes, the exchange energy naturally increases with the number of particles. Once particles are allocated in the L = 1 level, too, the reduced spatial overlap of their orbitals with those of the particles in L = 0 leads to a reduction of the per-particle exchange energy. As the number of particles in L = 1 increases, their intra-level exchange energy also grows and overcompensates the initial drop (as seen in the crossing of the full and the dashed curve).

For g = 2, the initial drop is much more pronounced than for g < 2, because for this value occupation of the spin-down subband of the L = 1 level occurs simultaneously with that of the energetically degenerate spin-up subband of the L = 0 level, so that the exchange energy of some of the aditional particles is not only reduced by small spatial overlap but strictly zero due to zero spin overlap with the spin-down particles already occupying L = 0. The resulting larger drop may be so large that for fields and densities compatible with  $L \leq 1$ ,  $e_x(n)$ does not recover the hypothetical value obtained by filling only the L = 0 level, as illustrated in the upper set of two curves in Fig. 1.

This complex behaviour of  $e_x(n)$  implies that magnetic-field Thomas-Fermi-Dirac theory,<sup>1,14,15</sup>



FIG. 1: Full curves: Exchange energy per particle, obtained from Eq. (1) by dividing by the density, as a function of  $r_s^{-1} \propto n^{1/3}$ . Dashed curves: (Erroneous) continuation of the L = 0expression into the L = 1 regime. Upper set of two curves: g = 2 and  $B = 1.4448 \times 10^4 T$ . Lower set of two curves: g = 0and  $B = 1.1474 \times 10^4 T$ .

current-density-functional theory<sup>6,7</sup> and magnetic-fielddensity-functional theory<sup>8,9</sup> should have a much richer solution space than have ordinary Thomas-Fermi-Dirac theory and spin-density-functional theory, where  $e_x(n)$ is strictly monotonous. The density  $n_d$  where the discontinuous drop occurs, and the critical density  $n_c$ where  $e_x(n)$  goes through its minimum, depend on the magnetic field. For sufficiently large fields both fall in the metallic density range, *i.e.*, both regimes, as well as the crossover between them, should be observable in the laboratory.

### **III. HIGHER FIELDS AND LIMITING CASES**

In the quantum limit, B is so high that only the L = 0level is occupied. In this limit, the exchange energy is given by keeping only  $e_x^{(0)}$  in Eq. (1). If the free-electron value g = 2 is employed, an accidental degeneracy between the spin-up subband of L = 0 level and the spindown subband of L = 1 occurs. Restriction to L = 0is thus only rigorously possible if either g < 2 or if the system is fully polarized, so that the spin-up subband of L = 0 level is empty, too. Moreover, the conditions guaranteeing L = 0 are stricter than those limiting occupation to  $L \leq 1$ , given above, and read

$$\frac{l}{r_s a_0} < \left(\frac{2\sqrt{2}}{3\pi}\right)^{1/3} \tag{7}$$

for g = 2,

$$\frac{l}{r_s a_0} < \left(\frac{4\sqrt{2}}{3\pi}\right)^{1/3} \tag{8}$$

for g = 0, and

$$\frac{l}{r_s a_0} < \left[\frac{2}{3\pi} \left(\sqrt{2 - |g|} + \sqrt{2}\right)\right]^{1/3} \tag{9}$$

for generic  $g \in (-2, 2)$ . These conditions are derived in Appendix B. As long as these conditions are satisfied, Eq. (1) with the contribution from just  $e_x^{(0)}$  can be used for arbitrary occupation of the up and down subbands, *i.e.* for arbitrary spin polarization.

The further restriction to full spin polarization, *i.e.* an empty spin-up subband, then leads to

$$e_x(n,B) = \frac{e^2}{8\pi^3} \left(\frac{m\omega_c}{\hbar}\right)^2 \left[C + \ln\left(p\right) - e^p Ei\left(-p\right) - e^{p^{1/4}} G_{23}^{22} \left(p \left| \begin{array}{c} 3/4, 5/4\\ 3/4, 3/4, 1/4 \end{array} \right) \right], (10)$$

which is the result obtained by DG.<sup>10</sup> Here the particle density *n* is equal to the occupation  $n_{L=0}^{\sigma=\downarrow}$  and we defined, following DG,  $p(n, B) := 8\pi^4 \hbar^3 n^2 / (m^3 \omega_c^3)$ . As long as no higher levels are occupied and g = 2, we also have  $p(n, B) = 4\epsilon_F/\hbar\omega_c$ , where  $\epsilon_F$  is the Fermi energy. (This equality breaks down if other spin states or Landau levels are involved, or  $g \neq 2$ .)

If  $\epsilon_F/\hbar\omega_c \ll 1$ , then only the bottom of the lowest subband is occupied, and the preceding equation reduces to

$$e_x(n,B) = \frac{\pi e^2 \hbar}{m\omega_c} n^2 \left[ \ln(p) - 3 + C \right].$$
(11)

In units in which  $\hbar = c = 1$ , Eq. (11) can be written as

$$e_x(n,B) = \frac{2\pi e^2}{eB} n^2 \left[ \ln\left(\frac{n}{(eB)^{3/2}}\right) + 2.11788 \right],$$
 (12)

which was first obtained by  $DG^{10}$  and can be directly compared with the corresponding equation (47) of BCR,<sup>11</sup>

$$e_x(n,B)^{BCR} = \frac{2\pi e^2}{eB} n^2 \left[ \ln\left(\frac{n}{(eB)^{3/2}}\right) + 2.32918 \right],$$
(13)

featuring a different numerical value inside the brackets. This difference was also noticed in Refs. 13,14.

For ultra-strong magnetic fields (quantified in Fig. 2) or for extremely low densities, that value, be it 2.11788 or 2.32918, becomes negligible relative to the logarithmic term, and the DG and BCR expressions become identical, both reducing to

$$\frac{e_x(r_s,B)}{Ry} = -\frac{27}{4\pi} \frac{1}{a_0^3 r_s^6} \left(\frac{\hbar\omega_c}{Ry}\right)^{-1} \ln\left(0.141 r_s^2 \frac{\hbar\omega_c}{Ry}\right),\tag{14}$$

where  $e_x/Ry$  and  $\hbar\omega_c/Ry$  denote exchange energy and cyclotron energy measured in Rydberg  $[1Ry = e^2/2a_0 = 13.6eV]$ . In DG this limit appears as their Eq. (1.1),

$$\frac{e_x(r_s,B)^{DG}}{Ry} = -\frac{27}{16\pi} \frac{1}{a_0^3 r_s^6} \left(\frac{\hbar\omega_c}{Ry}\right)^{-1} \ln\left(0.282 r_s^2 \frac{\hbar\omega_c}{Ry}\right),\tag{15}$$



FIG. 2: Applicability of the four expressions discussed here. The curves represent the *lowest* magnetic fields for which the indicated expression is valid, for the free-electron value of m and g = 2.

which differs in two ways from Eq. (14): the numerical factor inside the logarithm is  $0.282 = 2 \times 0.141$  and the prefactor is 27/16, instead of 27/4.

Our general expression (1) contains all these limits as special cases, and we can therefore verify which of the conflicting expressions is correct. In the high-field limit of the quantum limit, we find that the DG Eq. (12) is correct, while the BCR Eq. (13) is not. In the ultra-strong field limit, we find that Eq. (14) is correct, while the DG Eq. (15) is not. In this latter case, we suspect that DG inadvertedly used Hartree units instead of Rydberg units  $(1H = e^2/a_0 = 2Ry)$  is the atomic unit of energy), but denoted them as Rydberg, as this would explain both the factor of 2 inside and the factor of 1/4 outside the logarithm.<sup>24</sup>

Figure 2 illustrates the magnetic field and density regimes for which each of the above equations is valid. The curves represent the *lowest* magnetic fields for which the indicated expression is valid, for the free-electron value of m and g = 2. For all  $B(r_s)$  above the full curve,  $L \leq 1$  and Eq. (1) with arbitrary values of  $n_0^{\downarrow}$ ,  $n_0^{\uparrow}$ , and  $n_1^{\downarrow}$  may be applied. For g = 2,  $n_1^{\uparrow}$  must be zero to avoid degeneracy with the L = 2 Landau level, not included in our formula. For g < 2,  $n_1^{\uparrow}$  is also arbitrary.

For values of  $B(r_s)$  above the dashed curve, Eq. (10) with arbitrary values of  $n = n_0^{\downarrow}$  may be applied. For g = 2,  $n_0^{\uparrow}$  must be zero to avoid degeneracy with the L = 1 Landau level, not included in the DG formula. For g < 2,  $n_0^{\uparrow}$  is also arbitrary. The relevant expression in this case is our Eq. (1), but with only the  $e_x^{(0)}$  term kept.

The restrictions L = 0 and  $L \leq 1$  are precisely defined and easily applied. The restriction  $p \ll 1$ which leads from Eq. (10) to Eq. (11) and the condition  $\left|\ln\left(n/(eB)^{3/2}\right)\right| \gg 2.11788$ , which leads from Eq. (11) to Eq. (14), are less precisely defined, and we simply adopt as validity criterium that  $p \leq 0.05$  and that 2.11788 is less than 5% of the logarithmic term. From these criteria, we find that for  $B(r_s)$  above the dotted curve, the high-field limit (11) becomes valid. The inset shows the values of  $B(r_s)$  above which the ultra-high-field limit (14) is valid.

To generate Figs. 1 and 2 we used the free-electron value of the electron mass m, requiring very high magnetic fields to satisfy conditions (5) to (8). Such high fields are not without physical relevance. Continuous fields ~ 40T and pulsed fields ~ 10<sup>4</sup>T can be produced in the laboratory, white dwarfs have surface fields of order ~ 10<sup>4</sup>T, and neutron stars can have fields in excess of  $10^8 - 10^9T$ . For effective masses smaller than the free-electron mass, the required fields are considerably lower: the use of effective masses  $m^* = \gamma m$  rescales all B values by  $B \rightarrow \gamma^2 B$ , so that a reduction of m by a factor of 10 allows a reduction of B by a factor of 100. Small effective masses and low densities thus bring these fields into the laboratory range.

#### **IV. CONCLUSIONS**

Our key result is Eq. (1), for the exchange energy of the three-dimensional Fermi gas in magnetic fields for which both the lowest and the second-lowest Landau level contribute, and the fermions may have arbitrary spin polarization. This equation predicts the existence of two physically distinct regimes, one (entered with a discontinuous derivative) in which occupation of the L = 1 level lowers the per-particle exchange energy, and one (entered through a gradual crossover) in which it increases it. For high, but achievable, fields both the discontinuous drop and the gradual crossover between the two regimes occur in the metallic density range, so both regimes should be experimentally observable. We predict that similar separation in two regimes occurs every time a new Landau level is included in the calculation.

The availability of an analytical expression for the exchange energy of the electron gas in high magnetic fields facilitates the construction of density functionals for magnetic-field Thomas-Fermi-Dirac theory,<sup>1,14,15</sup>

current-density-functional theory,<sup>6,7</sup> and magnetic-field density-functional theory,<sup>8,9</sup> but the intricate structure of  $e_x(n, B)$  implies that their local approximations must display a much more complex behaviour than those of ordinary Thomas-Fermi-Dirac and spin-density-functional theory.

This work was supported by FAPESP and CNPq. We thank G. Vignale for useful remarks on an earlier version of this manuscript.

## APPENDIX A: DETAILS ON THE DERIVATION OF EQS. (1) TO (4)

From Eq. (2.1) of DG, and following their procedure until their Eq. (2.5), we find the per volume exchange energy

$$e_x = -\frac{e^2}{4\pi^3} \left(\frac{m\omega_c}{\hbar}\right)^2 \sum_{\sigma} \int_{-\infty}^{\infty} dr_z \int_{0}^{\infty} d\overline{r} \frac{\overline{r}e^{-\frac{m\omega_c}{2\hbar}\overline{r}^2}}{\left(\overline{r}^2 + r_z^2\right)^{1/2} r_z^2} \\ \times \left\{ \sum_{L=0}^{\infty} \Theta\left(C_L^{\sigma}\right) \sin\left(r_z D_L^{\sigma}\right) L_L\left(\frac{m\omega_c}{2\hbar}\overline{r}^2\right) \right\}^2 (A1)$$

where  $\Theta(x)$  is the step function,  $L_L(x)$  are Laguerre polynomials, and we defined

$$C_L^{\sigma} = \mu - \left(L + \frac{1}{2}\right)\hbar\omega_c - g\mu_0 B\sigma, \qquad (A2)$$

$$D_L^{\sigma} = \sqrt{\frac{2m}{\hbar^2} \left(\mu - \left(L + \frac{1}{2}\right) \hbar \omega_c - g\mu_0 B\sigma\right)}, \quad (A3)$$

where  $\mu_0 = e\hbar/2mc$  is the Bohr magneton.

Keeping the contributions from the L = 0 and L = 1 levels and performing the change of variable

$$\bar{r}^2 = \frac{2\hbar}{m\omega_c} \left( r_z t + \frac{\hbar}{2m\omega_c} t^2 \right) \tag{A4}$$

we obtain

$$e_{x} = -\frac{e^{2}}{2\pi^{3}} \frac{m\omega_{c}}{\hbar} \sum_{\sigma} \int_{0}^{\infty} dt \, e^{-\frac{\hbar t^{2}}{2m\omega_{c}}} \int_{0}^{\infty} dr_{z} \, \frac{e^{-r_{z}t}}{r_{z}^{2}} \left\{ \Theta(C_{0}^{\sigma}) \sin^{2}\left(D_{0}^{\sigma}r_{z}\right) + \Theta(C_{1}^{\sigma}) \sin^{2}\left(D_{1}^{\sigma}r_{z}\right) \left(1 - tr_{z} - \frac{\hbar t^{2}}{2m\omega_{c}}\right)^{2} + 2 \Theta(C_{0}^{\sigma})\Theta(C_{1}^{\sigma}) \sin\left(D_{0}^{\sigma}r_{z}\right) \sin\left(D_{1}^{\sigma}r_{z}\right) \left(1 - tr_{z} - \frac{\hbar t^{2}}{2m\omega_{c}}\right) \right\}$$
(A5)

Using Laplace Transforms to perform the  $r_z$  integral,<sup>25</sup> we find

$$e_x = -\frac{e^2}{2\pi^3} \frac{m\omega_c}{\hbar} \sum_{\sigma} \int_0^\infty dt \, e^{-\frac{\hbar t^2}{2m\omega_c}} \left\{ \Theta(C_0^{\sigma}) \left[ D_0^{\sigma} \tan^{-1}\left(\frac{2D_0^{\sigma}}{t}\right) - \frac{1}{4}t \ln\left(1 + \frac{4D_0^{\sigma^2}}{t^2}\right) \right] \right\}$$

$$+\Theta(C_{0}^{\sigma})\Theta(C_{1}^{\sigma})\left[\left(t-\frac{\hbar}{4m\omega_{c}}t^{3}\right)\ln\left(\frac{t^{2}+(D_{0}^{\sigma}-D_{1}^{\sigma})^{2}}{t^{2}+(D_{0}^{\sigma}+D_{1}^{\sigma})^{2}}\right)\right.\\\left.+\left(\frac{\hbar}{2m\omega_{c}}t^{2}-1\right)\left(D_{0}^{\sigma}-D_{1}^{\sigma}\right)\tan^{-1}\left(\frac{D_{0}^{\sigma}-D_{1}^{\sigma}}{t}\right)\right.\\\left.+\left(1-\frac{\hbar}{2m\omega_{c}}t^{2}\right)\left(D_{0}^{\sigma}+D_{1}^{\sigma}\right)\tan^{-1}\left(\frac{D_{0}^{\sigma}+D_{1}^{\sigma}}{t}\right)\right]\right.\\\left.+\Theta(C_{1}^{\sigma})\left[D_{1}^{\sigma}\left(1-\frac{\hbar}{2m\omega_{c}}t^{2}\right)^{2}\tan^{-1}\left(\frac{2D_{1}^{\sigma}}{t}\right)\right.\\\left.+\left(-\frac{3t}{4}+\frac{\hbar}{2m\omega_{c}}t^{3}-\frac{\hbar^{2}}{16m^{2}\omega_{c}^{2}}t^{5}\right)\ln\left(1+\frac{4D_{1}^{\sigma^{2}}}{t^{2}}\right)+2D_{1}^{\sigma^{2}}\frac{t}{t^{2}+4D_{1}^{\sigma^{2}}}\right]\right\}$$
(A6)

The integral over the first term, involving only contributions from L = 0, can be calculated following the procedure of Appendix A of the DG paper.<sup>10</sup> The other integrals are calculated by integration by parts and Laplace transforms. In particular, the integrals involving  $\tan^{-1}(x)$  can be calculated by using the result of the first integral.

 $C_L^{\sigma}$  and  $D_L^{\sigma}$  can be written in terms of the occupation number  $n_L^{\sigma}$  as  $C_L^{\sigma} = (2\pi^4\hbar^2 l^4/m)n_L^{\sigma^2}$  and  $D_L^{\sigma} = 2\pi^2 l^2 n_L^{\sigma}$ . The step functions turn out to be unnecessary, because the term arrising from L = 1, and the mixed term containing contributions from L = 0 and L = 1, are automatically zero when  $n_1^{\sigma} = 0$ . We thus find the exchange energy as given Eqs. (1) - (4). More details of the calculation are available through Ref. 26.

An extension of these analytical calculations to higher L values seems extraordinarily cumbersome. To include, e.g., L = 2 one must keep one more term in the sum of Eq. (A1), which gives rise to not just one more intra-level integral in Eq. (A5), but also to many new inter-level integrals. In the general case, for arbitrary L, there is no guarantee that the resulting integrals can all be solved by Laplace transforms, and reduced to some known special function. The three-dimensional case is, in this regard, more complicated than the two-dimensional one,<sup>27</sup> because of the additional  $k_z$  dependence of the single-particle energies and the resulting Landau-level dispersion. Numerical work on the general case is under way.

## APPENDIX B: DETAILS ON THE DERIVATION OF EQS. (5) TO (8)

The conditions specifying the fields and densities for which restriction to L = 0 or  $L \leq 1$  is valid depend on the Landau-level degeneracy. The single-particle dispersion in the spin  $\sigma$  sublevel of Landau level L is

$$\epsilon(k_z, L, \sigma) = \frac{\hbar^2}{2m} k_z^2 + \left(L + \frac{1}{2}\right) \hbar \omega_c + g\mu_0 B\sigma. \quad (B1)$$

In each subband, the Fermi momentum is related to the partial density by  $k_{FL}^{\sigma} = 2\pi^2 l^2 n_L^{\sigma}$ . The degeneracy of these levels depends on the value of g. For the freeelectron value g = 2, the spin-up subband of level L and the spin-down subband of level L+1 are degenerate. For g = 0, on the other hand, the spin-up and spin-down sublevels of Landau level L are degenerate. We first deal with g = 2.

The restriction to L = 0 implies

$$\epsilon(k_z = 0, L = 1, \downarrow) = \epsilon(k_z = 0, L = 0, \uparrow) \tag{B2}$$

$$> \epsilon_F = \epsilon(k_{F0}^{\downarrow}, L = 0, \downarrow).$$
 (B3)

From Eq. (B1), and using the fact that for L = 0 and  $g = 2, n_0^{\downarrow} = n$ , we immediately find

$$n^2 < \frac{1}{2\pi^4 l^6},$$
 (B4)

which is equivalent to Eq. (7) of the main text. Similarly, the restriction to  $L \leq 1$  implies

$$\epsilon(k_z = 0, L = 2, \downarrow) = \epsilon(k_z = 0, L = 1, \uparrow)$$
 (B5)

$$> \epsilon_F = \epsilon(k_{F1}^{\downarrow}, 1, \downarrow) = \epsilon(k_{F0}^{\uparrow}, 0, \uparrow) = \epsilon(k_{F0}^{\downarrow}, 0, \downarrow).$$
(B6)

The total density is  $n = n_0^{\downarrow} + n_0^{\uparrow} + n_1^{\downarrow}$ . Solving this set of equations for n yields

$$n < \frac{2 + \sqrt{2}}{\sqrt{2}\pi^2 l^3},\tag{B7}$$

which is equivalent to Eq. (5) of the main text.

The corresponding conditions for g = 0 follow in the same way. The case g = 0 is occasionally used as a methodological device in theoretical work, because it allows to cleanly separate the effects of spin magnetism from those of orbital magnetism. Experimentally, g = 0 occurs, *e.g.*, in systems studied in the context of *g*-factor engineering of devices.<sup>28</sup>

When |g| < 2 and  $g \neq 0$ , no such degeneracy occurs. The restriction to L = 0 then implies

$$n < \frac{\sqrt{2 - |g|} + \sqrt{2}}{2\pi^2 l^3},$$
 (B8)

where  $n = n_0^{\uparrow} + n_0^{\downarrow}$ .

- \* Electronic address: capelle@if.sc.usp.br
- <sup>1</sup> D. Lai, Rev. Mod. Phys. **73**, 629 (2001).
- <sup>2</sup> High Magnetic Fields in the Physics of Semiconductors, eds. G. Landwehr and W. Ossau (World Scientific, New York, 1997).
- <sup>3</sup> High Magnetic Fields in Semiconductor Physics III: Quantum Hall Effect, Transport and Optics, ed. G. Landwehr, (Springer Series in Solid-State Sciences, New York, 1992).
- <sup>4</sup> H. Ruder, G. Wunner, H. Herold, and F. Geyer, Atoms in Strong Magnetic Fields (Springer, Berlin, 1994).
- <sup>5</sup> Atoms and Molecules in Strong Magnetic Fields, eds. P. Schmelcher and W. Schweizer (Plenum, New York, 1998).
- <sup>6</sup> G. Vignale and M. Rasolt, Phys. Rev. Lett. **59**, 2360 (1987).
- <sup>7</sup> G. Vignale and M. Rasolt, Phys. Rev. B **37**, 10685 (1988).
- <sup>8</sup> C. J. Grayce and R. A. Harris, Phys. Rev. A **50**, 3089 (1994).
- <sup>9</sup> F. R. Salsbury and R. A. Harris, J. Chem. Phys. **107**, 7350 (1997).
- <sup>10</sup> R. W. Danz and M. L. Glasser, Phys. Rev. B 4, 94 (1971).
- <sup>11</sup> B. Banerjee, D. H. Constantinescu and P. Rehak, Phys. Rev. D 10, 2384 (1974).
- <sup>12</sup> N. J. M. Horing, R. W. Danz, and M. L. Glasser, Phys. Rev. A 6, 2391 (1972).
- <sup>13</sup> K. Yonei and T. Motomochi, J. Phys. Soc. Japan **59**, 3571 (1990).
- <sup>14</sup> I. Fushiki, E. H. Gudmundsson and C. J. Pethick, Astrophys. J. **342**, 958 (1989).
- <sup>15</sup> I. Fushiki, E. H. Gudmundsson, C. J. Pethick and J. Yngvason, Ann. Phys. **216**, 29 (1992).
- <sup>16</sup> M. Steinberg and J. Ortner, Phys. Rev. B **58**, 15460 (1998).
- <sup>17</sup> Y. Takada and H. Goto, J. Phys. Condens. Matter **10**, 11315 (1998).
- <sup>18</sup> P. Skudlarski and G. Vignale, Phys. Rev. B 48, 8547 (1993).

- <sup>19</sup> G. Vignale, P. Skudlarski and M. Rasolt, Phys. Rev. B 45, 8494 (1992).
- <sup>20</sup> A. K. Dhara and S. K. Ghosh, Phys. Rev. A **41**, 4653 (1990).
- <sup>21</sup> P. B. Jones, Phys. Rev. Lett. **55**, 1338 (1985).
- <sup>22</sup> K. Capelle and E. K. U. Gross, Phys. Rev. Lett. **78**, 1872 (1997).
- <sup>23</sup> Y. L. Luke, The special functions and their approximations Vol.I (Academic Press, New York, 1969).
- <sup>24</sup> After completion of this work we became aware of footnote 22 of A. R. P. Rau, R. O. Mueller and L. Spruch, Phys. Rev. A **11**, 1865 (1975), which confirms our suspicion regarding Eq. (15). Some other equations of Ref. 10 are also in need of minor corrections: in their Eq. (2.3), the exponent of the last exponential should be  $-\hbar n\omega_c s$  instead of  $-\hbar n\omega_c s/2$ , and in their Eq. (2.6)  $(\bar{r}^2 + r_z^2)^{1/2}$  should be  $(\bar{r}^2 + r_z^2)^{-1/2}$ . These problems seem to be mere misprints, however, since their correction leads to Eq. (11) [= Eq.(2.9.) of DG] as published in DG and reproduced by us. The wrong factors in Eq. (15) [=Eq. (1.1) of DG], on the other hand, are completely independent of these additional corrections, as is the difference between the DG and BCR expressions.
- <sup>25</sup> A. Erdélyi et al., *Tables of Integral Transforms* Vol. I (McGraw-Hill, New York, 1954).
- <sup>26</sup> J. M. Morbec, Ph.D. thesis, University of São Paulo, 2008 (in preparation, copy available upon request).
- <sup>27</sup> A. H. MacDonald, H. C. A. Oji, and K. L. Liu, Phys. Rev. B **34**, 2681 (1986).
- <sup>28</sup> V. F. Sapega, M. Cardona, K. Ploog, E. L. Ivchenko and D. N. Mirlin, Phys. Rev. B **45**, 4320 (1992). X. W. Zhang, Appl. Phys. Lett. **90**, 193111 (2007). S. V. Zaitsev et al, JETP Lett. **84**, 436 (2006). H. Kosaka et al., Elect. Lett. **37**, 464 (2001).