

MARKOV JUMP PROCESSES APPROXIMATING A NON-SYMMETRIC GENERALIZED DIFFUSION*

Nedžad Limić †

February 6, 2020

Abstract

Consider a non-symmetric generalized diffusion $X(\cdot)$ in \mathbb{R}^d determined by the differential operator $A(\mathbf{x}) = -\sum_{ij} \partial_i a_{ij}(\mathbf{x}) \partial_j + \sum_i b_i(\mathbf{x}) \partial_i$. In this paper the diffusion process is approximated by Markov jump processes $X_n(\cdot)$ in homogeneous and isotropic grids $G_n \subset \mathbb{R}^d$ which converge in distribution to diffusion. The generators of $X_n(\cdot)$ are constructed explicitly. Due to the homogeneity and isotropy of grids the proposed method for $d \geq 3$ can be applied to processes for which the diffusion tensor $\{a_{ij}(\mathbf{x})\}_{11}^{dd}$ fulfills an additional condition. The proposed construction offers a simple method for simulation of sample paths of non-symmetric generalized diffusion. Simulations are carried out in terms of jump processes $X_n(\cdot)$. For $d = 2$ the construction can be easily implemented into a computer code.

AMS subject classification: 60H35, 60J22, 60J27, 60J60, 65C30

Key words: Symmetric diffusion, Approximation of diffusion, Simulation of diffusion, Divergence form operators

1 INTRODUCTION

A symmetric tensor valued function $\mathbf{x} \mapsto a(\mathbf{x}) = \{a_{ij}(\mathbf{x})\}_{11}^{dd}$ which is measurable, bounded and strictly positive definite on \mathbb{R}^d defines a second order differential operator in divergence form on \mathbb{R}^d , $A_0(\mathbf{x}) = -\sum_{ij} \partial_i a_{ij}(\mathbf{x}) \partial_j$. Each $A_0(\mathbf{x})$ determines a symmetric diffusion $X(\cdot)$ in \mathbb{R}^d . In [SZ] the process $X(\cdot)$ is approximated by a sequences of Markov jump processes $X_n(\cdot)$ (MJP), i.e. processes in continuous time and discrete state space. The object of the present analysis is a non-symmetric diffusion determined by $A(\mathbf{x}) = A_0(\mathbf{x}) + B(\mathbf{x})$, where $B(\mathbf{x}) = \sum_i b_i(\mathbf{x}) \partial_i$, and a construction of its approximations by MJPs. In this way we offer an efficient method for simulation of sample paths of a non-symmetric generalized diffusion by using MJPs. Each $X_n(\cdot)$ can be simulated by well-known methods.

In the case of classical diffusion determined by an elliptic operator of form $A(\mathbf{x}) = -\sum_{ij} a_{ij}(\mathbf{x}) \partial_i \partial_j + \sum_i b_i(\mathbf{x}) \partial_i$, where a_{ij}, b_i are Hölder continuous on \mathbb{R}^d , approximations by MJPs can be efficiently used to simulate the first exit from a bounded set of \mathbb{R}^d . However, such an approach is one of existing possibilities and the motivation for the mentioned construction happen to be of lesser importance since the process has a representation in terms of SDE which can be simulated straightforwardly. For one-dimensional generalized diffusion there exist representations in terms of SDE as described by Étoré [Et] and Lejay & Martinez [LM], so that such representations can be used for simulation. In the case of a

*Supported by grant 0037014 of the Ministry of Science, Higher Education and Sports, Croatia.

†Dept. of Mathematics, University of Zagreb, Bijenička 30, 10002 Zagreb, Croatia, e-mail: nlimic@math.hr

process defined by a differential operator in divergence form on \mathbb{R}^d , $d > 1$, there is no such natural representation, so approximations by MJPs become essential.

A class of convergent approximations $X_n(\cdot)$ in [SZ] is constructed by using discretizations of the corresponding Dirichlet form $(v, u) \mapsto a(v, u)$. The functions $\partial_i v, \partial_i u$ in $a(v, u)$ are approximated by forward difference operators in local basis which generally varies. This approach is anticipated in [MW] without any remarks on the convergence of constructed MJPs $X_n(\cdot)$. In our approach the MJPs $X_n(\cdot)$ are constructed in terms of generators $A_n(\text{gen})$ on grids,

$$G_n = \left\{ h_n \sum_{i=1}^d k_i \mathbf{e}_i : k_i \in \mathbb{Z} \right\}, \quad h_n = 2^{-n}, \quad (1)$$

with a fixed basis $\{\mathbf{e}_i\}_1^d$. The generators $A_n(\text{gen})$ are constructed explicitly from a general principle which is not directly related to forward difference operators. Then the discretizations $h^d a_n(v, u)$ of original Dirichlet form are associated to the constructed generators $A_n(\text{gen})$. It happens that $h^d a_n(v, u)$ cannot be simply obtained from discretizations of original form by using forward/backward difference operators. The obtained class of $A_n(\text{gen})$ cannot be found among generators constructed in [SZ].

Advantages and drawbacks of the present construction can be briefly described as follows.

Advantages: The convergence of MJPs is proved for a non-symmetric generalized diffusion. The generators $A_n(\text{gen})$ are explicitly given in terms of values of functions a_{ij}, b_i . For the case of $d = 2$ the matrix entries of $A_n(\text{gen})$ can be easily implemented into a computer code. For $d = 2$ the construction covers a general matrix valued function $a(\cdot)$ on \mathbb{R}^2

Disadvantages: For $d \geq 3$ the proposed construction is not valid for all $a(\cdot)$ on \mathbb{R}^d . Let us define an auxiliary matrix valued function $\hat{a}(\cdot)$ by:

$$\hat{a}_{ii} = a_{ii}, \quad \hat{a}_{ij} = -|a_{ij}|, \quad i \neq j. \quad (2)$$

The here proposed construction of $A_n(\text{gen})$ is valid only if $\hat{a}(\cdot)$ is strictly positive definite on \mathbb{R}^d . There are simple examples for $d \geq 3$ in which $\hat{a}(\cdot)$ is indefinite. It happens that changes of local basis for $d \geq 3$ is necessary in the case of an indefinite $\mathbf{x} \mapsto \hat{a}(\mathbf{x})$.

Now we can describe basic steps in construction and proofs. Let $U(\cdot)$ be the strongly continuous semigroup in the Banach spaces $\dot{C}(\mathbb{R}^d)$ (continuous functions vanishing at infinity) which is associated with the diffusion process $X(\cdot)$ [EK]. Discretizations of $\dot{C}(\mathbb{R}^d)$ in terms of grid-functions on G_n are denoted by $\dot{l}_\infty(G_n)$. The generators $A_n(\text{gen})$ determine semigroups $U_n(\cdot)$ in $\dot{l}_\infty(G_n)$. There exist continuous mappings $\Phi_n : \dot{l}_\infty(G_n) \mapsto \dot{C}(\mathbb{R}^d)$ and $\Phi_n^{-1} : \dot{C}(\mathbb{R}^d) \mapsto \dot{l}_\infty(G_n)$ having the following properties: $\|\Phi_n\| = 1, \|\Phi_n^{-1}\| \leq 1$ and $\Phi_n^{-1}\Phi_n = I$ on $\dot{l}_\infty(G_n)$. For $f \in \dot{C}(\mathbb{R}^d)$ the grid function $\mathbf{f}_n = \Phi_n^{-1}f$ is called the discretization of f . Now, for each $t \geq 0$ the grid functions $\mathbf{u}_n(t) = \Phi_n^{-1}U(t)f$ and $U_n(t)\mathbf{f}_n$ can be compared. We need the following relation:

$$\lim_{n \rightarrow \infty} \sup \{ \|U_n(t)\mathbf{f}_n - \Phi_n^{-1}U(t)f\|_\infty : t \in [0, 1] \} = 0, \quad (3)$$

where $\|\cdot\|_\infty$ is the $l_\infty(G_n)$ -norm. By Theorem 2.11 of [EK] this relation is a sufficient condition for the convergence in distribution of MJPs $X_n(\cdot)$ to the diffusion process $X(\cdot)$.

Definitions of various objects are given in Sections 2. In Section 3. a class of generators $A_n(\text{gen})$ is constructed explicitly in terms of entries $(A_n(\text{gen}))_{\mathbf{k}\mathbf{l}}, \mathbf{k}, \mathbf{l} \in \mathbb{Z}^d$. Grid-functions $\mathbf{u}_n(t) = \exp(A_n(\text{gen})t)\mathbf{f}_n$ and their images $u(n, t) = \Phi_n \mathbf{u}_n(t)$ in $\dot{C}(\mathbb{R}^d)$ are studied in Section 4. The convergence $u(n, t) \mapsto U(t)f$ in $\dot{C}(\mathbb{R}^d)$ is proved first for the case of smooth coefficients a_{ij}, b_i and then for general coefficients a_{ij}, b_i . A numerical example is given in the last section.

2 PRELIMINARIES

The Euclidean norm in \mathbb{R}^d is denoted by $|\cdot|$. The only open subsets of \mathbb{R}^d , considered in this work, are bounded and connected open sets with Lipschitz boundary [Ma, Se]. We call a subset of this kind a *Lipshitz domain*, denote it by D , and its boundary by ∂D .

The Banach spaces of functions $C^{(k)}(\mathbb{R}^d)$, $C_0^{(k)}(\mathbb{R}^d) = C_0(\mathbb{R}^d) \cap C^{(k)}(\mathbb{R}^d)$ are defined as usual, $C_0(\mathbb{R}^d)$ being a linear space of continuous functions with compact supports. Their norms are denoted by $\|\cdot\|_\infty^{(k)}$. The closure of functions in $C(\mathbb{R}^d)$ with compact supports determines the Banach space $\dot{C}(\mathbb{R}^d)$. The Hölder space of parameter $k+\alpha$, $k \in \mathbb{N}_0$, $\alpha \in (0, 1)$, is denoted by $C^{(k+\alpha)}(\mathbb{R}^d)$ and defined as the completion of $C_0^{(\infty)}(\mathbb{R}^d)$ in the norm:

$$\|u\|^{(k+\alpha)} = \|u\|_\infty^{(k)} + \sup \left\{ \frac{|\partial^k u(\mathbf{x} + \mathbf{z}) - \partial^k u(\mathbf{x})|}{|\mathbf{z}|^\alpha} : \mathbf{x} \in \mathbb{R}^d, 0 < |\mathbf{z}| \leq 1 \right\}.$$

The L_p -spaces as well as Sobolev W_p^1 -spaces are defined in a standard way [Ma, Se]. Their norms are denoted by $\|\cdot\|_p$ and $\|\cdot\|_{p,1}$, respectively. For each p , $1 \leq p \leq \infty$, the norm of $W_p^1(\mathbb{R}^d)$ is defined by $\|u\|_{p,1} = (\|u\|_p^2 + \|\nabla u\|_p^2)^{1/2}$, where $\|\nabla u\|_p = (\sum_{j=1}^d \|\partial_j u\|_p^2)^{1/2}$. We say that a function f on \mathbb{R}^d belongs to a class $C^{(k+\alpha)}$ on \mathbb{R}^d if $\|f\|^{(k+\alpha)} < \infty$.

In this article we consider a 2nd-order elliptic operator on \mathbb{R}^d ,

$$A(\mathbf{x}) = - \sum_{i,j=1}^d \partial_i a_{ij}(\mathbf{x}) \partial_j + \sum_{j=1}^d b_j(\mathbf{x}) \partial_j, \quad (4)$$

for which the coefficients must fulfill the following:

Assumption 2.1 *The functions $a_{ij} = a_{ji}$, b_i , $i, j = 1, 2, \dots, d$, are measurable on \mathbb{R}^d and have the following properties:*

a) *There are positive numbers $\underline{M}, \overline{M}$, $0 < \underline{M} \leq \overline{M}$, such that the strict ellipticity is valid:*

$$\underline{M} |\mathbf{z}|^2 \leq \sum_{i,j=1}^d a_{ij}(\mathbf{x}) z_i \bar{z}_j \leq \overline{M} |\mathbf{z}|^2, \quad \mathbf{x} \in \mathbb{R}^d. \quad (5)$$

b) *The functions b_i are bounded on \mathbb{R}^d .*

In our analysis we regularly use the notation $A_0(\mathbf{x}) = -\sum_{i,j=1}^d \partial_i a_{ij}(\mathbf{x}) \partial_j$, $B(\mathbf{x}) = \sum_{j=1}^d b_j(\mathbf{x}) \partial_j$ and $A(\mathbf{x}) = A_0(\mathbf{x}) + B(\mathbf{x})$. The following real bilinear form on $W_q^1(D) \times W_p^1(D)$, $1/p + 1/q = 1$, defined by:

$$a(v, u) = \sum_{i,j=1}^d \int a_{ij}(\mathbf{x}) \partial_i v(\mathbf{x}) \partial_j u(\mathbf{x}) dx - \sum_{i=1}^d \int b_i(\mathbf{x}) v(\mathbf{x}) \partial_i u(\mathbf{x}) dx \quad (6)$$

is associated with the differential operator $A(\mathbf{x})$.

A basic result towards a proof of (3) is the following theorem [Sr]:

THEOREM 2.1 *Let the differential operator $A(\mathbf{x})$ be defined by (4) and Assumption 2.1. Then $-A(\mathbf{x})$ has the closure in $\dot{C}(\mathbb{R}^d)$ generating a Feller semigroup in $\dot{C}(\mathbb{R}^d)$.*

The linear space of grid-functions on G_n is denoted by $l(G_n)$. Elements of $l(G_n)$ are also called columns. Columns with a finite number of components determine a linear space $l_0(G_n)$. The corresponding l_p -spaces are denoted by $l_p(G_n)$ and their norms by $\|\cdot\|_p$. For $p < \infty$ this norm is defined by $\|\mathbf{u}\|_p = [\sum_{\mathbf{k} \in I_n} |u_{\mathbf{k}}|^p]^{1/p}$, and for $p = \infty$ by $\|\mathbf{u}\|_\infty = \sup\{|u_{\mathbf{k}}| : \mathbf{k} \in \mathbb{Z}^d\}$. The scalar product in $l_2(G_n)$ is denoted by $\langle \cdot | \cdot \rangle$ and sometimes by $(\cdot | \cdot)$.

Let $f \in C(\mathbb{R}^d)$ and the column $\mathbf{f}_n \in l(G_n)$ be defined by its components $(\mathbf{f}_n)_{\mathbf{k}} = f(h\mathbf{k})$. Then the mappings $C(\mathbb{R}^d) \ni f \mapsto \mathbf{f}_n \in l(G_n)$ are called the discretizations of f .

The shift operator $Z(\mathbf{x})$, $\mathbf{x} \in \mathbb{R}^d$, acting on functions $f : \mathbb{R}^d \mapsto \mathbb{R}$, is defined by $(Z(\mathbf{x})f)(\mathbf{x}) = f(\mathbf{x} + \mathbf{z})$. Similarly we define the discretized shift operator by $(Z_n(r, i)\mathbf{u}_n)_{\mathbf{k}} =$

$(\mathbf{u}_n)_l$, where $l = \mathbf{k} + r\mathbf{e}_i$. With respect to a grid step h , the partial derivatives of $u \in C^{(1)}(\mathbb{R}^d)$ are discretized by forward/backward finite difference operators in the usual way,

$$\begin{aligned}\bar{\partial}_i(t)u(\mathbf{x}) &= \frac{1}{t}(u(\mathbf{x} + t\mathbf{e}_i) - u(\mathbf{x})), \\ \widehat{\partial}_i(t)u(\mathbf{x}) &= \frac{1}{t}(u(\mathbf{x}) - u(\mathbf{x} - t\mathbf{e}_i)),\end{aligned}\quad \mathbf{x} \in \mathbb{R}^d, t \neq 0. \quad (7)$$

Let $r \in \mathbb{Z} \setminus \{0\}$. Discretizations of the functions $\partial_i u$ on G_n , denoted by $U_i(r)\mathbf{u}_n, V_i(r)\mathbf{u}_n$, are defined by:

$$(U_i(r)\mathbf{u}_n)_m = \bar{\partial}_i(rh)u(\mathbf{x}_m), \quad (V_i(r)\mathbf{u}_n)_m = \widehat{\partial}_i(rh)u(\mathbf{x}_m),$$

where $\mathbf{x} \in G_n$. Then

$$\begin{aligned}U_i(r) &= (rh)^{-1}(Z_n(r, i) - I), \\ V_i(r) &= (rh)^{-1}(I - Z_n(-r, i)) = U_i(-r) = -U_i(r)^T.\end{aligned}\quad (8)$$

Therefore we have $U_i(-r) = U_i(r)Z_n(-r, i) = Z_n(-r, i)U_i(r)$, and similarly for $V_i(r)$. In the case of $r = 1$ we use abbreviations U_i, V_i .

A matrix A_n on G_n is called a matrix of positive type if the diagonal entries of A_n are positive, off-diagonal entries are non-positive and the row sums are non-negative. If $A_n(\text{gen})$ is the generator of a MJP in G_n then $-A_n(\text{gen})$ is a matrix of positive type.

3 CONSTRUCTION OF GENERATORS OF MJPs

To discretize $A(\mathbf{x})$ means to associate to $A(\mathbf{x})$ a sequence of matrices A_n on $G_n, n \in \mathbb{N}$, with the following properties:

$$a(v, u) = \lim_{n \rightarrow \infty} h^d \langle \mathbf{v}_n | A_n \mathbf{u}_n \rangle, \quad v, u \in C_0^{(1)}(\mathbb{R}^d).$$

The convergence analysis is postponed to the next sections. Therefore, the terminology "discretizations" of $A(\mathbf{x})$ instead of approximations of $A(\mathbf{x})$ seems to be more suitable at the beginning of construction.

It is important to underline at the beginning that discretizations A_n of differential operator $A_0(\mathbf{x})$ are derived from a general principle which is not based on finite difference formulas [LR2]. Nevertheless, bilinear forms must be associated to A_n so that A_n can be derived from the corresponding variational equalities. The constructed bilinear forms are considered as discretizations of the original form (6). These forms are basic objects in our proof of convergence of approximate solutions.

Discretizations to be considered in this section are possible if certain conditions on a_{ij} are fulfilled. The required conditions are stronger than in Assumption 2.1. By relaxing them gradually as $n \rightarrow \infty$ we obtain discretizations for a general $A(\mathbf{x})$ given by Assumption 2.1.

To each pair $\mathbf{v} \in G_n, \mathbf{p} \in \mathbb{N}^d$, there is associated a rectangle $C_n(\mathbf{p}, \mathbf{v}) = \prod_{i=1}^d [v_i, v_i + hp_i]$ with the "lower left" vertex \mathbf{v} and edges of size hp_i . These rectangles define a partition of \mathbb{R}^d . Apart from these rectangles we need the closed rectangles,

$$S_n(\mathbf{p}, \mathbf{v}) = \prod_{i=1}^d [v_i - hp_i, v_i + hp_i], \quad (9)$$

which are defined by central grid-knots \mathbf{v} . Apparently, $S_n(\mathbf{p}, \mathbf{v})$ is the union of closures of those rectangles $C_n(\mathbf{p}, \mathbf{x})$ which share the grid-knot \mathbf{v} .

Discretizations A_n are defined in terms of its matrix entries $(A_n)_{\mathbf{k}l}$, where $h\mathbf{k}, h\mathbf{l} \in G_n$. For a fixed $\mathbf{x} = h\mathbf{k} \in G_n$ the set of all the grid-knots $\mathbf{y} = h\mathbf{l}$ such that $(A_n)_{\mathbf{k}l} \neq 0$ is denoted by $\mathcal{N}(\mathbf{x})$ and called the *numerical neighborhood* of A_n at $\mathbf{x} \in G_n$. A set $\mathcal{N}(\mathbf{x})$ contains always a "cross" consisting of \mathbf{x} and $2d$ elements $\pm h\mathbf{e}_i$. Additional elements of $\mathcal{N}(\mathbf{x})$ depend on the sign of $a_{ij}, i \neq j$. In terms of of MJP $X_n(\cdot)$ a set $\mathcal{N}(\mathbf{x})$ is the subset of states of possible jumps from the state \mathbf{x} .

General scenery

In order to give a comprehensive insight into the proposed construction of discretizations it is convenient to begin the construction with a differential operator $A_0 = -\sum_{ij} a_{ij} \partial_i \partial_j$ with a constant diffusion tensor $a = \{a_{ij}\}_{11}^{dd}$. In this case the matrices A_n must have a property which is called the consistency conditions. Let $\mathbf{x} \mapsto p(\mathbf{x}) = p_0 + \sum_i p_i x_i + \sum_{ij} p_{ij} x_i x_j$, $p_0, p_i, p_{ij} \in \mathbb{R}$, be a second order polynomial in arguments x_i , and let \mathbf{p}_n be its discretizations on grids G_n . Then the following identities are valid:

$$\delta(A, n, \mathbf{x})p(\mathbf{x}) := A(\mathbf{x})p(\mathbf{x}) - (A_n \mathbf{p}_n)_{\mathbf{k}} = 0, \quad \mathbf{x} = h\mathbf{k} \in G_n. \quad (10)$$

These consistency conditions are sufficient for a proof of the convergence of $U_n(t)\mathbf{f}_n$ to $\Phi_n U(t)f$ in Banach spaces of continuous functions on \mathbb{R}^d as specified in (3). Actually, the first step of construction of matrices A_n begins with a search for those matrices A_n which are simultaneously of positive type and fulfill the conditions $\delta(A, n, \mathbf{x})p(\mathbf{x}) = 0$ on grids [LR1].

To a sequence of vectors $\mathbf{w}_i, \mathbf{w}_j, \dots, \mathbf{w}_m \in \mathbb{R}^d$ we associate the cone $M(\mathbf{w}_i, \mathbf{w}_j, \dots, \mathbf{w}_m)$ consisting of elements $t_i \mathbf{w}_i + t_j \mathbf{w}_j + \dots + t_m \mathbf{w}_m$ with all possible $t_i \geq 0, t_j \geq 0, \dots, t_m \geq 0$. We demand that the discretizations A_n have the following properties:

- All the numerical neighborhood $\mathcal{N}(\mathbf{x}) \subset G_n$ are alike and the numerical neighborhood $\mathcal{N}(\mathbf{0})$ can have at most one non-trivial branch in a cone $M(\pm \mathbf{e}_i, \pm \mathbf{e}_j, \dots, \pm \mathbf{e}_m)$.
- The numerical neighborhoods $\mathcal{N}(\mathbf{0}) \subset G_n, n \in \mathbb{N}$, are alike, i.e. if $\mathbf{y} = h(n)\mathbf{l} \in \mathcal{N}(\mathbf{0}) \subset G_n$ then $\mathbf{y}' = h(m)\mathbf{l} \in \mathcal{N}(\mathbf{0}) \subset G_m$.
- The entries $(A_n)_{\mathbf{k}\mathbf{l}}$ are linear functions of $a_{ij}, 1 \leq i, j \leq d$.
- The matrices A_n are symmetric.
- The matrices A_n are consistent discretizations of A_0 .

Now we have the following result. There can be constructed matrices A_n of positive type fulfilling the conditions a)-e) if and only if \hat{a} of (2) is positive definite. The matrix A_n has the following structure:

$$\begin{aligned} (A_n)_{\mathbf{k}\mathbf{k}} &= -\sum_{\mathbf{l} \in \mathcal{N}(\mathbf{k}), \mathbf{l} \neq \mathbf{k}} (A_n)_{\mathbf{k}\mathbf{l}}, \\ (A_n)_{\mathbf{k}\mathbf{k} \pm r_i \mathbf{e}_i} &= \frac{1}{h^2 r_i^2} \left[a_{ii} - \sum_{m \neq i} \frac{r_i}{r_m} |a_{im}| \right], \\ (A_n)_{\mathbf{k}\mathbf{k} \pm \mathbf{z}} &= \frac{1}{h^2 r_i r_j} |a_{ij}|, \end{aligned} \quad (11)$$

where $\mathbf{z} = r_i \mathbf{e}_i \pm r_j \mathbf{e}_j$.

A proof of this result is simple. One starts from a general structure of matrix A_n ,

$$(A_n)_{\mathbf{k}\mathbf{l}} = (A_n)_{\mathbf{0}\mathbf{l}-\mathbf{k}} = h^{-2} \sum_{ij} \kappa(\mathbf{l} - \mathbf{k}, i, j) a_{ij},$$

where $\kappa(\mathbf{k}, i, j)$ are real numbers, which must be calculated from the conditions a)-e). Then the assertion follows from the Perron-Frobenius theorem.

In the case of $d = 2$ either both a, \hat{a} are positive definite or not. For $d = 3$ there are symmetric positive definite matrices a for which \hat{a} are indefinite. For instance, the symmetric matrix a of order $d = 3$ defined by $a_{ii} = 1, a_{12} = a_{23} = -1/\sqrt{2}$ has positive eigenvalues for $a_{13} > 0$ and a negative eigenvalue for $a_{13} < 0$.

Construction for $d = 2$

In a methodological approach one considers first constant coefficients a_{ij} and constructs the forms $a_n(v, u) = \langle \mathbf{v}_n | A_n \mathbf{u}_n \rangle$ for which the matrices A_n coincide with (11). Then the obtained expression of $a_n(v, u)$ is generalized to the case of non-constant coefficients a_{ij} . The corresponding matrices A_n are obtained naturally from the variational method. These intermediate steps of construction are omitted in the present discussion and we can write down the result of analysis.

Assumption 3.1 Let there exist a finite index set \mathcal{L} , a partition $\mathbb{R}^2 = \cup_l D_l$ and a diffusion tensor a on \mathbb{R}^2 satisfying the strict ellipticity conditions (5) and the following additional discretizations conditions:

- a) There exist $\mathbf{q} \in \mathbb{N}^2$, $n_0 \in \mathbb{N}$ and the corresponding $h_0 = 2^{-n_0}$ so that each set D_l is a connected union of cubes $\prod_{i=1}^d [x_i, x_i + q_i h_0)$. The tensor-valued function $\mathbf{x} \mapsto a(\mathbf{x})$ is continuous on D_l and the function a_{12} does not change its sign on D_l .
- b) To each D_l there is associated parameter $\mathbf{r}(l) \in \mathbb{N}^2$, such that the following inequality is valid:

$$\omega(a) = \inf_n \min_l \inf_i \left\{ \inf_{\mathbf{z} \in S_n(\mathbf{r}(l), \mathbf{x}) \cap D_l} a_{ii}(\mathbf{z}) - \sum_{m \neq i} \frac{r_i(l)}{r_m(l)} \sup_{\mathbf{z} \in S_n(\mathbf{r}(l), \mathbf{x}) \cap D_l} |a_{im}(\mathbf{z})| : \mathbf{x} \in G_n \right\} > 0. \quad (12)$$

Condition b) is crucial in our construction of discretizations A_n with the structure of matrices of positive type. In the case of $d = 2$ Condition b) can be always fulfilled.

The forms $a_n(\cdot, \cdot)$ are expressed in terms of $a_{ij}(\mathbf{x} + \mathbf{y})$, $\mathbf{x} \in G_n$, where \mathbf{y} are certain elements in \mathbb{R}^2 . The set \mathcal{L} of Assumption 3.1 is partitioned into the subsets \mathcal{L}_\mp , where $l \in \mathcal{L}_-$ means that $a_{ij} \leq 0$ on D_l and $l \in \mathcal{L}_+$ means $a_{ij} \geq 0$ on D_l . It is convenient to use a representation $a_n(v, u) = a_n^{(-)}(v, u) + a_n^{(+)}(v, u)$, where the forms $a_n^{(\mp)}(v, u)$ are related to the respective index sets \mathcal{L}_\mp . Let us define

$$\begin{aligned} \mathbf{t}^{(\pm+)}(\mathbf{r}) &= \frac{h}{2} (\pm r_1(l) \mathbf{e}_1 + r_2(l) \mathbf{e}_2) \in S_n(\mathbf{r}, \mathbf{0}), \\ \mathbf{t}^{(\pm-)}(\mathbf{r}) &= \frac{h}{2} (\pm r_1(l) \mathbf{e}_1 - r_2(l) \mathbf{e}_2) \in S_n(\mathbf{r}, \mathbf{0}). \end{aligned}$$

Then the form $a_n^{(-)}(\cdot, \cdot)$ is defined by:

$$\begin{aligned} a_n^{(-)}(v, u) &= \sum_{l \in \mathcal{L}_-} \sum_{\mathbf{x} \in G_n(l)} \left(\sum_{i=1}^2 a_{ii}(l, \mathbf{x} + \mathbf{t}^{(++)}(\mathbf{1})) (\partial_i(h)v)(\mathbf{x}) (\partial_i(h)u)(\mathbf{x}) \right. \\ &+ \sum_{i \neq j} a_{ij}(l, \mathbf{x} + \mathbf{t}^{(++)}(\mathbf{r})) (\partial_i(r_i(l)h)v)(\mathbf{x}) (\partial_j(r_j(l)h)u)(\mathbf{x}) \\ &+ \sum_{i \neq j} a_{ij}(l, \mathbf{x} + \mathbf{t}^{(++)}(\mathbf{r})) \frac{r_i(l)}{r_j(l)} \left[(\partial_i(h)v)(\mathbf{x}) (\partial_i(h)u)(\mathbf{x}) \right. \\ &\left. \left. - (\partial_i(r_i(l)h)v)(\mathbf{x}) (\partial_i(r_i(l)h)u)(\mathbf{x}) \right] \right). \end{aligned} \quad (13)$$

The form $a_n^{(+)}(v, u)$ is obtained from $a_n^{(-)}(v, u)$ formally by replacing a_{12} with $-a_{12}$, \mathcal{L}_- with \mathcal{L}_+ and $\mathbf{t}^{(++)}$ with $\mathbf{t}^{(+-)}$. The forms $a_n^{(\mp)}(v, u)$ are not the second degree polynomials of $\partial_i(h)$ with simple structure. They can be ultimately represented as forms depending on $\partial_i(q_i h)$ with various q_i .

We say that $\mathbf{x} \in G_n$ is an internal grid-knot if $\mathcal{N}(\mathbf{x}) \subset G_n \cap D_l$ for some l . All the other grid-knots are called boundary grid-knots. For an internal grid-knot \mathbf{x} the expressions of $(A_n)_{\mathbf{k}\mathbf{l}}$ follow directly from the definition of corresponding discrete forms $\mathbf{v}, \mathbf{u} \mapsto \langle \mathbf{v} | A_n \mathbf{u} \rangle$. For a boundary grid-knot $\mathbf{x} = h\mathbf{k}$ the expressions are complex and worse, the calculated $(A_n)_{\mathbf{k}\mathbf{l}}$ can lack the structure of matrices of positive type. Therefore, one seeks for possibilities of simpler procedure of construction of entries $(A_n)_{\mathbf{k}\mathbf{l}}$ for boundary grid-knots $\mathbf{x} = h\mathbf{k}$. The results of construction must be matrices A_n which determine MJPs and the convergence of MJPs determined by A_n must be ensured.

In order to write down the entries of A_n corresponding to internal grid-knots we need the following abbreviations:

$$\begin{aligned} \mathbf{w}^{(\pm)}(l) &= r_i(l) \mathbf{e}_i \pm r_j(l) \mathbf{e}_j \in I_n, \\ a_{ij}^{(\alpha\beta)}(\mathbf{r}) &= a_{ij}(\mathbf{x} + \mathbf{t}^{(\alpha\beta)}(\mathbf{r})), \quad \alpha, \beta \in \{+, -\}, \\ \hat{a}_{12}^{(-+)}(\mathbf{r}) &= a_{12}(\mathbf{x} + \mathbf{t}^{(++)}(\mathbf{r}) - h\mathbf{e}_1), \\ \hat{a}_{12}^{(+-)}(\mathbf{r}) &= a_{12}(\mathbf{x} + \mathbf{t}^{(++)}(\mathbf{r}) - h\mathbf{e}_2), \\ \hat{a}_{ii}^{(++)}(\mathbf{r}) &= a_{ii}^{(++)}(\mathbf{r}), \quad \hat{a}_{ii}^{(--) }(\mathbf{r}) = a_{ii}^{(--) }(\mathbf{r}). \end{aligned}$$




Figure 1: Numerical neighborhoods at grid-knots of D_l

Then we have the following nontrivial off-diagonal entries of A_n :

$$\begin{aligned}
 (A_n)_{\mathbf{k}\mathbf{k}\pm\mathbf{e}_1} &= -\frac{1}{h^2} \begin{cases} a_{11}^{(\pm+)}(\mathbf{1}) - \frac{r_1(l)}{r_2(l)} |\hat{a}_{12}^{(\pm+)}(\mathbf{r})| & \text{for } a_{12} \leq 0, \\ a_{11}^{(\pm-)}(\mathbf{1}) - \frac{r_1(l)}{r_2(l)} |\hat{a}_{12}^{(\pm-)}(\mathbf{r})| & \text{for } a_{12} \geq 0, \end{cases} \quad (14) \\
 (A_n)_{\mathbf{k}\mathbf{k}\pm\mathbf{e}_2} &= -\frac{1}{h^2} \left[a_{22}^{(+\pm)}(\mathbf{1}) - \frac{r_2(l)}{r_1(l)} |\hat{a}_{12}^{(+\pm)}(\mathbf{r})| \right].
 \end{aligned}$$

The entries in the plane spanned by $\mathbf{e}_1, \mathbf{e}_2$ have the structure:

$$\begin{aligned}
 (A_n)_{\mathbf{k}\mathbf{k}\pm\mathbf{w}^{(-)}(l)} &= -\frac{1}{h^2 r_1(l) r_2(l)} |a_{12}^{(\pm\mp)}(\mathbf{r})| \quad \text{for } a_{12} \leq 0 \text{ on } D_l, \\
 (A_n)_{\mathbf{k}\mathbf{k}\pm\mathbf{w}^{(+)}(l)} &= -\frac{1}{h^2 r_1(l) r_2(l)} |a_{12}^{(\pm\pm)}(\mathbf{r})| \quad \text{for } a_{12} \geq 0 \text{ on } D_l,
 \end{aligned} \quad (15)$$

where $+\mathbf{w}^{(\pm)}$ is associated with the upper and $-\mathbf{w}^{(\pm)}$ with the lower indices of $a^{(\alpha\beta)}$, respectively.

The numerical neighborhoods $\mathcal{N}(\mathbf{x})$ are illustrated in Figure 3.

If the quantities $\hat{a}_{ij}^{(\alpha\beta)}$ in (14) are replaced with $a_{ij}^{(\alpha\beta)}$, the convergence is still preserved. However, the quantities $a_{ij}^{(\alpha\beta)}$ must not be replaced with $a_{ij}(h\mathbf{k})$ since the resulting $(A_n)_{\mathbf{k}\mathbf{l}}$ would be discretizations of $-\sum_{ij} a_{ij} \partial_i \partial_j$.

Now we can describe a construction which gives a satisfactory result for both kinds of grid-knots, internal and boundary grid-knots. For each $\mathbf{x} = h\mathbf{k} \in \text{cls}(D_l)$ the entries $(A_n)_{\mathbf{k}\mathbf{l}}$ are constructed by the rules (14), (15). If $\mathcal{N}(\mathbf{x}) \subset \text{cls}(D_l)$ there is nothing to adjust. If $\mathcal{N}(\mathbf{x}) \cap \text{cls}(D_l)$ is less than $\mathcal{N}(\mathbf{x})$ and the constructed entries $(A_n)_{\mathbf{k}\mathbf{l}}, h\mathbf{l} \in \mathcal{N}(\mathbf{x})$, have the property $(A_n)_{\mathbf{k}\mathbf{l}} > 0, \mathbf{k} \neq \mathbf{l}$, the entries are accepted. If $\mathcal{N}(\mathbf{x}) \cap \text{cls}(D_l)$ is less than $\mathcal{N}(\mathbf{x})$ and some of $a_{12}^{(\alpha\beta)}(\mathbf{r})$ causes $(A_n)_{\mathbf{k}\mathbf{l}} < 0, \mathbf{k} \neq \mathbf{l}$, then these $a_{12}^{(\alpha\beta)}(\mathbf{r})$, all $\alpha, \beta \in \{\pm\}$, must be replaced with zero in all the expressions (14), (15).

In this way we determined the rules of construction of discretizations of a generalized diffusion for which the diffusion tensor satisfy Assumption 3.1. It is necessary to justify the described construction of entries $(A_n)_{\mathbf{k}\mathbf{l}}$ at boundary grid-knots. The justification is based on the convergence of generalized diffusions $X^{(n)}(\cdot)$ to the original diffusion $X(\cdot)$.

The set $\Gamma = \cup_l \partial D_l$ contains boundary grid-knots. A numerical neighborhood $\mathcal{N}(\mathbf{x}), \mathbf{x} \in D_l \cap G_n$ is contained in the closed rectangle

$$S_n(\mathbf{r}(l), \mathbf{x}) = \prod_{i=1}^d [x_i - hr_i(l), x_i + hr_i(l)]. \quad (16)$$

The union of all such rectangles centered at boundary grid-knots $\mathbf{x} \in \Gamma \cap G_n$ is denoted by $S_n(\Gamma)$. This is a closed set covering Γ . In addition, $\Gamma = \cap_n S_n(\Gamma)$. For each $n \in \mathbb{N}$ we

approximate the original diffusion tensor a by the tensor $a^{(n)}$ defined as follows:

$$\begin{aligned} a_{ij}^{(n)}(\mathbf{x}) &= a_{ij}(\mathbf{x}) \quad \text{for } \mathbf{x} \in \mathbb{R}^2 \setminus S_n(\Gamma), \quad i, j \in \{1, 2\}, \\ a_{ii}^{(n)}(\mathbf{x}) &= a_{ii}(\mathbf{x}) \quad \text{for } \mathbf{x} \in S_n(\Gamma), \\ a_{12}^{(n)}(\mathbf{x}) &= 0 \quad \text{for } \mathbf{x} \in S_n(\Gamma). \end{aligned}$$

The defined diffusion tensors $a^{(n)}$ determine differential operators $A_0^{(n)}(\mathbf{x})$ which approximate the differential operator $A_0(\mathbf{x})$. Each $A_0^{(n)}(\mathbf{x})$ determines a generalized diffusion $X^{(n)}(\cdot)$. The generalized diffusions $X^{(n)}(\cdot)$ converge in distributions to the original diffusion $X(\cdot)$. The matrices A_n are now constructed by the expressions (14), (15) where the values $a_{ij}(\mathbf{z})$ are replaced with the values $a_{ij}^{(n)}(\mathbf{z})$. The obtained A_n are matrices of positive type.

3.1 Construction for $d > 2$

The goal of overall analysis are those discretizations A_n of differential operator $A(\mathbf{x})$ which have the structure of matrices of positive type. Here we describe a general approach to the constructions of discretizations A_n with the structure of matrices of positive type which is based on reduction to the two-dimensional problems.

The index set of pairs $I(d) = \{\{ij\} : i < j, i, j = 1, 2, \dots, d, i \neq j\}$ has the cardinal number $m(d) = d(d-1)/2$. To each index $\{kl\} \in I(d)$ we associate three coefficients,

$$a_{kk}^{\{kl\}} = \frac{1}{d-1} a_{kk}, \quad a_{ll}^{\{kl\}} = \frac{1}{d-1} a_{ll}, \quad a_{kl}^{\{kl\}} = a_{kl}, \quad (17)$$

and a bilinear form $a^{\{kl\}}(\cdot, \cdot)$,

$$a^{\{kl\}}(v, u) = \sum_{i,j \in \{r,s\}} \int_{\mathbb{R}^d} a_{ij}^{\{kl\}}(\mathbf{x}) \partial_i v(\mathbf{x}) \partial_j u(\mathbf{x}) d\mathbf{x}.$$

Apparently, for each pair $v, u \in C_0^{(1)}(\mathbb{R}^d)$ the following equality is valid:

$$a(v, u) = \sum_{\{kl\} \in I(d)} a^{\{kl\}}(v, u). \quad (18)$$

To each of the forms $a^{\{kl\}}(\cdot, \cdot)$ we must associate a sequence of forms $a_n^{\{kl\}}(\cdot, \cdot)$ and matrices $A_n^{\{kl\}}$ constructed by using schemes in two dimensions. Then the matrix

$$A_n = \sum_{\{kl\} \in I} A_n^{\{kl\}}, \quad (19)$$

is a discretization of $A_0(\mathbf{x})$. If each $A_n^{\{kl\}}$ has the structure of a matrix of positive type then A_n is also a matrix of positive type. However, A_n can have the structure of a matrix of positive type although no $A_n^{\{kl\}}$ is a such matrix. Let us demonstrate the validity if this important property which enables a construction of matrices A_n of positive type for the case $d > 0$.

At the beginning we have to describe general structure of constructed discretizations of $A(\mathbf{x}) = A_0(\mathbf{x}) + B(\mathbf{x})$. It is sufficient to consider $A_0(\mathbf{x})$. From the definition of bilinear forms $a_n(u, v) = \langle \mathbf{v} | A_n \mathbf{u} \rangle$ the following property is obvious: If $\mathbf{v} = \mathbf{1}$ and \mathbf{u} are with compact supports on G_n there must be $a_n(\mathbf{1}, u) = a_n(u, \mathbf{1}) = 0$, implying that the row sums and column sums of A_n have zero values. Hence, if the off-diagonal entries of A_n are non-positive then A_n and A_n^T are simultaneously matrices of positive type.

It is convenient to utilize the quantities:

$$\omega_n(a_{ii}, \mathbf{x}) = \frac{1}{d-1} \sum_{s \neq i}^d a_{ii}(l, h\mathbf{k} + h\mathbf{m}_{ii}(l, s)), \quad (20)$$

where $\mathbf{m}_{ii}(l, s)$ are defined by the rules of construction of (19).

Discretization procedure 3.1 Let there exist a partition $\mathbb{R}^d = \cup_l D_l$ into a finite number of subsets D_l and let a parameter $\mathbf{r}(l) \in \mathbb{N}^d$ be associated to each D_l . Let matrices A_n on G_n be constructed by the rules (14), (15) and (19). Then their entries have the following properties:

1. Entries of $(A_n)_{\mathbf{k}\mathbf{l}}$, $\mathbf{x} = h\mathbf{k}$, $\mathbf{k} \in \mathbb{Z}^d$, $h\mathbf{l} \in \mathcal{N}(\mathbf{x})$ are linear combinations of $a_{ij}(l, \mathbf{x}_{ij}(n, \mathbf{x}, l))$ where $\mathbf{x}_{ij}(n, \mathbf{x}, l) = h\mathbf{k} + h\mathbf{m}_{ij}(l, s)$, $\mathbf{m}_{ij}(l, s)$ are n -independent elements of \mathbb{R}^d and $l \in \mathcal{L}$, $s \in \{1, 2, \dots, d\}$.
2. For each grid-knot $\mathbf{x} = h\mathbf{k}$: $(A_n)_{\mathbf{k}\mathbf{k}} = -\sum_l (A_n)_{\mathbf{k}\mathbf{l}}$.
3. For each $\mathbf{x} = h\mathbf{k} \in \text{cls}(D_l)$ entries on the "cross branches" $\mathbf{x} \pm h\mathbf{e}_i$, i.e. $(A_n)_{\mathbf{k}\mathbf{k} \pm h\mathbf{e}_i}$, are defined by:

$$(A_n)_{\mathbf{k}\mathbf{k} \pm \mathbf{e}_i} = -\frac{1}{h^2} \left[\omega_n(a_{ii}, \mathbf{x}) - \sum_{m \neq i} \frac{r_i(l)}{r_m(l)} |a_{im}(l, \mathbf{x}_{im}(n, \mathbf{x}, l))| \right].$$

4. Entries of A_n in the ij -plane are defined by using elements $\mathbf{z}_{ij}(l) = r_i(l)\mathbf{e}_i - r_j(l)\mathbf{e}_j \in \mathbb{Z}^d$ or elements $\mathbf{z}_{ij}(l) = r_i(l)\mathbf{e}_i + r_j(l)\mathbf{e}_j \in \mathbb{Z}^d$:

$$(A_n)_{\mathbf{k}\mathbf{k} \pm \mathbf{z}_{ij}(l)} = -\frac{1}{h^2 r_i(l) r_j(l)} |a_{ij}(l, \mathbf{x}_{ij}(n, \mathbf{x}, l))|.$$

Some peculiar features regarding the structure of sets $\mathcal{N}(\mathbf{x})$, $\mathbf{x} = h\mathbf{k} \in G_n$, must be pointed out. If $a_{ij} \neq 0$, $i \neq j$, then the maximal number of elements in $\mathcal{N}(\mathbf{x})$ is $1 + d + d^2$. In this case the set $\mathcal{N}(\mathbf{x})$ consists of its center, $2d$ -grid-knots on the d -dimensional cross $\{\pm \mathbf{e}_i : i = 1, 2, \dots, d\}$ and 2 grid-knots in each two-dimensional plane. Since there can be only two grid-knots at most in the two-dimensional planes the entries of $A_n^{(rs)}$ have the following property. Let the pairs $\mathbf{e}_r, \mathbf{e}_s$ and $\mathbf{e}_s, \mathbf{e}_t$ define two-dimensional planes and let $A_n^{(rs)}, A_n^{(st)}$ be the corresponding discretizations which are constructed by using parameters $\mathbf{r}^{(rs)}, \mathbf{r}^{(st)}$. Then there must hold $(\mathbf{r}^{(rs)})_s = (\mathbf{r}^{(st)})_s$. In such case the off-diagonal entries of A_n have the structure as described in item 4. of Discretization procedure 3.1.

3.2 Lower order differential operators

Discretizations of differential operator $-B(\mathbf{x}) = -\mathbf{b}(\mathbf{x})\nabla$ must be constructed by using an upwind scheme. Let the obtained discretizations be denoted by B_n . To each $\mathbf{x} \in G_n$ for which $\mathbf{b}(\mathbf{x}) \neq \mathbf{0}$ there is associated a non-trivial row of B_n with a positive diagonal entry and non-positive off diagonal entries. Let us define sets $\mathcal{K}(i, -) = \{\mathbf{x} \in G_n : b_i(\mathbf{x}) < 0\}$ and analogously $\mathcal{K}(i, +) = \{\mathbf{x} \in G_n : b_i(\mathbf{x}) > 0\}$. Then discretizations of $(v|Bu)$ are defined by

$$b_n(v, u) = \sum_i \left[\sum_{\mathbf{x} \in \mathcal{K}(i, -)} b_i(\mathbf{x}) v(\mathbf{x}) (\partial_i(-h)u)(\mathbf{x}) + \sum_{\mathbf{x} \in \mathcal{K}(i, +)} b_i(\mathbf{x}) v(\mathbf{x}) (\partial_i(h)u)(\mathbf{x}) \right].$$

These forms must be summed with the forms (18) in order to get discretizations of the original form (6). If discretizations $(A_0)_n$ of $A_0(\mathbf{x})$ have the structure of matrices of positive type then obviously $(A_0)_n + B_n$ preserve this structure.

The constructed forms a_n of this section are discretizations of the form (6). One can be easily convinced that the terminology "a discretization of the original form (6)" is not artificial. At the present level of analysis it is easy to check $a(v, u) = \lim_n h^d a_n(v, u)$ for any pair $v, u \in C_0^{(1)}(\mathbb{R}^d)$.

3.3 Results of construction

THEOREM 3.1 *Let there exist a partition $\mathbb{R}^d = \cup_l D_l$ into a finite number of connected sets D_l , each being the union of $C_m(\mathbf{p}, \mathbf{x})$ with some fixed m , so that a_{ij} are uniformly continuous on D_l and $a_{ij}, i \neq j$, do not change sign on D_l . Let (2) be valid. There exist discretizations A_n which are constructed by the rules of Discretization procedure 3.1, such that A_n are matrices of positive type.*

PROOF: For each D_l we have to choose the parameters $r_i(l)$ so that the entries $(A_n)_{\mathbf{k}\mathbf{k} \pm \mathbf{e}_i}$ of item 3. in Discretization procedure 3.1 have negative values. Then the rules of construction (15) ensure the existence of A_n with the structure of matrices of positive type. **QED**

The uniform continuity of functions a_{ij} on D_l and the inequality (12) imply another important property of matrices A_n which are constructed by described procedure. There exists $\sigma_0 > 0$, independent of n , such that

$$(A_n)_{\mathbf{k}\mathbf{k} \pm \mathbf{e}_i} \geq \frac{\sigma_0}{h^2} a_{ii}(h\mathbf{k}), \quad \mathbf{x} = h\mathbf{k} \in G_n. \quad (21)$$

The described construction of matrices A_n for which (21) holds is called *admissible method* anticipating that the obtained A_n have all the necessary properties for the convergence of constructed MVPs to generalized diffusion.

LEMMA 3.1 *Let the matrices A_n on G_n be discretizations of $A_0 = -\sum \partial_i a_{ij} \partial_j$ by an admissible method. Then the matrices A_n are irreducible and there exist positive numbers $\underline{M}, \overline{M}$ such that the following inequalities*

$$\underline{M} \sum_i \|U_i \mathbf{u}\|_2^2 \leq \langle \mathbf{u} | A_n \mathbf{u} \rangle \leq \overline{M} \sum_i \|U_i \mathbf{u}\|_2^2,$$

are valid uniformly with respect to $n \in \mathbb{N}$.

For a differential operator $A(\mathbf{x}) = A_0(\mathbf{x}) + \mathbf{b}(\mathbf{x})\nabla$ with non-constant coefficients the functions $G_n \ni \mathbf{x} \mapsto \delta(A, n, \mathbf{x})p(\mathbf{x})$ of (10) are not zero on G_n . For the proof of convergence in Section 4 the following weaker result is used:

LEMMA 3.2 *Let the differential operator $A(\mathbf{x})$ satisfy the conditions of Theorem 3.1 and let the following additional conditions be valid:*

- a) *Functions a_{ij} belong to the class $C^{(1+\alpha)}$ on \mathbb{R}^d .*
- b) *Functions b_i belong to the class $C^{(\alpha)}$ on \mathbb{R}^d .*

Then there exists a positive constant κ , independent of n , depending on $\overline{M}, \|\mathbf{b}\|_\infty, \|a_{ij}\|^{(1+\alpha)}$ and $\|b_i\|^{(\alpha)}$, such that

$$\sup_{\mathbf{x} \in G_n} |\delta(A, n, \mathbf{x})f(\mathbf{x})| \leq \kappa h^\alpha \|f\|^{(2+\alpha)}$$

for any $f \in C^{(2+\alpha)}(\mathbb{R}^d)$.

This result is proved by calculating $A_n \mathbf{f}_n$ directly.

Let us mention that generally $\lim_n \delta(A, n, \mathbf{x})f(\mathbf{x})$ on grid-knots $\mathbf{x} \in \text{int}(D_l)$ behave like h^α and $\delta(A, n, \mathbf{x})p(\mathbf{x})$ is bounded elsewhere. This property is sufficient for a proof of the convergence in W_2^1 -spaces [LR2].

4 CONVERGENCE OF MJPS

The convergence of MJPs to generalize diffusion is analyzed here in terms of the criterion (3). Therefore, we have to define explicitly the mappings $\Phi_n : l_\infty(G_n) \mapsto \dot{C}(\mathbb{R}^d)$.

An element (column) $\mathbf{u}_n \in l(G_n)$ can be associated to a continuous function on \mathbb{R}^d in various ways. Here is utilized a mapping $l(G_n) \mapsto C(\mathbb{R}^d)$ which is defined in terms of hat functions. Let χ be the canonical hat function on \mathbb{R} , centered at the origin and having the support $[-1, 1]$. Then $z \mapsto \phi(h, x, z) = \chi(h^{-1}(z - hx))$ is the hat function on \mathbb{R} , centered at $x \in \mathbb{R}$ with support $[x - h, x + h]$. The functions $\mathbf{z} \mapsto \phi_{\mathbf{k}}(\mathbf{z}) = \prod_{i=1}^d \phi(h, x_i, z_i)$, $x_i = hk_i$, define d -dimensional hat functions with supports $S_n(\mathbf{1}, \mathbf{x}) = \prod_i [x_i - h, x_i + h]$. The functions $\phi_{\mathbf{k}}(\cdot) \in G_n$, span a linear space, denoted by $E(n, \mathbb{R}^d)$. Let $\mathbf{u}_n \in l(G_n)$ have the entries $u_{n\mathbf{k}} = (\mathbf{u}_n)_{\mathbf{k}}$. Then the function $u(n) = \sum_{\mathbf{k} \in I_n} u_{n\mathbf{k}} \phi_{\mathbf{k}}$ belongs to $E(n, \mathbb{R}^d)$ and defines imbedding of grid-functions into the space of continuous functions. We denote the corresponding mapping by $\Phi_n : l(G_n) \mapsto E(n, \mathbb{R}^d)$ and write

$$u(n) = \Phi_n \mathbf{u}_n = \sum_{\mathbf{k}} (\mathbf{u}_n)_{\mathbf{k}} \phi_{\mathbf{k}}. \quad (22)$$

It is obvious that there exists $\Phi_n^{-1} : E(n, \mathbb{R}^d) \mapsto l(G_n)$ and the spaces $l(G_n)$ and $E(n, \mathbb{R}^d)$ are isomorphic with respect to the pair of mappings Φ_n, Φ_n^{-1} . It is clear that $E(n, \mathbb{R}^d) \subset E(n+1, \mathbb{R}^d)$ and the space of functions $\cup_n E(n, \mathbb{R}^d)$ is dense in $L_p(\mathbb{R}^d)$, $p \in [1, \infty)$, as well as in $\dot{C}(\mathbb{R}^d)$. Let us mention that $\sum_{\mathbf{k}} \phi_{\mathbf{k}} = 1$ on \mathbb{R}^d . For two functions $v(n), u(n)$ we have $(v(n)|u(n)) = h^d \sum_{\mathbf{k}, \mathbf{l}} s_{\mathbf{k}\mathbf{l}} v_{\mathbf{k}} u_{\mathbf{l}}$ where $s_{\mathbf{k}\mathbf{l}} = \|\phi_{\mathbf{k}}\|_1^{-1} (\phi_{\mathbf{k}} | \phi_{\mathbf{l}})$. Let us notice that $\sum_{\mathbf{l}} s_{\mathbf{k}\mathbf{l}} = 1$. We have $\Phi_n^{-1} \Phi_n = I$ in $l(\mathbb{R}^d)$ and $\Phi_n \Phi_n^{-1} = I$ in $E(n, \mathbb{R}^d)$, while $\Phi_n \Phi_n^{-1} = P(n)$ in $\dot{C}(\mathbb{R}^d)$, where $P(n)$ is the projector defined by $f \mapsto P(n)f = \Phi_n \mathbf{f}_n$.

If F_n is a matrix on G_n then $F(n) = \Phi_n F_n \Phi_n^{-1}$ is a linear operator in the linear space $E(n, \mathbb{R}^d)$. It is easy to verify $\|F(n)\|_p = \|F_n\|_p$ for $p = 1, \infty$, where the norm $\|\cdot\|_p$ is induced by the restriction of $L_p(\mathbb{R}^d)$ to $E(n, \mathbb{R}^d)$. By applying Rietz-Thorin theorem we get $\|F(n)\|_p \leq \|F_n\|_p$ for $p \in [1, \infty]$.

The objective of analysis in this section is the comparison of Feller semigroup $U(\cdot)$ in $\dot{C}(\mathbb{R}^d)$ and matrix semigroups $U(n, t) = \Phi_n \exp(-A_n t) \Phi_n^{-1}$ in $\dot{E}_n(\mathbb{R}^d) \subset \dot{C}(\mathbb{R}^d)$ and a proof of limit (3). For this purpose we consider two basic objects. The first one is the collection of equations:

$$\begin{aligned} (\lambda I - A)u &= f, \\ (\lambda I_n - A_n)\mathbf{u}_n &= \mathbf{f}_n, \quad n \in \mathbb{N}, \end{aligned} \quad (23)$$

consisting of a PDE on \mathbb{R}^d and its discretizations on grids G_n . The second object are the IVPs associated to (23)

$$\begin{aligned} (\partial_t - A(\mathbf{x}))u(t, \mathbf{x}) &= 0, \quad u(0, \mathbf{x}) = u_0(\mathbf{x}), \\ \dot{\mathbf{u}}_n(t) - A_n \mathbf{u}_n(t) &= \mathbf{0}, \quad \mathbf{u}_n(0) = \mathbf{u}_{0n}. \end{aligned} \quad (24)$$

By using standard methods in the Sobolev space $W_2^1(\mathbb{R}^d)$ we can prove a result which is weaker than (3). Let $A(\mathbf{x})$ be defined by (4) and Assumption 2.1 and let A_n be its discretizations on G_n constructed by the rules of Section 4. We consider $u(t) = U(t)f$ and $u(n, t) = U(n, t)f(n)$. For each $f \in W_2^1(\mathbb{R}^d)$ there must be valid [LR2]: $\lim_{n \rightarrow \infty} \|u(t) - u(n, t)\|_{2,1} = 0$ uniformly with respect to t in segments of $[0, \infty)$. Here the symbol \lim_n means that each infinite subsequence of functions $u(n)$ contains a convergent subsequence and each convergent subsequence has the same limit.

The function $u(t)$ is continuous on \mathbb{R}^d . Each function $u(n, t)$ is also continuous on \mathbb{R}^d and the sequence $\mathfrak{U} = \{u(n, t) : n \in \mathbb{N}\}$ is uniformly bounded. This sequence converges a.e. to u on \mathbb{R}^d , while we need the uniform convergence. As usually in such problems the uniform convergence can be proved eventually for a subsequence of \mathfrak{U} . In our approach the plan of proof of (3) is slightly changed. The original differential operator $A(\mathbf{x})$ is approximated by differential operators $A^{(m)}(\mathbf{x})$ with smooth coefficients. The corresponding semigroups are

denoted by $U^{(m)}(\cdot)$ and $U^{(m)}(n, \cdot)$. The limit (3) must be proved for each $m \in \mathbb{N}$. Then by applying the diagonalization argument to the sequence $\{u^{(m)}(n, t) : m, n \in \mathbb{N}\}$ we get the desired result.

Convergence for smooth coefficients

Let $E_2 = \dot{C}^{(2+\alpha)}(\mathbb{R}^d)$, and $E_2(n)$ be the closed subspace of all elements in E_2 vanishing at points of G_n . If the space $F_2(n) = E_2/E_2(n)$ is endowed with a quotient norm, it becomes isomorphic to a subspace of l_∞ . Analogously, the spaces $E_0 = \dot{C}^{(\alpha)}(\mathbb{R}^d)$, $E_0(n)$ and $F_0(n)$ are defined. Because of the mentioned isomorphism, we identify $F_2(n)$ with a subspace of $F_0(n)$. The norms of E_k are denoted by $\|\cdot\|_{k+\alpha}$, while the norms of $F_k(n)$ are denoted by $\|\cdot\|_\infty$. The natural embedding from E_k into F_k , $k = 2, 0$, is denoted by $\Psi_n(k)$.

We say that an element $u_n \in F_k(n)$ approximates $u \in E_k$ with the order $\beta > 0$ if there exists an n -independent positive number κ such that $\|\Psi_n(k)u - u_n\|_\infty < \kappa h^\beta$. Let $\mathfrak{U}(k) = \{u_n : n \in \mathbb{N}\}$ be a sequence of such elements. We say in this case that $\mathfrak{U}(k)$ factor converges to u with the order β . We say that a sequence of operators $A_n \in \mathbf{L}(F_2(n), F_0(n))$ approximates an operator $A \in \mathbf{L}(E_2, E_0)$ with the order of approximation β , if there exists an n -independent positive number κ such that

$$\|\Psi_n(0)A - A_n\Psi_n(2)\|_{\mathbf{L}(E_2, F_0(n))} \leq \kappa h^\beta. \quad (25)$$

A sequence of approximations, $\mathfrak{A} = \{A_n : n \in \mathbb{N}\} \subset \mathbf{L}(E_2, F_0(n))$, is called stable if there exists an n -independent $\rho > 0$ such that $\|(A_n)^{-1}\|_{\mathbf{L}(F_0(n), F_2(n))} \leq \rho$. Because of $F_2(n) \subset F_0(n) \subset l_\infty$ the operator A_n is considered as an infinite order matrix with matrix entries $(A_n)_{ij}$.

For a differential operator $A(\mathbf{x}) : E_2 \mapsto E_0$, $\Re \lambda > 0$ and $f \in E_0$, we consider a PDE of the form $(\lambda I - A(\mathbf{x}))u(\mathbf{x}) = f(\mathbf{x})$ on \mathbb{R}^d . Let us assume that the differential operator $A(\mathbf{x})$ is discretized on grids G_n by matrices A_n and let us define linear algebraic systems of (23) where \mathbf{f}_n are the discretizations on G_n of $f \in E_0$. Within the formulated abstract framework of solving PDE in terms of discretized equations of (23) the following result is valid [Kr]:

THEOREM 4.1 *Let the following conditions be satisfied:*

- There exist a unique solution u to the first equation of (23).*
- The sequence of matrices $\mathfrak{A} = \{\lambda I_n - A_n : n \in \mathbb{N}\}$ approximates $\lambda I - A$ on G_n with the order α .*
- \mathfrak{A} is stable.*
- The elements \mathbf{f}_n factor converge to the element f with the order α .*

Then the sequence $\mathbf{u}_n = (\lambda I_n - A_n)^{-1}\mathbf{f}_n$ factor converges with the order α to the solution u of first equation of (23).

Now we have a straightforward application of this theorem:

COROLLARY 4.1 *Let $t \mapsto U(t)$ is a strongly continuous semigroup in E_2 , $\|U(t)\|_{2+\alpha} \leq \exp(\sigma t)$ with some $\sigma \geq 0$, such that $u(t) = U(t)u_0$ solves the first IVP of (24). Let $t \mapsto U_n(t)$, $\|U_n(t)\|_\infty \leq 1$, be semigroups generated by $-A_n$ in $F_2(n) \subset F_0(n)$, such that $\mathbf{u}_n(t) = U_n(t)\mathbf{u}_{0n}$ solve the IVPs for ODE of (24). If the differential operator $A(\mathbf{x})$ on \mathbb{R}^d fulfills the condition b) of Theorem 4.1 then the following assertion is valid: For each $T > 0$ there exists a positive number $\rho(T)$ such that*

$$\|\Psi_n(2)u(t) - \mathbf{u}_n(t)\|_\infty \leq \rho(T)\|u_0\|_{2+\alpha} h^\alpha,$$

for all $u_0 \in E_2 = \dot{C}^{(2+\alpha)}(\mathbb{R}^d)$, uniformly with respect to $0 \leq t \leq T$.

PROOF: The following identity [Kr, Tb] is used in this proof:

$$\Psi_n(2)U(t) - U_n(t)\Psi_n(2) = \int_0^t U_n(t-s) (\Psi_n(0)A - A_n\Psi_n(2)) U(s) ds.$$

By using (25) we get

$$\begin{aligned} \left\| \left(\Psi_n(2)U(t) - U_n(t)\Psi_n(2) \right) u_0 \right\|_\infty &\leq \\ \int_0^t \left\| U_n(t-s) \right\|_\infty \|A_n\Psi_n(2) - \Psi_n(0)A\|_{\mathbf{L}(E_2, F_0(n))} \|U(s)\|_{2+\alpha} \|u_0\|_{2+\alpha} &\leq \\ T \exp(\sigma T) \kappa h^\alpha \|u_0\|_{2+\alpha}, & \end{aligned}$$

and the statement follows from Theorem 4.1. **QED**

Now we can turn to the differential operator $A(\mathbf{x})$ of (4). If the differential operator (4) has coefficients b_i belonging to the class $C^{(\alpha)}$ and coefficients a_{ij} belonging to the class $C^{(1+\alpha)}$ then it can be represented in another form,

$$A(\mathbf{x}) = - \sum_{ij=1}^d a_{ij}(\mathbf{x}) \partial_{ij}^2 + \sum_{i=1}^d b'_i(\mathbf{x}) \partial_i,$$

where $b'_i(\mathbf{x}) = b_i(\mathbf{x}) - \sum_j \partial_j a_{ij}(\mathbf{x})$. Hence, $A(\mathbf{x})$ can be represented as an elliptic operator in non-divergence form with coefficients belonging to the class $C^{(\alpha)}$ on \mathbb{R}^d . This form of $A(\mathbf{x})$ supplies us with a possibility to use results on the existence of a strongly continuous semigroup in $E_2 = \dot{C}^{(2+\alpha)}(\mathbb{R}^d)$ as developed by Solonnikov [So] (a concise exposition of results can be found in [GM]). The existence of a strongly continuous semigroup in E_2 is demonstrated from the representation of $U(t)$ in terms of integral operators for which kernels are fundamental solutions. By using $\|U(t)\|_{\mathbf{L}(E_2, F_0(n))} \leq \exp(\sigma t)$ and the representation $T(\lambda, A) = \int_0^\infty \exp(-\lambda t) U(t) dt$ with $\lambda > \sigma$ we conclude that the elliptic problem in (23) has solutions for any $f \in E_0$. In order to apply Corollary 4.1 to $A(\mathbf{x})$ we have to prove that the corresponding discretizations A_n of Section 3 satisfy the conditions of Theorem 4.1.

The matrices $A_n \in \mathbf{L}(F_2(n), F_0(n))$ of Section 4 approximate $A(\mathbf{x})$ with the order α as demonstrated in Lemma 3.2. Each matrix A_n is of positive type so that the equations $(\lambda I_n - A_n)\mathbf{u}_n = \mathbf{f}_n$, $\Re \lambda > 0$, have unique solutions in $F_k(n)$ satisfying $\|\mathbf{u}_n\|_\infty < \|\mathbf{f}_n\|_\infty / \Re \lambda$. Thus the approximations $\lambda I_n - A_n$ of $\lambda I - A(\mathbf{x})$ are stable.

COROLLARY 4.2 *Let $A(\mathbf{x})$ be defined by (4), Assumption 2.1 and fulfil conditions of Theorem 3.1. If the coefficients b_i belong to the class $C^{(\alpha)}$ on \mathbb{R}^d and a_{ij} belong to the class $C^{(1+\alpha)}$ on \mathbb{R}^d then:*

- (i) *The operators $U(n, t)P(n)$ converge strongly in $\dot{C}(\mathbb{R}^d)$ to $U(t)$ uniformly with respect to t in segments of $[0, \infty)$.*
- (ii) *The limit (3) is valid.*

Convergence in general case

Let us defines a molifier $\mathbf{x} \mapsto \vartheta(n, \mathbf{x}) = h^{-d} \vartheta(h^{-1} \mathbf{x})$ in terms of a non-negative function $\vartheta(\cdot)$ of class $C^{(2)}$ on \mathbb{R}^d with the support equal to $B_1(0)$ and $\int \vartheta(\mathbf{x}) d\mathbf{x} = 1$. A smoothing procedure of coefficients a_{ij}, b_i is determined by replacing the coefficients with sequences of coefficients $a_{ij}^{(m)} = \vartheta(m) \circ a_{ij}, b_i^{(m)} = \vartheta(m) \circ b_i$, $m \in \mathbb{N}$. The resulting differential operators $A^{(m)}(\mathbf{x})$ have the closures in $\dot{C}(\mathbb{R}^d)$ and generate Feller semigroups $U^{(m)}(\cdot)$. Now we consider the mappings:

$$\begin{aligned} A(\mathbf{x}) &\longmapsto \{A^{(m)}(\mathbf{x}) : m \in \mathbb{N}\}, \\ A_n &\longmapsto \{A_n^{(m)} : m \in \mathbb{N}\}, \end{aligned} \tag{26}$$

and the associated operators $U^{(m)}(t)$ and $U^{(m)}(n, t)P(n)$ in the Banach space $\dot{C}(\mathbb{R}^d)$, where semigroups $U^{(m)}(\cdot)$ are generated by the closures of $A^{(m)}(\mathbf{x})$ and $U^{(m)}(n, t) = \exp(-A_n^{(m)}t)$. For the double sequence of operators $U^{(m)}(n, t)P(n)$, $m, n \in \mathbb{N}$, we have to prove the following limits:

$$U^{(m)}(n, t)P(n) \xrightarrow{\mathfrak{n}} U^{(m)}(t) \xrightarrow{\mathfrak{m}} U(t) \quad (27)$$

uniformly with respect to t in segments of $[0, \infty)$. By applying the diagonalization argument to the sequence $\{U^{(m)}(n, \cdot)P(m) : (m, n) \in \mathbb{N}^2\}$ and using (27) we get the main result of this article:

THEOREM 4.2 *Let the differential operator $A(\mathbf{x})$ be defined by (4) and Assumption 2.1. There exists a sequence of pairs $(m, n(m)) \in \mathbb{N}^2$ such that*

$$\lim_m \|U_{n(m)}^{(m)}(t)P_{n(m)} - U(t)\|_\infty = 0,$$

uniformly with respect to t in segments of $[0, \infty)$. The limit (3) is valid for the sequence $U_{n(m)}^{(m)}(\cdot)$.

PROOF: Since the coefficients $b_i^{(m)}$ belong to the class $C^{(\alpha)}$ on \mathbb{R}^d and the coefficients $a_{ij}^{(m)}$ belong to the class $C^{(1+\alpha)}$ on \mathbb{R}^d the first limit in (27) follows from Corollary 4.2.

It remains to prove the second limit in (27), i.e. the convergence of the semigroups $U^{(m)}(\cdot)$ in $\dot{C}(\mathbb{R}^d)$ to $U(\cdot)$, uniformly with respect to t in segments of $[0, \infty)$. The strong convergence of $U^{(m)}(\cdot)$ to $U(t)$ in $L_2(\mathbb{R}^d)$ can be easily obtained (for instance, by using Theorem 6.1, Chapter 1. in [EK]). The strong convergence of $U^{(m)}(t)$ to $U(t)$ in $\dot{C}(\mathbb{R}^d)$, uniformly with respect to t in segments of $[0, \infty)$, would follow from such convergence on a dense subspace in $\dot{C}(\mathbb{R}^d)$. We choose the subspace $C_0^{(3)}(\mathbb{R}^d)$ which is dense in both, the space $L_2(\mathbb{R}^d)$ and the space $\dot{C}(\mathbb{R}^d)$. The operators $U(t)$ and $U^{(m)}(t)$ map $C_0^{(3)}(\mathbb{R}^d)$ into the space $\dot{C}^{(2+\alpha)}(\mathbb{R}^d)$ uniformly with respect to t in segments of $[0, \infty)$, due to $\|U^{(m)}(t)\|^{(2+\alpha)} \leq (\sigma t)$. Thus we come to the following conclusion. The operators $U(t) - U^{(m)}(t)$ are continuous mappings from $C_0^{(3)}(\mathbb{R}^d)$ into $\dot{C}^{(2+\alpha)}(\mathbb{R}^d)$ and they converge strongly to zero in $L_2(\mathbb{R}^d)$, uniformly with respect to t in segments of $[0, \infty)$.

Now we have to apply the following auxiliary result to $u_n(t) = U(t) - U^{(m)}(t)v$, $v \in C_0^{(3)}(\mathbb{R}^d)$. Let $\mathfrak{U} = \{u_n : n \in \mathbb{N}\}$ be a sequence of continuous functions on $[0, 1] \times \mathbb{R}^d$ such that:

- a) $u_n(t) \in L_2(\mathbb{R}^d)$ and $\sup\{\lim_n \|u_n\|_2 : t \in [0, 1]\} = 0$.
- b) The functions u_n are uniformly Hölder continuous in the following sense. For each $t \in [0, 1]$ and $\mathbf{x} \in \mathbb{R}^d$ there holds $u_n(t)|_{B_1(\mathbf{x})} \in C^{(\alpha)}(B_1(\mathbf{x}))$, $\|u_n(t)\|_\infty^{(\alpha)} \leq c_\alpha$, where α and c_α do not depend on t and \mathbf{x} .

Then $\lim_n u_n(t) = 0$ in $\dot{C}(\mathbb{R}^d)$, uniformly with respect to $t \in [0, 1]$.

A proof of this auxiliary result is simple. If the assertion of lemma were not valid there would exist a positive number δ and a sequence of pairs $(t_k, \mathbf{x}_k) \in [0, 1] \times \mathbb{R}^d$, $\lim_k |\mathbf{x}_k| = \infty$, such that $u_k(t_k, \mathbf{x}_k) \geq \delta$. Due to b) the following must be also valid: $|u_k(t_k)| \geq \delta/2$ on the ball $B_r(\mathbf{x}_k)$ where $r = (\delta/2c_\alpha)^{1/\alpha}$. A consequence of these inequalities would be $\|u_k(t_k)\|_2 \geq 2^{-1}\delta\sqrt{|B_r(\mathbf{0})|}$, contradicting a). **QED**

From a computational point of view it is natural to assume that the coefficients a_{ij}, b_i have extensions on $\overline{(D_l)}$ belonging to the class $C^{(1)}$. By using the decomposition of unit we can realize the first mapping of (26) in such way that $A_m^{(m)}$ and A_m coincide, while the matrices $A_n^{(m)}$, $n > m$, differ from the matrices A_n of Section 3. Entries $(A_n^{(m)})_{kl}$ and $(A_n)_{kl}$, $n > m$, eventually differ for $\mathbf{y} = hl \in (G_n \setminus G_m) \cap S_n(\Gamma)$ where $S_n(\Gamma)$ is defined in Section 3.

5 SIMULATION OF SAMPLE PATHS

Here we demonstrate the efficiency of simulation of sample paths of a generalized diffusion by using MJPs. We intend to estimate the expectation and variance of the first exit time from a Lipschitz domain. The differential operator $A(\mathbf{x}) = -\sum_{ij=1}^2 \partial_i a_{ij}(\mathbf{x}) \partial_j$ on \mathbb{R}^2 is defined by its diffusion tensor, being a piecewise constant tensor-valued function of the form,

$$a(\mathbf{x}) = \begin{bmatrix} \sigma^2 & \alpha(\mathbf{x}) \\ \alpha(\mathbf{x}) & 1 \end{bmatrix}, \quad \alpha(\mathbf{x}) = \rho \mathbb{1}_{D_0}(\mathbf{x}), \quad \rho^2 < \sigma^2,$$

where σ^2 is a positive number, ρ is a real number and $D_0 = (1/4, 3/4)^2$. Let $X(\cdot)$ be diffusion determined by $A(\mathbf{x})$ starting from $\mathbf{x}_0 = (1/2, 1/2)$. For a bounded Lipschitz domain D the expectation and variance of first exit time from D are given by expressions:

$$\mathbf{E}[\theta] = \|u\|_1, \quad \mathbf{Var}[\theta] = 2\|v\|_1 - \mathbf{E}[\theta]^2, \quad (28)$$

where u, v are solutions of the following boundary value problems,

$$\begin{aligned} A(\mathbf{x})^\dagger u(\mathbf{x}) &= \delta(\mathbf{x} - \mathbf{x}_0), \quad \mathbf{x} \in D, & A(\mathbf{x})^\dagger v(\mathbf{x}) &= u(\mathbf{x}), \quad \mathbf{x} \in D, \\ u|_{\partial D} &= 0, & v|_{\partial D} &= 0, \end{aligned} \quad (29)$$

and $\delta(\mathbf{x})$ is Dirac δ -function at $\mathbf{0}$. We intend to compute $\mathbf{E}[\theta], \mathbf{Var}[\theta]$ by simulations and by using deterministic methods formulated in terms of Expressions (28), (29). In order to simulate sample paths of $X(\cdot)$ we shall approximate the diffusion by a MJP $X_n(\cdot)$ and simulate sample paths of $X_n(\cdot)$. In this example we choose $\sigma^2 = 0.1, \rho = 0.02, D = (0, 1)^2$ and two cases of discretizations, $h = 1/200$ and $h = 1/400$.

The generator of process $X_n(\cdot)$ in G_n is denoted by $A_n(\text{gen})$. Since the entries a_{12} are non-trivial on D_0 we have to use construction of Section 3 in order to get $A_n(\text{gen}) = -A_n$. The parameters $r_1 = 3, r_2 = 1$ of construction are illustrated in Figure 3. These values of parameters ensure A_n to have the structure of a matrix of positive type.

Two boundary value problems of (29) have unique solutions $u, v \in L_1(D)$ as proved in [BO, LR1]. An efficient numerical method is constructed and the convergence in $L_1(D)$ is proved [LR2]. This numerical method is based on the construction of A_n which is described in Section 3. Efficiency of constructed methods is demonstrated by examples in which solutions in closed forms are compared with numerical solutions.

Results of computation are expressed in terms of relative errors:

$$\varepsilon_{exp} = \frac{\langle \theta \rangle_{det} - \langle \theta \rangle_{sim}}{\langle \theta \rangle_{det}}, \quad \varepsilon_{var} = \frac{\ll \theta \gg_{det} - \ll \theta \gg_{sim}}{\ll \theta \gg_{det}},$$

where $\langle \theta \rangle_{det}, \langle \theta \rangle_{sim}$ are the estimates of $\mathbf{E}[\theta]$ obtained by using deterministic methods (28) and Monte Carlo simulations, respectively. Analogously, $\ll \theta \gg_{det}, \ll \theta \gg_{sim}$ are the corresponding estimates of $\mathbf{Var}[\theta]$. Some results of computations are given in the table bellow. The last column contains the ratios, $r = t_{det}/t_{sim}$, of computational times t_{det} and t_{sim} of deterministic and Monte Carlo method, respectively. Sample paths are simulated 20000 times.

| $h = 1/n$ | ε_{exp} | ε_{var} | r |
|-----------|---------------------|---------------------|------|
| $n = 200$ | -0.039 | 0.018 | 15.7 |
| $n = 400$ | -0.044 | 0.007 | 20.3 |

Comparison of results obtained by deterministic and Monte Carlo methods

As expected, the first two statistical moments of first exit time can be estimated by Monte Carlo simulations dozen times faster than by using the deterministic method formulated by (28) and (29).

References

- [BO] BOTTARO G. & OPPEZZI P., *Elliptic equations in divergence form, with right hand side a measure, on unbounded domains*, Ann. Mat. Pura Appl., **163**, 223–246, 1993
- [EK] ETHIER S. N. & KURTZ T. G., *Markov Processes, characteristics and Convergence*, Wiley, New York, 1986.
- [Et] ÉTORÉ P., *On random walk simulation of one-dimensional diffusion process with discontinuous coefficients*, Electronic Journal of Probability, **11**, 2006, 249-275.
- [GM] GARRONI M.G. & MENALDI J.L., *Second order elliptic integro-differential problems*, Chapman, 2002.
- [Kr] KREIN S.G. *Linear Differential Equations in Banach Space*, Translations of Mathematical Monographs, 1992.
- [LM] LEJAY A. & MARTINEZ M., *A scheme for simulating one-dimensional diffusion process with discontinuous coefficients*, The Annales of Applied Probability **16:1**, 107-139, 2006.
- [LR1] LIMÍĆ N. & ROGINA M., *Explicit stable methods for second order parabolic systems*, Math. Commun. Vol 5, 97–115, 2000
- [LR2] ROGINA M. & LIMÍĆ N., *Monotone numerical schemes for a Dirichlet problem for elliptic operators in divergence form*, to be published in Mathematical Methods in Applied Sciences, 2009
- [Ma] MAZ'YA V.G., *Sobolev spaces*, Springer, N.Y., 1985
- [MW] MOTZKYN T.S. & WASOV W., *On the approximation of linear elliptic differential equations by difference equations with positive coefficients*, J. Mathematical Physics **31**, 253-259 1953.
- [Se] STEIN F.M., *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press, New Jersey, 1970
- [So] SOLONNIKOV V.A., *A priori estimates for second order equations of parabolic type*, Trudy Math. Inst. Steklov, **70**, 133-212, 1964; *On a boundary value problem for linear general parabolic system of differential equations*, Trudy Math. Inst. Steklov, **83**, 1965.
- [Sr] STROOCK D.W., *Diffusion semigroups corresponding to uniformly elliptic divergence form operators*, Séminaire de Probabilité XXII, Springer Verlag LNMS # 1321, 316-348, 1988.
- [SZ] STROOCK D.W. & W. ZHENG, *Markov chain approximations to symmetric diffusions*, Annal, Section B, **33**, 619-649 1997.
- [Tb] TANABE H., *Equations of Evolution*, Pitman, 1975.