

Weibel Instabilities in a Completely Degenerate Electron Fermi Gas

Levan N.Tsintsadze

Department of Plasma Physics, E.Andronikashvili Institute of Physics, Tbilisi, Georgia

(Dated: November 16, 2018)

We study the Weibel instability in a degenerate Fermi plasma. We disclose new type of quantum Weibel instabilities. In particular, a novel oscillatory Weibel instability is found and its growth rate is obtained. We reveal a transverse zero-sound in a quantum degenerate electron gas, which has no counterpart in the classical consideration.

In 1959, Weibel showed [1] that a self-excited transverse electromagnetic (EM) waves can exist in an unmagnetized plasma, when the electron distribution in the momentum space is sufficiently anisotropic. The classical Weibel instability [2]-[5] is a purely growing transverse EM mode excited in an unmagnetized plasma by surplus the perpendicular (relative to the propagation of EM waves) mean energy to the parallel mean energy, i.e., $\varepsilon_{\perp} > \varepsilon_{\parallel}$. In particular case it reads the temperature anisotropy ($T_{\perp} > T_{\parallel}$).

The Weibel instability, which leads to the generation of large-scale magnetic fields, has wide range of applicability to astrophysical and space plasmas, as well as laboratory plasmas. The relativistic theory was developed for it in Refs.[6]-[12] in order to interpret the astrophysical data on gamma-ray bursts, relativistic jets from active galactic nuclei, and clusters of galaxies. The Wigner's type of quantum kinetic equation for a plasma was derived by Klimontovich and Silin [13], and in the linear approximation obtained dispersion relations for longitudinal plasma waves and transverse EM waves in an isotropic plasma.

In this Brief Report, we consider the Weibel instability in fully degenerate electron gas, supposing that a plasma consists of degenerate electrons and nondegenerate ions (ions are assumed to be immobile), such as those in neutron stars, supernovae, black holes and white dwarfs. The purpose of the present work is to investigate detailed properties of the Weibel instability in fully degenerate electron plasma for a particular choice of anisotropic distribution function.

As is well known in an electron gas at the temperature of absolute zero (completely degenerate Fermi gas) the electrons occupy all states with momentum from zero to $p = p_F = (3\pi^2)^{1/3}\hbar n^{1/3}$, which is the radius of the Fermi sphere in momentum space (\hbar is the Planck constant divided by 2π). Hence, the possible electron trajectories in momentum space are closed. In general the topology of Fermi surface can be more complex with close electron trajectory [14, 15]. As it was indicated in Ref.[16] exists also the open Fermi surface, which means that in some direction of momentum space the electron trajectory is not closed. In our study we assume that the Fermi surface is closed, but it has a cylindrical type of shape.

Let us consider one dimensional propagation of EM waves along the z-axis. Relative to this propagation, we introduce the perpendicular, p_{\perp} , and the parallel, p_{\parallel} , momentums.

The distribution function adopted in the present calculation assumes that some electrons occupy states with the perpendicular momentum from zero to the limit $p_{\perp} = p_{F\perp} = \sqrt{p_{Fx}^2 + p_{Fy}^2}$, and some of the electrons occupy states with parallel momentum in the range $p_{\parallel} = -p_{F\parallel}$ and $p_{\parallel} = p_{F\parallel}$. So, the anisotropic distribution function of completely degenerate electrons can be expressed as

$$f_0 = H(p_{F\perp} - p_{\perp}) \cdot H(p_{F\parallel}^2 - p_{\parallel}^2), \quad (1)$$

where $H(x)$ is the Heaviside step function, that is $H(x) = 1$ for $x > 0$, and $H(x) = 0$ for $x < 0$.

The number of quantum states in an element of volume dV with values of electron momentum from p_{\perp} to $p_{\perp} + dp_{\perp}$ and p_{\parallel} to $p_{\parallel} + dp_{\parallel}$ equals $2\pi p_{\perp} dp_{\perp} dp_{\parallel} dV / (2\pi\hbar)^3$. This quantum number we must multiply by 2 due to the spin of electrons. Thus, the N electrons, contained in the volume V , fill all the lowest energy states with momentum ranging from zero to $p_{F\perp}$ and $-p_{F\parallel}$ to $p_{F\parallel}$. Therefor, the density of electrons can be presented as

$$n = \frac{N}{V} = 2 \int_0^{p_{F\perp}} \frac{2\pi p_{\perp} dp_{\perp}}{(2\pi\hbar)^2} \int_{-p_{F\parallel}}^{p_{F\parallel}} \frac{dp_{\parallel}}{2\pi\hbar}. \quad (2)$$

Based on the expression (2) we can introduce the number of electrons that fill the energy states of perpendicular and parallel directions

$$n_{\perp} = \frac{p_{F\perp}^2}{2\pi^2\hbar^2} \text{ cm}^{-2} \quad \text{and} \quad n_{\parallel} = \frac{p_{F\parallel}}{\pi\hbar} \text{ cm}^{-1}. \quad (3)$$

With these expressions of n_{\perp} and n_{\parallel} at hand, we write the kinetic energy as

$$\varepsilon_{F\perp} = \frac{p_{F\perp}^2}{2m_0} = \frac{\pi^2\hbar^2}{m_0} n_{\perp} \quad \text{and} \quad \varepsilon_{F\parallel} = \frac{p_{F\parallel}^2}{2m_0} = \frac{\pi^2\hbar^2}{2m_0} n_{\parallel}^2,$$

where m_0 is the electron rest mass.

Correspondingly for the degeneracy temperatures we can write

$$T_{F\perp} = \frac{\varepsilon_{F\perp}}{K_B} \quad \text{and} \quad T_{F\parallel} = \frac{\varepsilon_{F\parallel}}{K_B}, \quad (4)$$

where K_B is the Boltzmann constant.

In order to investigate the Weibel instability at quantum scales, we employ the expression of dielectric permeability, $\varepsilon^{tr}(\omega, k)$, given in Ref.[17], which reads

$$\varepsilon^{tr}(\omega, k) = 1 - \frac{\omega_p^2}{\omega^2} \left(1 - \frac{1}{\hbar m_0 n_0} \int \frac{p_{\perp}^2}{\omega - \vec{k} \cdot \vec{v}} \left[f_0 \left(\vec{p} + \frac{\hbar \vec{k}}{2} \right) - f_0 \left(\vec{p} - \frac{\hbar \vec{k}}{2} \right) \right] \frac{d\vec{p}_{\perp} dp_{\parallel}}{(2\pi\hbar)^3} \right), \quad (5)$$

where n_0 is the total density of electrons and equals $n_0 = n_{\perp} \cdot n_{\parallel}$, k is the wave-vector, ω is the wave frequency, ω_p is the electron plasma frequency.

Using the dispersion relation $k^2 c^2 / \omega^2 = \varepsilon^{tr}$ (c is the speed of light in vacuum) for the transverse EM waves, and taking into account the fact that we consider the propagation of EM waves along the z-axis, we replace $\vec{p} + \hbar \vec{k} / 2$ by the \vec{p} in the first integral of Eq.(5), and in the second integral $\vec{p} - \hbar \vec{k} / 2 \rightarrow \vec{p}$, and use the newly introduced distribution function (1) in the expression (5) to obtain

$$\frac{k^2 c^2}{\omega^2} = 1 - \frac{\omega_p^2}{\omega^2} \left(1 + \frac{n_{\perp}}{k n_{\parallel}} \left[\ln \frac{\omega + \eta - k v_{F\parallel}}{\omega + \eta + k v_{F\parallel}} - \ln \frac{\omega - \eta - k v_{F\parallel}}{\omega - \eta + k v_{F\parallel}} \right] \right), \quad (6)$$

where $\eta = \hbar k^2 / 2 m_0$. Note that the ratio $\beta = n_{\perp} / k n_{\parallel}$ can in general be arbitrary.

We now examine the dispersion relation (6) for different range of frequencies. First, for the weak quantum limit, when $|\omega - k v_{F\parallel}| \gg \eta$, from Eq.(6) follows

$$\varepsilon^{tr} = 1 - \frac{\omega_p^2}{\omega^2} \left(1 + \frac{2 n_{\perp}}{m_0 n_{\parallel}} \frac{\hbar k \cdot k v_{F\parallel}}{\omega^2 - k^2 v_{F\parallel}^2} \right). \quad (7)$$

In the case $\omega^2 \ll \omega_p^2$, expressing $v_{F\parallel}$ through $n_{F\parallel}$, we get the following dispersion equation

$$\omega^2 = k^2 \frac{\pi^2 \hbar^2}{4 m_0^2} \left[n_{F\parallel}^2 - \frac{4}{\pi} \frac{\omega_p^2}{\omega_p^2 + k^2 c^2} n_{F\perp} \right]. \quad (8)$$

We note here that if $\omega_p \sim kc$, then the condition for the generation of low frequency EM field is $n_{F\perp} > n_{F\parallel}^2$, which is similar to the one for the classical Weibel instability, $T_{\perp} > T_{\parallel}$.

Next, we consider the range of frequencies $|\omega - \eta| \gg k v_{F\parallel}$. In this case the dispersion relation (6) reduces to

$$\omega^2 = \eta \left(\eta - \frac{4\pi\omega_p^2}{\omega_p^2 + k^2 c^2} \frac{\hbar n_{\perp}}{m_0} \right). \quad (9)$$

Equation (9) predicts a purely growing quantum Weibel instability for the wavelengths $\lambda \gg \sqrt{2\pi/n_{\perp}}$. Here we assumed that $\omega_p^2 \gg k^2 c^2$.

In the frequency region $|\omega - \eta| < k v_{F\parallel} < \omega + \eta$, use of the definition $\ln x = \ln |x| - i\pi$ for $x < 0$ in Eq.(6) yields

$$\omega^2 = \omega_p^2 + k^2 c^2 + i\omega_p^2 \pi \beta. \quad (10)$$

The solution of which we can write as

$$\omega = \pm \sqrt{\omega_p^2 + k^2 c^2 + \omega_p^2 \pi \beta} \left(\cos \frac{\varphi}{2} + i \sin \frac{\varphi}{2} \right), \quad (11)$$

where $\varphi = \arctg \frac{\omega_p^2 \pi \beta}{\omega_p^2 + k^2 c^2}$.

If we now suppose that $\omega_p^2 \pi \beta \ll \omega_p^2 + k^2 c^2$, then $\cos \frac{\varphi}{2} \approx 1$ and $\sin \frac{\varphi}{2} \sim \frac{\pi}{2} \beta \frac{\omega_p^2}{\omega_p^2 + k^2 c^2}$, and from Eq.(11) follows that $Re\omega \gg Im\omega$, which indicates that there is the possibility of an oscillatory Weibel instability in the degenerate

electron gas, that has no counterpart in classical plasma. The growth rate of this novel oscillatory Weibel instability is

$$Im\omega = \frac{\pi\omega_p^2\beta}{2\sqrt{\omega_p^2 + k^2c^2}}. \quad (12)$$

In the opposite case, i.e., $\omega_p^2\pi\beta \gg \omega_p^2 + k^2c^2$, one can immediately see that $Re\omega \simeq Im\omega$ because $\varphi \sim \pi/2$, $\cos \frac{\varphi}{2} \simeq \sin \frac{\varphi}{2}$, and we have

$$\omega \approx \omega_p\sqrt{\pi\beta}(1 + i). \quad (13)$$

Finally, we shall demonstrate the existence of transverse zero-sound in a quantum degenerate electron gas. As is well known in a Fermi liquid [15], as well as in a plasma the longitudinal waves, known as the zero-sound exist in the range of short wavelengths, i.e., $k^2r_D^2 = k^2\frac{3v_F^2}{\omega_p^2} \gg 1$ (r_D is the Debye length). Note that this wave is the continuation of electron Langmuir waves. It should be emphasized that the transverse zero-sound cannot arise in the classical consideration. However, as we show here undamped transverse EM zero-sound appears in quantum electron gas in the frequency region $\omega \simeq kv_{F\parallel} - \eta$. To this end, we replace the frequency by $\omega = kv_{F\parallel} - \eta + \gamma$ (where $|kv_{F\parallel} - \eta| \gg \gamma$) in Eq.(6) to obtain

$$\omega = kv_{F\parallel} - \eta \left(1 + 2 \exp\left\{-\frac{k^2c^2}{\omega_p^2\beta}\right\} \right). \quad (14)$$

Note that for these waves to be true the inequalities $k^2c^2 \gg \omega_p^2\beta$ and $v_{F\parallel} \gg \hbar k/2m_0$ should hold. These inequalities can be satisfied in most practical cases.

In summary, we have investigated the Weibel instability in fully degenerate electron gas, assuming that a plasma consists of degenerate electrons and nondegenerate ions, which are immobile. A purely growing quantum Weibel instabilities, as well as oscillatory Weibel instability are found. We have discovered the transverse zero-sound in a quantum degenerate electron gas. We specifically note here that this wave does not exist in classical plasmas. The new quantum Weibel instabilities may be responsible for the generation of non-stationary magnetic fields in compact astrophysical objects as well as in the forthcoming intense laser-solid density plasma experiments.

-
- [1] E.S.Weibel, Phys. Rev. Lett. **2**, 83 (1959).
 - [2] B.D.Fried, Phys. Fluids **2**, 337 (1959).
 - [3] R.L.Morse and C.W.Nielson, Phys. Fluids **14**, 830 (1971).
 - [4] R.C.Davidson, D.Hammer, I.Haber, and C.E.Wagner, Phys. Fluids **15**, 317 (1972).
 - [5] M.Lazar, R.Schlickeiser and P.K.Shukla, Phys. Plasmas **13**, 102107 (2006).
 - [6] P.H.Yoon and R.C.Davidson, Phys. Rev. A **35**, 2718 (1987).
 - [7] P.H.Yoon, Phys. Fluids B **1**, 1336 (1989).
 - [8] T.Y.B.Yang, J.Arons and A.B.Langdon, Phys. Plasmas **1**, 3059 (1994).
 - [9] P.H.Yoon, Phys. Plasmas **14**, 024504 (2007).
 - [10] M.V.Medvedev and A.Loeb, Astrophys. J. **526**, 697 (1999).
 - [11] R.A. Fonseca, L.O.Silva, J.W.Tonge, W.M.Mori and J.M.Dawson, Phys. Plasmas **10**, 1979 (2003).
 - [12] U.Schaefer-Rolffs and I.Lerche, Phys. Plasmas **13**, 062303 (2006).
 - [13] Yu.L.Klimontovich and V.P.Silin, Zh. Eksp. Teor. Fiz. **23** 151 (1952).
 - [14] R.L.Liboff, *Kinetic Theory*, 3rd ed. (Springer, 2003).
 - [15] E.M.Lifshitz and L.P.Pitaevskii, *Statistical Physics*, part 2 (Butterworth - Heinemann, Oxford, 1997), 13.
 - [16] I.M.Lifshitz and V.G.Peschanskii, Sov. Phys. JETP **35**, 1251 (1958).
 - [17] M.V.Kuzelev and A.A.Rukhadze, Phys. Usp. **42** 603 (1999).