

# Asymptotic stability of certain sets of associated prime ideals of local cohomology modules

Nguyen Tu Cuong<sup>a</sup>, Nguyen Van Hoang<sup>b</sup> and Pham Huu Khanh<sup>c</sup>

Institute of Mathematics  
18 Hoang Quoc Viet Road, 10307 Hanoi, Vietnam

**Abstract**<sup>1</sup>. Let  $(R, \mathfrak{m})$  be a Noetherian local ring,  $I, J$  two ideals of  $R$  and  $M$  a finitely generated  $R$ -module. It is first shown that for  $k \geq -1$  the integer  $r_k = \text{depth}_k(I, J^n M/J^{n+1}M)$ , it is the length of a maximal  $(J^n M/J^{n+1}M)$ -sequence in dimension  $> k$  in  $I$  defined by M. Brodmann and L. T. Nhan [4], becomes for large  $n$  independent of  $n$ . Then we prove in this paper that the sets  $\bigcup_{j \leq r_k} \text{Ass}_R(H_I^j(J^n M/J^{n+1}M))$  with  $k = -1$  or  $k = 0$ , and  $\bigcup_{j \leq r_1} \text{Ass}_R(H_I^j(J^n M/J^{n+1}M)) \cup \{\mathfrak{m}\}$  are stable for large  $n$ . We also obtain similar results for modules  $M/J^n M$ .

## 1 Introduction

Let  $(R, \mathfrak{m})$  be a Noetherian local ring,  $I, J$  two ideals of  $R$  and  $M$  a finitely generated  $R$ -module. This paper is concerned the asymptotic stability of sets of associated prime ideals  $\text{Ass}_R(H_I^j(J^n M/J^{n+1}M))$  and  $\text{Ass}_R(H_I^j(M/J^n M))$ . There are two starting points for our investigation as follows.

Firstly, In 1979, M. Brodmann [1] had proved that the sets  $\text{Ass}_R(J^n M/J^{n+1}M)$  and  $\text{Ass}_R(M/J^n M)$  are stable for large  $n$ . Based on this result he showed in [2, Theorem 2 and Proposition 12] that the integers  $\text{depth}(I, J^n M/J^{n+1}M)$  and  $\text{depth}(I, M/J^n M)$  take constant values for large  $n$ . Recently, M. Brodmann and L.T. Nhan introduced the notion of  $M$ -sequence in dimension  $> k$  in [4]: Let  $k$  be an integer  $k \geq -1$ . A sequence  $x_1, \dots, x_r$  of elements of  $\mathfrak{m}$  is called an  $M$ -sequence in dimension  $> k$  if  $x_i \notin \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Ass}_R(M/(x_1, \dots, x_{i-1})M)$  with  $\dim(R/\mathfrak{p}) > k$  and all  $i = 1, \dots, r$ . They also showed that every maximal  $M$ -sequence in dimension  $> k$  in  $I$  have the same length,

---

<sup>1</sup>*Key words and phrases:* associated prime, local cohomology, depth, filter depth, generalized depth  
*2000 Subject Classification:* 13D45, 13E05  
This work is supported in part by the National Basis Research Programme in Natural Science of Vietnam.

<sup>a</sup> E-mail: ntcuong@math.ac.vn

<sup>b</sup> E-mail: nguyenvanhoang1976@yahoo.com

<sup>c</sup> E-mail: huukhanh75@yahoo.com

and the common length is the least integer  $i$  such that there exists  $\mathfrak{p} \in \text{Supp}(H_I^i(M))$  with  $\dim(R/\mathfrak{p}) > k$ . For convenience, we denote this common length by  $\text{depth}_k(I, M)$ . It should be mentioned that  $\text{depth}_{-1}(I, M)$  is just  $\text{depth}(I, M)$  the length of a maximal  $M$ -sequence in  $I$ ,  $\text{depth}_0(I, M)$  is the filter depth  $f\text{-depth}(I, M)$  with respect to the ideal  $I$  defined by R. Lü and Z. Tang [14], and  $\text{depth}_1(I, M)$  is the generalized depth  $g\text{depth}(I, M)$  with respect to the ideal  $I$  defined by L.T. Nhan [20]. The first result of this paper is a generalization of Brodmann's Theorem.

**Theorem 1.1.** *For all  $k \geq -1$ ,  $\text{depth}_k(I, J^n M/J^{n+1}M)$  and  $\text{depth}_k(I, M/J^n M)$  take constant values  $r_k$  and  $s_k$ , respectively, for large  $n$ .*

Secondly, in 1990, C. Huneke [9, Problem 4] asked whether the set of associated primes of  $H_I^i(M)$  is finite for all generated modules  $M$  and all ideals  $I$ . Affirmative answers were given by G. Lyubeznik [15] and Huneke-R.Y. Sharp [10] for equicharacteristic regular local rings. Although, A. Singh [21] and M. Katzman [11] provided examples of finitely generated modules having some local cohomology with infinite associated prime ideals, the problem is still true in many situations ( see [3] [12], [13], [16], [20]...). Especially, it was proved in [13] and [20] that  $\text{Ass}_R(H_I^j(M))$  is finite for all  $j \leq \text{depth}_1(I, M)$ . From this and Theorem 1.1 we obtain for all  $j \leq r_1 = \text{depth}_1(I, J^n M/J^{n+1}M)$  and  $i \leq s_1 = \text{depth}_1(I, M/J^n M)$  that the sets  $\text{Ass}_R(H_I^j(J^n M/J^{n+1}M))$  and  $\text{Ass}_R(H_I^i(M/J^n M))$  are finite for large  $n$ . So it is natural to ask the following question:

*Are the sets  $\text{Ass}_R(H_I^j(J^n M/J^{n+1}M))$  and  $\text{Ass}_R(H_I^i(M/J^n M))$  for  $j \leq r_1$  and  $i \leq s_1$  stable for large  $n$ ?*

The main result of this paper is the following theorem which shows that a weakening of the above questions may have a positive answer.

**Theorem 1.2.** *Let  $k \geq -1$  be an integer, and  $r_k, s_k$  as in Theorem 1.1. Then the following statements are true.*

- (i)  $\text{Ass}_R(H_I^{r-1}(J^n M/J^{n+1}M))$  and  $\text{Ass}_R(H_I^{s-1}(M/J^n M))$  are stable for large  $n$ .
- (ii)  $\bigcup_{j \leq r_0} \text{Ass}_R(H_I^j(J^n M/J^{n+1}M))$  and  $\bigcup_{i \leq s_0} \text{Ass}_R(H_I^i(M/J^n M))$  are stable for large  $n$ .
- (iii)  $\bigcup_{t \leq j} \text{Ass}_R(H_I^t(J^n M/J^{n+1}M)) \cup \{\mathfrak{m}\}$  and  $\bigcup_{t \leq i} \text{Ass}_R(H_I^t(M/J^n M)) \cup \{\mathfrak{m}\}$  for all  $j \leq r_1$  and  $i \leq s_1$  are stable for large  $n$ .

Notice here that since  $r_{-1} \leq r_0 \leq r_1$  as well as  $s_{-1} \leq s_0 \leq s_1$  and the modules  $H_I^j(J^n M/J^{n+1}M)$ ,  $H_I^i(M/J^n M)$  are either vanished or Artinian for all  $j < r_0$  and  $i < s_0$  by [14], the question is trivial at these  $j, i$ . So the conclusion (i) of Theorem 1.2 is the first non-trivial answer to the above question at  $j = r_{-1}$  and  $i = s_{-1}$  when  $r_{-1} = r_0$  and  $s_{-1} = s_0$ . Moreover, the conclusion (ii) (the conclusion (iii), respectively) of Theorem 1.2 says that except the maximal ideal  $\mathfrak{m}$  (a finite set, respectively) the sets in the question above are stable for all  $j \leq r_0$  and  $i \leq s_0$  ( $j \leq r_1$  and  $i \leq s_1$ , respectively).

This paper is divided into 5 sections. In Section 2 we study some properties concerning generalizations of the depth of a finite generated module over a local ring. Theorem

1.1 will be also showed in this section. Sections 3, 4 and 5 are devoted to prove the statements (i), (ii) and (iii), respectively, of Theorem 1.2.

## 2 Asymptotic stability of generalizations of depth

Throughout this paper,  $(R, \mathfrak{m})$  is a Noetherian local ring with the maximal ideal  $\mathfrak{m}$ ,  $I, J$  two ideals of  $R$  and  $M$  is a finitely generated  $R$ -module.

**Definition 2.1.** ([4, Definition 2.1]) Let  $k \geq -1$  be an integer. A sequence  $x_1, \dots, x_r$  of elements of  $\mathfrak{m}$  is called an  $M$ -sequence in dimension  $> k$  if  $x_i \notin \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Ass}_R(M/(x_1, \dots, x_{i-1})M)$  with  $\dim(R/\mathfrak{p}) > k$  and all  $i = 1, \dots, r$ .

It is clear that  $x_1, \dots, x_r$  is an  $M$ -sequence in dimension  $> -1$  if and only if it is a regular sequence of  $M$ ; and  $x_1, \dots, x_r$  is an  $M$ -sequence in dimension  $> 0$  if and only if it is a filter regular sequence of  $M$  introduced by P. Schenzel, N. V. Trung and the first author in [8]. Moreover,  $x_1, \dots, x_r$  is an  $M$ -sequence in dimension  $> 1$  if and only if it is a generalized regular sequence of  $M$  defined by L. T. Nhan in [20].

**Remark 2.2.** (i) Let  $k$  be a non negative integer. Assume that  $\dim(M/IM) > k$ . Then any  $M$ -sequence in dimension  $> k$  in  $I$  is of finite length, and all maximal  $M$ -sequence in dimension  $> k$  in  $I$  have the same length which is equal to the least integer  $i$  such that there exists  $\mathfrak{p} \in \text{Supp}(H_I^i(M))$  with  $\dim(R/\mathfrak{p}) > k$  ([4, Lemma 2.4]). We denote in this case by  $\text{depth}_k(I, M)$  the length of a maximal  $M$ -sequence in dimension  $> k$  in  $I$ . Moreover, if  $x_1, \dots, x_r$  is a maximal  $M$ -sequence in dimension  $> k$  in  $I$ , then  $x_1, \dots, x_r$  is a part of system of parameters of  $M$ , and thus  $\text{depth}_k(I, M) \leq \dim(M) - \dim(M/IM)$ . Note that  $\text{depth}_{-1}(I, M)$  is the usual depth  $\text{depth}(I, M)$  of  $M$  in  $I$ ,  $\text{depth}_0(I, M)$  is the filter depth  $f\text{-depth}(I, M)$  of  $M$  in  $I$  denoted by Lü and Tang in [14] and  $\text{depth}_1(I, M)$  is just the generalized depth  $\text{gdepth}(I, M)$  of  $M$  in  $I$  defined by Nhan [20].

(ii) If  $\dim(M/IM) \leq k$ , then we can choose an  $M$ -sequence in dimension  $> k$  in  $I$  of length  $r$  for every positive integer  $r$ , and we set in this case that  $\text{depth}_k(I, M) = \infty$ .

Let  $S$  be a subset of  $\text{Spec}(R)$  and  $i \geq 0$  an integer, we set

$$S_{\geq i} = \{\mathfrak{p} \in S \mid \dim(R/\mathfrak{p}) \geq i\} \text{ and } S_{> i} = \{\mathfrak{p} \in S \mid \dim(R/\mathfrak{p}) > i\}.$$

**Lemma 2.3.** Let  $k \geq -1$  be an integer. Then

$$\begin{aligned} \text{depth}_k(I, M) &= \inf\{j \mid \dim(\text{Ext}_R^j(R/I, M)) > k\} \\ &= \inf\{\text{depth}_{k-i}(I_{\mathfrak{p}}, M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Supp}(M/IM)_{\geq i}\} \end{aligned}$$

for all  $0 \leq i \leq k + 1$ , where we use the convenience that  $\inf(\emptyset) = \infty$ .

*Proof.* Since  $\text{depth}_k(I, M) = \infty$  if and only if  $\dim(M/IM) \leq k$ , the statement is clear for the case  $\text{depth}_k(I, M) = \infty$ . Assume that  $r = \text{depth}_k(I, M)$  is a non negative integer. By [4, Lemma 2.4], we get

$$r = \inf\{i \mid \exists \mathfrak{p} \in \text{Supp}(H_I^i(M)), \dim(R/\mathfrak{p}) > k\}.$$

Moreover, by [7, Lemma 2.8], we have

$$\bigcup_{j \leq l} \text{Supp}(H_I^j(M)) = \bigcup_{j \leq l} \text{Supp}(\text{Ext}_R^j(R/I, M))$$

for all  $l \geq 0$ . It follows that  $r = \inf\{j \mid \dim(\text{Ext}_R^j(R/I, M)) > k\}$ . To prove the second equality, let  $x_1, \dots, x_r \in I$  be an  $M$ -sequence in dimension  $> k$ , and  $i \in \{0, \dots, k+1\}$ . For each  $\mathfrak{p} \in \text{Supp}(M/IM)_{\geq i}$ ,  $x_1/1, \dots, x_r/1$  is an  $M_{\mathfrak{p}}$ -sequence in dimension  $> k-i$  in  $I_{\mathfrak{p}}$ ; and thus  $r \leq \text{depth}_{k-i}(I_{\mathfrak{p}}, M_{\mathfrak{p}})$ . On the other hand, by the first equality, there exists  $\mathfrak{q} \in \text{Supp}(\text{Ext}_R^r(R/I, M))$  with  $\dim(R/\mathfrak{q}) > k$ . Hence, we can choose a prime ideal  $\mathfrak{p}'$  such that  $\mathfrak{p}' \supseteq \mathfrak{q}$ ,  $\dim(R/\mathfrak{p}') = i$  and  $\dim(R_{\mathfrak{p}'}/\mathfrak{q}R_{\mathfrak{p}'}) > k-i$ . Therefore  $\mathfrak{q}R_{\mathfrak{p}'} \in \text{Supp}_{R_{\mathfrak{p}'}}(\text{Ext}_{R_{\mathfrak{p}'}}^r(R_{\mathfrak{p}'}/I_{\mathfrak{p}'}, M_{\mathfrak{p}'})_{>k-i})$ . It follows that  $r \geq \text{depth}_{k-i}(I_{\mathfrak{p}'}, M_{\mathfrak{p}'})$ , and so  $r = \inf\{\text{depth}_{k-i}(I_{\mathfrak{p}}, M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Supp}(M/IM)_{\geq i}\}$  as required.  $\square$

Let  $\mathcal{R} = \bigoplus_{n \geq 0} R_n$  be finitely generated a standard graded algebra over  $R_0 = R$ , and  $\mathcal{M} = \bigoplus_{n \geq 0} M_n$  is a finitely generated graded  $\mathcal{R}$ -module. For the convenience, we use  $N_n$  to denote from now on either the  $R$ -module  $M_n$  or  $M/J^n M$ . Then Brodmann's Theorem on the asymptotic stability of associated prime ideals [1] (see also [17, Theorem 3.1]) can be stated as follows.

**Lemma 2.4.** *The set  $\text{Ass}_R(N_n)$  is stable for large  $n$ .*

**Lemma 2.5.** *Let  $k \geq -1$  and  $r \geq 1$  be integers. If  $\dim(\text{Ext}_R^i(R/I, N_n)) \leq k$  for infinitely many  $n$  and all  $i < r$ . Then there always exists a sequence of  $r$  elements  $x_1, \dots, x_r \in I$  which is an  $N_n$ -sequence in dimension  $> k$  for all large  $n$ .*

*Proof.* Assume that  $T$  is an infinite subset of the set of integers such that for all  $n \in T$  and  $i < r$  we have  $\dim(\text{Ext}_R^i(R/I, N_n)) \leq k$ . We proceed by induction on  $r$ . If  $r = 1$ , then  $\dim(\text{Hom}(R/I, N_n)) \leq k$  for all  $n \in T$ . It implies  $I \not\subseteq \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Ass}_R(N_n)_{>k}$  and all  $n \in T$ . Since  $\text{Ass}_R(N_n)$  is stable for large  $n$  by Lemma 2.4, there exists an integer  $a \in T$  such that  $I \not\subseteq \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Ass}_R(N_n)_{>k}$  and all  $n \geq a$ . Hence, we can choose an element  $x_1 \in I$  which is an  $N_n$ -sequence in dimension  $> k$  for all  $n \geq a$ . Assume that  $r > 1$ . Note that  $\dim(0 :_{N_n} x_1) \leq k$  and  $\dim(\text{Ext}_R^i(R/I, N_n)) \leq k$  for all  $n \in T$  with  $n \geq a$  and all  $i < r$ . From this, by using the following exact sequences

$$\text{Ext}_R^i(R/I, N_n) \rightarrow \text{Ext}_R^i(R/I, N_n/(0 :_{N_n} x_1)) \rightarrow \text{Ext}_R^{i+1}(R/I, (0 :_{N_n} x_1))$$

and

$$\text{Ext}_R^{i-1}(R/I, N_n) \rightarrow \text{Ext}_R^{i-1}(R/I, N_n/x_1 N_n) \rightarrow \text{Ext}_R^i(R/I, N_n/(0 :_{N_n} x_1)),$$

we obtain that  $\dim(\text{Ext}_R^i(R/I, N_n/x_1 N_n)) \leq k$  for all  $n \in T$  with  $n \geq a$  and all  $i < r-1$ . Hence, by inductive hypothesis, there exists an  $N_n/x_1 N_n$ -sequence in dimension  $> k$   $x_2, \dots, x_r \in I$  for all  $n \geq a$  for some  $a \in T$  as required.  $\square$

**Proof of Theorem 1.1.** By Lemma 2.4, there is an integer  $u > 0$  such that  $d = \dim(N_n)$  and  $d' = \dim(N_n/IN_n)$  for all  $n \geq u$ . If  $d' \leq k$ , then  $\text{depth}_k(I, N_n) = \infty$  for

$n \geq u$ , and the conclusion follows.

Assume that  $d' > k$ . Then  $0 \leq \text{depth}_k(I, N_n) \leq d - d'$  for all  $n \geq u$  by Remark 2.2, (i). Thus there exists an infinite subset  $T$  of  $\mathbb{Z}$ , and an integer  $r \in \{0, \dots, d - d'\}$  such that

$$r = \inf\{i \mid \dim(\text{Ext}_R^i(R/I, N_n)) > k\}$$

for all  $n \in T$ . We will show that  $\text{depth}_k(I, N_n) = r$  for all  $n$  large. Indeed, if  $r = 0$ , then  $\dim(\text{Hom}(R/I, N_n)) > k$  for all  $n \in T$ . Since  $\text{Ass}_R(\text{Hom}(R/I, N_n)) = \text{Ass}_R(N_n) \cap V(I)$ , we get by Lemma 2.4 that the set  $\text{Ass}_R(\text{Hom}(R/I, N_n))$  is stable for all  $n \geq a$  for some  $a \in T$ . It follows that  $\text{depth}_k(I, N_n) = 0$  for all  $n \geq a$ . If  $r \geq 1$ , then we get by Lemma 2.5 that  $\text{depth}_k(I, N_n) \geq r$  for all  $n \geq v$  for some  $v > 0$ . Therefore  $\dim(\text{Ext}_R^i(R/I, N_n)) \leq k$  for all  $n \geq v$  and all  $i < r$ . Assume that  $\dim(\text{Ext}_R^r(R/I, N_n)) \leq k$  for all  $n$  in an infinite subset  $S$  of  $\mathbb{Z}$ . Then there exists by Lemma 2.5 an integer  $b \geq v$  such that  $\text{depth}_k(I, N_n) \geq r + 1$ , and so  $\dim(\text{Ext}_R^r(R/I, N_n)) \leq k$  for all  $n \geq b$  by Lemma 2.3. This contradicts the definition of  $r$ . Thus there exists an enough large integer  $c$  such that  $\dim(\text{Ext}_R^r(R/I, N_n)) > k$  and  $\dim(\text{Ext}_R^i(R/I, N_n)) \leq k$  for all  $n \geq c$  and  $i < r$ . Therefore by Lemma 2.3 again we get  $\text{depth}_k(I, N_n) = r$  for all  $n \geq c$ .  $\square$

In view of Theorem 1.1, if  $r = \text{depth}_k(I, N_n)$  for all  $n$  large, we call that  $r$  is the eventual value of  $\text{depth}_k(I, N_n)$ . By Lemma 2.5 and the proof of Theorem 1.1, we get immediately the following consequence.

**Corollary 2.6.** *Let  $k \geq -1$  be an integer and  $r$  the eventual value of  $\text{depth}_k(I, N_n)$ . Then*

$$r = \inf\{i \mid \dim(\text{Ext}_R^i(R/I, N_n)) > k \text{ for infinitely many } n\}.$$

**Lemma 2.7.** *Let  $k \geq -1$  be an integer. Assume that  $r$  and  $s$  are the eventual values of  $\text{depth}_k(I, J^n M/J^{n+1}M)$  and  $\text{depth}_k(I, M/J^n M)$ , respectively. Then we have  $r \geq s$ .*

*Proof.* Let  $n > 0$ . Set  $r(n) = \text{depth}_k(I, J^n M/J^{n+1}M)$  and  $s(n) = \text{depth}_k(I, M/J^n M)$ . By the short exact sequence

$$0 \rightarrow J^n M/J^{n+1}M \rightarrow M/J^{n+1}M \rightarrow M/J^n M \rightarrow 0,$$

we get the long exact sequence

$$\begin{aligned} \cdots \rightarrow \text{Ext}_R^{j-1}(R/I, M/J^n M) &\rightarrow \text{Ext}_R^j(R/I, J^n M/J^{n+1}M) \\ &\rightarrow \text{Ext}_R^j(R/I, M/J^{n+1}M) \rightarrow \cdots \end{aligned}$$

Therefore for any  $j < \min\{s(n) + 1, s(n + 1)\}$  we get by Lemma 2.3 that

$$\dim(\text{Ext}_R^j(R/I, J^n M/J^{n+1}M)) \leq k.$$

It implies that  $r(n) \geq \min\{s(n) + 1, s(n + 1)\}$  by Lemma 2.3. Note that, by the hypothesis of  $r$  and  $s$ , there exists an integer  $a > 0$  such that  $r = r(n)$  and  $s = s(n)$  for all  $n \geq a$ . It follows that  $r \geq s$ .  $\square$

Recall that a sequence  $x_1, \dots, x_r \in I$  is called an  $I$ -filter regular sequence of  $M$  if  $x_i \notin \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Ass}_R(M/(x_1, \dots, x_{i-1})M) \setminus V(I)$  and all  $i = 1, \dots, r$ , where  $V(I)$  is the set of all prime ideals of  $R$  containing  $I$ .

**Lemma 2.8.** *Let  $k \geq -1$  be an integer and  $r$  the eventual value of  $\text{depth}_k(I, N_n)$ . Assume that  $1 \leq r < \infty$ . Then there is a sequence  $x_1, \dots, x_r$  in  $I$  which is at the same time a permutable  $N_n$ -sequence in dimension  $> k$  and a permutable  $I$ -filter regular sequence of  $N_n$  for all large  $n$ .*

*Proof.* Let  $l$  be an integer that such  $1 \leq l \leq r$ . We will show by induction on  $l$  the existence of a sequence of  $l$  elements in  $I$  satisfying the conclusion of the lemma. If  $l = 1$ , then by Lemma 2.4 we may assume that  $\text{Ass}_R(N_n)$  is stable for all  $n \geq t$  for some integer  $t > 0$ . Since  $r \geq 1$ ,  $I \not\subseteq \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Ass}_R(N_n)_{>k}$  and all  $n \geq t$ . From this, by the Prime Avoidance Theorem,  $I \not\subseteq \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Ass}_R(N_n)_{>k} \cup (\text{Ass}_R(N_n) \setminus V(I))$  and all  $n \geq t$ . Therefore, there is an element  $x_1 \in I$  satisfying our lemma.

Assume that  $1 < l \leq r$  and there already exists a sequence  $x_1, \dots, x_{l-1} \in I$  which is at the same time a permutable  $N_n$ -sequence in dimension  $> k$  and a permutable  $I$ -filter regular sequence of  $N_n$  for all  $n \geq t$  for some integer  $t > 0$ . Let  $\Omega$  be the set of all subsets of  $\{1, \dots, l-1\}$ . Hence, since  $\Omega$  is finite, we get by Lemma 2.4 that the set

$$T = \bigcup_{\Lambda \in \Omega} (\text{Ass}_R(N_n/\sum_{j \in \Lambda} x_j N_n)_{>k} \cup (\text{Ass}_R(N_n/\sum_{j \in \Lambda} x_j N_n) \setminus V(I)))$$

is finite and stable for all  $n$  large. Hence there exists an element  $x_l \in I$  such that  $x_l \notin \mathfrak{p}$  for all  $\mathfrak{p} \in T$  by the Prime Avoidance Theorem. Then with the same method that used in the proof of Brodmann-Nhan in [4, Proposition 2.5], we can show that  $x_1, \dots, x_l$  is at the same time a permutable  $N_n$ -sequence in dimension  $> k$  and a permutable  $I$ -filter regular sequence of  $N_n$  for all large  $n$ .  $\square$

Finally, we recall the following fact which will be used in the sequel.

**Lemma 2.9.** (see [12, 1.2], [19, 3.4]) *If  $x_1, \dots, x_r$  is an  $I$ -filter regular sequence of  $M$  then we have*

$$H_I^j(M) = \begin{cases} H_{(x_1, \dots, x_r)}^j(M) & \text{if } j < r \\ H_I^{j-r}(H_{(x_1, \dots, x_r)}^r(M)) & \text{if } j \geq r. \end{cases}$$

### 3 Proof of Theorem 1.2, (i)

In the rest of this paper, instead of  $\text{depth}_{-1}(I, M)$ ,  $\text{depth}_0(I, M)$  and  $\text{depth}_1(I, M)$  we denote, by tradition, by  $\text{depth}(I, M)$ ,  $f\text{-depth}(I, M)$  and  $g\text{depth}(I, M)$ , respectively. We first recall a known fact on local cohomology modules.

**Lemma 3.1.** (cf. [6, Theorem 2.4]) *Let  $r = \text{depth}(I, M)$ . Assume that  $1 \leq r < \infty$  and  $x_1, \dots, x_r$  a regular sequence of  $M$  in  $I$ . Then we have*

$$\text{Ass}_R(H_I^r(M)) = \text{Ass}_R(M/(x_1, \dots, x_r)M) \cap V(I).$$

The statement (i) of Theorem 1.2 is then an immediate consequence of following theorems.

**Theorem 3.2.** *Let  $r$  be the eventual value of  $\text{depth}(I, M_n)$ . Then the set  $\text{Ass}_R(H_I^r(M_n))$  is stable for large  $n$ .*

*Proof.* If  $r = \infty$  then  $\text{Ass}_R(H_I^r(M_n)) = \emptyset$  for all large  $n$ . If  $r = 0$  then  $\text{Ass}_R(H_I^0(M_n)) = \text{Ass}_R(M_n) \cap V(I)$  is stable for large  $n$  by Lemma 2.4. It remains to consider the case  $1 \leq r < \infty$ . Then we get from Theorem 1.1, Lemma 2.4 and Lemma 2.8 that there exists a positive integer  $a$  such that for all  $n \geq a$  the following are true: (i)  $r = \text{depth}(I, M_n)$ ; (ii) there exists a sequence  $x_1, \dots, x_r \in I$  which is a regular sequence of  $M_n$ ; and (iii) the set  $\text{Ass}_R(M_n/(x_1, \dots, x_r)M_n)$  is independent of  $n$ . Therefore, we get by Lemma 3.1 that

$$\text{Ass}_R(H_I^r(M_n)) = \text{Ass}_R(M_n/(x_1, \dots, x_r)M_n) \cap V(I)$$

for all  $n \geq a$ , and so that  $\text{Ass}_R(H_I^r(M_n))$  is stable for large  $n$ .  $\square$

**Theorem 3.3.** *Let  $s$  be the eventual value of  $\text{depth}(I, M/J^n M)$ . Then the set  $\text{Ass}_R(H_I^s(M/J^n M))$  is stable for large  $n$ .*

*Proof.* Let  $r$  be the eventual value of  $\text{depth}(I, J^n M/J^{n+1}M)$ . Then  $r \geq s$  by Lemma 2.7. Moreover, we can choose an integer  $a > 0$  such that  $r = \text{depth}(I, M/J^n M)$  and  $s = \text{depth}(I, J^n M/J^{n+1}M)$  for all  $n \geq a$ . From the short exact sequence

$$0 \rightarrow J^n M/J^{n+1}M \rightarrow M/J^{n+1}M \rightarrow M/J^n M \rightarrow 0,$$

we get the following long exact sequence

$$\begin{aligned} \dots \rightarrow H_I^{j-1}(M/J^n M) \rightarrow H_I^j(J^n M/J^{n+1}M) \rightarrow \\ \rightarrow H_I^j(M/J^{n+1}M) \rightarrow H_I^j(M/J^n M) \rightarrow \dots \end{aligned}$$

for all  $n > 0$ . We consider three cases as follows.

*Case 1:*  $r - 2 \geq s$ . By the long exact sequence, we have the following isomorphisms

$$H_I^s(M/J^{n+1}M) \cong H_I^s(M/J^n M)$$

for all  $n \geq a$ . Thus  $\text{Ass}_R(H_I^s(M/J^n M))$  is stable for large  $n$ .

*Case 2:*  $r - 1 = s$ . By the long exact sequence above, we get the exact sequence

$$0 \rightarrow H_I^s(M/J^{n+1}M) \rightarrow H_I^s(M/J^n M) \rightarrow H_I^{s+1}(J^n M/J^{n+1}M)$$

for all  $n \geq a$ . So  $\text{Ass}_R(H_I^s(M/J^{n+1}M)) \subseteq \text{Ass}_R(H_I^s(M/J^n M))$  for all  $n \geq a$ . It follows that  $\text{Ass}_R(H_I^s(M/J^n M))$  is stable for large  $n$ , since  $\text{Ass}_R(H_I^s(M/J^a M))$  is a finite set by Lemma 3.1.

*Case 3:*  $r = s$ . By the long exact sequence, we have the following exact sequence

$$0 \rightarrow H_I^s(J^n M/J^{n+1}M) \rightarrow H_I^s(M/J^{n+1}M) \rightarrow H_I^s(M/J^n M)$$

for all  $n \geq a$ . There exists by Theorem 3.2 an integer  $b \geq a$  such that

$$\text{Ass}_R(H_I^s(J^n M/J^{n+1} M)) = \text{Ass}_R(H_I^s(J^b M/J^{b+1} M))$$

for all  $n \geq b$ . Set  $X = \text{Ass}_R(H_I^s(J^b M/J^{b+1} M))$ , then we get by the previous exact sequence that

$$X \subseteq \text{Ass}_R(H_I^s(M/J^{n+1} M)) \subseteq \text{Ass}_R(H_I^s(M/J^n M)) \cup X$$

for all  $n \geq b$ . Hence, for any  $n \geq b$ , we have

$$\text{Ass}_R(H_I^s(M/J^{n+2} M)) \subseteq \text{Ass}_R(H_I^s(M/J^{n+1} M)) \cup X = \text{Ass}_R(H_I^s(M/J^{n+1} M)).$$

Since  $\text{Ass}_R(H_I^s(M/J^{b+1} M))$  is finite by Lemma 3.1, it follows that  $\text{Ass}_R(H_I^s(M/J^n M))$  is stable for large  $n$ , this finishes the proof of the theorem.  $\square$

## 4 Proof of Theorem 1.2, (ii)

We first need a characterization of filter depth by the Artinianness of local cohomology modules as follows.

**Lemma 4.1.** ([18, Theorem 3.1])  $\text{f-depth}(I, M) = \inf\{i \mid H_I^i(M) \text{ is not Artinian}\}$ .

Now, we apply Theorems 3.2 and 3.3 to prove the statement (ii) of Theorem 1.2. We will do this in the two following theorems corresponding to  $N_n = M_n$  in the graded case  $\mathcal{M} = \bigoplus_{n \geq 0} M_n$  and  $N_n = M/J^n M$ .

**Theorem 4.2.** *Let  $r_0$  be the eventual value of  $\text{f-depth}(I, M_n)$ . Then the set  $\bigcup_{j \leq r_0} \text{Ass}_R(H_I^j(M_n))$  is stable for large  $n$ .*

*Proof.* By virtue of Lemmas 2.4 and 4.1 the theorem is trivial for the case of either  $r_0 = 0$  or  $r = \infty$ . Therefore it remains to prove the theorem in the case  $1 \leq r_0 < \infty$ . Then by Theorem 1.1 and Lemmas 2.4 and 2.8 there exists an integer  $a$  such that for all  $n \geq a$  the following conditions are true: (i)  $r_0 = \text{f-depth}(I, M_n)$ ; (ii) there is a sequence  $x_1, \dots, x_{r_0} \in I$  which is at the same time a permutable filter regular sequence of  $M_n$  and a permutable  $I$ -filter regular sequence of  $M_n$ ; and (iii) the set  $\text{Ass}_R(M_n/(x_1, \dots, x_{r_0})M_n)$  is independent of  $n$ . We first prove that  $\bigcup_{n \geq a} \text{Ass}_R(H_I^{r_0}(M_n))$  is a finite set. Let  $n \geq a$ . We have  $H_I^{r_0}(M_n) \cong H_I^0(H_{(x_1, \dots, x_{r_0})}^{r_0}(M_n))$  by Lemma 2.9, and

$$H_{(x_1, \dots, x_{r_0})}^{r_0}(M_n) \cong \varinjlim_t (M_n/(x_1^t, \dots, x_{r_0}^t)M_n)$$

by [5, Theorem 5.2.9]. Thus by virtue of [4, Proposition 2.6] we obtain

$$\text{Ass}_R(H_I^{r_0}(M_n)) \subseteq \bigcup_{t > 0} \text{Ass}_R(M_n/(x_1^t, \dots, x_{r_0}^t)M_n) = \text{Ass}_R(M_n/(x_1, \dots, x_{r_0})M_n).$$

Therefore the set  $\bigcup_{n \geq a} \text{Ass}_R(H_I^{r_0}(M_n))$  is finite by the choice of  $a$ . From this we may assume without loss of generality that  $a$  large enough such that the set  $S =$



$\bigcup_{n \geq a} \text{Ass}_R(H_I^{r_0}(M_n))$  only consists of all prime ideals  $\mathfrak{p}$  which belong to  $\text{Ass}_R(H_I^{r_0}(M_n))$  for infinitely many  $n \geq a$ .

Next, we claim that  $\text{Ass}_R(H_I^{r_0}(M_n)) \setminus \{\mathfrak{m}\}$  is stable for large  $n$ . Indeed, for any  $\mathfrak{p} \in S$  there is an infinite set  $T$  of integers such that  $\mathfrak{p} \in \text{Ass}_R(H_I^{r_0}(M_n))$  for all  $n \in T$ . Hence  $H_{\mathfrak{p}}^{r_0}((M_n)_{\mathfrak{p}}) \neq 0$  so that  $\text{depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}}) \leq r_0$ . As  $\mathfrak{p} \in \text{Supp}(M_n/IM_n) \setminus \{\mathfrak{m}\}$ ,  $\text{depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}}) \geq r_0$  by Lemma 2.3. Thus  $\text{depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}}) = r_0$  for all  $n \in T$ . Keep in mind that  $\text{depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}})$  is stable for large  $n$  by Theorem 1.1. Hence  $\text{depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}}) = r_0$  for all large  $n$ . It follows by Theorem 3.2 that  $\text{Ass}_{R_{\mathfrak{p}}}(H_{I_{\mathfrak{p}}}^{r_0}((M_n)_{\mathfrak{p}}))$  is stable for large  $n$ . Therefore  $\text{Ass}_R(H_I^{r_0}(M_n)) \setminus \{\mathfrak{m}\}$  is stable for large  $n$ , and the claim follows.

Let  $r$  be the eventual value of  $\text{depth}(I, M_n)$ . It is clear that  $r \leq r_0$ . If  $r = r_0$ , then  $\bigcup_{j \leq r_0} \text{Ass}_R(H_I^j(M_n)) = \text{Ass}_R(H_I^{r_0}(M_n))$ , and this set is stable for large  $n$  by Theorem 3.2. If  $r < r_0$ , there exists an enough large  $n$  such that  $H_I^r(M_n) \neq 0$ , which is an Artinian module by Lemma 4.1. It follows that

$$\bigcup_{j \leq r_0} \text{Ass}_R(H_I^j(M_n)) = \text{Ass}_R(H_I^{r_0}(M_n)) \cup \{\mathfrak{m}\}$$

is stable for large  $n$ , and the theorem is proved.  $\square$

Before proving the statement (ii) of Theorem 1.2 for  $M/J^n M$ , we need the following lemma.

**Lemma 4.3.** *Let  $A$  be a submodule of an  $R$ -module  $K$ . Then we have*

$$\text{Ass}_R(K/A) \setminus \text{Supp}(A) = \text{Ass}_R(K) \setminus \text{Supp}(A).$$

*Proof.* Let  $\mathfrak{p} \in \text{Ass}_R(K/A) \setminus \text{Supp}(A)$  then  $\mathfrak{p} = (A : \nu)_R$  for some  $\nu \in K$ . It yields that  $\mathfrak{p}\nu \subseteq A$ . So as  $A_{\mathfrak{p}} = 0$  we have  $(\mathfrak{p}\nu)_{\mathfrak{p}} = 0$ . From this, since  $\mathfrak{p}\nu$  is a finitely generated  $R$ -module, it easy to verify that  $\mathfrak{p} \subseteq \text{ann}(r\nu)$  for some  $r \notin \mathfrak{p}$ . For any  $x \in \text{ann}(r\nu)$  then  $rx \in \text{ann}(\nu) \subseteq (A : \nu)_R = \mathfrak{p}$ ; so  $x \in \mathfrak{p}$ , since  $r \notin \mathfrak{p}$ . Thus  $\mathfrak{p} = \text{ann}(r\nu)$ , so that  $\mathfrak{p} \in \text{Ass}_R(K)$ . It follows that  $\text{Ass}_R(K/A) \setminus \text{Supp}(A) \subseteq \text{Ass}_R(K) \setminus \text{Supp}(A)$ . The converse inclusion is trivial.  $\square$

**Theorem 4.4.** *Let  $s_0$  be the eventual value of  $\text{f-depth}(I, M/J^n M)$ . Then the set  $\bigcup_{j \leq s_0} \text{Ass}_R(H_I^j(M/J^n M))$  is stable for large  $n$ .*

*Proof.* By Theorem 1.1, there is an integer  $a$  such that  $r_0 = \text{f-depth}(I, J^n M/J^{n+1} M)$  and  $s_0 = \text{f-depth}(I, M/J^n M)$  for all  $n \geq a$ . By Theorem 3.3 the result is trivial for the case  $s_0 = \infty$ . Assume now that  $s_0 < \infty$ . Using the last part in the proof of Theorem 4.2 we need only to show that the set  $\text{Ass}_R(H_I^{s_0}(M/J^n M)) \setminus \{\mathfrak{m}\}$  is stable for large  $n$ . From the short exact sequence  $0 \rightarrow J^n M/J^{n+1} M \rightarrow M/J^{n+1} M \rightarrow M/J^n M \rightarrow 0$ , we get the following long exact sequence of local cohomology modules

$$\begin{aligned} \cdots \rightarrow H_I^{j-1}(M/J^n M) &\rightarrow H_I^j(J^n M/J^{n+1} M) \\ &\rightarrow H_I^j(M/J^{n+1} M) \rightarrow H_I^j(M/J^n M) \rightarrow \cdots \end{aligned}$$

for all  $n > 0$ . Note that  $r_0 \geq s_0$  by Lemma 2.7. Thus we consider two cases as follows.

*Case 1:*  $r_0 - 1 \geq s_0$ . By the long exact sequence, we get the following exact sequences

$$H_I^{s_0}(J^n M/J^{n+1}M) \rightarrow H_I^{s_0}(M/J^{n+1}M) \rightarrow H_I^{s_0}(M/J^n M),$$

for all  $n \geq a$ , in which  $H_I^{s_0}(J^n M/J^{n+1}M)$  is Artinian by Lemma 4.1. Thus we have

$$\text{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\} \subseteq \text{Ass}_R(H_I^{s_0}(M/J^n M)) \setminus \{\mathfrak{m}\}$$

for all  $n \geq a$ . Since  $\text{Ass}_R(H_I^{s_0}(M/J^n M))$  is finite for  $n = a$  by [6, Theorem 2.5], the set  $\text{Ass}_R(H_I^{s_0}(M/J^n M)) \setminus \{\mathfrak{m}\}$  is stable for large  $n$ .

*Case 2:*  $r_0 = s_0$ . Note by Theorem 4.2 that  $\text{Ass}_R(H_I^{s_0}(J^n M/J^{n+1}M)) \setminus \{\mathfrak{m}\}$  is stable for all  $n \geq b$  for some integer  $b \geq a$ . Set  $X = \text{Ass}_R(H_I^{s_0}(J^b M/J^{b+1}M)) \setminus \{\mathfrak{m}\}$ . Thus we get by the exact sequence

$$H_I^{s_0-1}(M/J^n M) \rightarrow H_I^{s_0}(J^n M/J^{n+1}M) \rightarrow H_I^{s_0}(M/J^{n+1}M) \rightarrow H_I^{s_0}(M/J^n M)$$

for all  $n \geq a$ , in which  $H_I^{s_0-1}(M/J^n M)$  is Artinian by Lemma 4.1, and Lemma 4.3 that

$$X \subseteq \text{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\} \subseteq (\text{Ass}_R(H_I^{s_0}(M/J^n M)) \setminus \{\mathfrak{m}\}) \cup X$$

for all  $n \geq b$ . Therefore

$$\begin{aligned} \text{Ass}_R(H_I^{s_0}(M/J^{n+2}M)) \setminus \{\mathfrak{m}\} &\subseteq (\text{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\}) \cup X \\ &= \text{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\} \end{aligned}$$

for all  $n \geq b$ . Note that  $\text{Ass}_R(H_I^{s_0}(M/J^{n+1}M)) \setminus \{\mathfrak{m}\}$  is a finite set for  $n = b$  by [6, Theorem 2.5]. It follows that  $\text{Ass}_R(H_I^{s_0}(M/J^n M)) \setminus \{\mathfrak{m}\}$  is stable for large  $n$  as required.  $\square$

## 5 Proof of Theorem 1.2, (iii)

To prove the statement (iii) of Theorem 1.2 we first need the following known results.

**Lemma 5.1.** ([7, Corollary 4.4]) *The following equality holds true*

$$\text{gdepth}(I, M) = \inf\{i \mid \text{Supp}(H_I^i(M)) \text{ is not finite}\}.$$

**Lemma 5.2.** ([7, Corollary 4.6]) *The set  $\text{Ass}(H_I^j(N))$  is finite for all  $j \leq \text{gdepth}(I, N)$ .*

The first conclusion of Theorem 1.2, (iii) is a special case of the following result.

**Theorem 5.3.** *Let  $r_1$  be the eventual value of  $\text{gdepth}(I, M_n)$ . Then for each  $l \leq r_1$  the set  $\bigcup_{j \leq l} \text{Ass}_R(H_I^j(M_n)) \cup \{\mathfrak{m}\}$  is stable for large  $n$ .*

*Proof.* By theorem 1.1 and Lemmas 2.4 and 2.8 we can choose a positive integer  $a$  such that  $d = \dim(M_n)$ ,  $d' = \dim(M_n/IM_n)$  and  $r_1 = \text{gdepth}(I, M_n)$  for all  $n \geq a$ . We first claim that there is an integer  $b \geq a$  such that  $S = \bigcup_{n \geq b} \bigcup_{j \leq l} \text{Ass}_R(H_I^j(M_n))$  is finite. In order to do this we consider three cases:  $r_1 = 0$ ,  $r_1 = \infty$  and  $1 \leq r_1 < \infty$ . If  $r_1 = 0$ , then  $S \subseteq \bigcup_{n \geq a} \text{Ass}_R(M_n)$  is finite by Lemma 2.4. If  $r_1 = \infty$ , then  $d' \leq 1$ , so that  $S \subseteq \bigcup_{n \geq a} \text{Supp}_R(M_n/IM_n)$  is finite by Lemma 2.4. Now we consider the case  $1 \leq r_1 < \infty$ . Then, by Lemma 2.8 we can choose an integer  $b \geq a$  such that the following statements are true for all  $n \geq b$ : (i) there is a sequence  $x_1, \dots, x_{r_1} \in I$  which is at the same time a permutable generalized regular sequence of  $M_n$  and a permutable  $I$ -filter regular sequence of  $M_n$ ; and (ii) the sets  $\text{Ass}_R(M_n/(y_1, \dots, y_j)M_n)$  are stable for all  $j \leq r_1$ . From this and [4, Proposition 2.6] we have  $S \subseteq \bigcup_{n \geq b} \bigcup_{j \leq l} \text{Ass}_R(M_n/(y_1, \dots, y_j)M_n)$ . Therefore  $S$  is finite by Lemma 2.4 again, and the claim is proved.

Since  $S$  is finite, we can choose the integer  $b$  enough large such that  $S$  only consists of all prime ideals  $\mathfrak{p}$ , which belong to  $\bigcup_{j \leq l} \text{Ass}_R(H_I^j(M_n))$  for infinitely many  $n \geq b$ . For any  $\mathfrak{p} \in S_{\geq 1}$ , by Theorem 1.1, there is an integer  $n(\mathfrak{p}) \geq b$  such that  $r(\mathfrak{p}) = \text{depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}})$  and  $r_0(\mathfrak{p}) = \text{f-depth}(I_{\mathfrak{p}}, (M_n)_{\mathfrak{p}})$  for all  $n \geq n(\mathfrak{p})$ . Then, by the choice of  $S$  and Lemma 2.3, we obtain that  $r(\mathfrak{p}) \leq l \leq r_1 \leq r_0(\mathfrak{p})$ . If  $l < r_0(\mathfrak{p})$ ,  $\bigcup_{j \leq l} \text{Ass}_{R_{\mathfrak{p}}}(H_{I_{\mathfrak{p}}}^j((M_n)_{\mathfrak{p}})) = \{\mathfrak{p}R_{\mathfrak{p}}\}$  for all  $n \geq n(\mathfrak{p})$  by Lemma 4.1. If  $l = r_0(\mathfrak{p})$ , it follows from Theorem 4.2 that  $\bigcup_{j \leq l} \text{Ass}_{R_{\mathfrak{p}}}(H_{I_{\mathfrak{p}}}^j((M_n)_{\mathfrak{p}}))$  is stable for all  $n \geq m(\mathfrak{p})$  for some integer  $m(\mathfrak{p}) \geq b$ . Therefore  $\bigcup_{j \leq l} \text{Ass}_R(H_I^j(M_n)) \cup \{\mathfrak{m}\}$  is stable for all  $n \geq \max\{n(\mathfrak{p}), m(\mathfrak{p}) \mid \mathfrak{p} \in S_{\geq 1}\}$  as required.  $\square$

Finally, by applying Theorem 5.3 we can prove the statement (iii) of Theorem 1.2 for the  $R$ -modules  $M/J^n M$ .

**Theorem 5.4.** *Let  $s_1$  be the eventual value of  $\text{gdepth}(I, M/J^n M)$ . Then for each  $l \leq s_1$  the set  $\bigcup_{j \leq l} \text{Ass}_R(H_I^j(M/J^n M)) \cup \{\mathfrak{m}\}$  is stable for large  $n$ .*

*Proof.* By Theorem 1.1, there is an integer  $a$  such that  $r_1 = \text{gdepth}(I, J^n M/J^{n+1} M)$  and  $s_1 = \text{gdepth}(I, M/J^n M)$  for all  $n \geq a$ . Note that  $s_1 \leq r_1$  by Lemma 2.7. For any  $0 \leq l \leq s_1$  we set

$$X_l(n) = \bigcup_{j \leq l} \text{Ass}_R(H_I^j(M/J^n M)) \cup \{\mathfrak{m}\} \text{ and } S_l(n) = \bigcup_{j \leq l} \text{Ass}_R(H_I^j(J^n M/J^{n+1} M)) \cup \{\mathfrak{m}\}.$$

We will show by induction on  $l$  that the set  $X_l(n)$  is stable for large  $n$ . It is nothing to prove for the case  $l = 0$ . Let  $l > 0$ . From the short exact sequence

$$0 \rightarrow J^n M/J^{n+1} M \rightarrow M/J^{n+1} M \rightarrow M/J^n M \rightarrow 0$$

we get the following long exact sequence

$$\begin{aligned} \cdots \rightarrow H_I^{j-1}(M/J^n M) \rightarrow H_I^j(J^n M/J^{n+1} M) \\ \xrightarrow{f_j} H_I^j(M/J^{n+1} M) \rightarrow H_I^j(M/J^n M) \rightarrow \cdots \end{aligned}$$

for all  $n > 0$ . From this, for any  $j \leq l$  and  $n \geq a$  we have

$$\text{Ass}_R(H_I^j(M/J^{n+1} M)) \subseteq \text{Ass}_R(H_I^j(M/J^n M)) \cup \text{Ass}_R(\text{Im}(f_j)).$$

Note that

$$\text{Ass}_R(\text{Im}(f_j)) \subseteq \text{Ass}_R(H_I^j(J^n M/J^{n+1}M)) \cup \text{Supp}_R(\text{Ker}(f_j))$$

by Lemma 4.3. Since  $j - 1 < s_1$ ,  $\text{Supp}_R(H_I^{j-1}(M/J^n M))$  is a finite set by Lemma 5.1, so that

$$\text{Supp}_R(\text{Ker}(f_j)) \subseteq \text{Supp}_R(H_I^{j-1}(M/J^n M)) \subseteq \text{Ass}_R(H_I^{j-1}(M/J^n M)) \cup \{\mathfrak{m}\}.$$

Therefore we get

$$X_l(n+1) \subseteq X_l(n) \cup S_l(n).$$

On the other hand, by the long exact sequence above and Lemma 5.1, we have

$$\begin{aligned} \text{Ass}_R(H_I^j(J^n M/J^{n+1}M)) &\subseteq \text{Ass}_R(H_I^j(M/J^{n+1}M)) \cup \text{Supp}_R(H_I^{j-1}(M/J^n M)) \\ &\subseteq \text{Ass}_R(H_I^j(M/J^{n+1}M)) \cup \text{Ass}_R(H_I^{j-1}(M/J^n M)) \cup \{\mathfrak{m}\}. \end{aligned}$$

Hence

$$S_l(n) \subseteq X_l(n+1) \cup X_{l-1}(n)$$

for all  $n \geq a$ . By Theorem 5.3 and the inductive hypothesis, there exists an integer  $b \geq a$  such that  $S_l(n) = S$  and  $X_{l-1}(n) = X$  for all  $n \geq b$ . Therefore, since  $X = X_{l-1}(n+1) \subseteq X_l(n+1)$ ,

$$S \subseteq X_l(n+1) \cup X = X_l(n+1)$$

for all  $n \geq b$ . Finally, we obtain the following inclusions

$$X_l(n+2) \subseteq X_l(n+1) \cup S \subseteq X_l(n+1)$$

for all  $n \geq b$ . Thus  $X(n)$  is stable for large  $n$ , since  $X_l(b+1)$  is finite by Lemma 5.2, and the proof of Theorem 5.4 is complete.  $\square$

## References

- [1] M. Brodmann, *Asymptotic stability of  $\text{Ass}_R(M/I^n M)$* , Proc. Amer. Math. Soc., (1) **74** (1979), 16 - 18.
- [2] M. Brodmann, *The asymptotic nature of the analytic spread*, Math. Proc. Camb. Phil. Soc., **86** (1979), 35 - 39.
- [3] M. Brodmann and A. L. Faghani, *A finiteness result for associated primes of local cohomology modules*, Proc. Amer. Math. Soc., (10) **128** (2000), 2851 - 2853.
- [4] M. Brodmann and L. T. Nhan, *A finiteness result for associated primes of certain Ext-modules*, to appear in Comm. Algebra.
- [5] M. Brodmann and R. Y. Sharp, "Local cohomology: an algebraic introduction with geometric applications", Cambridge University Press, 1998.

- [6] N. T. Cuong and N. V. Hoang, *Some finite properties of generalized local cohomology modules*, East-West J. Math., (2) **7** (2005), 107 - 115.
- [7] N. T. Cuong and N.V. Hoang, *On the vanishing and the finiteness of supports of generalized local cohomology modules*, to appear in Manuscripta Math., 2008.
- [8] N. T. Cuong, P. Schenzel and N. V. Trung, *Verallgemeinerte Cohen-Macaulay Moduln*, Math. Nachr., **85** (1978), 57 - 73.
- [9] C. Huneke, *Problems on local cohomology*, Free resolutions in commutative algebra and algebraic geometry (Sundance, Utah, 1990), Res. Notes Math., **2** (1992), 93 - 108.
- [10] C. Huneke and R. Y. Sharp, *Bass numbers of local cohomology modules*, Trans. Amer. Math. Soc., **339** (1993), 765 - 779.
- [11] M. Katzman, *An example of an infinite set of associated primes of a local cohomology module*, J. Algebra, **252** (2002), 161 - 166.
- [12] K. Khashyarmanesh and Sh. Salarian, *Filter regular sequences and the finiteness of local cohomology modules*, Comm. Algebra, (8) **26** (1998), 2483 - 2490.
- [13] K. Khashyarmanesh and Sh. Salarian, *On the associated primes of local cohomology modules*, Comm. Algebra, **27** (1999), 6191 - 6198.
- [14] R. Lü and Z. Tang, *The  $f$ -depth of an ideal on a module*, Proc. Amer. Math. Soc., (7) **130** (2001), 1905 - 1912.
- [15] G. Luybeznik, *Finiteness properties of local cohomology modules (an application of  $D$ -modules to commutative algebra)*, Invent. Math., **113** (1993), 41 - 55.
- [16] T. Marley, *Associated primes of local cohomology module over rings of small dimension*, Manuscripta Math., (4) **104** (2001), 519 - 525.
- [17] L. Melkersson, *On asymptotic stability for sets of prime ideals connected with the powers of an ideal*, Math. Proc. Camb. Phil. Soc., **107** (1990), 267 - 271.
- [18] L. Melkersson, *Some applications of a criterion for artinianness of module*, J. Pure Appl. Algebra, **101** (1995), 291 - 303.
- [19] U. Nagel and P. Schenzel, *Cohomological annihilators and Castelnuovo-Mumford regularity*, In: Commutative Algebra: Syzygies, Multiplicities, and Birational Algebra (South Hadley, 1992). Contemp. Math., 159. Amer. Math. Soc., (1994), 307 - 328.
- [20] L. T. Nhan, *On generalized regular sequences and the finiteness for associated primes of local cohomology modules*, Comm. Algebra, **33** (2005), 793 - 806.
- [21] A. Singh,  *$p$ -torsion elements in local cohomology modules*, Math. Res. Lett., **7** (2000), 165 - 176.