

Metal-insulator transition in the two-dimensional fully polarized homogeneous electron gas from Hartree-Fock solutions

B. Bernu,¹ F. Delyon,² M. Duneau,² and M. Holzmann^{1,3}

¹*LPTMC, UMR 7600 of CNRS, Université P. et M. Curie, Paris, France*

²*CPHT, UMR 7644 of CNRS, École Polytechnique, Palaiseau, France*

³*LPMMC, UMR 5493 of CNRS, Université J. Fourier, Grenoble, France*

(Dated: July 19, 2022)

We determine the ground state of the two-dimensional, fully polarized electron gas within the Hartree-Fock approximation without imposing any particular symmetries on the solutions. At low electronic densities, the Wigner crystal solution is stable, but for higher densities (r_s less than ~ 3) we obtain a ground state of different symmetry: the charge density forms a triangular lattice with about 11% more sites than electrons. We argue that this conducting state with broken translational symmetry remains the ground state of the high density region in the thermodynamic limit giving rise to a metal to insulator transition.

PACS numbers: 71.10.-w, 71.10.Ca, 71.10.Hf, 71.30.+h, 03.67.Ac

The two-dimensional homogeneous electron gas is one of the fundamental models in condensed matter physics. Despite its simplicity - the system consists of electrons interacting with each other through a $1/r$ potential to which a uniform positive background is added for charge neutrality - the phase diagram at zero temperature is non trivial^{1,2,3}. In general, it is given in terms of the dimensionless parameter $r_s = (\pi n a_B^2)^{-1/2}$, where n is the electronic density and a_B the Bohr radius. At low density (large r_s), the potential energy dominates over the kinetic energy and the system forms a perfect triangular lattice, the Wigner crystal, whereas in the high density region ($r_s \rightarrow 0$) the kinetic energy favors a uniform Fermi liquid phase¹. The experimental observation⁴ of an apparent metal to insulator transition in two-dimensional Si MOSFETs is believed to be incompatible with this simple phase diagram. The scaling theory of localization⁵ suggests that for any disorder, the Fermi liquid solution will become an insulator in the high density region ($r_s \rightarrow 0$), but also the (pinned) Wigner crystal of the low density phase is a natural candidate for an insulator. The physics behind the apparent metallic phase in the intermediate region where roughly the transition between the Wigner crystal and the Fermi liquid is expected, is still puzzling⁶.

In this letter, we present results of a Hartree-Fock calculation of the disorder-free, polarized electron gas, and give evidence for a metal to insulator transition in the zero-temperature phase-diagram at $r_s \approx 2.6$. In contrast to previous Hartree-Fock calculations⁷ we do not impose any symmetry of the solution. Around $r_s \approx 2$, where the uniform Fermi liquid solution is energetically favorable compared to the hexagonal Wigner crystal for the completely polarized system, we find a metallic state with broken translational symmetry which minimizes the energy. The possibility of a metallic, translational symmetry breaking phase, as proposed by our Hartree-Fock solutions, opens a new perspective for the qualitative understanding of the experimental observed metal to insulator transition.

The Hartree-Fock approximation is certainly the sim-

plest approach for quantum many-body calculations. Based on the variational principle, the energy is minimized in the subspace of a single Slater determinant containing single particle orbitals; many-body correlation effects are completely neglected. The free Fermi gas with single particle plane wave states occupied up to the Fermi-wavevector, k_F , is an eigenstate of the Hartree-Fock Hamiltonian with energy $E_{FG} = r_s^{-2} - (8/3\pi)r_s^{-1}$ in Hartree units. In the high density limit ($r_s \rightarrow 0$), it is energetically preferred compared to the low-density Wigner crystal which is well described by gaussian orbitals around the lattice site of a hexagonal crystal. However, already Wigner⁸ argued that the Fermi gas is unstable even in the limit $r_s \rightarrow 0$. Later, Overhauser showed the instability of the unpolarized Fermi gas with respect to spin-density waves, even within the Hartree-Fock approximation⁹. The true ground state of the electron gas within Hartree-Fock is still not completely established¹⁰. Only recently, a Hartree Fock study of the unpolarized three-dimensional electron gas was performed which proposes a more complicated structure of a ground state with spin-density waves¹¹. Here, we consider the two-dimensional electron gas, and, for simplification, we concentrate on the Hartree-Fock ground state of the completely polarized system.

The N-body Hamiltonian, $H = K + V$, contains the kinetic energy K and the $1/r$ -periodic Coulomb potential V where a uniform positive charge background is subtracted. Within the Hartree-Fock approximation, the search of the true ground state of the quantum many-body system is reduced to the simpler problem of finding the lowest energy states in the subset of the Slater determinants. Let $\Phi = \varphi_1 \wedge \dots \wedge \varphi_N$ be the Slater determinant associated with the single particle states $\{\varphi_i\}$

$$\Phi(x_1, \dots, x_N) = \begin{vmatrix} \varphi_1(x_1) & \dots & \varphi_1(x_N) \\ \vdots & \ddots & \vdots \\ \varphi_N(x_1) & \dots & \varphi_N(x_N) \end{vmatrix}.$$

and $E(\varphi_1, \varphi_2, \dots, \varphi_N)$ the corresponding energy expecta-

tion value. The variation of E with respect to a variation of the single particle state $\delta\varphi_i$ is then given by

$$\delta E = \sum_i (h_\Phi \varphi_i, \delta\varphi_i) + \sum_i (\delta\varphi_i, h_\Phi \varphi_i).$$

where h_Φ , the so-called Hartree-Fock Hamiltonian, is a single particle operator depending on the full state Φ (not on the particular choice of the φ_i 's). Extremal states must satisfy the following equation

$$h_\Phi \varphi_i = \sum_j C_{ij} \varphi_j, \quad (1)$$

where C_{ij} are the Lagrange coefficients associated with the normalization constraint $(\varphi_i, \varphi_j) = \delta_{ij}$. Conversely, if $\Phi = \varphi_1 \wedge \dots \wedge \varphi_N$ is not an extremum, we have

$$h_\Phi \varphi_i = \sum_j C_{ij} \varphi_j + \theta_i \quad (2)$$

where the θ_i 's satisfy $(\varphi_i, \theta_j) = 0, \forall i, j$. Within the steepest descent method one chooses first a $N \times N$ unitary transformation $A = (a_{ij})$ such that one obtains

$$(\varphi'_i, \theta'_j) = 0, \quad (\theta'_i, \theta'_j) \propto (\varphi'_i, \varphi'_j) = \delta_{ij}, \quad \forall i, j \quad (3)$$

for the transformed single particle states $\theta'_i = \sum_j a_{ij} \theta_j$, $\varphi'_i = \sum_j a_{ij} \varphi_j$. Then, the energy $E(\varphi_1 + \lambda\theta_1, \dots, \varphi_N + \lambda\theta_N)$ can be expressed as a sum of rational fractions whose numerators and denominators are polynomials of order four, at most. Thus, it is possible to find the best λ and to iterate the process until a stationary state is reached.

In fact, this method has the same drawbacks as the steepest decent method in linear optimization problems; in general, it converges slowly. For linear problems, conjugate gradient methods are preferable^{12,13}. However, since the Hartree-Fock states do not form a linear space, the genuine conjugate gradient method does not apply here. We have therefore adapted a variant of this method to the non-linear case. Let η_i be the previous variation $\delta\varphi_i$, and θ_i is obtained by Eq.2. We then compute $E(\varphi_1 + \lambda\theta_1 + \mu\eta_1, \dots, \varphi_n + \lambda\theta_n + \mu\eta_n)$ for six values of the pairs $\{\lambda, \mu\}$ in order to approximate E by a polynomial of order two in λ and μ . Minimizing the polynomial with respect to λ and μ , we obtain the new changes of the single particle states, $\delta\varphi_i$, and the corresponding energy change. This process is iterated until the relative variation of the energy, $\delta E/E$, is sufficiently small.

Since we expect the electrons to crystallize on a triangular lattice at low densities, we have chosen periodic conditions compatible with this geometry. Thus, the unit cell of our periodized system is given in terms of two vectors $\{\mathbf{L}_1, \mathbf{L}_2\}$ of length L and with an angle of 60 degrees between both; the volume of the unit cell is $\Omega = L^2\sqrt{3}/2$. We have restricted our study to system sizes which are compatible both with the triangular lattice and with a closed shell occupation in k -space. Any triangular crystal with unit-cell vectors $\{\mathbf{e}_1, \mathbf{e}_2\}$ is compatible with the




FIG. 1: Extrapolated energies $E_\infty - E_{FG}$ (Hartree units) versus r_s . Points with error bars: present calculations, full line (red): data of Ref.⁷, full line (green): fit to present results for ($r_s \leq 2.67$), vertical dash line (blue): $r_s = 2.67$.

boundary conditions if it satisfies $\mathbf{L}_1 = l\mathbf{e}_1 + m\mathbf{e}_2$ and $\mathbf{L}_2 = -m\mathbf{e}_1 + (l+m)\mathbf{e}_2$, where (l, m) are two non-negative integers. The number of sites of the lattice is given by $N_c = \det(\mathbf{L}_1, \mathbf{L}_2) / \det(\mathbf{e}_1, \mathbf{e}_2) = l^2 + m^2 + lm$. For a commensurate lattice $N = N_c$.

We compute the wavefunction on a $N_g \times N_g$ grid, the fast Fourier transform is used to switch between real and reciprocal space¹⁴. We have systematically checked the convergence of the solution with respect to the grid size. For the Fermi gas ground state, convergence is reached once all k -vectors in the Fermi surface are represented in the grid ($N_g^2 \sim 3N$). At larger r_s , in the Wigner crystal phase, the wave functions are essentially gaussians¹⁵. The width δ of the gaussians scales as $\delta/L \propto (r_s N)^{-1/2}$. For a correct resolution of the gaussians we need $L/N_g \propto \delta$, so that the number of grid points increases at low densities, $N_g \propto (Nr_s)^{1/2}$. Convergence was reached for $N_g = 32$ (resp. 64, 128) for $N \leq 43$ (resp. $N \leq 200$, $N \leq 500$) up to $r_s = 30$. Whenever the number of grid points was chosen too small, solutions without any particular symmetries have been obtained.

We have further studied the influence of the initial state on the final solution, choosing different types of wavefunction for initialization: a Wigner crystal state, a converged state stored at larger or lower r_s , a state initialized with random numbers, or a ‘‘conducting state’’ as described below (this state has $N_c > N$ maxima in the charge density).

Typically, during the minimization procedure, the energies decrease exponentially; however, the rate of convergence depends strongly on the initial state. The decrease in energy during transitions to a different symmetry is in general much smaller than the convergence within the same symmetry. We have often seen energy plateaus with changes of energy $\delta E/E < 10^{-4}$ just before the occurrence of a transition to a completely different state. For sizes up to $N = 151$, we have con-




FIG. 2: Maxima of $\tau(\mathbf{q})$ (Eq. 6) for $N=151$. The black (resp. white) dots corresponds to $r_s = 3.33$ (resp. 2.67). The radii of the dashed circles are k_F and $2k_F$. The ratio of the radii of the solid circles is $\sqrt{157/151}$.

tinued the minimization until $\delta E/E \lesssim 10^{-12}$, whereas $\delta E/E \lesssim 10^{-5}$ was chosen for convergence for larger system sizes.

We have studied systems with up to 500 electrons at densities $r_s = 4/3 \times \{20, 10, 5, 3, 2.5, 2, 1.75, 1.5, 1.25, 1\}$. In Figure 1, we report the energies of the obtained Hartree-Fock ground state $E_\infty(r_s)$, extrapolated to the thermodynamic limit, as a function of r_s . In the high density region, for $r_s > 3$, we obtain good agreement with Trail *et al.* results⁷, which imposed the triangular Wigner crystal symmetry. For $r_s < 3$, we find ground states of different symmetries than the Wigner crystal which we describe below. These new states have been eventually found down to $r_s = 1$, at smaller r_s , the Fermi gas solution is stable for our finite system sizes with $N \leq 500$.

We now turn to a more detailed analysis of the different phases. It is instructive to follow the evolution of the Slater determinants in k -space as a function of r_s . In order to quantify the differences to the Fermi gas solution, we project each plane wave $\phi_{\mathbf{k}_0}$ of the Fermi gas ($k_0 \leq k_F$), on our Hartree-Fock solution $\{\varphi_i\}$,

$$\phi_{\mathbf{k}_0} = \phi_{\mathbf{k}_0}^{\parallel} + \phi_{\mathbf{k}_0}^{\perp}, \quad (4)$$

$$\phi_{\mathbf{k}_0}^{\parallel} = \sum_i (\varphi_i, \phi_{\mathbf{k}_0}) \varphi_i \quad (5)$$

where $\phi_{\mathbf{k}_0}^{\parallel}$ is the projection of $\phi_{\mathbf{k}_0}$ on the subspace spanned by the φ_i 's. The deviation of the Hartree-Fock solution from the Fermi gas can be measured by $\tau(\mathbf{q})$ as

$$\tau(\mathbf{q}) = \sum_{k_0 \leq k_F, k_0 \neq \mathbf{q}} |\phi_{\mathbf{k}_0}^{\perp}(\mathbf{q} - \mathbf{k}_0)|^2 \quad (6)$$

For the Fermi gas, $\tau(\mathbf{q})$ is 0. For the Wigner crystal, $\tau(\mathbf{q})$ is non zero on the points corresponding to the tri-




FIG. 3: Charge density $\rho/\langle\rho\rangle - 1$, with $\rho(x) = \sum_{i=1}^N |\varphi_i(x)|^2$. Top: the number of maxima is $N_c = 9^2 + 5^2 + 9 \times 5 = 151$. Bottom: the number of maxima is $N_c = 12^2 + 1^2 + 12 \times 1 = 157$. Gray levels corresponds to the same density in both figures.

angular reciprocal lattice: $a\mathbf{q}_1 + b\mathbf{q}_2$, with a, b integers and $\mathbf{q}_1 = l\mathbf{K}_1 - m\mathbf{K}_2$, $\mathbf{q}_2 = m\mathbf{K}_1 + (l+m)\mathbf{K}_2$ ($\mathbf{K}_\alpha \cdot \mathbf{L}_\beta = 2\pi\delta_{\alpha\beta}$)¹⁶. As r_s decreases from 26.7 down to 3, only the first layer of q -vectors corresponding to the triangular lattice remain important. In this region, only the hexagon of the 6 lowest q -vectors of the Wigner crystal remains, and the shape of the charge density in k -space is hexagonal instead of the circular shape at lower densities. This wavefunction has therefore a similar character as the hybrid wave functions studied in Ref.³. However, in contrast to Ref.³, we do not associate a phase transition, rather it is a continuous change from many layers of k -vectors compatible with the Wigner crystal at high r_s to only one layer (Figure 2) as r_s decreases. At the same time, $\tau(\mathbf{q})$ in Eq.6 is dominated by the contribution of the Fermi gas vectors \mathbf{k}_0 close to Fermi surface. For $|\mathbf{k}_0| \approx k_F$, $\phi_{\mathbf{k}_0}^{\perp}(\mathbf{k})$ has essentially only one non zero component at the vector $\mathbf{k} = \mathbf{k}_0 + \mathbf{q}$ closest to the Fermi surface, with \mathbf{q} in the reciprocal lattice of the Wigner crystal.

At $r_s \approx 2.6$, the system lowers its energy by delocalizing the electrons on a denser lattice with more sites than electrons (Figure 2 and 3). This denser lattice is characterized by integer numbers (l', m') different from the Wigner crystal lattice (l, m) . For some sizes, the maxima correspond to various couples of $\{l', m'\}$ leading to different number of lattice sites, $N' = l'^2 + m'^2 + l'm'$ (Fig.4). In this case, our system looks like a periodic crys-




FIG. 4: $N_c - N$ versus the number N of electrons, where N_c is the number of maxima in the charge density. Remark that N_c never decreases with r_s . The line is from Eq.7.

tal with an incomplete band in contrast to the Wigner crystal solutions of fully occupied bands, studied in Ref.⁷. Therefore, our translational symmetry breaking Hartree-Fock solutions below $r_s \approx 2.6$, have a genuine metallic character. However, as r_s approaches zero, the energy gain of this metallic crystal compared to the Fermi gas gets more and more tiny. At the same time, as one can see in Fig.4, N_c is either constant or increases when r_s decreases. For $r_s < 1$ and $N < 500$, the Fermi gas is always the ground state.

Now we show that the translational symmetry breaking, metallic states should exist at small r_s in the thermodynamic limit. Let us replace a plane wave state \mathbf{k} of the free Fermi gas ($\|\mathbf{k}\| \leq k_F$) by a superposition of two plane waves with wavevectors \mathbf{k} and $\mathbf{k}+\mathbf{q}$ ($\|\mathbf{k}+\mathbf{q}\| > k_F$). Choosing \mathbf{q} on the six-fold star of a triangular lattice

we certainly obtain a gain in potential energy. The increase of kinetic energy is minimized if $\|\mathbf{k}\| \sim k_F$ and $\|\mathbf{k}+\mathbf{q}\| \sim k_F$. Then, the number of solutions for \mathbf{k} is optimal if $\|\mathbf{q}\| \sim 2k_F$. This solution corresponds to a triangular lattice of length $L_c = 2\pi/(\sqrt{3}k_F)$ in real space leading to a unit cell of volume $\Omega_c = \sqrt{3}L_c^2/2$. Since our system is contained in the volume $\Omega = \sqrt{3}L^2/2$, we will obtain $N_c = \Omega_c/\Omega$ lattice sites, or

$$N_c = \frac{2\sqrt{3}}{\pi}N \approx 1.1N \quad (7)$$

where we have used $n = N/\Omega = k_F^2/(4\pi)$. From Fig. 4 we see, that, as r_s decreases, the number of sites of the Hartree-Fock ground state indeed approaches the limit of Eq. 7.

A simple model accounting for such a perturbation¹⁷ (actually far in norm from the Fermi gas) gives an energy difference from the Fermi gas varying as $\exp(-a/r_s)$ at the limit $r_s \rightarrow 0$. Our data are reasonably well represented by such a law (see Fig. 1).

In conclusion, we have computed the ground state energy of the two dimensional fully polarized periodic electron gas in the Hartree-Fock approximation. We recover the Wigner crystal at large r_s . For $r_s \lesssim 2.6$, the ground state changes symmetry. Still breaking translational symmetry, the crystal formed contains more lattice sites than particle, and should therefore have metallic properties. For our finite systems, this solution disappears at small r_s in favor of the Fermi gas. Post Hartree-Fock approaches, like the Quantum Monte Carlo method^{1,2,3}, have to be used in order to study the influence of correlation effects and to establish precisely the phase diagram of the electron gas.

Acknowledgment: We thank J. Trail and R. Needs for providing us numerical data of Ref.⁷.

¹ B. Tanatar and D. M. Ceperley, Phys. Rev. **B 39**, 5005 (1989); C. Attaccalite, S. Moroni, P. Gori-Giorgi, and G. B. Bachelet, Phys. Rev. Lett. **88**, 256601 (2002).
² B. Bernu, L. Cândido, and D. M. Ceperley, Phys. Rev. Lett. **86**, 870 (2001).
³ H. Falakshahi and X. Waintal, Phys. Rev. Lett. **94**, 046801 (2005), X. Waintal, Phys. Rev. **B 73**, 075417 (2006).
⁴ S.V. Kravchenko, G.V. Kravchenko, J.E. Furneaux, V.M. Pudalov, and M.D'Iorio, Phys. Rev. **B 50**, 8039 (1994).
⁵ E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
⁶ E. Abrahams, S. V. Kravchenko, and M. P. Sarachik, Rev. Mod. Phys. **73**, 251 (2001).
⁷ J. R. Trail, M. D. Towler, and R. J. Needs, Phys. Rev. **B 68**, 045107 (2003).
⁸ E. P. Wigner, Trans. Faraday Soc. **34**, 678 (1938); Phys. Rev. **46**, 1002 (1934).
⁹ A. W. Overhauser, Phys. Rev. Lett. **4**, 462 (1960); Phys. Rev. **128**, 1437 (1962).

¹⁰ G. F. Giuliani and G. Vignale, *Quantum Theory of the Electron Liquid*, Cambridge University Press, Cambridge (2005).
¹¹ S. Zhang and D. M. Ceperley, arXiv:0712.1194 (2007).
¹² G. H. Golub and C. Van Loan, *Matrix Computations*, Johns Hopkins University Press, 2nd edition (1989).
¹³ M. Duneau, F. Delyon, J. of Comput. Phys. **207**, 375 (2005).
¹⁴ In reciprocal space, the k -vectors of the Wigner crystal fall on the grid.
¹⁵ Indeed, a simple variational calculation with only one parameter, the width of the gaussians, gives the HF-energy with a relative precision better than 0.1%.
¹⁶ In the triangular triangular, the magnitude of the reciprocal lattice vectors are scaled $\|\mathbf{K}_1\| = \|\mathbf{K}_2\| = (2/\sqrt{3})2\pi/L$, compared to a rectangular lattice.
¹⁷ F. Delyon, M. Duneau, B. Bernu, and M. Holzmann, to be published.