A NOTE ON COMBINATORIAL SPLICING FORMULAS FOR HEEGAARD FLOER HOMOLOGY

EAMAN EFTEKHARY

ABSTRACT. We give a precise description of splicing formulas from a previous paper in terms of knot Flor complex associated with a knot in a homology sphere.

1. INTRODUCTION

Heegaard Floer homology has provided topologists with a quite powerful technique in the study of surgery along null-homologous knots inside three-manifolds. After the introduction of surgery formulas for Heegaard Floer homology by Ozsváth and Szabó ([OS3, OS4]), several interesting results have been proved (e.g. [OS4, Hed]). The author has considered another type of surgery formulas which are suggested from the combinatorial approach of Sarkar and Wang to Heegaard Floer homology (see [Ef3], also [SW], also [MOS, MOST]). More generally, there is a formula for Heegaard Floer homology of the three-manifold obtained by splicing two knot complements, provided in [Ef4]. In this paper, we will compare these formulas, and will bring the combinatorial surgery formulas into a form similar to the formulas of Ozsváth and Szabó. More interestingly, this will give a more explicit formula for Heegaard Floer homology of the three-manifold obtained by splicing two knot complements, which is described in terms of knot Floer complexes of the two knots.

Let K be a knot inside the homology sphere Y. We may remove a tubular neighborhood of K and glue it back to obtain the three-manifold $Y_{p/q} = Y_{p/q}(K)$, which is the result of p/q-surgery on K. The core of the solid torus, which is the tubular neighborhood of K, will represent a knot in $Y_{p/q}$ which will be denoted by $K_{p/q}$. We may denote (Y, K) by (Y_{∞}, K_{∞}) , as an extension of the above notation. Let $\mathbb{H}_{\bullet}(K)$ be the Heegaard Floer homology group $HFK(Y_{\bullet}, K_{\bullet})$ for $\bullet \in \mathbb{Q} \cup \{\infty\}$. Note that HFK is defined for knots inside rational homology spheres (see [OS4]), and that $\mathbb{H}_0(K) = \widehat{\mathrm{HFL}}(Y, K)$ is the longitude Floer homology of K from [Ef1]. In all these cases, we choose the coefficient ring to be $\mathbb{Z}/2\mathbb{Z}$ (for simplicity). If we choose a Heegaard diagram for Y - K and let λ_{\bullet} denote a longitude which has framing • $\in \mathbb{Z} \cup \{\infty\}$ (with $\lambda_{\infty} = \mu$ the meridian for K), one may observe that the pairs $(\lambda_{\infty}, \lambda_1)$ and (λ_1, λ_0) have a single intersection point in the Heegaard diagram. Let $(\bullet, \star) \in \{(\infty, 1), (1, 0)\}$ correspond to either of these pairs. There are four quadrants around the intersection point of λ_{\star} and λ_{\bullet} . If we puncture three of these quadrants and consider the corresponding holomorphic triangle map, we obtain an induced map $\mathbb{H}_{\bullet} \to \mathbb{H}_{\star}$. If the punctures are chosen as in figure 1, the result would be two

Key words and phrases. Heegaard Floer homology, rank.

maps $\phi, \overline{\phi} : \mathbb{H}_{\infty}(K) \to \mathbb{H}_{1}(K)$ and two other maps $\psi, \overline{\psi} : \mathbb{H}_{1}(K) \to \mathbb{H}_{0}(K)$ so that the following two sequences are exact:

$$\mathbb{H}_0(K) \xrightarrow{\psi} \mathbb{H}_1(K) \xrightarrow{\overline{\phi}} \mathbb{H}_\infty(K), \text{ and} \\ \mathbb{H}_0(K) \xrightarrow{\overline{\psi}} \mathbb{H}_1(K) \xrightarrow{\phi} \mathbb{H}_\infty(K).$$

The homology of the mapping cones of ϕ (or $\overline{\phi}$) and ψ (or $\overline{\psi}$) are $\mathbb{H}_0(K)$ and $\mathbb{H}_{\infty}(K)$ respectively (see [Ef4]). With the above notation fixed, we have proved the following surgery formula in [Ef4] (The orientation convention of [Ef4] is different from that of [Ef2] and [OS3], so we have changed the direction of all maps to compensate this difference):

Theorem 1.1. Suppose that K_1 and K_2 are two knots inside homology spheres Y_1 and Y_2 respectively. Define the maps $\phi^i, \overline{\phi}^i, \psi^i$ and $\overline{\psi}^i$ for K_i as above. Let $\mathbb{M} = \mathbb{M}(K_1, K_2)$ denote the following cube of maps



where $\mathbb{H}^1_{\star} \otimes \mathbb{H}^2_{\bullet} = \mathbb{H}_{\star}(K_1) \otimes_{\mathbb{Z}/2\mathbb{Z}} \mathbb{H}_{\bullet}(K_2)$ for $\star, \bullet \in \{\infty, 1, 0\}$. The differential $d_{\mathbb{M}}$ of the complex \mathbb{M} is defined to be the sum of all the maps that appear in this cube.



FIGURE 1. For defining chain maps between $C_{\bullet}(K)$ and $C_{\star}(K)$, the punctures around the intersection point of λ_{\bullet} and λ_{\star} should be chosen as illustrated in the above diagrams.

Then the Heegaard Floer homology of the three-manifold Y, obtained by splicing knot complements $Y_1 - K_1$ and $Y_2 - K_2$, is given by

$$\operatorname{HF}(Y; \mathbb{Z}/2\mathbb{Z}) = H_*(\mathbb{M}, d_{\mathbb{M}}),$$

where $H_*(\mathbb{M}, d_{\mathbb{M}})$ denotes the homology of the cube \mathbb{M} .

On the other hand, the groups $\mathbb{H}_{\bullet}(K_i)$ for $\bullet \in \mathbb{Z}$ are described in terms of knot Floer complex for (Y_i, K_i) in [Ef2]. We have shown that the following is true:

Theorem 1.2. Suppose that (Y, K) is a homology sphere and \mathbb{B} is the complex $\widehat{\operatorname{CF}}(Y, \mathbb{Z}/2\mathbb{Z})$, equipped by the filtration induced by K. Let $\mathbb{B}(\geq s)$ denote the quotient-complex generated by those generators whose relative Spin^c class in $\mathbb{Z} = \operatorname{Spin}^c(Y, K)$ is greater than or equal to $s \in \mathbb{Z}$. The homology group $\widehat{\operatorname{HFK}}(Y_n(K), K_n; s; \mathbb{Z}/2\mathbb{Z})$ are isomorphic to the homology of the following chain complex

$$\mathbb{B}(>s-n) \xrightarrow{i} \mathbb{B} \xleftarrow{i} \mathbb{B}(\geq -s),$$

where i is the inclusion map.

The question we would like to address in this paper is the description of the maps $\phi, \overline{\phi}, \psi$ and $\overline{\psi}$ in terms of a presentation of $\mathbb{H}_{\bullet}(K)$ given by the above theorem. This would give surgery and splicing formulas which will look similar to the surgery formulas of Ozsváth and Szabó in [OS3, OS4].

Namely, if we denote the part of $\mathbb{H}_{\bullet}(K)$ in relative $\operatorname{Spin}^{c} \operatorname{class} s \in \mathbb{Z} = \underline{\operatorname{Spin}}^{c}(Y, K)$ by $\mathbb{H}_{\bullet}(K, s)$ we will prove

Theorem 1.3. Under the identification of $\mathbb{H}_{\infty}(K,s)$ with $\mathbb{B}\{s\}$, $\mathbb{H}_{1}(K,s)$ with the homology of the complex

$$C_1(s) = \left(\mathbb{B}\{\geq s\} \xrightarrow{i} \mathbb{B} \xleftarrow{i} \mathbb{B}\{\geq -s\} \right),$$

and $\mathbb{H}_0(K,s)$ with the homology of the complex

$$C_0(s) = \left(\mathbb{B}\{\geq s+1\} \xrightarrow{i} \mathbb{B} \xleftarrow{i} \mathbb{B}\{\geq -s\} \right),$$

the map $\phi : \mathbb{H}_1(K) \to \mathbb{H}_{\infty}(K)$ is induced by the sum of maps ϕ_s which take the quotient complex $\mathbb{B}\{\geq s\}$ of the complex $C_1(s)$ to it quotient $\mathbb{B}\{s\} = \mathbb{H}_{\infty}(K, s)$. The map $\overline{\phi}$ is the sum of maps $\overline{\phi}_s$ which are induced by using the map ϕ_{-s} followed by the isomorphism $\mathbb{B}\{s\} \simeq \mathbb{B}\{-s\}$. Moreover, the map ψ is a sum of inclusion maps ψ_s from $C_0(s-1)$ to $C_1(s)$ and the map $\overline{\psi}$ is induced by the sum of inclusion maps $\overline{\psi}_s$ from $C_0(s)$ to $C_1(s)$.

2. Surgery on null-homologous knots; A short review

In this section we will review the construction of [Ef2], at least to the extent relevant for the purposes of this paper. The reader is referred to [Ef2] for the proofs.

Consider a Heegaard diagram for the pair (Y, K). Suppose that the curve $\mu = \beta_g$ in the Heegaard diagram

$$H = (\Sigma, \boldsymbol{\alpha} = \{\alpha_1, ..., \alpha_g\}, \boldsymbol{\beta} = \{\beta_1, ..., \beta_g\}, p)$$

corresponds to the meridian of K and that the marked point p is placed on β_g . One may assume that the curve β_g cuts α_g once and that this is the only element of α

that has an intersection point with β_g . Suppose that λ represents a longitude for the knot K (i.e. it cuts β_g once and stays disjoint from other elements of β) such that the Heegaard diagram

$$(\Sigma, \boldsymbol{\alpha}, \{\beta_1, ..., \beta_{g-1}, \lambda\})$$

represents the three-manifold $Y_0(K)$ obtained by zero surgery on K. Winding λ around β_{g} - if it is done n times- would produce a Heegaard diagram for the three-manifold $Y_n(K)$. More precisely, if the resulting curve is denoted by λ_n , the Heegaard diagram

$$H_n = (\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}_n = \{\beta_1, ..., \beta_{g-1}, \lambda_n\}, p_n)$$

would give a diagram associated with the knot $(Y_n(K), K_n)$, where $y = p_n$ is a marked point at the intersection of λ_n and β_g .

The curve λ_n intersects the α -curve α_g in *n*-points which appear in the winding region (there may be other intersections outside the winding region). Denote these points of intersection by

$$\dots, x_{-2}, x_{-1}, x_0, x_1, x_2, \dots,$$

where x_1 is the intersection point with the property that three of its four neighboring quadrants belong to the regions that contain either p_n as a corner. Any generator which is supported in the winding region is of the form

$$\{x_i\} \cup \mathbf{y}_0 = \{y_1, \dots, y_{q-1}, x_i\},\$$

and it is in correspondence with the generator

$$\mathbf{y} = \{x\} \cup \mathbf{y}_0 = \{y_1, ..., y_{q-1}, x\}$$

for the complex associated with the knot (Y, K), where x denotes the unique intersection point of α_g and β_g . Denote the former generator by $(\mathbf{y})_i$, keeping track of the intersection point x_i among those in the winding region.

By taking an intersection point \mathbf{x} in relative $\operatorname{Spin}^c \operatorname{class} s - k \in \operatorname{Spin}^c(Y, K) = \mathbb{Z}$ to $(\mathbf{x})_k$ when $k \leq 0$ and $(\mathbf{x})_{k+n}$ if k > 0 we obtain an identification of the complex $\widehat{\operatorname{CFK}}(Y_n(K), K_n; s)$ and the complex

$$\mathbb{U}(s) = \mathbb{B}\{\geq s\} \oplus \mathbb{B}\{>n-s\} = \mathbb{B}\{\geq s\}$$

for large values of n, having fixed $s \in \mathbb{Z}$.

Choose an intersection point between the curves λ_n and λ_{m+n} in the middle of the winding region, denoted by q. We will assume that $m = \ell n$ for an integer ℓ which is chosen to be appropriately large. We continue to assume that α_g is the unique α -curve in the winding region. From the 4 quadrants around the intersection point q, two of them are parts of small triangles Δ_0 and Δ_1 between α, β_n and β_{m+n} . We may assume that the intersection points between α_g and λ_{m+n} in the winding region are

$$\dots, y_{-2}, y_{-1}, y_0, y_1, y_2, \dots,$$

and that the intersection points between λ_n and α_g are

$$\dots, x_{-2}, x_{-1}, x_0, x_1, x_2, \dots$$

as before. We may also assume that the domain Δ_i for i = 0, 1 is the triangle with vertices q, x_i and y_i , and Δ_1 is one of the connected domains in the complement of curves $\Sigma \setminus C$ where

$$C = \boldsymbol{\alpha} \cup \boldsymbol{\beta}_n \cup \boldsymbol{\beta}_{m+n}.$$

Other that Δ_0 and Δ_1 there are two other domains which have q as a corner. One of them is on the right-hand-side of both λ_n and λ_{m+n} , denoted by D_1 , and the other one is on the left-hand-side of both of them, denoted by D_2 . The domains D_1 and D_2 are assumed to be connected regions in the complement of the curves $\Sigma \setminus C$. We may assume that the meridian μ passes through the regions D_1, D_2 and Δ_0 , cutting each of them into two parts: $\Delta_0 = \Delta_0^R \cup \Delta_0^L$, $D_1 = D_1^R \cup D_1^L$ and $D_2 = D_2^R \cup D_2^L$. Here $\Delta_0^R \subset \Delta_0$ is the part on the right-hand-side of μ and Δ_0^L is the part on the left-hand-side. Similarly for the other partitions. Choose the marked points so that u is in D_1^R , v is in D_2^R , w is in D_2^L and z is in D_1^L (see figure 2). We obtain the Heegaard diagram

$$R_{m,n} = (\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\beta}_n, \boldsymbol{\beta}_{m+n}; u, v, w, z),$$

which determines three chain complexes associated with the pairs $(\boldsymbol{\alpha}, \boldsymbol{\beta}), (\boldsymbol{\alpha}, \boldsymbol{\beta}_n)$, and $(\boldsymbol{\alpha}, \boldsymbol{\beta}_{m+n})$ once we use the marked points u, v, w, z. The complexes associated with the last two pairs are simply $\widehat{\operatorname{CFK}}(Y_n(K), K_n)$ and $\widehat{\operatorname{CFK}}(Y_{m+n}(K), K_{m+n})$, obtained by puncturing the surface at these marked points. The complex associated with the first pair is constructed by making punctures at u, v, counting holomorphic disks, and twisting according to the intersection number of the disks with either of the marked points w, z. As we have already seen, $\widehat{\operatorname{CFK}}(K_{m+n}; s)$ may be identified with the complex $\widehat{\mathbb{U}}(s)$ if m is large enough. In fact, the holomorphic triangle map gives a long sequence for each relative Spin^c class $s \in \mathbb{Z}$ as

$$\dots \xrightarrow{f^s} \widehat{\operatorname{CF}}(Y) \xrightarrow{h^s} \widehat{\operatorname{CFK}}(K_n, s) \xrightarrow{g^s} \widehat{\mathbb{E}}(s) \xrightarrow{f^s} \widehat{\operatorname{CF}}(Y) \xrightarrow{h} \dots$$

such that the induced maps in homology form a long exact sequence, if $m = \ell . n$ and ℓ is large enough. Here $\widehat{\mathbb{E}}(s)$ is the complex $\mathbb{B}\{\geq s\} \oplus \mathbb{B}\{>n-s\}$. This implies



FIGURE 2. The Heegaard diagram $R_{m,n}$. The shaded triangles are Δ_0 and Δ_1 . The marked points u, v, w and z are placed in D_1^R, D_2^R, D_2^L and D_1^L respectively.

EAMAN EFTEKHARY

that $\widehat{HFK}(Y_n(K), K_n; s)$ may be computed as the homology of the mapping cone of $f^s : \widehat{\mathbb{E}}(s) \to \mathbb{B}$, which is the sum of two inclusion maps. One particular case, is the case where n = 1 and this later mapping cone is the same as the complex $C_1(s)$ mentioned before.

3. UNDERSTANDING THE MAPS

In this section we prove theorem 1.3.

Proof. In the construction of the previous section, note that $A(s) = \widehat{\operatorname{CFK}}(Y_n(K), K_n; s)$

may be written as the mapping cone of some chain map $\rho: A_1(s) \to A_2(s)$, where $A_1(s)$ is the part of A(s) generated by the generators of the form $(\mathbf{x})_1$. The differential of A(s) induces a differential on the quotient complex $A_1(s)$ and the subcomplex $A_2(s)$ of it. There is a similar decomposition of $B = \widehat{\operatorname{CFK}}(Y_{m+n}(K), K_{m+n})$ as the mapping cone of $\sigma: B_1 \to B_2$, and the image of A_1 under g is in B_1 . Under the identification of B(s) with $\widehat{\mathbb{U}}(s)$, the quotient complex $B_1(s)$ corresponds to the quotient complex $\mathbb{B}\{s\}$ of $\mathbb{B}\{\geq s\} = \widehat{\mathbb{U}}(s)$.

We will use the diagram S_1 , together with the marked points u, w and z to define the map $\phi : C_1(s) = \widehat{\operatorname{CFK}}(Y_1(K), K_1, s) \to \widehat{\operatorname{CFK}}(Y, K, s) = C_{\infty}(s)$. Note that puncturing the surface Σ at these three marked points implies that this chain map will respect the relative Spin^c classes on the two sides. One can easily see from the Heegaard diagram that the map ϕ is defined on the generators by $\phi(\mathbf{y}) = \mathbf{x}$, if $\mathbf{y} = (\mathbf{x})_1 + b_2$ for some generator \mathbf{x} of C_{∞} and some $b_2 \in B_2$. This means that ϕ is trivial on the sub-complex A_2 of $A = C_1$. The image of the generators $(\mathbf{x})_1$ of A_1 under the map g is the generator $(\mathbf{x})_1$, as a generator for $\widehat{\operatorname{CFK}}(Y_{m+1}(K), K_{m+1}, s)$, i.e. g identifies $A_1(s)$ with $B_1(s)$, via the natural isomorphism of both of them with $\mathbb{B}\{s\}$.

On the other hand, since the map h uses punctures at u and v, its image does not contain any generator of the form $(\mathbf{x})_1$ in $\widehat{\operatorname{CFK}}(Y_1(K), K_1, s)$, i.e. the image of h is in the sub-complex A_2 of A. This implies that if an element a in $\operatorname{Ker}(g_*)=\operatorname{Im}(h_*)$ is of the form $a_1 \oplus a_2$ in $A = A_1 \oplus A_2$, and $a_1 = (\mathbf{x})_1$, we should have $(\mathbf{x})_1 = d(\mathbf{y})_1 = (dy)_1$ in A_2 for some generator \mathbf{y} of C_{∞} , i.e. \mathbf{x} is trivial in $\mathbb{H}_{\infty}(K)$.

Finally, note that by the exactness of the long sequence in homology we have

$$\begin{aligned} \mathrm{HFK}(K_1, s) &= \mathrm{Ker}(g_*^s) \oplus \mathrm{Im}(g_*^s) \\ &= \mathrm{Im}(h_*^s) \oplus \mathrm{Ker}(f_*^s) \\ &= \frac{H_*(\mathbb{B})}{\mathrm{Im}(f_*^s)} \oplus \mathrm{Ker}(f_*^s) \oplus H_*(\frac{\widehat{\mathbb{E}}(s)}{\mathrm{Ker}(f^s)} \xrightarrow{\simeq} \mathrm{Im}(f^s)) \\ &= H_*(\frac{\mathbb{B}}{\mathrm{Im}(f^s)} \xrightarrow{d_*} \mathrm{Im}(f^s) \xleftarrow{\simeq} \frac{\widehat{\mathbb{E}}(s)}{\mathrm{Ker}(f^s)} \xrightarrow{d_*} \mathrm{Ker}(f^s)) \\ &= H_*(\widehat{\mathbb{E}}(s) \xrightarrow{f^s} \mathbb{B}). \end{aligned}$$

In these equalities, $H_*(\bullet)$ denotes the homology of the complex \bullet . The above discussion shows that $\operatorname{Ker}(g_*)$ is in the kernel of the induced map ϕ . We thus need to understand the image of $\operatorname{Ker}(f_*^s)$ under the map ϕ in the above description, which are represented in the last line of the above equality by the closed elements e in $\widehat{\mathbb{E}}(s)$ such that $f^s(e)$ is exact in \mathbb{B} . Any such element is the sum of an exact element in $\widehat{\mathbb{E}}(s)$ and an element of the form $g(a_1 \oplus a_2)$ in B_1 , where $a_1 \in A_1$ is closed and $\rho(a_1) = da_2$ for some $a_2 \in A_2$. Under the identification of B_1 with A_1 , this image is of the form $a_1 \oplus e_2 \in B_1 \oplus B_2$. The discussion implies that for a closed element $e \oplus b \in \widehat{\mathbb{E}} \oplus \mathbb{B}$ (i.e. such that e is closed and f(e) = db) such that $e = a_1 \oplus e_2 \in B_1 \oplus B_2$ we have $\phi_*(e \oplus b) = a_1 \in \widehat{\operatorname{HFK}}(Y, K) = \mathbb{H}_{\infty}(K)$. This proves the first assertion in theorem 1.3. The other claims follow similarly.

References

- [Ef1] Eftekhary, E., Longitude Floer homology for knots and the Whitehead double, Alg. and Geom. Topology 5 (2005), also available at math.GT/0407211
- [Ef2] Eftekhary, E., Heegaard Floer homology and knot surgery, preprint, available at math.GT/0603171
- [Ef3] Eftekhary, E, A combinatorial approach to surgery formulas in Heegaard Floer homology, preprint, available at math.GT/0802.3623
- [Ef4] Eftekhary, E., Floer homology and splicing knot complements, *preprint, available* atmath.GT/0802.2874
- [Ef5] Eftekhary, E., Floer homology and existence of incompressible tori in homology spheres, preprint
- [Hed] Hedden, M., On Floer homology and Berge conjecture on knots admitting lens space surgeries, preprint, available at math.GT/0710.0357v2
- [MOS] Manolescu, C., Ozsváth, P., Sarkar, S., A combinatorial description of knot Floer homology, preprint, available at math.GT/0607691
- [MOST] Manolescu, C., Ozsváth, P., Szabó, Z., Thurston, D., On combinatorial link Floer homology, preprint, available at math.GT/0610559
- [Ni] Ni, Y., Link Floer homology detects the Thurston norm, preprint, available at math.GT/0604360
- [OS1] Ozsváth, P., Szabó, Z., Holomorphic disks and knot invariants, Advances in Math. 189 (2004) no.1, also available at math.GT/0209056
- [OS2] Ozsváth, P., Szabó, Z., Holomorphic disks and genus bounds, Geom. Topol. 8 (2004), 311-334
- [OS3] Ozsváth, P., Szabó, Z., Knot Floer homology and integer surgeries, preprint, available at math.GT/0410300
- $[{\rm OS4}] \qquad {\rm Ozsváth, P., Szabó, Z., Knot Floer homology and rational surgeries, , preprint, available at math.GT/0504404 }$
- [SW] Sarkar, S., Wang, J., A combinatorial description of some Heegaard Floer homologies, preprint, available at math.GT/0607777

School of Mathematics, Institute for Studies in Theoretical Physics and Mathematics (IPM), P. O. Box 19395-5746, Tehran, Iran

E-mail address: eaman@ipm.ir