

Dyonic Instantons in 5-dim Yang-Mills Chern-Simons Theories

Seok Kim

*Theoretical Physics Group, Blackett Laboratory,
Imperial College, London SW7 2AZ, U.K.*
s.kim@imperial.ac.uk

Ki-Myeong Lee

*School of Physics
Korea Institute for Advanced Study
Seoul 130-012, KOREA*
klee@kias.re.kr

Sungjay Lee

*School of Physics
Korea Institute for Advanced Study
Seoul 130-012, KOREA*
sjlee@kias.re.kr

ABSTRACT: We consider the BPS dyonic instantons in the 5-dim supersymmetric Yang-Mills Chern-Simons theories. Its field theoretic structures and the moduli space dynamics in term of the ADHM data have been explored in detail. We find that the field theoretic Chern-Simons term leads to an effective magnetic field on instanton moduli space.

Contents

1. Introduction and Conclusion	1
2. 5-dim $\mathcal{N} = 1$ Supersymmetric Gauge Theories	4
2.1 the Lagrangian	4
2.2 Brane Configuration	7
2.3 Vacuum Moduli	9
3. BPS Configurations	10
3.1 BPS Energy Bound and BPS Equations	10
3.2 BPS Configurations	12
3.3 A Single Dyonic $SU(N)$ Instanton	14
4. Low Energy Dynamics	16
4.1 One-dimensional $U(K)$ Gauge Theory	16
4.2 Moduli Space Dynamics	19
4.3 A Single Chern-Simons Instanton Moduli Dynamics	21
A. the ADHM Formalism and the Higgs Solutions	25

1. Introduction and Conclusion

Dyonic instantons in 5-dim Yang-Mills theories in the Coulomb phase have been explored in many different point of view. Electric Coulomb repulsion balances the force to shrink instantons in the Coulomb phase [1, 2, 3]. Dyonic instantons in the Coulomb phase represents a typical example of supertubes realized in a field theory. In the Coulomb phase, it has been argued that dyonic instantons has magnetic monostrings imbedded interior along which distributed are instanton electric charge and linear momentum [4, 5, 6, 7, 8].

Recently dyonic instantons configurations are studied in 5-dim $\mathcal{N} = 1$ supergravity model with nonabelian gauge group [9]. A new feature is the 5-dim Chern-Simons term in the action, which leads nonabelian electric charge on instantons even in the symmetric phase without costing additional energy.

In this work we investigate dyonic instantons in the 5-dim $\mathcal{N} = 1$ supersymmetric Yang-Mills theories with Chern-Simons term. While the kinetic energy is not positive definite, we were able to characterize the 1/2 BPS supersymmetric dyonic instanton configurations which saturate a ‘pseudo’ BPS energy bound. The low energy dynamics of these dyonic instantons is described by the gauged matrix mechanics theory corrected by a simple background charge term, which reduces the number of supersymmetries to four. The moduli space dynamics of these dyonic instantons is a modification of the standard instanton moduli space dynamics by a term which is first order in time derivative and so can be attributed as a background magnetic field on the moduli space. We work out several examples to illuminate the various aspects of the dyonic instantons and also the quantum aspect of nonabelian charge for a single instanton in the symmetric phase.

The 16 supersymmetric 5-dim $U(N)$ Yang-Mills theories can arise from the low energy dynamics of $D4$ branes. While these theories are not renormalizable, there exists an ultraviolet completion in terms of the field theory, the so-called $(2, 0)$ theory compactified on a circle of radius $g^2/4\pi$ with the gauge coupling constant g , and the instantons play the KK mode on the circle. In the Coulomb phase, the instantons collapse. However, there exist regular dyonic instanton configurations in the Coulomb phase, which can be regarded as composite of F1 and D0 branes on D4 branes. In the Coulomb phase, there is a potential induced on the moduli space dynamics without breaking the supersymmetry [1]. There is a natural induced D2 brane dipole charge which can be regarded as the magnetic monopoles inside the dyonic instantons, and so these dyonic instantons in the Coulomb phase is a typical example of a supertube. These dyonic instantons have been studied in detail [4, 5, 6, 7, 8].

The 8 supersymmetric 5-dim $U(N)$ Yang-Mills theories with flavor hypermultiplets have been approached in somewhat differently as their brane counterparts are more complicated. When the flavor number is small, there can be a ultraviolet fixed point with enhanced global symmetries. The Chern-Simons term can be also induced by the massive hypermultiplet in the symmetric phase and by the massive gauginos and hypermultiplets in the Coulomb phase [10]. The kinetic energy of the classical theory with the Chern-Simons term is not positive definite anymore when the scalar field is large. But one expects there is an ultraviolet complete theory for a small flavor case, for which the kinetic energy remains nonnegative [10, 11]. A complete classification of all possible nontrivial five-dimensional fixed points of gauge-theoretic origin as been given in Ref. [12].

Such 8 supersymmetric 5-dim Yang-Mills theories can appear in the brane dynamics as low energy dynamics of the D5- NS5 web dynamics [13, 14]. The additional massive fundamental hypermultiplets can be introduced either by spectator D7 branes or D5 branes.

The integration of the massive hypermultiplets would lead to the Chern-Simons term. The instantons in 5-dim gauge theories with Chern-Simons term have been discussed briefly in relation to skyrmions in Ref. [15, 16].

The 5-dim Chern-Simons term is possible for $SU(N)$ gauge theories with $N \geq 3$, and for those gauge theories with $U(1)$ factor. The Chern-Simons term can be induced by the massive fermions and its coefficient κ should be quantized for the consistency of the path integral. On noncommutative four-space, the natural gauge group is $U(N)$ and the Chern-Simons term is possible for all N . The instantons cannot collapse on noncommutative space, and the low energy dynamics described by the gauged sigma model is modified by the FI term, which makes the instanton moduli space smooth. While our primary focus is on dyonic instantons on commutative space, we will make some discussion for the noncommutative case also. (See ref. [17] for a review on the noncommutative field theory.)

The 1/2 BPS dyonic instanton configuration is characterized by (anti-)selfdual instanton configurations. In the symmetric phase, the Gauss law leads to the electric field profile for a given instanton configuration. For a single instanton configuration in the $SU(N)$ gauge theory we find the scalar field profile explicitly which is singular when the instanton size is less than of order κg^2 . This shows that the ultraviolet cutoff length scale of the Yang-Mills theory with the Chern-Simons term is somewhat larger than the natural scale g^2 for large κ . We do not know the stringy origin of this gap.

The low energy dynamics of the K number of instantons for the 16 supersymmetric theories have 8 supersymmetric gauged sigma model with hypermultiplets in the adjoint and fundamental representation of $U(K)$ gauge symmetry. The noncommutative space can be taken into account by the F-I term in this gauge theory without breaking any supersymmetry. We argue that the 5-dim Chern-Simons term in the field theory leads to an 1-dim Chern-Simons term in the gauged linear sigma model, which is basically an interaction term between the $U(K)$ gauge field and the background charge. The coefficients of two Chern-Simons terms are identical up to an obvious multiplicative factor. This 1-dim Chern-Simons term also reduces the supersymmetry of the gauged sigma model by half. While one needs an ultraviolet completion of the 5-dim theory, the 1-dim gauged sigma model for dyonic instantons is complete by its own. This leads to a tiny gap for the electric charge energy calculated in the field theory and the gauged sigma model in the Coulomb phase.

One can reduce further the gauged sigma model to nonlinear sigma model on the instanton moduli space by solving the vacuum equation of this sigma model and the gauge field equation. The 1-dim Chern-Simons term leads to a background magnetic field on the instanton moduli space. We find this magnetic field on the instanton moduli space explicitly for a single instanton on noncommutative space. For a single instanton on the commutative space, we quantize the moduli space dynamics and show what kind of the ‘nonabelian’ electric charge a single instanton can carry in the symmetric phase.

Our theory allows also 1/2 BPS magnetic monostrings in the Coulomb phase. As in the

maximally supersymmetric case, instantons and W-bosons are attracted to monostings. In addition, one can get the linear momentum be induced on the monostings. Similar to Ref. [6], we can easily obtain the BPS energy bound on magnetic monostings with both instanton and electric charge density. This shows again the supertube origin of our dyonic instantons in the Coulomb phase. There is some investigation of the magnetic monostings in the context of anomaly [18]. Our work should shed some further light on the subject.

There are a couple of directions one can explore starting from our work. The role of the instanton solitons in 5-dim Yang-Mills theories is not fully explored. In the Coulomb phase, there is a 1-loop correction to the Chern-Simons term due to the massive gauginos and hypermultiplets. This would lead to quantum modification of the characteristics of the dyonic instantons.

When one of the spatial direction is compact, instantons or calorons can be decomposed into magnetic monopoles. Our dyonic instantons would be also decomposed into dyons. When the hypermultiplets are present and when the theory is in the Higgs phase, one could have the magnetic flux vortices and magnetic monopoles would be in trapped between flux vortices. [19, 20] We expect similar composite objects in the theory with Chern-Simons term.

The plan of this paper is as follows. In Sec. 2, we review the basic properties of the 5-dim $\mathcal{N} = 1$ supersymmetric Yang-Mills theories with the Chern-Simons term. In Sec. 3, we consider the 1/2 BPS dyonic instanton configurations from the energy bound point of view and solve the Gauss law explicitly for a single instanton in $U(N)$ theory. In Sec.4, we discuss the gauged sigma model with 1-dim Chern-Simons term and also the moduli space dynamics with a background magnetic field. In Appendix, we review some aspects of the ADHM formalism for instantons.

2. 5-dim $\mathcal{N} = 1$ Supersymmetric Gauge Theories

2.1 the Lagrangian

Let us fix the Lagrangian for 5-dim $\mathcal{N} = 1$ supersymmetric gauge theories with the 5-dim Chern-Simons term by using the 4-dim $\mathcal{N} = 2$ supersymmetric Yang-Mills theories. To find the supersymmetric counterpart for the Chern-Simons term, we consider the $U(1)$ theory on a circle. The low energy 4-dim Lagrangian theory would have terms at most quadratic in the spacetime derivatives. As the 4- dim theory has the eight supersymmetries, it is decided by the prepotential $\mathcal{F}(A)$. Especially the A_5 plays the role of scalar field and the 5-dim Chern-Simons term would be come the 4-dim axion-photon coupling term. In addition, five-dimensional gauge invariance highly restricts the prepotential to be at most cubic polynomial; Invariance under $A_5 \rightarrow A_5 + a$ with a arbitrary real constants translates to invariance under the shift $\mathcal{A} \rightarrow \mathcal{A} + ia$ of the chiral field $\mathcal{A} = \phi + iA_5$. Assuming the

prepotential to have the cubic term \mathcal{A}^3 so that

$$\mathcal{F} \sim \mathcal{A}^3 \quad (2.1)$$

the 4-dim bosonic Lagrangian becomes

$$\mathcal{L} \sim -\phi \left((\partial_\mu \phi)^2 + (\partial_\mu A_5)^2 + \frac{1}{2} F_{\mu\nu} F^{\mu\nu} \right) + A_5 \left(\frac{1}{4} F_{\mu\nu} \tilde{F}^{\mu\nu} \right) + \dots, \quad (2.2)$$

which is invariant under the constant shift $A_5 \rightarrow A_5 + a$. When the prepotential has terms higher than cubic order, the Lagrangian is no longer invariant under the constant shift. For instance, let us consider the quartic prepotential $\mathcal{F} = \frac{1}{4!} \mathcal{A}^4$. The Lagrangian now contains

$$\mathcal{L} \sim (\phi^2 - A_5^2) F_{\mu\nu} F^{\mu\nu} + (2\phi A_5) F_{\mu\nu} \tilde{F}^{\mu\nu} + \dots, \quad (2.3)$$

which is not invariant obviously under the shift $A_5 \rightarrow A_5 + a$.

We therefore conclude that the most general five dimensional supersymmetric $SU(N)$ gauge theory can be described by the cubic prepotential

$$\begin{aligned} \mathcal{F}(\mathcal{A}) &= \text{tr}_N \left(\frac{1}{g^2} \mathcal{A}^2 + \frac{\kappa}{3} \mathcal{A}^3 \right) \\ &= \frac{1}{2g^2} \mathcal{A}^a \mathcal{A}^a + \frac{\kappa}{3} d_{abc} \mathcal{A}^a \mathcal{A}^b \mathcal{A}^c. \end{aligned} \quad (2.4)$$

Here we used the convention for the traceless $SU(N)$ generators T_a as

$$\{T_a, T_b\} = \frac{1}{N} \delta_{ab} \mathbf{1}_N + 4d_{abc} T_c. \quad (2.5)$$

so that $\text{tr} T_a T_b = \delta_{ab}/2$. The bosonic part of the action is therefore

$$\begin{aligned} S &= \int d^5x \text{tr} \left(-\frac{1}{2g_{\text{eff}}^2} (F_{\mu\nu} F^{\mu\nu} + 2D_\mu \phi D^\mu \phi) \right) + S_{CS} \\ &= \int d^5x \left(-\frac{1}{4} \left(\frac{1}{g_{\text{eff}}^2} \right)_{ab} (F_{\mu\nu}^a F^{b\mu\nu} + 2D_\mu \phi^a D^\mu \phi^b) \right) + S_{CS} \end{aligned} \quad (2.6)$$

with

$$\frac{1}{g_{\text{eff}}^2} = \frac{1}{g^2} \mathbf{1}_N + \kappa \phi, \quad \left(\frac{1}{g_{\text{eff}}^2} \right)_{ab} = \frac{1}{g^2} \delta_{ab} + 2\kappa d_{abc} \phi^c. \quad (2.7)$$

The 5-dim Chern-Simons term is defined as

$$\begin{aligned} S_{CS} &= \frac{\kappa}{3} \int \text{tr} \left(A \wedge F \wedge F + \frac{i}{2} A \wedge A \wedge A \wedge F - \frac{1}{10} A \wedge A \wedge A \wedge A \wedge A \right) \\ &= \frac{\kappa}{12} \int d^5x \epsilon^{\mu\nu\rho\sigma\chi} \left(d^{abc} A_\mu^a F_{\nu\rho}^b F_{\sigma\chi}^c + \dots \right), \end{aligned} \quad (2.8)$$

which allows only eight real supersymmetries rather than sixteen. The Gauss law constraint for all physical configurations is given by the traceless part of the following equation

$$-D_m \left\{ \frac{1}{g_{\text{eff}}^2}, E_m \right\} - i \left[\phi, \left\{ \frac{1}{g_{\text{eff}}^2}, D_0 \phi \right\} \right] + \frac{\kappa}{2} F_{mn} \tilde{F}_{mn} = 0, \quad (2.9)$$

where $E_m \equiv F_{m0}$ and the dual of the spatial field strength is

$$\tilde{F}_{mn} \equiv \frac{1}{2} \epsilon_{mnpq} F_{pq} . \quad (2.10)$$

Of course the 5-dim Yang-Mills theory is not renormalizable as the coupling constant g^2 has the dimension of the length. We thus consider the field theory with cut-off while it is possible to imagine a consistent quantum completion of the theory such that additional degrees of freedom appears in the ultraviolet region in the energy scale larger than $1/g^2$. With the Chern-Simons term, another length scale $\kappa g^2 > g^2$ could be introduced. Later we will see the classical instanton configuration breaks down in some cases when the instanton size is less than the order of κg^2 .

The Chern-Simons coefficient is quantized for the consistency of the path integral. Note that the Chern-Simons term vanishes for the $SU(2)$ gauge group as there is no nontrivial d_{abc} coefficient. When one makes a large gauge transformation with nontrivial homotopy as $\pi_5(SU(N)) = \mathbf{Z}$ for $N \geq 3$, the Chern-Simons term transforms so that the topological quantity for this homotopy group appears. The quantum consistency requires the quantization,

$$\kappa = \frac{n}{8\pi^2} , \quad (n \in \mathbf{Z}) , \quad (2.11)$$

where we call the integer quantity n as the *Chern-Simons level*.

One important feature of the Chern-Simons theories is that the Chern-Simons couplings can be induced by the radiative quantum corrections by a massive Dirac fermion[21]. Let us consider an amplitude \mathcal{A} involving three external non-abelian gauge bosons of momentum p , q and $-(q+p)$ together with gauge and Lorentz indices (a, μ) , (b, ν) and (c, ρ) , respectively. The parity violating term can be expressed as

$$\begin{aligned} \mathcal{A}_{\mu\nu\rho}^{abc} &= - \int \frac{d^5 k}{(2\pi)^5} \text{tr} \left(\gamma_\mu \frac{1}{k \cdot \gamma + im} \gamma_\nu \frac{1}{(k-q) \cdot \gamma + im} \gamma_\rho \frac{1}{(k+p) \cdot \gamma + im} \right) \text{tr} \left(T^a \{ T^b, T^c \} \right) \\ &= \frac{i}{16\pi^2} \frac{m}{|m|} \epsilon_{\mu\nu\rho\gamma\delta} q^\gamma p^\delta \text{tr} \left(T^a \{ T^b, T^c \} \right) , \end{aligned} \quad (2.12)$$

where the gamma matrices γ_μ here should satisfy the Dirac-Clifford algebra relation $\{\gamma_\mu, \gamma_\nu\} = 2\eta_{\mu\nu}$ with $\gamma_{01234} = i$. One can therefore obtain the induced couplings after integrating out the N_f massive hypermultiplets

$$\begin{aligned} \mathcal{L}_{\text{induced}} &= \text{sign}(m) \frac{N_f}{48\pi^2} \int d^5 x \epsilon^{\mu\nu\rho\gamma\delta} A_\mu^a \partial_\gamma A_\nu^b \partial_\delta A_\rho^c \text{tr} \left(T^a T^b T^c \right) \\ &= -\text{sign}(m) \frac{N_f}{2} \frac{1}{24\pi^2} \int \text{tr} (A \wedge F \wedge F) . \end{aligned} \quad (2.13)$$

Taking account of four and five external gauge bosons diagram in additions, one can find the induced shift of the Chern-Simons level

$$\Delta n = -\frac{N_f}{2} \quad (2.14)$$

If the zeroth order Chern-Simons term vanishes, one requires the even numbers of hypermultiplets for the gauge invariances. It agrees with the result in [12].

When the gauge group is $U(N)$, the gauge coupling constant for $U(1)$ theory and that for the $SU(N)$ theory can be different in general. There could be also Chern-Simons terms for the purely $U(1)$ theory and the mixed term between $SU(N)$ and the $U(1)$ parts with different coefficients. On noncommutative 4-dim space, the gauge group however should be $U(N)$ instead of the $SU(N)$, and so there exist only one gauge coupling constant and also only one Chern-Simons coefficient. It is noteworthy here that, when we consider the noncommutative five-dimensional $U(N)$ gauge theory, all traceless condition for $SU(N)$ gauge symmetry should be dropped out.

2.2 Brane Configuration

In this section, we will present certain brane configurations describing the five-dimensional $\mathcal{N} = 1$ supersymmetric gauge theory [13, 14] with the Chern-Simons couplings. These configurations leads later a useful way to consider the low energy dynamics of dyonic instantons in terms of the dynamics of the D- branes corresponding to instantons.

Let us start with two parallel solitonic (NS) fivebranes, N D5-branes suspended between two NS5-branes. Their configurations are summarized in the table 1. It is noteworthy here that the brane configuration above preserves quarter of maximal supersymmetries, that is, eight real supersymmetries. Since D5-branes are finite in x^6 direction, their macroscopic description is five-dimensional $\mathcal{N} = 1$ $G = SU(N)$ super Yang- Mills theories. The five-dimensional gauge coupling can be now described as

$$\frac{1}{g^2} = \frac{\Delta x^6}{(4\pi g_s)(2\pi l_s)^2} \quad \rightarrow \quad \left[\frac{1}{g^2} \right] = \text{mass} , \quad (2.15)$$

where l_s and g_s are the string length and coupling respectively. When D5 branes meets NS5 branes, they form a (p,q) web of 5-branes, and so the $U(1)$ group factor which is responsible for the motion of the center of the mass of D5 brane segments are frozen as discussed in [22]. It would be interesting find out why this is not case when the noncommutativity on D5 branes are introduced.

For the field theory decoupling limit, we should require the physical states in the field theory not to excite the Kaluza-Klein modes and stringy states. In other words, one must require the characteristic energy scale $\frac{1}{g^2}$ in the field theory to be smaller than

$$\frac{1}{g^2} \ll \frac{1}{l_s} , \frac{1}{\Delta x^6} , \quad (2.16)$$

while taking the $g_s \rightarrow 0$. One can therefore conclude that the decoupling limit is given by

$$\Delta x^6 \ll g_s l_s . \quad (2.17)$$

When NS5-branes are far apart than l_s , one can recover six-dimensional theories with various stringy effects.

	0	1	2	3	4	5	6	7	8	9
D5	○	○	○	○	○		○			
NS5	○	○	○	○	○	○				
D7	○	○	○	○	○			○	○	○

Table 1: Brane configuration for 5-dimensional $\mathcal{N} = 1$ supersymmetric gauge theories

In five-dimensional gauge theories, there is an additional $U(1)_I$ symmetry of interests, whose conserved current takes the form

$$j = \frac{1}{8\pi^2} *_5 \text{tr}(F \wedge F). \quad (2.18)$$

Its charge is the instanton number. In our brane picture, an instanton whose $U(1)_I$ charge is unit can be identified as a D1-brane stretched along the x^6 direction. It is depicted in Fig. One simple way to check this identification follows from the agreement in the mass

$$\frac{8\pi^2}{g^2} = \frac{\Delta x^6}{g_s(2\pi l_s)} = \tau_{D1} \Delta x^6, \quad (2.19)$$

where τ_{D1} denotes the tension of a D1-brane.

In order to introduce massive N_f hypermultiplets, there are two options, of which (1) one is to add the N_f D7-branes whose world volume is parameterized by 01234789 coordinates and (2) another is to add N_f flavor semi-infinite D5-branes. Their configurations are also summarized in the table 1, and depicted in Fig.1. As discussed in [23], the open string modes connecting D5 with D7 branes for the case (1), or D5 with flavor fivebranes for the case (2) give rise to massive hypermultiplets of masses $m = \frac{\delta x^6}{2\pi l_s^2}$. While introducing D7-branes changes the global property of the space-time geometry, which requires a more careful analysis in general, we ignore it here. For our discussion, either picture is fine[13].

The Chern-Simons coupling can be finally introduced after integrating out these massive fermions in the hypermultiplets. From D5 brane point of the view, it is exactly the induced Chern-Simons term by the massive Dirac fermions as discussed in the previous subsection.




Figure 1: Brane configurations for 5d supersymmetric gauge theory

As a final comment, let us consider the Coulomb branch of the theories in the presence of the Chern-Simons couplings. In our brane setting, the Chern-Simons terms are in fact induced by integrating out the massive hypermultiplets. When we try to analyze the Coulomb branch of theory with those brane configurations, we therefore have to suppose that Higgs vevs, or distance between several D5-branes are much smaller than the masses of hypermultiplets or δx^6 .

2.3 Vacuum Moduli

We now in turn discuss the Coulomb branch of the present theories and non-trivial fixed points. In five- dimensional $\mathcal{N} = 1$ supersymmetric gauge theories, the Coulomb branch is parameterized by real scalar fields of vector multiplets. For the gauge group $SU(N)$ of rank $N - 1$, the gauge symmetry is spontaneously broken down to $U(1)^r$ at a generic point on the moduli space. Weyl reflection symmetries implies that the Coulomb branch can therefore be identified with a Weyl chamber $\mathbf{R}^{N-1}/\mathcal{W}$.

We can naively expect from the gauge coupling of negative mass dimension that the low-energy effective theories on the Coulomb branch are trivial. Seiberg however shown in [10] that under one-loop corrections there exist certain strongly-coupled non-trivial fixed point. Although those topics are not relevant in our discussion later, basic ideas will be presented briefly.

For simplicity, let us consider the $SU(2)$ gauge theory with N_f hypermultiplets of masses m_i ($i = 1, 2, \dots, N_f$). For a generic value of ϕ , a scalar field of vector multiplet, the induced Chern-Simons level is proportional to

$$n_{\text{induced}} = 2 \text{sign}(\phi) - \frac{1}{8} \sum_{i=1}^{N_f} \left[\text{sign}(\phi - m_i) + \text{sign}(\phi + m_i) \right], \quad (2.20)$$

where the former contribution is from massive gauginos and the latter from massive hypermultiplets. This implies that the effective gauge coupling (2.7), or metric on the Coulomb branch, takes the form as

$$\frac{1}{g_{\text{eff}}^2} = \frac{1}{g^2} + \frac{1}{16\pi^2} \left(2|\phi| - \frac{1}{8} \sum_{i=1}^{N_f} \left[|\phi - m_i| + |\phi + m_i| \right] \right), \quad (2.21)$$

after finite constant shift of $\frac{1}{g^2}$. We will assume from now on that all mass parameters are turned off.

This metric implies that for $N_f > 8$, there is a singularity at a finite point $\phi \sim \frac{1}{g^2}$ in the moduli space, reflecting the non-renormalizability of the present model. As discussed in [10], non-existence of the singular points on the Coulomb branch for $N_f < 8$ implies the non-trivial fixed point together with the enhancement of global symmetries in the strong coupling limit $g \rightarrow \infty$. Possibility of those non-trivial RG fixed points for various

models with higher rank gauge groups and the classification of them in accordance with the classification of singularities of Calabi-Yau threefolds have been presented extensively in [11, 12].

3. BPS Configurations

3.1 BPS Energy Bound and BPS Equations

We now in turn consider the BPS bound for the present model. The energy functional now takes the form

$$\mathcal{E} = \int d^4\mathbf{x} \operatorname{tr}_N \left(\frac{1}{g_{\text{eff}}^2} \left[E_m^2 + \frac{1}{2} F_{mn}^2 + (D_0\phi)^2 + (D_m\phi)^2 \right] \right). \quad (3.1)$$

In order to make the energy positive definite, we will consider the family of the field configurations such that $g_{\text{eff}}^2 \gg 0$. Thus we consider the region where the value of $1/g^2$ is much larger than the value of $\kappa\phi$ for the sake of argument. We rearrange the above energy functional as

$$\begin{aligned} \mathcal{E} &= \int d^4\mathbf{x} \operatorname{tr}_N \left(\frac{1}{g_{\text{eff}}^2} \left[(E_m - \alpha D_m\phi)^2 + \frac{1}{4} (F_{mn} - \alpha \tilde{F}_{mn})^2 \right. \right. \\ &\quad \left. \left. + (D_0\phi)^2 + \alpha \{E_m, D_m\phi\} + \frac{\alpha}{4} \{F_{mn}, \tilde{F}_{mn}\} \right] \right) \\ &\geq \alpha \int d^4\mathbf{x} \operatorname{tr}_N \left(\frac{1}{g_{\text{eff}}^2} \left[\{E_m, D_m\phi\} + \frac{1}{4} \{F_{mn}, \tilde{F}_{mn}\} \right] \right) \end{aligned}$$

where $\alpha = \pm 1$. One can show from the Gauss law (2.9) that the total energy is bounded by

$$\mathcal{E} \geq \left| \frac{8\pi^2}{g^2} K + Q_E \right|, \quad (3.2)$$

where the topological charge and the electric charge are defined respectively as

$$K = \frac{1}{16\pi^2} \int d^4x \operatorname{tr}_N (F_{mn} \tilde{F}_{mn}), \quad Q_E = \int_{S_\infty^3} dS^m \operatorname{tr}_N \left(\phi \left\{ \frac{1}{g_{\text{eff}}^2}, E_m \right\} \right). \quad (3.3)$$

It should be noted here that the energy due to the electric charge vanishes in the symmetric phase where $\phi_\infty = 0$, even though the instanton can carry nontrivial ‘nonabelian’ electric charge. The bound is saturated by the BPS configurations satisfying the Gauss law (2.9) and the BPS equations

$$F_{mn} - \alpha \tilde{F}_{mn} = 0, \quad E_m - \alpha D_m\phi = 0, \quad D_0\phi = 0. \quad (3.4)$$

In order to see that these BPS configurations preserve four supersymmetries, let us consider the supersymmetry transformation rules of the theory. As discussed in section 2, the theory at hands has eight supersymmetries whose the supersymmetric parameter satisfies the 6-dimensional Weyl condition

$$\Gamma^{012345} \epsilon = \epsilon, \quad (3.5)$$

together with the 10-dimensional Majorana and Weyl condition. (Here we use the 10-dimensional notation for the Gamma matrices, and the gaugino field λ also satisfies all the same conditions as ϵ .) For the BPS configurations, the supersymmetric transformation of the gaugino field

$$\delta\lambda = \frac{1}{2}\Gamma^{MN}F_{MN}\epsilon, \quad (3.6)$$

where $A_5 = \phi$, is required to vanish. Imposing a supersymmetric condition compatible with the Weyl condition (3.5)

$$\Gamma^{1234}\epsilon = \alpha\epsilon, \quad \text{or, equivalently} \quad \Gamma^{05}\epsilon = \alpha\epsilon, \quad (3.7)$$

one can recover the BPS equations (3.4)

$$F_{mn} - \alpha\tilde{F}_{mn} = 0, \quad F_{m0} - \alpha D_m\phi = 0, \quad D_0\phi = 0. \quad (3.8)$$

This implies that the BPS configuration preserves the half of the supersymmetries. Unlike the $\mathcal{N} = 2$ theories, there is a correlation between the sign of the instanton charge and that of electric charge in the $\mathcal{N} = 1$ theory, regardless the presence of the Chern-Simons term.

Once we choose the gauge $A_0 = \alpha\phi$, the field configuration becomes static in time and the Gauss law (2.9) can be reduced to

$$-D_m \left\{ \frac{1}{g_{\text{eff}}^2}, D_m\phi \right\} + \frac{\kappa}{2}F_{mn}^2 = 0. \quad (3.9)$$

One can further reduce the Eqn. (3.9) by using the BPS equations as the traceless part of the following equation,

$$D_m^2\Phi = \frac{\kappa}{4}F_{mn}^2, \quad (3.10)$$

where

$$\Phi \equiv \frac{1}{g^2}\phi + \frac{\kappa}{2} \left(\phi^2 - \frac{\text{tr}(\phi^2)}{N}\mathbf{1}_N \right) \quad (3.11)$$

or, explicitly

$$D_m^2\Phi^a = \frac{\kappa}{2}d_{abc}F_{mn}^b F_{mn}^c, \quad (3.12)$$

where $\Phi = \Phi^a T^a$ and $\Phi^a = \frac{\partial}{\partial\phi^a}\mathcal{F}(\phi) = \frac{1}{g^2}\phi^a + \kappa d_{abc}\phi^b\phi^c$. This scalar equation does not depend on the choice of α . Here we want to note that BPS equations (3.8,3.12) are exactly the same to those obtained from the SUGRA analysis[9].

Before closing this subsection, we now in turn present the physical charges of our BPS solutions. We have to define the electric charge more carefully in the present model. We can usually read off the electric charge from the asymptotic behavior of the displacement vector $\mathcal{D}_m = \frac{\delta\mathcal{L}}{\delta A_m}$

$$\mathcal{D}_m = \left\{ \frac{1}{g_{\text{eff}}^2}, E_m \right\} - (\text{trace part}) = \frac{x_m}{4\pi^2 r^4} Q_N + \mathcal{O}\left(\frac{1}{r^4}\right), \quad (3.13)$$

where Q_N is the ‘nonabelian’ electric charge of the system. Assuming that the BPS configuration is localized in the space and that the scalar field will fall off as

$$\phi \approx v' + \mathcal{O}\left(\frac{1}{r^2}\right), \quad (3.14)$$

where we choose the gauge where v' is constant, one can determine the asymptotic behavior of Φ from the Eqn. (3.13) as

$$\Phi = \frac{v}{g^2} - \frac{Q_N}{8\pi^2 r^2} + \mathcal{O}\left(\frac{1}{r^3}\right). \quad (3.15)$$

The electric energy therefore becomes

$$Q_E = 2 \int d^4 \mathbf{x} \partial_m \text{tr}_N(\phi D_m \Phi) = \text{tr}(v' Q_N). \quad (3.16)$$

Taking into account of the positive kinetic energy, we should consider the parameter region $|v'| \ll (\kappa g^2)^{-1}$ for the Coulomb branch so that

$$v = v' + \frac{\kappa g^2}{2}(v')^2 - \frac{\kappa g^2 \mathbf{1}_N}{2N} \text{tr}_N(v')^2 \simeq v'. \quad (3.17)$$

Our BPS configurations can also carry the conserved angular momentum

$$\begin{aligned} J_{mn} &= \int d^4 \mathbf{x} (x_m P_n - x_n P_m) \\ &= 2 \int_{S_\infty^3} r^3 dS^p (x_m \text{tr}(\Phi F_{np}) - x_n \text{tr}(\Phi F_{mp})) + 4 \int d^4 \mathbf{x} \text{tr}(\Phi F_{mn}), \end{aligned} \quad (3.18)$$

where we used for the last equality the expression of the linear momentum density

$$P_m = \text{tr}(F_{mp}\{E_p, 1/g_{\text{eff}}^2\}) = 2\partial_p \text{tr}(F_{mp}\Phi). \quad (3.19)$$

The second term in Eqn. (3.18) can also be translated to the surface term as

$$\int d^4 \mathbf{x} \text{tr}(\Phi F_{mn}) = \int_{S_\infty^3} r^3 (dS^m \alpha_n - dS^n \alpha_m + \epsilon_{mnpq} dS^p \alpha_q), \quad (3.20)$$

once the one-form $\alpha = \alpha_m dx^m$ satisfy the differential equation

$$(1 + *_4) d\alpha = \text{tr}(\Phi F). \quad (3.21)$$

The solution of the above equation for the BPS configurations will be presented in the next subsection.

3.2 BPS Configurations

To find out the BPS configurations, we need to first solve the antiselfdual equation with $\alpha = -1$,

$$F_{mn} + \tilde{F}_{mn} = 0 \quad (3.22)$$

The general solutions of the antiselfdual equations are given the ADHM method. In a given instanton background, one needs to solve the second equation (3.10). Fortunately, the general solutions of the second equations can be also found in terms of the ADHM data.

Let us first summarize the ADHM method briefly. For details, it is referred to [24]. Starting with the constant matrices $(\bar{q}_i)_{N,K}$ and $(a_m)_{K,K}$, one can construct a $(N+2K) \times 2K$ matrix

$$\Delta = \left((\bar{q}_1 \ \bar{q}_2), M_m e^m \right) \quad (3.23)$$

where $e_m = (\sigma_a, i)$ and $M_m = a_m + \mathbf{1}_K x_m$. Note that $\bar{e}_m e_n = \delta_{mn} + i\eta_{mn}^a \sigma^a$ where selfdual t'Hooft tensor η_{mn}^a satisfies, for instances, $\eta_{12}^3 = \eta_{34}^1 = 1$. We introduce $V_{N+2N,N}(x)$ such that

$$\Delta^\dagger V = 0, \quad V^\dagger V = \mathbf{1}_N. \quad (3.24)$$

The condition on the ADHM data requires that

$$\Delta^\dagger \Delta = f^{-1}(x) \mathbf{1}_2 \quad (3.25)$$

with an invertible $K \times K$ matrix $f(x)$. This implies the ADHM condition

$$i\eta_{mn}^a [a_m, a_n] + \sigma_{ij}^a q_j \bar{q}_i = 0, \quad (3.26)$$

where $a = 1, 2, 3$. The ADHM method leads to the hermitian gauge field to be

$$A_\mu = iV^\dagger \partial_\mu V \quad (3.27)$$

for the anti-instantons.

As shown in appendix, the solution of Eqn. (3.10) for the traceless scalar field is given by

$$\Phi(x) = \bar{V} \begin{pmatrix} \frac{v}{g^2} & 0 \\ 0 & (\frac{\varphi}{g^2} - \frac{\kappa}{2} f(x)) \otimes \mathbf{1}_2 \end{pmatrix} V - \frac{\kappa}{4N} (\partial_m^2 \ln \det f) \mathbf{1}_N \quad (3.28)$$

where the constant matrices v and φ satisfy $\text{tr}_N v + \text{tr}_K \varphi = 0$ and an inhomogeneous linear matrix equation,

$$\frac{8\pi^2}{g^2} (\mathbf{L}\varphi + q_i v \bar{q}_i) - 8\pi^2 \kappa \mathbf{1}_K = 0 \quad (3.29)$$

where $\mathbf{L}\varphi = -[a_m, [a_m, \varphi]] - \frac{1}{2} \{q_i \bar{q}_i, \varphi\}$.

We now in turn express the physical charges of those BPS objects, dyonic instantons in terms of the ADHM data. From the leading order behavior of the scalar field Φ (3.28) at infinity

$$\Phi = \frac{v}{g^2} + \frac{1}{8\pi^2 r^2} \left(\frac{8\pi^2}{g^2} (\bar{q}_i \varphi q_i - \frac{1}{2} \{\bar{q}_i q_i, v\}) - \frac{nK}{N} \right) + \mathcal{O}\left(\frac{1}{r^3}\right), \quad (3.30)$$

together with Eqn. (3.15), one can read off the following electric charge of the $N \times N$ matrix

$$Q_N = \frac{8\pi^2}{g^2} \left(\frac{1}{2} \{\bar{q}_i q_i, v\} - q_i \varphi \bar{q}_i \right) + \frac{nK}{N} \mathbf{1}_N. \quad (3.31)$$

One can show from Eqn. (3.29) that the electric charge matrix Q_N is traceless. We now in turn consider the angular momentum of the dyonic instantons. As shown in [9], one can solve Eqn. (3.21) using the various identities in the ADHM method

$$\alpha_m = \frac{i}{4} \text{tr} \left(\mathcal{J}(P\partial_m P - \partial_m P P) + 4[\varphi, a_m]f - \frac{\kappa}{6} \epsilon_{mnpq} \partial_n f^{-1} f \partial_p f^{-1} f \partial_q f^{-1} f \right), \quad (3.32)$$

where $P = VV^\dagger$ and $\mathcal{J} = \frac{1}{g^2} \text{diag}(v_N, \varphi_K \mathbf{1}_2)$. For the BPS configurations, the angular momentum (3.18,3.20) therefore takes the form as

$$J_{mn} = -\frac{8\pi^2}{g^2} i \left\{ \frac{1}{2} \text{tr}_k \left(q_i v \bar{q}_j (\bar{\sigma}_{mn})_{ji} \right) + (1 + *_4) \text{tr}_k \left(\varphi [a_m, a_n] \right) \right\}. \quad (3.33)$$

3.3 A Single Dyonic $SU(N)$ Instanton

To be concrete, let us consider the profile of a single dyonic instanton in the $SU(N)$ gauge theory. The solution can be obtained either by using the general solution presented in the previous section, or simply solving it directly in the single instanton case. While the former is discussed in appendix, we follow the latter.

The $4N$ -dimensional moduli space of an instanton is composed of 4-parameters for the center of the instanton position and $4N - 4$ parameters for the internal degrees of freedom. The internal part is made of a single scale and a symmetric space $SU(N)/U(N-2)$. The internal $4N - 4$ dimensional moduli space is a hyper-Kähler cone over the quaternionic space $\frac{SU(N)}{SU(2) \times U(N-2)}$. The moduli space is spanned by moduli of the single $SU(2)$ 't Hooft instanton, plus additional moduli which determines the way how $SU(2)$ is embedded in $SU(N)$. In this section we work in the 'regular' gauge, opposed to the singular gauge which is convenient for the ADHM instantons.

The $SU(N)$ single instanton gauge field is given as

$$A_m = T^a A_m^a = T^a \bar{\eta}_{mn}^a \frac{2x_n}{r^2 + \rho^2}, \quad (3.34)$$

where T^a generates arbitrary $SU(2)$ sub-algebra of $SU(N)$ and $r = \sqrt{x_l x_l}$. We can take it to be the following:

$$T^a = U^\dagger \left(\frac{1}{2} \sigma^a \right) U, \quad [T^a, T^b] = i \epsilon^{abc} T^c, \quad (3.35)$$

where

$$U_{2,N} = \begin{pmatrix} e_{11} & e_{12} & \cdots & e_{1N} \\ e_{21} & e_{22} & \cdots & e_{2N} \end{pmatrix} \equiv \begin{pmatrix} \bar{\mathbf{e}}_1 \\ \bar{\mathbf{e}}_2 \end{pmatrix}, \quad \bar{\mathbf{e}}_i \cdot \mathbf{e}_j = \delta_{ij}. \quad (3.36)$$

In addition to the $SU(2)$ algebra, the generators T^a satisfy the anti-commutation relation

$$\{T^a, T^b\} = \frac{1}{2} \delta^{ab} U^\dagger U = \frac{\delta^{ab}}{N} \mathbf{1}_N - \frac{\delta^{ab}}{2} T, \quad (3.37)$$

where

$$T \equiv \frac{2}{N} \mathbf{1}_N - U^\dagger U . \quad (3.38)$$

The mutually orthogonal complex unit vectors \mathbf{e}_i contain $4N - 4$ real components. Among these, the $U(1)$ phase rotation $\mathbf{e}_i \rightarrow \mathbf{e}_i e^{i\alpha}$ does not affect T^a or A_m , so we may regard α as a ‘gauge’ degree. Together with the remaining $4N - 5$ gauge-invariant degrees, the scalar λ provides the desired $4N - 4$ zero modes apart from the 4 translations that we ignore here.

We are interested in the solution of the traceless part of Eqn. (3.10) for Φ . For a single instanton (3.34), the field strength can be described as

$$F_{mn} = \frac{4\rho^2}{(r^2 + \rho^2)^2} \bar{\eta}_{mn}^a T^a , \quad (3.39)$$

from which one can obtain

$$F_{mn}^2 = \frac{16\rho^4}{(r^2 + \rho^2)^4} 4T^a T^a = \frac{16\rho^4}{(r^2 + \rho^2)^4} \left(\frac{6}{N} \mathbf{1}_N - 3T \right) . \quad (3.40)$$

Taking the ansatz for Φ field as $\Phi = F(r) T$, the equation for Φ can be simply reduced to

$$\partial_m^2 F(r) = - \frac{12\kappa\rho^4}{(r^2 + \rho^2)^4} = \frac{\kappa}{8} (\partial^2)^2 (\log(r^2 + \rho^2)) . \quad (3.41)$$

The solution for $F(r)$ is therefore given by

$$F(r) = \frac{\kappa}{2} \frac{r^2 + 2\rho^2}{(r^2 + \rho^2)^2} , \quad (3.42)$$

in perfect agreement with that obtained from the ADHM method.

Now we would like to obtain the scalar $\phi = \phi^a T^a$ from the above Φ . Our claim is that ϕ is also spanned by the generator T only as $T^2 - \text{tr}_N T^2 / N \propto T$. Therefore we try $\phi = h(r)T$, and obtain the following equation for $h(r)$:

$$\frac{1}{g^2} h(r) + \frac{\kappa}{2} \frac{4 - N}{N} h(r)^2 = F(r) . \quad (3.43)$$

The solution $h_N(r)$ for the $SU(N)$ gauge theory is

$$h_4(r) = g^2 F(r) \quad (\text{for } N = 4) \quad (3.44)$$

$$h_N(r) = - \frac{1}{\kappa g^2} \frac{N}{4 - N} \pm \sqrt{\left(\frac{1}{\kappa g^2} \frac{N}{4 - N} \right)^2 + \frac{2}{\kappa} \frac{N}{4 - N} F(r)} , \quad (N \neq 4) \quad (3.45)$$

where the sign in front of the square-root for the second case is fixed by requiring that $\phi = 0$ at $r = \infty$.

To summarize, the scalar solution, and therefore the whole instanton solution in the presence of Chern-Simons term is classified into the following three cases:

1. For $SU(3)$, the scalar solution is given as

$$h_3(r) = -\frac{3}{\kappa g^2} + \sqrt{\left(\frac{3}{\kappa g^2}\right)^2 + \frac{3r^2 + 6\rho^2}{(r^2 + \rho^2)^2}}. \quad (3.46)$$

The solution is well-defined for all $\rho > 0$.

2. For $SU(4)$, the scalar is given as

$$h_4(r) = \frac{\kappa g^2}{2} \frac{r^2 + 2\rho^2}{(r^2 + \rho^2)^2}. \quad (3.47)$$

3. For $SU(N)$ with $N \geq 5$, the solution is given as

$$h_N(r) = \frac{1}{\kappa g^2} \frac{N}{N-4} - \sqrt{\left(\frac{1}{\kappa g^2} \frac{N}{N-4}\right)^2 - \frac{N}{N-4} \frac{r^2 + 2\rho^2}{(r^2 + \rho^2)^2}}. \quad (3.48)$$

The solution is well-defined at all $r \geq 0$ only in the background of large enough instanton, namely,

$$\rho \geq \rho_c \equiv \sqrt{\frac{2(N-4)}{N}} \kappa g^2. \quad (3.49)$$

For ρ smaller than this value ρ_c , there is no *real solution* $\phi(r)$ for $r < r_0$, where

$$4r_0^2 = \rho_c^2 - 4\rho^2 + \sqrt{\rho_c^4 + 8\rho^4}. \quad (3.50)$$

For large κ , there is however no meaning of the classical instanton of size of order $\kappa g^2 = ng^2/8\pi^2$, which is quite bigger than the ultraviolet scale g^2 . Although the UV completion of theory, or the reliable UV description of those solutions, seems to be important, we will not discuss about it any more.

4. Low Energy Dynamics

4.1 One-dimensional $U(K)$ Gauge Theory

The 16 supersymmetric $SU(N)$ gauge theory arises from the low energy dynamics of N D4 branes. It is well-known that the four-dimensional K instanton solitons are realized as K D0 branes on D4 branes. The low energy dynamics of instantons can be described by the one-dimensional $U(K)$ gauge theory with eight supersymmetries. The bosonic part of the Lagrangian is

$$\begin{aligned} \mathcal{L}_0 = \frac{M}{2} \text{tr}_K \left\{ (D_0 a_m)^2 + |D_0 q_i|^2 + [\varphi, a_m]^2 - |(\varphi q_i - q_i v)|^2 \right. \\ \left. + c(D_0 \varphi)^2 + c(D^a)^2 + D^a (\sigma_{ij}^a q_j \bar{q}_i + i \eta_{mn}^a [a_m, a_n]) - \zeta^a D^a \right\}, \quad (4.1) \end{aligned}$$

where $M = 8\pi^2/g^2$ denotes the instanton mass parameters. This Lagrangian also has a $SU(N)$ flavor symmetry, understood as the gauge symmetry in the five-dimensional gauge

symmetries. Under the flavor symmetry, the hypermultiplet scalars $q_i, i = 1, 2$ ($K \times N$ matrices) transform as

$$q_i \rightarrow q'_i = q_i U \quad (U \in SU(N)) . \quad (4.2)$$

Another hypermultiplet scalars $a_m, m = 1, 2, 3, 4$ denote the matrix position of K instantons on $D4$ branes. There are two controllable parameters of the present model of which one is the $N \times N$ hermitian matrix v , representing the positions of N D4-branes, and another is the Fayet-Iliopoulos parameter ζ , necessary for the noncommutative field theory. As well as the gauge symmetry $U(K)$ and flavor symmetry $SU(N)$, the theory has $SO(4) \simeq SU(2)_R \times SU(2)$ global symmetry. For instances, a_m furnish a vector representation under $SO(4)$ while q_i are in a fundamental representation under $SU(2)_R$. The arbitrary parameter c of dimension $mass^4$ can be determined from this D0–D4 system by $c = 2\pi/(2\pi l_s)^4$.

In this work, our 5-dim theory has 8 supersymmetries and a Chern-Simons term. Instantons of our theory is again 1/2 BPS with now four supersymmetries and so their property will be different from those from 16 supersymmetric theories. One may wonder what the low energy dynamics of instantons with four supersymmetries. As far as the classical feature is concerned, our analysis of the BPS equations tells us that the gauge field configuration does not change. Only our instantons now carry some electric charge due to the Chern-Simons term. Thus we expect a simple modification of the above Lagrangian may suffice. Indeed, here we argue that it is sufficient to add to the Lagrangian (4.1) the following 1-dim Chern-Simons term

$$\mathcal{L}_{cs} = n \operatorname{tr}_K(A_0 - \varphi) . \quad (4.3)$$

Let us first look at the supersymmetry. Note that a_m, q_i are hypermultiplet and A_0, φ , are part of the vector multiplet. In 10-dim notation for the vector multiplet, the above Chern-Simons term is invariant under supersymmetric transformation

$$\delta(A_0 - \varphi) = \bar{\lambda}(\Gamma_0 - \Gamma_5)\epsilon = \bar{\lambda}\Gamma_0(1 - \Gamma^{05})\epsilon = 0 \quad (4.4)$$

if $\Gamma^{05}\epsilon = \epsilon$. Thus imposing additional condition on the spinor parameter breaks the supersymmetry to half. It is to be noted that the 1-dim Chern-Simons term introduces a background electric charge to $U(K)$ theory. With the presence of the fundamental representation of $U(K)$, the invariance of the path integral amplitude under a local $U(1)$ gauge transformation $e^{i2\pi\Lambda(t)T}$ such that $\Lambda(\infty) - \Lambda(-\infty) = 2\pi l$ where $T = \operatorname{diag}(1, 0, 0, \dots, 0)$, leads to the quantization $n \in \mathbf{Z}$.

We then consider the stringy origin of the one-dimensional Chern-Simons term (4.3). From our D-brane picture in section 2, the instantons were identified as D1-branes stretched between NS5-branes. The open strings connecting flavor D7 or D5 branes to D1-instantons leads to the massive fundamental hypermultiplets in the matrix model (4.1). Their one-loop effect indeed induces the very 1-dimensional Chern-Simons term.

The ground state should have the lowest energy. First of all the D-term equation implies that

$$D^a = \frac{1}{2c}(\zeta^a - i\bar{\eta}_{mn}^a[a_m, a_n] - \sigma_{ij}^a q_j \bar{q}_i) . \quad (4.5)$$

Since the Chern-Simons term does not affect the energy, the energy of the Lagrangian $L_0 + L_{cs}$ can be rearranged as

$$\begin{aligned} \mathcal{E} = & \frac{M}{2} \text{tr}_K \left\{ (D_0 a_m + i[\varphi, a_m])^2 + |D_0 q_i + i(\varphi q_i - q_i v)|^2 + c(D_0 \varphi)^2 + c(D^a)^2 \right\} \\ & + M \text{tr}_K \left\{ \varphi \left(-i[a_m, D_0 a_m] - \frac{i}{2}(q_i D_0 \bar{q}_i - D_0 q_i \bar{q}_i) \right) + \frac{n}{M} \right\} \\ & + \frac{M}{2} \text{tr}_N \left\{ i(D_0 \bar{q}_i q_i - \bar{q}_i D_0 q_i) v \right\} \end{aligned} \quad (4.6)$$

After using the Gauss law constraint

$$M \left(-i[a_m, D_0 a_m] - \frac{i}{2}(q_i D_0 \bar{q}_i - D_0 q_i \bar{q}_i) \right) + n \mathbf{1}_K = 0 , \quad (4.7)$$

we can obtain a BPS bound on the energy

$$\mathcal{E} \geq \text{tr}_N(v Q_N) , \quad (4.8)$$

where the $SU(N)$ flavor charge is given by

$$Q_N = M \left(\frac{i}{2}(D_0 \bar{q}_i q_i - \bar{q}_i D_0 q_i) - \frac{i}{2N} \text{tr}_N(D_0 \bar{q}_i q_i - \bar{q}_i D_0 q_i) \mathbf{1}_N \right) . \quad (4.9)$$

The BPS configurations which saturate the bound should satisfy

$$cD^a = \zeta - \bar{\eta}_{mn}^a[a_m, a_n] - \sigma_{ij}^a q_j \bar{q}_i = 0 \quad (4.10)$$

which is nothing but the ADHM constraint. We started with $4K^2$ -parameterized a_m and $4NK$ parameterized q_i . Imposing the $3K^2$ ADHM constraint and taking out $K^2 U(K)$ gauge degrees of freedom leads to $4KN$ moduli-space coordinates $z_A, A = 1, 2, \dots, 4KN$. The solutions of the above equation would be given by $a_m(z_A)$ and $q_i(z_A)$. The rest of the BPS equations are

$$D_0 a_m + i[\varphi, a_m] = 0, \quad D_0 q_i + i(\varphi q_i - q_i v) = 0, \quad D_0 \varphi = 0 . \quad (4.11)$$

In the gauge $A_0 = \varphi$, the configuration is static in time, besides $q_i \sim q_{i0} e^{ivt}$. For the BPS configurations, the Gauss law becomes

$$M \left(-[a_m, [a_m, \varphi]] - \frac{1}{2} \{q_i \bar{q}_i, \varphi\} + q_i v \bar{q}_i \right) - 8\pi^2 \kappa \mathbf{1}_K = 0 , \quad (4.12)$$

which is identical to Eqn. (3.29) in the field theory, once we noting that $8\pi^2 \kappa = n$. This leads to a further support for the validity of our low energy Lagrangian for the instanton solitons. As the kinetic part for φ is positive-definite, the solution would be a linear combination,

$$\varphi = \varphi_v(z_A) + \varphi_\kappa(z_A) \quad (4.13)$$

where $\mathbf{L}\varphi_v + q_i v \bar{q}_i = 0$ and $M\mathbf{L}\varphi_\kappa - 8\pi^2\kappa = 0$.

Let us now in turn consider the various charges that the BPS configurations can carry. The traceless $SU(N)$ flavor charge (4.9) becomes

$$Q_N = -M \left(\bar{q}_i \varphi q_i - \frac{1}{2} \{ \bar{q}_i q_i, v \} \right) + \frac{nK}{N} \mathbf{1}_N , \quad (4.14)$$

which is in the agreement with the electric charge (3.31) of the dyonic instantons. We then compute the angular momentum of this half BPS solution

$$\begin{aligned} J_{mn} &= -iM \text{tr} \left(\frac{1}{4} (q_i D_0 \bar{q}_j - D_0 q_i \bar{q}_j) (\bar{\sigma}_{mn})_{ji} + (D_0 a_m a_n - a_m D_0 a_n) \right) \\ &= -iM \text{tr} \left(\frac{1}{2} (q_i v \bar{q}_j - \varphi q_i \bar{q}_j) (\bar{\sigma}_{mn})_{ji} + 2\varphi [a_m, a_n] \right) \\ &= -iM \text{tr} \left(\frac{1}{2} (q_i v \bar{q}_j) (\bar{\sigma}_{mn})_{ji} + (1 + *_4) (\varphi [a_m, a_n]) \right) , \end{aligned} \quad (4.15)$$

where for the last equality we used the ADHM constraints (4.10). This angular momentum agrees again with the result (3.33) in field theory analysis. It is therefore conceivable to identify the dyonic instantons with the BPS configurations of the present 1-dim gauge theory.

4.2 Moduli Space Dynamics

The vacuum configuration modulo the gauge equivalence is the moduli space. The low energy dynamics is described by the nonlinear sigma model of the moduli coordinates. For the K instantons of the $SU(N)$ gauge theory, the nonlinear sigma model is characterized by the $4KN$ -dimensional hyper-Kähler space, with several Killing vectors for symmetry of the instanton moduli space. The Coulomb phase leads the natural potential on the moduli space. We argue that the 1-dim Chern-Simons term leads to the magnetic field on the instanton moduli space. We may wonder how the Chern-Simons effect can be realized in the moduli space dynamics. In terms of the moduli space coordinates z^A , the dynamics can be described by the Lagrangian

$$L = \frac{M}{2} g_{AB}(z) \dot{z}^A \dot{z}^B + n \dot{z}^A \mathcal{A}_A(z) - U(z) . \quad (4.16)$$

(A similar magnetic field has appeared in the moduli space of magnetic flux vortices in the Abelian Higgs model with a background charge and a Chern-Simons terms [25].)

The metric for the moduli space can be determined by the standard procedure. We start with ignoring the effect of the Chern-Simons term κ and the vacuum parameter v . One way to approach is to put the initial field configuration,

$$a_m(z), q_i(z), \quad (4.17)$$

determined by the moduli coordinate, and its time derivative

$$\begin{aligned}\hat{D}_0 a_m &= \partial_0 a_m - i[\hat{A}_0, a_m] = \dot{z}^A \hat{D}_A a_m = \dot{z}^A \left(\frac{\partial a_m}{\partial z^A} - i[\Lambda_A, a_m] \right), \\ \hat{D}_0 q_i &= \partial_0 q_i - i\hat{A}_0 q_i = \dot{z}^A \hat{D}_A q_i = \dot{z}^A \left(\frac{\partial q_i}{\partial z^A} - i(\Lambda_A q_i - q_i) \right),\end{aligned}\quad (4.18)$$

which is given by the velocity of the moduli coordinate. The initial configuration should satisfies the Gauss law constraint (4.7), implying that

$$\mathbf{L}\hat{A}_0 = -i[a_m \partial_0 a_m] - \frac{i}{2}(q_i \partial_0 \bar{q}_i - \partial_0 q_i \bar{q}_i). \quad (4.19)$$

Inserting this initial field configuration back to the $U(K)$ gauge theory Lagrangian (4.1), one can obtain the metric on the moduli space

$$g_{AB}(z) = \text{tr}_K \left(\hat{D}_A a_m \hat{D}_B a_m + \frac{1}{2}(\hat{D}_A q_i \hat{D}_B \bar{q}_i + \hat{D}_B q_i \hat{D}_A \bar{q}_i) \right). \quad (4.20)$$

With the Chern-Simons term and the vacuum expectation value v , we modify the field velocity of the previous initial field configurations to be

$$D_0 a_m = \hat{D}_0 a_m - i[\Delta A_0, a_m], \quad D_0 q_i = \hat{D}_0 q_i - i\Delta A_0 q_i \quad (4.21)$$

and suppose the scalar field φ relaxes to the lowest energy configuration $\varphi = \varphi_\kappa + \varphi_v$ as before. The Gauss law and the scalar field equation tells us $\Delta A_0 = \varphi_\kappa$ and the $U(K)$ gauge theory Lagrangian now can take the form as

$$L = \frac{M}{2} g_{AB} \dot{z}^A \dot{z}^B + 8\pi^2 \kappa (\hat{A}_0 - \varphi_v) - U(z), \quad (4.22)$$

from which one can find various corrections to the instanton moduli space dynamics. Defining the gauge field in the moduli space to be

$$\mathcal{A}_A(z) = \text{tr}_K \mathbf{L}^{-1} \left(-i \left[a_m, \frac{\partial a_m}{\partial z^A} \right] - \frac{i}{2} \left(q_i \frac{\partial \bar{q}_i}{\partial z^A} - \frac{\partial q_i}{\partial z^A} \bar{q}_i \right) \right), \quad (4.23)$$

the Chern-Simons term becomes

$$8\pi^2 \kappa \text{tr}_K(\hat{A}_0) = 8\pi^2 \kappa \sum_A \dot{z}^A \mathcal{A}_A(z). \quad (4.24)$$

Finally the potential term

$$U(z) = \frac{M}{2} \text{tr}_K \left(-[a_m, \varphi_v]^2 + (\varphi_v q_i - q_i v)(\bar{q}_i \varphi_v - v \bar{q}_i) \right) = \frac{M}{2} \text{tr}_K \left(-\varphi_v q_i v \bar{q}_i + q_i v^2 \bar{q}_i \right) \quad (4.25)$$

is of order $\mathcal{O}(v^2)$ correction. As well as $\mathcal{O}(\kappa)$ correction $8\pi^2 \kappa \hat{A}_0$, there is of $\mathcal{O}(\kappa v)$ correction $-8\pi^2 \kappa \varphi_v$.

We can find the Killing vector related to the $SU(N)$ gauge symmetry and the potential more explicitly. The $SU(N)$ phase rotations of the theory lead to the cyclic coordinates

with Killing vectors forming $SU(N)$ algebra. We choose the $N - 1$ commuting direction ζ^a so that we can choose $\dot{\zeta}^a T^a$ to be diagonal. The Killing vector becomes

$$K_a^A \frac{\partial}{\partial z^A} = \frac{\partial}{\partial \zeta^a} \quad (4.26)$$

The moduli space dynamics becomes

$$L = \frac{M}{2} g_{AB} \dot{z}^A \dot{z}^B + 8\pi^2 \kappa (\dot{z}^A - G^A) \mathcal{A}_A - \frac{M}{2} g_{AB} G^A G^B \quad (4.27)$$

where $G^A = K_a^A v^a$. This moduli space dynamics is of order $\dot{z}^2, \kappa \dot{z}, \kappa v, v^2$. We ignore other higher order corrections. Calling the rest of the coordinates y^i , we can re-express the above low energy dynamics as

$$L = \frac{M}{2} h_{ij}(y) \dot{y}^i \dot{y}^j + \frac{M}{2} k_{ab}(y) (\dot{\zeta}^a + \dot{y}^i \omega_i^a(y)) (\dot{\zeta}^b + \dot{y}^j \omega_j^b(y)) \quad (4.28)$$

$$+ 8\pi^2 \kappa \dot{y}^i \mathcal{C}_i(y) + 8\pi^2 \kappa (\dot{\zeta}^a - v^a) \mathcal{V}_a(y) - \frac{M}{2} k_{ab}(y) v^a v^b \quad (4.29)$$

One can easily give a BPS bound on the energy related to the above Lagrangian and can obtain the ground state energy as

$$E_{ground} = v^a p_a \quad (4.30)$$

where p_a is the canonical momentum related to the cyclic coordinates. This ground state energy is equivalent to the minimum of the bound (4.8). The BPS ground state configuration satisfies

$$\dot{y}^i = 0, \quad \dot{\zeta}^a = v^a \quad (4.31)$$

It would be interesting to find the supersymmetric extension of the moduli space dynamics (4.27).

4.3 A Single Chern-Simons Instanton Moduli Dynamics

Let us consider the moduli space dynamics of a single instanton. As discussed before, its moduli space is a $4N$ -dimensional hyper-Kähler manifold. For generality, we will turn on every possible parameters κ and ζ except the vev v . It implies that we will analyze the moduli space of a single instanton in the $U(N)$ gauge theory on noncommutative R^4 space rather than commutative space. We can of course take the commutative limit to the $SU(N)$ gauge theory instanton by letting $\zeta \rightarrow 0$ together with decoupling $U(1)$ factor. The single instanton moduli space for $U(N)$ gauge group is known to be the Calabi space of dim $4N$ which is complete nonsingular hyperKähler space with cohomogeneity one.

Before the detailed analysis, we first summarize the conventions used in Ref. [26] to find the moduli space metric. For the invariant one-forms L^i_j of $SU(N) (\subset U(N))$ defined as

$$U dU^\dagger = -i L^i_j t_i^j, \quad [t_i^j, t_k^l] = \delta_k^j t_i^l - \delta_i^l t_k^j, \quad (4.32)$$

with $(t_i^j)^\dagger = t_j^i$ (or $(L_i^j)^\dagger = L_j^i$) and $L^i_i = 0$, the Maurer-Cartan equations are

$$dL^i_j = iL^i_k \wedge L^k_j. \quad (4.33)$$

Here i run over $(1, 2, a)$, where a denote the $SU(N-2)$ indices. One can then define the generators of the coset group $SU(N)/(SU(N-2) \times U(1))$ as

$$\sigma_a = L^1_a, \quad \Sigma_a = L^2_a, \quad \mu = L^1_1 - L^2_2, \quad \nu = L^1_2, \quad (4.34)$$

and those of $SU(N-2) \times U(1)$ as

$$\tilde{L}^a_b = L^a_b + \frac{1}{n} Q \delta^a_b, \quad Q = L^1_1 + L^2_2. \quad (4.35)$$

With those new basis, the Maurer-Cartan equations of our interests, for examples, take the form as

$$d\mu = 2i\nu \wedge \bar{\nu} + i\sigma_a \wedge \bar{\sigma}^a - i\Sigma_a \wedge \bar{\Sigma}^a, \quad dQ = i\sigma_a \wedge \bar{\sigma}^a + i\Sigma_a \wedge \bar{\Sigma}^a. \quad (4.36)$$

Let us first consider the D-term conditions. Since for a single instanton a_m are simple c-numbers, we can solve explicitly the D-term condition with $\zeta^a = \delta^{a3} \zeta^2$ as

$$\begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{u^2 + \zeta^2} & 0 & \cdots & 0 \\ 0 & \sqrt{u^2 - \zeta^2} & \cdots & 0 \end{pmatrix} U \equiv q_0 U \quad (4.37)$$

where U is an $U(N)$ element. The two useful quantities in our discussion are summarized as

$$\begin{aligned} \text{tr} \left\{ (\bar{q}_0 q_0) U dU^\dagger \right\} &= -\frac{i}{2} (\zeta^2 \mu + u^2 Q) - i d\alpha \\ \text{tr} \left\{ (\bar{q}_0 q_0) U dU^\dagger U dU^\dagger \right\} &= - \left(u^2 \left(\frac{Q^2 + \mu^2}{4} + \nu \bar{\nu} \right) + \frac{\zeta^2}{2} Q \mu + \frac{u^2 + \zeta^2}{2} \sigma_a \bar{\sigma}^a + \frac{u^2 - \zeta^2}{2} \Sigma_a \bar{\Sigma}^a \right), \end{aligned} \quad (4.38)$$

where $d\alpha$ denotes the one-form corresponding to $U(1) (\subset U(N))$ generator that rotates overall phase of q_i . For the single instanton, the moduli space metric (4.20) of $U(N)$ non-commutative theory can be described as

$$ds^2 = \left(1 - \frac{\zeta^4}{u^4} \right)^{-1} du^2 + \frac{u^2}{4} \left(1 - \frac{\zeta^4}{u^4} \right) \mu^2 + u^2 \nu \bar{\nu} + \frac{u^2 + \zeta^2}{2} \sigma_a \bar{\sigma}^a + \frac{u^2 - \zeta^2}{2} \Sigma_a \bar{\Sigma}^a, \quad (4.39)$$

which is known to be Calabi metric [26]. The one-form gauge field (4.23) on the moduli space now becomes

$$\mathcal{A}_A dz^A = \frac{\zeta^2}{2u^2} \mu + \frac{1}{2} Q + d\alpha, \quad (4.40)$$

whose field strength is therefore

$$F = d\mathcal{A} = \frac{1}{2u^2} \left(-2\zeta^2 \frac{du}{u} \wedge \mu + 2i\zeta^2 \nu \wedge \bar{\nu} + i(u^2 + \zeta^2) \sigma_a \wedge \bar{\sigma}^a + i(u^2 - \zeta^2) \Sigma_a \wedge \bar{\Sigma}^a \right). \quad (4.41)$$

As mentioned before, the single Chern-Simons instanton can carry the electric charge (4.14) even in the symmetric phase $v = 0$

$$Q_N = -8\pi^2\kappa \left[\frac{1}{N} \begin{pmatrix} \frac{N-2}{2}\mathbf{1}_2 & 0 \\ 0 & -\mathbf{1}_{N-2} \end{pmatrix} - \frac{\zeta^2}{2u^2} \begin{pmatrix} \sigma_3 & 0 \\ 0 & 0 \end{pmatrix} \right], \quad (4.42)$$

and no angular momentum. We note that, for commutative $SU(2)$ gauge theory, the charge matrix Q_N identically vanishes. It agrees with the fact that there is no non-abelian Chern-Simons coupling for $SU(2)$ group.

Especially for a single instanton in $U(2)$ gauge theory on noncommutative space, the moduli space metric is Eguchi-Hanson space with the metric the moduli space metric of the non-commutative $U(2)$ single instanton now becomes

$$ds^2 = \left(1 - \frac{\zeta^4}{u^4}\right)^{-1} du^2 + \frac{u^2}{4} (\sigma_1^2 + \sigma_2^2) + \frac{u^2}{4} \left(1 - \frac{\zeta^4}{u^4}\right) \sigma_3^2, \quad (4.43)$$

which is nothing but the Eguchi-Hanson metric. Here, σ_i are the standard $SU(2)$ invariant one-forms

$$\begin{aligned} \sigma_1 &= -\sin\psi d\theta + \cos\psi \sin\theta d\varphi \\ \sigma_2 &= \cos\psi d\theta + \sin\psi \sin\theta d\varphi \\ \sigma_3 &= d\psi + \cos\theta d\varphi, \end{aligned} \quad (4.44)$$

which satisfy $d\sigma_i = \frac{1}{2}\epsilon_{ijk}\sigma_j \wedge \sigma_k$. The gauge field \mathcal{A}_A can be expressed as

$$\mathcal{A} = \frac{\zeta^2}{2u^2}\sigma_3 + d\alpha, \quad (4.45)$$

whose field strength now becomes the unique normalizable self-dual harmonic two-form on the Eguchi-Hanson space[26]

$$\mathcal{F} = d\mathcal{A} = \frac{\zeta^2}{2u^2} \left(-\frac{2}{u} du \wedge \sigma_3 + \sigma_1 \wedge \sigma_2 \right). \quad (4.46)$$

When we take the commutative limit $\zeta \rightarrow 0$ with ignoring overall $U(1)$ factor, the moduli space is reduced to the orbifold C^2/Z_2 and the gauge field A_A disappears. This disappearance is again consistent to the non-existence of the single Chern-Simons instanton for $SU(2)$.

For two identical instantons in the $U(1)$ theory on noncommutative space, the moduli space metric is again Eguchi-Hanson and the 1-form \mathcal{A} is again identical to that of the single instanton in the $U(2)$ theory as shown in the previous paragraphs [27].

We then restrict our attention only on the commutative five-dimensional field theories, i.e., $\zeta \rightarrow 0$ as well as ignoring $d\alpha$ ($U(1)$) effect. Since the asymptotic Higgs vev is not developed, non-abelian gauge symmetry $SU(N)$ is not broken. We therefore expect that a single Chern-Simons instanton can carry charges in a certain representation under the gauge

symmetry $SU(N)$. In order to specify the representation, we will discuss the quantization of the single Chern-Simons instanton moduli space and then find out the ground state degeneracies.

In the $\zeta \rightarrow 0$ limit, the metric of the moduli space becomes

$$ds^2 = du^2 + \frac{u^2}{4}\mu^2 + u^2\nu\bar{\nu} + \frac{u^2}{2}\sigma_a\bar{\sigma}^a + \frac{u^2}{2}\Sigma_a\bar{\Sigma}^a, \quad (4.47)$$

while the Chern-Simons effect on the moduli space for the present model is given by

$$8\pi^2\kappa\mathcal{A}_Adz^A = \frac{8\pi^2\kappa}{2}Q. \quad (4.48)$$

One can show that the uniform magnetic flux F , proportional to the Kähler two-form on the coset space $\mathcal{G} = G/H \equiv \frac{SU(N)}{SU(2)\times SU(N-2)\times U(1)}$, is now turned on in the moduli space

$$8\pi^2\kappa\mathcal{F} = \frac{8\pi^2i\kappa}{2}(\sigma_a \wedge \bar{\sigma}^a + \Sigma_a \wedge \bar{\Sigma}^a), \quad (4.49)$$

from which the quantization of the Chern-Simons coefficient

$$8\pi^2\kappa = n, \quad (4.50)$$

can be also understood as the Dirac quantization. The problem in considerations can be thus reduced to as the Landau problem of a charged particle moving on the coset space \mathcal{G} under the background $U(1)$ gauge field, analyzed extensively in [28, 29]. As a shortcut, we will just sketch the basic idea how to analyze the representation, or degeneracy of the ground states. For details, it is referred to [28, 29].

When a particle moves on the coset space $\mathcal{G} = G/H$ (with the background gauge fields), energy eigenstates of the system should be in certain representations under G . Since the background gauge field \mathcal{A} (4.48) is turned on only along the $U(1)$ direction in H , the eigenstates are singlet under $SU(2) \times SU(N-2)$, implying that they can be represented as \square , its complex conjugate $\bar{\square}$ and their symmetric tensor products without any trace part. We will denote such representations by (\mathbf{p}, \mathbf{q}) (p -times symmetric products of \square and q -times products of $\bar{\square}$). One can show from Eqn. (4.48) that a state in (\mathbf{p}, \mathbf{q}) carries $U(1)$ charge $p - q$, which has to be identified as the background charge $8\pi^2\kappa = n$, that is

$$p - q = n. \quad (4.51)$$

Assuming $n > 0$ without loss of generality, one can conclude that q can play the role of the Landau level index because higher dimensional representation states are in the higher energy levels. For the lowest Landau level ($q = 0$, or $p = n$), the degenerate ground states are therefore in the representation of κ -times symmetric products of \square , i.e.,

$$\underbrace{\begin{array}{|c|c|c|c|c|c|c|} \hline \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}}_n \quad (SU(N) \text{ Young tableaux}) . \quad (4.52)$$

As non-trivial cross checks of these results, let us consider the case of $N = 2$. Since there is no Chern- Simons terms for the gauge group $SU(2)$ in the five-dimensional theories, the instanton should be in a singlet whatever value the coefficient κ is. The result above in (4.52) agrees with this expectation. For the case of $N = 3$, the coset space can be identified as the projective space CP^2 whose lowest Landau levels under the n-unit magnetic field are well-known to be

$$\underbrace{\begin{array}{|c|c|c|c|c|c|c|} \hline & & & & & & \\ \hline \end{array}}_n \quad (SU(3) \text{ Young tableaux }) , \quad (4.53)$$

nothing but the complex conjugate of the representation above. One can therefore again see the agreement.

Acknowledgment

We would like to thank Sangmin Lee, Ho-Ung Yee and Piljin Yi for valuable discussions. This work is supported in part by the KOSEF SRC Program through CQUeST at Sogang University (KML), KRF Grants No. KRF-2005-070-C00030 (KML), and the KRF National Scholar program (SL,KML).

A. the ADHM Formalism and the Higgs Solutions

Using the ADHM method, the field strength can be expressed as

$$F_{mn} = -2\bar{\eta}_{mn}^a \bar{V} b f \sigma^a b^\dagger V , \quad (A.1)$$

where $b^\dagger = (0_{2K,N}, \mathbf{1}_{2K})$. As $\bar{\eta}_{mn}^a \bar{\eta}_{mn}^b = 4\delta^{ab}$, we get

$$\begin{aligned} \frac{F_{mn}^2}{16} &= \bar{V} b f \sigma^a b^\dagger V \bar{V} b f \sigma^a b^\dagger V \\ &= 3\bar{V} b f^2 b^\dagger V - 3\bar{V} b f M_m f M_m f b^\dagger V + i\bar{\eta}_{mn}^a \bar{V} b f \sigma^a M_m f M_n f b^\dagger V \end{aligned} \quad (A.2)$$

Note that

$$\partial_m f = -f(\partial_m f^{-1})f = -2fM_m f \quad (A.3)$$

and

$$D_m(\bar{V} b f b^\dagger V) = -4\bar{V} b f M_m f b^\dagger V \quad (A.4)$$

Thus

$$D_m^2(\bar{V} b f b^\dagger V) = 8\bar{V} b f^2 b^\dagger V - \frac{1}{2}F_{mn}^2 \quad (A.5)$$

Thus, the hermitian field strength satisfies

$$\text{tr}_N(F_{mn}^2) = \partial_m^2 \partial_n^2 \ln \det(f) \quad (A.6)$$

We present an ADHM solution for the equation (3.10)

$$\mathcal{D}^2\Phi = \frac{\kappa}{4}F_{mn}^2 \quad (\text{A.7})$$

where fields are in $N \times N$ matrix representation. We now replace F^2 by using Eq. (A.5). This leads to

$$D_m^2\Phi = \frac{\kappa}{2}\left(8\bar{V}bf^2b^\dagger V - D_m^2(\bar{V}bfb^\dagger V)\right) \quad (\text{A.8})$$

or

$$D_m^2(\Phi + \frac{\kappa}{2}\bar{V}bfb^\dagger V) = 4\kappa\bar{V}f^2b^\dagger V \quad (\text{A.9})$$

For simplicity we introduce

$$\Psi = \Phi + \frac{\kappa}{2}\bar{V}bfb^\dagger V \quad (\text{A.10})$$

which satisfy

$$D_m^2\Psi = 4\kappa\bar{V}bf^2b^\dagger V \quad (\text{A.11})$$

We solve this equation with the ansatz

$$\Psi = \frac{1}{g^2}\bar{V}\begin{pmatrix} v & 0 \\ 0 & \varphi \otimes \mathbf{1}_2 \end{pmatrix}V \quad (\text{A.12})$$

$$D_m^2\Psi = \frac{4}{g^2}\bar{V}bf\left(\mathbf{L}\varphi + q_iv\bar{q}_i\right)fb^\dagger V \quad (\text{A.13})$$

where a linear operator \mathbf{L} is defined so that

$$\mathbf{L}\varphi \equiv -[a_m, [a_m, \varphi]] - \frac{1}{2}\{q_i\bar{q}_i, \varphi\} \quad (\text{A.14})$$

Thus the equation $D_m^2\Psi = 2\kappa\bar{V}bf^2b^\dagger V$ can be solved if

$$\frac{8\pi^2}{g^2}\left(\mathbf{L}\varphi + q_iv\bar{q}_i\right) - 8\pi^2\kappa\mathbf{1}_K = 0 \quad (\text{A.15})$$

Note that the above equation is invariant under $(q, \varphi) \rightarrow (q, \varphi) + c(\mathbf{1}_N, \mathbf{1}_K)$. Once the solution is known, the traceless scalar field configuration would be

$$\Phi(x) = \bar{V}\begin{pmatrix} \frac{v}{g^2} & 0 \\ 0 & (\frac{\varphi}{g^2} - \frac{\kappa}{2}f(x)) \otimes \mathbf{1}_2 \end{pmatrix}V - \frac{\kappa}{4N}\partial_m^2 \ln \det f \quad (\text{A.16})$$

where the traceless condition implies that $\text{tr}_N q + \text{tr}_K \varphi = 0$.

References

- [1] N.D. Lambert and D. Tong, ‘‘Dyonic instantons in five-dimensional gauge theories,’’ Phys. Lett. B **462** (1999) 89 [arXiv:hep-th/9907014].
- [2] M. Zamaklar, ‘‘Geometry of the nonabelian DBI dyonic instanton,’’ Phys. Lett. B **493** (2000) 411 [arXiv:hep-th/0006090].

- [3] E. Eyras, P. K. Townsend and M. Zamaklar, “The heterotic dyonic instanton,” JHEP **0105** (2001) 046 [arXiv:hep-th/0012016].
- [4] D. s. Bak and K. M. Lee, “Supertubes connecting D4 branes,” Phys. Lett. B **544** (2002) 329 [arXiv:hep-th/0206185].
- [5] S. Kim and K.M. Lee, “Dyonic instanton as supertube between D4 branes,” JHEP **0309**, 035 (2003) [arXiv:hep-th/0307048].
- [6] S. Kim, K. M. Lee and H. U. Yee, “Supertubes in field theories,” Phys. Rev. D **75** (2007) 125011 [arXiv:hep-th/0603179].
- [7] H.Y. Chen, M. Eto and K. Hashimoto, “The shape of instantons: Cross-section of supertubes and dyonic instantons,” JHEP **0701**, 017 (2007) [arXiv:hep-th/0609142].
- [8] M.Y. Choi, K.K. Kim, C. Lee and K.M. Lee, “Higgs Structures of Dyonic Instantons,” arXiv:0712.0735 [hep-th].
- [9] S. Kim and S. Lee, “The geometry of dyonic instantons in 5-dimensional supergravity,” arXiv:0712.0090 [hep-th].
- [10] N. Seiberg, “Five dimensional SUSY field theories, non-trivial fixed points and string dynamics,” Phys. Lett. B **388** (1996) 753 [arXiv:hep-th/9608111].
- [11] D.R. Morrison and N. Seiberg, “Extremal transitions and five-dimensional supersymmetric field theories,” Nucl. Phys. B **483** (1997) 229 [arXiv:hep-th/9609070].
- [12] K.A. Intriligator, D.R. Morrison and N. Seiberg, “Five-dimensional supersymmetric gauge theories and degenerations of Calabi-Yau spaces,” Nucl. Phys. B **497** (1997) 56 [arXiv:hep-th/9702198].
- [13] O. Aharony and A. Hanany, “Branes, superpotentials and superconformal fixed points,” Nucl. Phys. B **504** (1997) 239 [arXiv:hep-th/9704170].
- [14] O. Aharony, A. Hanany and B. Kol, “Webs of (p,q) 5-branes, five dimensional field theories and grid diagrams,” JHEP **9801** (1998) 002 [arXiv:hep-th/9710116].
- [15] M. Eto, M. Nitta, K. Ohashi and D. Tong, “Skyrmions from instantons inside domain walls,” Phys. Rev. Lett. **95** (2005) 252003 [arXiv:hep-th/0508130].
- [16] D. K. Hong, M. Rho, H. U. Yee and P. Yi, “Dynamics of Baryons from String Theory and Vector Dominance,” JHEP **0709** (2007) 063 [arXiv:0705.2632 [hep-th]].
- [17] M. R. Douglas and N. A. Nekrasov, “Noncommutative field theory,” Rev. Mod. Phys. **73** (2001) 977 [arXiv:hep-th/0106048].
- [18] A. Boyarsky, J. A. Harvey and O. Ruchayskiy, “A toy model of the M5-brane: Anomalies of monopole strings in five dimensions,” Annals Phys. **301** (2002) 1 [arXiv:hep-th/0203154].
- [19] D. Tong, “TASI lectures on solitons,” arXiv:hep-th/0509216.
- [20] M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, “Instantons in the Higgs phase,” Phys. Rev. D **72**, 025011 (2005) [arXiv:hep-th/0412048].
- [21] E. Witten, “Phase transitions in M-Theory and F-Theory,” Nucl. Phys. B **471** (1996) 195 [arXiv:hep-th/9603150].

- [22] E. Witten, “Solutions of four-dimensional field theories via M-theory,” Nucl. Phys. B **500** (1997) 3 [arXiv:hep-th/9703166].
- [23] A. Hanany and E. Witten, “Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics,” Nucl. Phys. B **492** (1997) 152 [arXiv:hep-th/9611230].
- [24] N. Dorey, T.J. Hollowood, V.V. Khoze and M.P. Mattis, “The calculus of many instantons,” Phys. Rept. **371** (2002) 231 [arXiv:hep-th/0206063].
- [25] Y. b. Kim and K. M. Lee, “First and second order vortex dynamics,” Phys. Rev. D **66**, 045016 (2002) [arXiv:hep-th/0204111].
- [26] M. Cvetič, G.W. Gibbons, H. Lu and C.N. Pope, “Hyper-Kähler Calabi metrics, L^2 harmonic forms, resolved M2-branes, and AdS_4/CFT_3 correspondence,” Nucl. Phys. B **617** (2001) 151 [arXiv:hep-th/0102185].
- [27] K.M. Lee, D. Tong and S. Yi, “The moduli space of two $U(1)$ instantons on noncommutative R^4 and $R^3 \times S^1$,” Phys. Rev. D **63** (2001) 065017 [arXiv:hep-th/0008092].
- [28] D. Karabali and V.P. Nair, “Quantum Hall effect in higher dimensions,” Nucl. Phys. B **641** (2002) 533 [arXiv:hep-th/0203264].
- [29] D. Karabali and V.P. Nair, “Quantum Hall effect in higher dimensions, matrix models and fuzzy geometry,” J. Phys. A **39** (2006) 12735 [arXiv:hep-th/0606161].