

Universal star products

Mourad Ammar^{1,2}

¹ *Université du Luxembourg*
avenue de la faïencerie
mourad.ammar@uni.lu

Véronique Chloup²

² *Université Paul Verlaine - Metz, LMAM*
Ile du Saulcy, F-57045 Metz Cedex 01, France
chloup@poncelet.univ-metz.fr

Simone Gutt^{3,2}

Membre de l'Académie Royale de Belgique
³ *Université Libre de Bruxelles, Campus Plaine CP 218,*
Bvd du Triomphe, B-1050 Brussels, Belgium
sgutt@ulb.ac.be, gutt@poncelet.univ-metz.fr

Abstract

One defines the notion of universal deformation quantization: given any manifold M , any Poisson structure Λ on M and any torsionfree linear connection ∇ on M , a universal deformation quantization associates to this data a star product on (M, Λ) given by a series of bidifferential operators whose corresponding tensors are given by universal polynomial expressions in the Poisson tensor Λ , the curvature tensor R and their covariant iterated derivatives. Such universal deformation quantization exist. We study their unicity at order 3 in the deformation parameter, computing the appropriate universal Poisson cohomology.

Mathematics Subject Classifications (2000): 53D55, 81S10

Keywords: star products, formality, universal differential operators.

1 Introduction

Let (M, Λ) be a d -dimensional Poisson manifold. Let $C^\infty(M)$ be the commutative algebra of \mathbb{K} -valued smooth function on M , where \mathbb{K} is \mathbb{R} or \mathbb{C} . The Poisson bracket of functions on (M, Λ) is denoted by $\{f, g\} := \Lambda(df, dg)$, $f, g \in C^\infty(M)$, and $m(f, g) = f \cdot g$ denotes the pointwise multiplication.

Definition 1 [1] Let $C^\infty(M)[[\mathbf{v}]]$ be the space of formal power series in the formal parameter \mathbf{v} with coefficients in $C^\infty(M)$. A **differential star product** on (M, Λ) is a bilinear map

$$*: C^\infty(M) \times C^\infty(M) \rightarrow C^\infty(M)[[\mathbf{v}]] \quad (f, g) \rightarrow f * g := f \cdot g + \sum_{r \geq 1} \mathbf{v}^r C_r(f, g),$$

such that

1. its $\mathbb{K}[[\mathbf{v}]]$ -bilinear extension is an associative product $(f * g) * h = f * (g * h)$;
2. $C_1(f, g) - C_1(g, f) = 2\{f, g\}$;
3. each C_r is a bidifferential operator vanishing on constants.

Kontsevich gave in [10] an explicit formula for a star product on \mathbb{R}^d endowed with any Poisson structure, as a special case of his formality theorem for the Hochschild complex of multidifferential operators.

Theorem 1 [10] *The formality theorem on \mathbb{R}^d is proven through the explicit formula of an L_∞ morphism from the differential graded Lie algebra of polyvectorfields on \mathbb{R}^d to the differential graded Lie algebra of polydifferential operators on \mathbb{R}^d . This consists in a collection of multilinear graded symmetric maps U_j associating to j multivectorfields $\alpha_k \in \Gamma(\mathbb{R}^d, \wedge^{m_k} T\mathbb{R}^d)$ a multidifferential operator*

$$U_j(\alpha_1, \dots, \alpha_j) : C^\infty(\mathbb{R}^d)^{\otimes r} \rightarrow C^\infty(\mathbb{R}^d)$$

with $r = \sum_k m_k - 2j + 2$, satisfying quadratic relations $(*)$ which translate the fact that they are the Taylor coefficients of a L_∞ morphism. The maps U_j are defined in terms of graphs; in particular the coefficients of the multidifferential operator $U_j(\alpha_1, \dots, \alpha_j)$ are given by multilinear universal expressions in the partial derivatives of the coefficients of the multivectorfields $\alpha_1, \dots, \alpha_j$.

The relations $(*)$ imply that, given any Poisson structure Λ on \mathbb{R}^d , the formula

$$f *_K^\Lambda g = f g + \sum_{n=1}^{\infty} \frac{\mathbf{v}^n}{n!} U_n(\Lambda, \dots, \Lambda)(f, g) = f g + \mathbf{v} \Lambda(df, dg) + O(\mathbf{v}^2) \quad (1.1)$$

defines a star product on (\mathbb{R}^d, Λ) . Observe that each $U_n(\Lambda, \dots, \Lambda)$ is a bidifferential operator of order maximum n in each argument whose coefficients are polynomials of degree n in the partial derivatives of the coefficients of the tensor Λ .

Kontsevich then obtained the existence of star products on a general Poisson manifold using abstract arguments.

A more direct construction of a star product on a d -dimensional Poisson manifold (M, P) , using Kontsevich's formality on \mathbb{R}^d , was given by Cattaneo, Felder and Tomassini in [3]. Given a torsion-free connexion ∇ on (M, P) one builds an identification of the commutative algebra $C^\infty(M)$ of smooth functions on M with the algebra of flat sections of the jet bundle $E \rightarrow M$, for the Grothendieck connection D^G . The next point is to "quantize" this situation: a deformed algebra structure on $\Gamma(M, E)[[\mathbf{v}]]$ is obtained through fiberwise quantization of the jet bundle using Kontsevich star product on \mathbb{R}^d , and

a deformed flat connection D which is a derivation of this deformed algebra structure is constructed “à la Fedosov”. Then one constructs an identification between the formal series of functions on M and the algebra of flat sections of this quantized bundle of algebras; this identification defines the star product on M .

Later, Dolgushev [7] gave in a similar spirit a construction for a Kontsevich’s formality quasi-isomorphism for a general smooth manifold. The construction starts again with a torsionfree linear connexion ∇ on M and the identification of the commutative algebra $C^\infty(M)$ of smooth functions on M with the algebra of flat sections of the jet bundle $E \rightarrow M$, for a connection D^F constructed “à la Fedosov”. This is extended to a resolution of the space $T_{poly}(M)$ of polyvectors on the manifold using the complexes of forms on M with values in the bundle of formal fiberwise polyvectorfields on E and a resolution of the space $D_{poly}(M)$ of polydifferential operators on the manifold using the complexes of forms on M with values in the bundle of formal fiberwise polydifferential operators on E . The fiberwise Kontsevich L_∞ morphism is then twisted and contracted to yield a L_∞ -morphism from $T_{poly}(M)$ to $D_{poly}(M)$.

Given a torsionfree linear connection ∇ on a manifold M , any multidifferential operator $\text{Op} : (C^\infty(M))^k \rightarrow C^\infty(M)$ writes in a unique way as

$$\text{Op}(f_1, \dots, f_k) = \sum_{J_1, \dots, J_k} \text{Op}^{J_1, \dots, J_k} \nabla_{J_1}^{sym} f_1 \dots \nabla_{J_k}^{sym} f_k \quad (1.2)$$

where the J_1, \dots, J_k are multiindices and $\nabla_J^{sym} f$ is the symmetrised covariant derivative of order $|J|$ of f :

$$\nabla_J^{sym} f = \sum_{\sigma \in S_m} \frac{1}{m!} \nabla_{i_{\sigma(1)} \dots i_{\sigma(m)}}^m f \quad \text{for } J = (i_1, \dots, i_m),$$

where $\nabla_{i_1 \dots i_m}^m f := \nabla^m f(\partial_{i_1}, \dots, \partial_{i_m})$ with $\nabla^m f$ defined inductively by $\nabla f := df$ and $\nabla^m f(X_1, \dots, X_m) = (\nabla_{X_1}(\nabla^{m-1} f))(X_2, \dots, X_m)$.

The tensors $\text{Op}^{J_1, \dots, J_k}$ are covariant tensors of order $|J_1| + \dots + |J_k|$ which are symmetric within each block of J_r indices; they are called **the tensors associated** to Op for the given connection.

Definition 2 A **universal Poisson-related multidifferential operator** will be the association to any manifold M , any torsionfree connection ∇ on M and any Poisson tensor P on M , of a multidifferential operator $\text{Op}^{(M, \nabla, \Lambda)} : (C^\infty(M))^k \rightarrow C^\infty(M)$, so that, the tensors associated to $\text{Op}^{(M, \nabla, \Lambda)}$ for ∇ are given by universal polynomials in Λ , the curvature tensor R and their covariant multiderivatives, involving concatenations.

We shall say that a universal Poisson-related multidifferential operator is **of no-loop type** if the concatenations only arise between different terms, not within a given term (i.e. $(\nabla_r \Lambda)^{is} (\nabla_s \Lambda)^{jr} \nabla_{ij}^{sym}$ is of no-loop type but $(\nabla_i \Lambda)^{ti} \nabla_i^{sym}$ or $R_{str}^r \Lambda^{si} \Lambda^{tj} \nabla_{ij}^{sym}$ are not).

We shall say that the universal Poisson-related operator $\text{Op}^{(M, \nabla, \Lambda)}$ is a **polynomial of degree r in the Poisson structure** if $\text{Op}^{(M, \nabla, t\Lambda)} = t^r \text{Op}^{(M, \nabla, \Lambda)}$.

Definition 3 A **universal star product** $* = m + \sum_{r \geq 1} v^r C_r$ will be the association to any manifold M , any torsionfree connection ∇ on M and any Poisson tensor P on M , of a differential star product $*^{(M, \nabla, \Lambda)} := m + \sum_{r \geq 1} v^r C_r^{(M, \nabla, \Lambda)}$ where each C_r is a universal Poisson-related bidifferential operator of no-loop type, which is a polynomial of degree r in the Poisson structure.

An example of a universal star product at order 3 is given in section 2.

Unicity at order 3 is studied in section 3 using universal Poisson cohomology. This, we compute for universal Poisson-related bidifferential operators of order 1 in each argument defined by low order

polynomials in the Poisson structure.

The existence of a universal star product is implied either by the globalisation proof of Cattaneo, Felder and Tomassini [3, 4] using the exponential map of a torsionfree linear connection, either by the globalisation of the formality given by Dolgushev [7]. We show this existence in section 5, stressing first the relations between the resolutions involved in the two constructions in section 4.

2 An example at order 3

Theorem 2 *There exists a universal star product up to order three, which associates to a Poisson manifold (M, Λ) and a torsionfree linear connection ∇ on M , the star product at order three defined by*

$$f \tilde{*}_3^{(M, \nabla, \Lambda)} g = f \cdot g + \nu \{f, g\} + \nu^2 \tilde{C}_2^{(M, \nabla, \Lambda)}(f, g) + \nu^3 \tilde{C}_3^{(M, \nabla, \Lambda)}(f, g), \quad f, g \in C^\infty(M) \quad (2.1)$$

for

$$\tilde{C}_2^{(M, \nabla, \Lambda)}(f, g) = \frac{1}{2} \Lambda^{kr} \Lambda^{ls} \nabla_{kl}^2 f \nabla_{rs}^2 g + \frac{1}{3} \Lambda^{kr} \nabla_r \Lambda^{ls} (\nabla_{kl}^2 f \nabla_s g + \nabla_s f \nabla_{kl}^2 g) + \frac{1}{6} \nabla_l \Lambda^{kr} \nabla_k \Lambda^{ls} \nabla_r f \nabla_s g, \quad (2.2)$$

and

$$\tilde{C}_3^{(M, \nabla, \Lambda)}(f, g) = \frac{1}{6} S_{\nabla}^{(M, \Lambda) 3}(f, g) = -\frac{1}{6} \Lambda^{ls} (\mathcal{L}_{X_f} \nabla)_{kl}^j (\mathcal{L}_{X_g} \nabla)_{js}^k \quad \text{with } X_f = i(df)\Lambda, \quad (2.3)$$

where $\mathcal{L}_{X_f} \nabla$ is the tensor defined by the Lie derivative of the connection ∇ in the direction of the Hamiltonian vector field X_f

$$(\mathcal{L}_{X_f} \nabla)_{kl}^j = \Lambda^{ij} \nabla_{ki}^3 f + \nabla_k \Lambda^{ij} \nabla_{li}^2 f + \nabla_l \Lambda^{ij} \nabla_{ki}^2 f + \nabla_{kl}^2 \Lambda^{ij} \nabla_i f + R_{ikl}^j \Lambda^{si} \nabla_s f.$$

This can be seen by direct computation.

Remark 1 The operator $S_{\nabla}^{(M, \Lambda) 3}$ was introduced by Flato, Lichnerowicz and Sternheimer [9]; it is a Chevalley-cocycle on (M, P) , i.e.

$$\bigoplus_{u, v, w} \left\{ S_{\nabla}^{(M, \Lambda) 3}(u, v), w \right\} + S_{\nabla}^{(M, \Lambda) 3}(\{u, v\}, w) = 0,$$

where $\bigoplus_{u, v, w}$ denotes the sum over cyclic permutations of u, v, w .

For this universal star product at order 3, there exists a universal Poisson-related-differential-operator-valued 1-form D defined as follows:

Proposition 1 *Given any Poisson manifold (M, Λ) , any torsionfree linear connection ∇ on M , and any vector field X on M , the differential operator $D_X^{(M, \nabla, \Lambda)}$ defined by*

$$D_X^{(M, \nabla, \Lambda)} g = Xg - \nu^2 \frac{1}{6} \Lambda^{ls} (\mathcal{L}_X \nabla)_{kl}^j (\mathcal{L}_{X_g} \nabla)_{js}^k, \quad g \in C^\infty(M)$$

verifies at order 3 in ν

$$D_X^{(M, \nabla, \Lambda)}(f \tilde{*}_3^{(M, \nabla, \Lambda)} g) - (D_X^{(M, \nabla, \Lambda)} f) \tilde{*}_3^{(M, \nabla, \Lambda)} g - f \tilde{*}_3^{(M, \nabla, \Lambda)} (D_X^{(M, \nabla, \Lambda)} g) = \frac{d}{dt} \Big|_{t=0} f \tilde{*}_3^{(M, \nabla, \phi_{tX}^* \Lambda)} g + O(\nu^4).$$

where ϕ_t^X denotes the flow of the vectorfield X .

If X is a Hamiltonian vector field corresponding to a function $f \in C^\infty(M)$, then $D_{X_f}^{(M,\nabla,\Lambda)}$ coincides with the inner derivation at order 3 of $\tilde{\ast}_3$ defined by the function f , i.e.

$$D_{X_f}^{(M,\nabla,\Lambda)} g = \frac{1}{2\nu} (f \tilde{\ast}_3^{(M,\nabla,\Lambda)} g - g \tilde{\ast}_3^{(M,\nabla,\Lambda)} f).$$

3 Equivalence of universal star products – Universal Poisson cohomology

Lemma 1 • Any universal star product $\ast = m + \sum_{r \geq 1} \nu^r C_r$ is a natural star product, i.e. each bidifferential operator C_r is of order at most r in each argument. Indeed C_r is a universal Poisson-related bidifferential operator defined by a polynomial of degree r in the Poisson structure; this implies, in view of the Bianchi's identities for the curvature tensor, that C_r is of order at most r in each argument.

- The universal Poisson-related bidifferential operator C_1 of any universal star product is necessarily the Poisson bracket $C_1^{(M,\nabla,\Lambda)} = \Lambda^{ij} \nabla_i \wedge \nabla_j$.
- The Gerstenhaber bracket $[\cdot, \cdot]_G$ of two universal Poisson-related multidifferential operator of degree k and l in Λ , is a universal Poisson-related multidifferential operator of degree $k+l$ in Λ .
- If a universal Poisson-related p -differential operator C is a Hochschild p -cocycle (where $\partial := \text{ad } m = [m, \cdot]_G$ denotes the Hochschild differential) then $C = A + \partial B$ where A a universal Poisson-related p -differential operator which is of order 1 in each argument and is the totally skewsymmetric part of C , and where B is a universal Poisson-related $(p-1)$ -differential operator.

The last point comes from the explicit formulas [2, 5] for the tensors associated to B in terms of those associated to C when one is given a connection.

Definition 4 A **universal Poisson p -cocycle** is a universal Poisson-related p -differential skewsymmetric operator C of order 1 in each argument which is a cocycle for the Chevalley cohomology for the adjoint representation of $(C^\infty(M), \{, \})$, i.e. with the coboundary defined by

$$\begin{aligned} \delta_P C(u_1, \dots, u_{m+1}) &= \sum_{i=1}^{m+1} (-1)^i \{u_i, C(u_1, \dots, \hat{u}_i, \dots, u_{m+1})\} \\ &\quad + \sum_{i < j} (-1)^{i+j} C(\{u_i, u_j\}, u_1 \dots \hat{u}_i \dots \hat{u}_j \dots, u_{m+1}). \end{aligned}$$

which can be written as a multiple of

$$\text{skew}[\Lambda, C^{(M,\nabla,\Lambda)}]_G$$

where *skew* indicates the skewsymmetrisation in all its arguments of an operator.

Equivalently, a universal Poisson p -cocycle C is defined by a universal Poisson related skewsymmetric p -tensor c (with $C(u_1, \dots, u_p) = c(du_1, \dots, du_p)$) so that

$$[\Lambda, c^{(M,\nabla,\Lambda)}]_{SN} = 0$$

where $[\cdot, \cdot]_{SN}$ denotes the Schouten-Nijenhuis bracket of skewsymmetric tensors (which is the extension, as a graded derivation for the exterior product \wedge of the usual bracket of vectorfields).

A universal Poisson p -cocycle C is a **universal Poisson coboundary** if there exists a universal Poisson-related skewsymmetric $(p-1)$ -differential operator C of order 1 in each argument so that

$$C^{(M, \nabla, \Lambda)} = \delta_p B^{(M, \nabla, \Lambda)} (= skew[\Lambda, B^{(M, \nabla, \Lambda)}]_G);$$

(equivalently, if there exists a universal Poisson related tensor b so that $c^{(M, \nabla, \Lambda)} = [\Lambda, b^{(M, \nabla, \Lambda)}]_{SN}$).

The **universal Poisson cohomology** H^p is the quotient of the space of universal Poisson p -cocycles by the space of universal Poisson p -coboundaries.

We can restrict ourselves to the space of universal Poisson p -cocycles defined by polynomials of degree k in the Poisson structures and make the quotient by the space of universal Poisson coboundaries defined by polynomials of degree $k-1$. We speak then of the universal Poisson p -cohomology of degree k in the Poisson structure and we denote it by H_{polk}^p . We can further restrict ourselves to universal Poisson related tensors (or operators of order 1 in each argument) of no-loop type.

Definition 5 If $* = m + \sum_{r \geq 1} v^r C_r$ is a universal star product and if $E = \sum_{r=2}^{\infty} v E_r$ is a formal series of universal differential operators vanishing on constants, of no-loop type, with each E_r a polynomial of degree r in the Poisson structure, then the series $*'$ defined by

$$*' = (\exp \text{ad} E) *$$

where $\text{ad} E \cdot = [E, \cdot]_G$, (i.e. $f *' g = \exp E((\exp -E)f * (\exp -E)g)$), is an equivalent universal star product. We say that $*$ and $*'$ are universally equivalent.

Lemma 2 If $*$ and $*'$ are universal star products which coincide at order k in the deformation parameter v , then, by the associativity relation at order k , $C'_k - C_k$ is a universal Hochschild 2-cocycle of no-loop type which is a polynomial of degree k in the Poisson structure. Furthermore, associativity at order $k+1$ implies that its skewsymmetric part p_2 is a universal Poisson 2-cocycle :

$$\bigoplus_{u,v,w} \left\{ p_2^{M, \nabla, \Lambda}(u, v), w \right\} + p_2^{M, \nabla, \Lambda}(\{u, v\}, w) = 0,$$

where $\bigoplus_{u,v,w}$ denotes the sum over cyclic permutations of u, v, w .

If it is a universal Poisson 2-coboundary of no-loop type, then there is a formal series E of universal differential operators vanishing on constants such that $(\exp \text{ad} E) *$ and $*'$ coincide at order $k+1$.

In particular, two universal star products are universally equivalent if $H_{(no-loop, pol)}^2 = \{0\}$. They are always equivalent at order k in the deformation parameter v if $H_{(no-loop) pol j}^2 = \{0\} \forall 1 \leq j \leq k$.

Consider now any universal star product $* = m + \sum_{r \geq 1} v^r C_r$. We automatically have that C_1 is the Poisson bracket. Associativity at order 2 yields $\partial C_2 = \partial \tilde{C}_2^{(M, \nabla, \Lambda)}$ so

$$C_2(f, g) = \tilde{C}_2^{(M, \nabla, \Lambda)}(f, g) + p_2(f, g) + \partial E_2(f, g)$$

and the skewsymmetric part of associativity at order 3 yields that p_2 is a universal Poisson 2-cocycle (which is a polynomial of degree 2 in the Poisson structure).

Proposition 2 The spaces $H_{pol2}^2(\Lambda)$ and $H_{(no-loop) pol2}^2(\Lambda)$ of universal Poisson 2-cohomology of degree 2 in the Poisson structure vanish.

PROOF The universal skewsymmetric 2-tensors of degree 2 in Λ are combinations of

$$\begin{aligned} & \nabla_s \Lambda^{ir} \nabla_r \Lambda^{js} \nabla_i \wedge \nabla_j, \\ & (\nabla_{rs}^2 \Lambda^{ir} \Lambda^{js} - \nabla_{rs}^2 \Lambda^{jr} \Lambda^{is}) \nabla_i \wedge \nabla_j, \\ & (\Lambda^{ir} \Lambda^{st} R_{rst}^j - \Lambda^{jr} \Lambda^{st} R_{rst}^i) \nabla_i \wedge \nabla_j, \\ & \Lambda^{ri} \Lambda^{sj} R_{rst}^t \nabla_i \wedge \nabla_j. \end{aligned}$$

The only universal cocycles are the multiples of

$$\Lambda^{ri} \Lambda^{sj} R_{rst}^t \nabla_i \wedge \nabla_j$$

and those are the boundaries of the multiples of $\nabla_r \Lambda^{ir} \partial_i$. Remark that there are no cocycles of no-loop type. ■

Thus, universal star product at order 2 are unique modulo equivalence and one can assume that $C_2 = \tilde{C}_2^{(M, \nabla, \Lambda)}$. Then the skewsymmetric part of the Hochschild 2-cocycle $C_3 - \tilde{C}_3^{(M, \nabla, \Lambda)}$ is a universal Poisson 2-cocycle of no-loop type which is a polynomial of degree 3 in the Poisson structure.

Proposition 3 *The space $H_{(no-loop)pol3}^2(\Lambda)$ of universal Poisson 2-cohomology of no-loop type and of degree 3 in the Poisson structure vanishes.*

PROOF We consider all possible universal Poisson 2-cochains of no loop type which are polynomials of degree 3 in the Poisson structure. They are defined by universal skewsymmetric 2-tensors of degree 3 in Λ which are combinations with constant coefficients of the different concatenations (with no loops) of

$$\begin{aligned} & \Lambda \wedge \Lambda \wedge (\nabla^2 R) \dots \quad \Lambda \wedge \Lambda \wedge R \wedge R \dots, \\ & \Lambda \wedge (\nabla \wedge \Lambda) \wedge (\nabla \wedge \Lambda) \wedge R \dots \quad \Lambda \wedge \Lambda \wedge (\nabla \wedge \Lambda) \wedge (\nabla R) \dots \\ & \Lambda \wedge \Lambda \wedge (\nabla^2 \Lambda) \wedge R \dots \quad \Lambda \wedge (\nabla^2 \Lambda) \wedge (\nabla^2 \Lambda) \wedge (\nabla^2 \Lambda) \wedge (\nabla \wedge \Lambda) \wedge (\nabla \wedge \Lambda) \wedge (\nabla \wedge \Lambda) \dots \end{aligned}$$

Using the symmetry properties of R , the Bianchi's identities and the fact that Λ is a Poisson tensor, one is left with a combination with constant coefficients of 49 independant terms.

Universal 2-coboundaries come from the boundaries of universal 1-tensors of degree 2 in Λ ; such 1-tensors are given by combinations with constant coefficients of concatenations of

$$\Lambda \wedge \Lambda \wedge (\nabla R) \dots \quad \Lambda \wedge (\nabla \wedge \Lambda) \wedge R \dots$$

Hence, modulo universal coboundaries, one can assume that the coefficients of 4 of the 49 terms in a universal cochain are zero.

The cohomology that we are looking for is then given by the combinations with constant coefficients of the remaining 45 terms which are cocycles (for all possible choices of manifold, Poisson structure Λ and connection ∇ .)

Let C be a combination of those 45 terms. The cocycle condition is $[\Lambda, C]_{SN} = 0$. We plug in examples of Poisson structures and connections and impose this cocycle condition. This shows that all 45 coefficients must vanish.

It is enough, for instance, to consider the example on \mathbb{R}^4 , with the non vanishing coefficients of the connection defined by

$$\begin{aligned} \Gamma_{12}^1 &= x_1^3, & \Gamma_{14}^1 &= x_4 & \Gamma_{11}^2 &= x_1^2, & \Gamma_{13}^2 &= 1, & \Gamma_{22}^2 &= 1, & \Gamma_{14}^2 &= x_3, & \Gamma_{13}^3 &= -x_4, \\ \Gamma_{33}^3 &= 1, & \Gamma_{44}^3 &= -x_2x_3x_4, & \Gamma_{11}^4 &= 1, & \Gamma_{13}^4 &= 1, & \Gamma_{22}^4 &= x_1, & \Gamma_{44}^4 &= -3. \end{aligned}$$

and the quadratic Poisson structure defined by

$$\Lambda = \sum_{1=i<j=4} x_i x_j \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$$

From this example, one gets that 41 of the 45 coefficients have to vanish. One is left with a combination with constant coefficients of four terms and an example with constant Poisson structure in dimension 7 shows that all those coefficients must vanish. The non vanishing coefficients of this example are:

$$\begin{aligned} \Lambda^{12} &= 1 & \Lambda^{15} &= 1 & \Lambda^{17} &= 1 & \Lambda^{25} &= 1 & \Lambda^{26} &= 2 & \Lambda^{27} &= 2 \\ \Lambda^{34} &= 1 & \Lambda^{37} &= 3 & \Lambda^{46} &= 1 & \Lambda^{47} &= 4 & \Lambda^{56} &= 1 & \Lambda^{57} &= 5 & \Lambda^{67} &= 6. \end{aligned}$$

$$\begin{aligned} \Gamma_{12}^1 &= 1 & \Gamma_{62}^1 &= x_7 & \Gamma_{77}^1 &= -1 & \Gamma_{17}^2 &= x_1 & \Gamma_{13}^3 &= x_6 & \Gamma_{11}^4 &= 1 & \Gamma_{22}^4 &= 1 & \Gamma_{33}^4 &= 2 \\ \Gamma_{77}^4 &= 3 & \Gamma_{12}^5 &= x_1x_5 & \Gamma_{33}^5 &= x_2 & \Gamma_{11}^6 &= 1 & \Gamma_{44}^6 &= x_2 & \Gamma_{44}^6 &= x_5 & \Gamma_{11}^7 &= x_7 & \Gamma_{44}^7 &= x_3 & \Gamma_{17}^7 &= x_1. \end{aligned}$$

■

Corollary 1 *Any universal star product is universally equivalent to one whose expression at order 3 is given by formula (2.1).*

4 Grothendieck- and Dolgushev-resolution of the space of functions

Our purpose in this section is to prove that the Fedosov-resolution of the algebra of smooth functions constructed in Dolgushev [7] coincides with its resolution given by Cattaneo, Felder and Tomassini in [3]. We also give explicitly the identification of smooth functions, tensorfields and differential operators on M with flat sections in the corresponding bundles.

Let M be a d -dimensional manifold and consider the **jet bundle** $E \rightarrow M$ (the bundle of infinite jet of functions) with fibers $\mathbb{R}[[y^1, \dots, y^d]]$ (i.e. formal power series in $y \in \mathbb{R}^d$ with real coefficients) and transition functions induced from the transition functions of the tangent bundle TM . Thus

$$E = F(M) \times_{\text{Gl}(d, \mathbb{R})} \mathbb{R}[[y^1, \dots, y^d]] \quad (4.1)$$

where $F(M)$ is the frame bundle. Remark that E can be seen as the formally completed symmetric algebra of the cotangent bundle T^*M ; a section $s \in \Gamma(M, E)$ can be written in the form

$$s = s(x; y) = \sum_{p=0}^{\infty} s_{i_1 \dots i_p}(x) y^{i_1} \dots y^{i_p}$$

with repeated indices varying from 1 to d , and where the $s_{i_1 \dots i_p}$ are components of symmetric covariant tensors on M . This bundle E is denoted $\mathcal{S}M$ by Dolgushev.

The construction of a star product on a d -dimensional Poisson manifold (M, P) given by Cattaneo, Felder and Tomassini in [3], using a linear torsionfree connection ∇ on the manifold M , starts

with the identification of the commutative algebra $C^\infty(M)$ of smooth functions on M with the algebra $\mathcal{L}^0(\Gamma(M, E), D^G)$ of flat sections of the jet bundle $E \rightarrow M$, for the Grothendieck connection D^G (which is constructed using ∇). Let us recall this construction.

The exponential map for the connection ∇ gives an identification

$$\exp_x : U \cap T_x M \rightarrow M \quad y \mapsto \exp_x(y) \quad (4.2)$$

at each point x , of the intersection of the tangent space $T_x M$ with a neighborhood U of the zero section of the tangent bundle TM with a neighborhood of x in M .

To a function $f \in C^\infty(M)$, one associates the section f_ϕ of the jet bundle $E \rightarrow M$ given, for any $x \in M$ by the Taylor expansion at $0 \in T_x M$ of the pullback $f \circ \exp_x$.

Lemma 3 *The section f_ϕ is given by:*

$$f_\phi(x; y) = f(x) + \sum_{n>0} \frac{1}{n!} \nabla_{i_1 \dots i_n}^n f(x) y^{i_1} \dots y^{i_n} = f(x) + \sum_{n>0} \frac{1}{n!} \nabla_{i_1 \dots i_n}^{n, \text{sym}} f(x) y^{i_1} \dots y^{i_n}. \quad (4.3)$$

PROOF In local coordinates x^i 's one has $\frac{d}{dt} f(\exp_x ty) = \sum_{k=1}^d (\partial_{x^k} f)(\exp_x ty) \frac{d}{dt} (\exp_x ty)^k$ and $\frac{d^2}{dt^2} f(\exp_x ty) = \sum_{k,l} (\partial_{x^k x^l}^2 f)(\exp_x ty) \frac{d}{dt} (\exp_x ty)^k \frac{d}{dt} (\exp_x ty)^l + \sum_{k=1}^d (\partial_{x^k} f)(\exp_x ty) \frac{d^2}{dt^2} (\exp_x ty)^k$. The definition of the exponential map imply that

$$\frac{d^2}{dt^2} (\exp_x ty)^k = - \sum_{r,s} \Gamma_{rs}^k(\exp_x ty) \frac{d}{dt} (\exp_x ty)^r \frac{d}{dt} (\exp_x ty)^s \quad (4.4)$$

hence $\frac{d^2}{dt^2} f(\exp_x ty) = \sum_{k,l} (\nabla_{kl}^2 f)(\exp_x ty) \frac{d}{dt} (\exp_x ty)^k \frac{d}{dt} (\exp_x ty)^l$. By induction, one gets

$$\frac{d^n}{dt^n} f(\exp_x ty) = \sum_{k_1, \dots, k_n} (\nabla_{k_1 \dots k_n}^n f)(\exp_x ty) \frac{d}{dt} (\exp_x ty)^{k_1} \dots \frac{d}{dt} (\exp_x ty)^{k_n}$$

and the result follows at $t = 0$. ■

Definition 6 [4] The Grothendieck connection D^G on E is defined by:

$$D_X^G s(x; y) := \frac{d}{dt} \Big|_{t=0} s(x(t); \exp_{x(t)}^{-1}(\exp_x(y))) \quad (4.5)$$

for any curve $t \rightarrow x(t) \in M$ representing $X \in T_x M$ and for any $s \in \Gamma(M, E)$. It is locally given by

$$D_X^G = \sum_{i=1}^d X^i \left(\partial_{x^i} + \sum_k \sum_j \left(\frac{\partial \phi_x^{-1}}{\partial y^j} \right)^k \frac{\partial \phi_x^j}{\partial x^i} \partial_{y^k} \right) \quad (4.6)$$

where $\phi_x(y) = \phi(x, y)$ is the Taylor expansion of $\exp_x y$ at $y = 0$:

$$\phi(x, y)^k = x^k + y^k - \frac{1}{2} \sum_{rs} \Gamma_{rs}^k(x) y^r y^s + \frac{1}{3!} \sum_{rst} \left(-(\partial_{x^r} \Gamma_{st}^k)(x) + 2 \sum_u \Gamma_{rs}^u(x) \Gamma_{ut}^k(x) \right) y^r y^s y^t + O(y^4).$$

Remark 2 • From the definition (4.5) it is clear that D^G is flat ($D_X^G \circ D_Y^G - D_Y^G \circ D_X^G = D_{[X,Y]}^G$).

• It is also obvious that $D^G(f_\phi) = 0 \forall f \in C^\infty(M)$.

Lemma 4 [4] *Introducing the operator on E -valued forms on M*

$$\delta = \sum_i dx^i \frac{\partial}{\partial y^i}, \quad (4.7)$$

one can write

$$D^G = -\delta + \nabla' + A, \quad (4.8)$$

where

$$\nabla' = \sum_i dx^i \left(\partial_{x^i} - \sum_{jk} \Gamma_{ij}^k y^j \partial_{y^k} \right) \quad (4.9)$$

is the covariant derivative on E associated to ∇ and where A is a 1-form on M with values in the fiberwise vectorfields on E ,

$$A(x;y) =: \sum_{ik} dx^i A_i^k(x;y) \partial_{y^k} = \sum_{ik} dx^i \left(-\frac{1}{3} \sum_{rs} R_{ris}^k(x) y^r y^s + 0(y^3) \right) \partial_{y^k}. \quad (4.10)$$

One extends as usual the operator D^G to the space $\Omega(M, E)$ of E -valued forms on M :

$$D^G = -\delta + \nabla' + A \quad \text{with} \quad \nabla' = d - \sum_{ijk} dx^i \Gamma_{ij}^k y^j \partial_{y^k}. \quad (4.11)$$

One introduces the operator $\delta^* = \sum_j y^j i(\frac{\partial}{\partial x^j})$ on $\Omega(M, E)$. Clearly $(\delta^*)^2 = 0, \delta^2 = 0$ and for any $\omega \in \Omega^q(M, E_p)$, i.e. a q -form of degree p in y , we have $(\delta \delta^* + \delta^* \delta) \omega = (p+q) \omega$.

Defining, for any $\omega \in \Omega^q(M, E_p)$

$$\begin{aligned} \delta^{-1} \omega &= \frac{1}{p+q} \delta^* \omega && \text{when } p+q \neq 0 \\ &= 0 && \text{when } p=q=0 \end{aligned}$$

we see that any δ -closed q -form ω of degree p in y , when $p+q > 0$, writes uniquely as $\omega = \delta \sigma$ with $\delta^* \sigma = 0$; σ is given by $\sigma = \delta^{-1} \omega$.

One proceeds by induction on the degree in y to see that the cohomology of D^G is concentrated in degree 0 and that any flat section of E is determined by its part of degree 0 in y . Indeed a q -form ω is D^G -closed iff $\delta \omega = (\nabla' + A) \omega$; this implies that $\delta \omega_p = 0$ for ω_p the terms of lowest order (p) in y . When $p+q > 0$ we can write $\omega_p = \delta(\delta^{-1} \omega_p)$ and $\omega - D^G(\delta^{-1} \omega_p)$ has terms of lowest order at least $p+1$ in y . Remark that given any section s of E then $s(x; y=0)$ determines a smooth function f on M . If $D^G s = 0$, then $s - f_\phi$ is still D^G closed. By the above, Its terms of lowest order in y must be of the form $\delta \sigma$ hence must vanish since we have a 0-form. Hence we have:

Lemma 5 [4] *Any section of the jet bundle $s \in \Gamma(E)$ is the Taylor expansion of the pullback of a smooth function f on M via the exponential map of the connection ∇ if and only if it is horizontal for the Grothendieck-connection D^G :*

$$s = f_\phi \text{ for a } f \in C^\infty(M) \Leftrightarrow s \in \Gamma_{hor}(E) := \{s' \in \Gamma(E) \mid D^G s' = 0\}. \quad (4.12)$$

Furthermore, the cohomology of D^G is concentrated in degree 0. In other word, one obtains a ‘‘Grothendieck-resolution’’ of the algebra of smooth functions, i.e.

$$H^\bullet(\Omega(M, E), D^G) = H^0(\Omega(M, E), D^G) = \Gamma_{hor}(E) \cong C^\infty(M).$$

Remark 3 When a q -form ω is D^G exact, we have written $\omega = D^G\sigma$ where the tensors defining σ are given by universal polynomials (with no-loop concatenations) in the tensors defining ω , the tensors defining A , the curvature of the connection, and their iterated covariant derivatives.

Lemma 6 *The 1-form A on M with values in the fiberwise vectorfields on E is given by $A(x;y) =: \sum_{ik} dx^i A_i^k(x;y) \partial_{y^k}$ where the A_i^k are universal polynomials given by (no-loop) concatenations of iterative covariant derivatives of the curvature; they are of the form*

$$\sum (\nabla \dots R)_{i \dots}^{j_1} (\nabla \dots R)_{j_1 \dots}^{j_2} \dots (\nabla \dots R)_{j_{s-1} \dots}^{j_s} y^1 \dots y^s. \quad (4.13)$$

In particular $\delta^{-1}A = 0$ since the curvature is skewsymmetric in its first two lower arguments. The 1-form A is uniquely characterized by the fact that $\delta^{-1}A = 0$. and the fact that $D^G = -\delta + \nabla' + A$ is flat, i.e.; $(D^G)^2 = 0$ which is equivalent to

$$\delta A = R^{\nabla'} + \nabla' A + \frac{1}{2}[A, A] \quad (4.14)$$

for $\frac{1}{2}[A, A](X, Y) := [A(X), A(Y)]$ and $R^{\nabla'} = -\frac{1}{2}R_{ijk}^l dx^i \wedge dx^j y^k \frac{\partial}{\partial y^l}$.

PROOF Any section $s \in \Gamma(M, E)$ writes $\sum_{p=0}^{\infty} s_{i_1 \dots i_p}^p(x) y^{i_1} \dots y^{i_p}$ with symmetric p -covariant tensors $s_{i_1 \dots i_p}^p$. Write $A(x;y) = \sum_{r \geq 2} dx^i (A^{(r)}(x))_{i, j_1 \dots j_r}^k y^{j_1} \dots y^{j_r} \partial_{y^k}$ with $(A^{(2)}(x))_{i, rs}^k = -\frac{1}{3} \sum_{rs} R_{ris}^k(x) y^r y^s$. Then the covariant tensors of $D_X^G s$ are given by the symmetrisation of

$$(D_X^G s)^p = -i(X)s^{p+1} + \nabla_X s^p + \sum_{r=0}^{p-2} (A^{(p-r)}(X))^k \partial_{y^k} s^{r+1}.$$

The fact that $D^G(f_\phi) = 0 \forall f \in C^\infty(M)$ implies the expression given in the lemma for A . Indeed, the symmetric tensors defining f_ϕ are given by $\frac{1}{p!} \nabla^{p, \text{sym}} f$ and we must have

$$\begin{aligned} 0 &= \left(D_{\partial_{y^i}}^G f_\phi \right)_{j_1 \dots j_p}^p = -\frac{1}{(p+1)!} (\nabla^{p+1, \text{sym}} f)_{ij_1 \dots j_p} + \frac{1}{p!} (\nabla (\nabla^{p, \text{sym}} f))_{ij_1 \dots j_p} \\ &\quad + \sum_{r=0}^{p-2} \left(A^{(p-r)}(x) \right)_{i, j_1 \dots j_{p-r}}^k \frac{1}{r!} (\nabla^{r+1, \text{sym}} f)_{kj_{p-r+1} \dots j_p} \end{aligned}$$

with the last terms symmetrized in the j 's. The commutation of covariant derivatives of a q -form ω gives

$$(\nabla^{p+2} f)_{klj_1 \dots j_p} - (\nabla^{p+2} f)_{lkj_1 \dots j_p} = -\sum_{r=0}^p R_{klj_r}^s (\nabla^p f)_{j_1 \dots j_{r-1} s j_{r+1} \dots j_p}$$

and implies by induction that $(\nabla (\nabla^{p, \text{sym}} f))_{ij_1 \dots j_p} - (\nabla^{p+1, \text{sym}} f)_{ij_1 \dots j_p}$ is a universal expression contracting covariant derivatives of the curvature tensor with lower covariant derivatives of f of the form

$$(\nabla \dots R)_{i \dots}^{t_1} (\nabla \dots R)_{t_1 \dots}^{t_2} \dots (\nabla \dots R)_{t_{s-1} \dots}^s (\nabla^{r+1, \text{sym}} f)_{s \dots}$$

with the j 's put in a symmetrised way at the \cdot 's, and for $0 \leq r \leq p-2$. Hence the expression for A .

Observe that $d\delta + \delta d = 0$ and also $\delta \nabla' + \nabla' \delta = 0$ since ∇ is torsionfree. Hence $(D^G)^2 = 0$ iff $-\delta A + R^{\nabla'} + \nabla' A + \frac{1}{2}[A, A]$ vanishes on all sections of E ; since it is a 2-form on M with values in the fiberwise vectorfields on E , this must vanish. ■

Dolgushev [7] gave in a similar spirit a construction for a Kontsevich's formality quasi-isomorphism for a general smooth manifold. The construction starts again with a torsionfree linear connexion ∇ on M . A resolution (called Fedosov's resolution in Dolgushev's paper) of the algebra of functions is given using the complex of algebras $(\Omega(M, E), D_F)$ for a flat connexion (differential) D_F defined by

$$D_F := \nabla' - \delta + A \quad (4.15)$$

where A is a 1-form on M with values in the fiberwise vectorfields on E , obtained by induction on the order in y by the equation

$$A = \delta^{-1} R^{\nabla'} + \delta^{-1} (\nabla' A + \frac{1}{2} [A, A]). \quad (4.16)$$

This implies that $\delta^{-1} A = 0$ and $\delta A = R^{\nabla'} + \nabla' A + \frac{1}{2} [A, A]$ so that A coincides with the 1-form already considered. Hence

Lemma 7 *The differential D^G and D_F coincide.*

Similarly, Dolgushev defined a resolution of polydifferential operators and polyvectorfields on M using the complexes $(\Omega(M, \mathcal{D}_{poly}), D_F^{\mathcal{D}_{poly}})$ and $(\Omega(M, \mathcal{T}_{poly}), D_F^{\mathcal{T}_{poly}})$ where \mathcal{T}_{poly} is the bundle of formal fiberwise polyvectorfields on E and \mathcal{D}_{poly} is the bundle of formal fiberwise polydifferential operators on E . A section of \mathcal{T}_{poly}^k is of the form

$$\mathcal{F}(x; y) = \sum_{n=0}^{\infty} \mathcal{F}_{i_1 \dots i_n}^{j_1 \dots j_{k+1}}(x) y^{i_1} \dots y^{i_n} \frac{\partial}{\partial y^{j_1}} \wedge \dots \wedge \frac{\partial}{\partial y^{j_{k+1}}}, \quad (4.17)$$

where $\mathcal{F}_{i_1 \dots i_n}^{j_1 \dots j_{k+1}}(x)$ are coefficients of tensors, symmetric in the covariant indices i_1, \dots, i_n and anti-symmetric in the contravariant indices j_1, \dots, j_{k+1} . A section of \mathcal{D}_{poly}^k is of the form

$$\mathcal{O}(x; y) = \sum_{n=0}^{\infty} \mathcal{O}_{i_1 \dots i_n}^{\alpha_1 \dots \alpha_{k+1}}(x) y^{i_1} \dots y^{i_n} \frac{\partial^{|\alpha_1|}}{\partial y^{\alpha_1}} \otimes \dots \otimes \frac{\partial^{|\alpha_{k+1}|}}{\partial y^{\alpha_{k+1}}}, \quad (4.18)$$

where the α_l are multi-indices and $\mathcal{O}_{i_1 \dots i_n}^{\alpha_1 \dots \alpha_{k+1}}(x)$ are coefficients of tensors symmetric in the covariant indices i_1, \dots, i_n . and symmetric in each block of α_i contravariant indices.

The spaces $\Omega(M, \mathcal{T}_{poly})$ and $\Omega(M, \mathcal{D}_{poly})$ have a formal fiberwise DGLA structure. Namely, the degree of an element in $\Omega(M, \mathcal{T}_{poly})$ (resp. $\Omega(M, \mathcal{D}_{poly})$) is defined by the sum of the degree of the exterior form and the degree of the polyvector field (resp. the polydifferential operator), the bracket on $\Omega(M, \mathcal{T}_{poly})$ is defined by $[\omega_1 \otimes \mathcal{F}_1, \omega_2 \otimes \mathcal{F}_2]_{SN} := (-1)^{k_1 q_2} \omega_1 \wedge \omega_2 \otimes [\mathcal{F}_1, \mathcal{F}_2]_{SN}$ for ω_i a q_i form and \mathcal{F}_i a section in $\mathcal{T}_{poly}^{k_i}$ and similarly for $\Omega(M, \mathcal{D}_{poly})$ using the Gerstenhaber bracket. The differential on $\Omega(M, \mathcal{T}_{poly})$ is 0 and the differential on $\Omega(M, \mathcal{D}_{poly})$ is defined by $\partial := [m_{pf}, \cdot]_G$ where m_{pf} is the fiberwise multiplication of formal power series in y of E .

Definition 7 [7] The differential $D_F^{\mathcal{T}_{poly}}$ is defined on $\Omega(M, \mathcal{T}_{poly})$ by

$$D_F^{\mathcal{T}_{poly}} \mathcal{F} := \nabla^{\mathcal{T}_{poly}} \mathcal{F} - \delta^{\mathcal{T}_{poly}} \mathcal{F} + [A, \mathcal{F}]_{SN} \quad (4.19)$$

where $\nabla^{\mathcal{T}_{poly}} \mathcal{F} = d\mathcal{F} - \left[\sum_{ijk} dx^i \Gamma_{ij}^k y^j \partial_{y^k}, \mathcal{F} \right]_{SN}$ and where $\delta \mathcal{F} = \left[\sum_i dx^i \frac{\partial}{\partial y^i}, \mathcal{F} \right]_{SN}$. Similarly $D_F^{\mathcal{D}_{poly}}$ is defined on $\Omega(M, \mathcal{D}_{poly})$ by

$$D_F^{\mathcal{D}_{poly}} \mathcal{O} := \nabla^{\mathcal{D}_{poly}} \mathcal{O} - \delta^{\mathcal{D}_{poly}} \mathcal{O} + [A, \mathcal{O}]_G \quad (4.20)$$

with $\nabla^{\mathcal{D}_{poly}}$ and $\delta^{\mathcal{D}_{poly}}$ defined as above with the Gerstenhaber bracket.

Again the cohomology is concentrated in degree 0 and a flat section $\mathcal{F} \in \mathcal{T}_{poly}$ or $\mathcal{O} \in \mathcal{D}_{poly}$ is determined by its terms \mathcal{F}_0 or \mathcal{O}_0 of order 0 in y ; it is defined inductively by

$$\mathcal{F} = \mathcal{F}_0 + \delta^{-1} \left(\nabla^{\mathcal{T}_{poly}} \mathcal{F} + [A, \mathcal{F}]_{SN} \right) \quad \mathcal{O} = \mathcal{O}_0 + \delta^{-1} \left(\nabla^{\mathcal{D}_{poly}} \mathcal{O} + [A, \mathcal{O}]_G \right).$$

On the other hand, if $s_1 \dots s_{k+1}$ are sections of E , we have for a $\mathcal{F} \in \Gamma(M, \mathcal{T}_{poly})$:

$$D_F(\mathcal{F}(s_1, \dots, s_{k+1})) = (D_F^{\mathcal{T}_{poly}} \mathcal{F})(s_1, \dots, s_{k+1}) + \mathcal{F}(D_F s_1, \dots, s_{k+1}) + \dots + \mathcal{F}(s_1, \dots, D_F s_{k+1}) \quad (4.21)$$

and similarly for a $\mathcal{O} \in \Gamma(M, \mathcal{D}_{poly})$.

Definition 8 [3] As in Cattaneo et al. we associate to a polyvector field $F \in T_{poly}^k(M)$ a section $F_\phi \in \Gamma(M, \mathcal{T}_{poly})$: for a point $x \in M$ one considers the Taylor expansion (infinite jet) $F_\phi(x; y)$ at $y = 0$ of the push-forward $(\exp_x)_*^{-1} F(\exp_x y)$. Clearly this definition implies that $X_\phi(f_\phi) = (Xf)_\phi$ so that F_ϕ is uniquely determined by the fact that

$$F_\phi(f_\phi^1, \dots, f_\phi^{k+1}) = \left(F(f^1, \dots, f^{k+1}) \right)_\phi \quad \forall f^j \in C^\infty(M). \quad (4.22)$$

Similarly we associate to a differential operator $O \in D_{poly}^k(M)$ a section $O_\phi \in \Gamma(M, \mathcal{D}_{poly})$ determined by the fact that

$$O_\phi(f_\phi^1, \dots, f_\phi^{k+1}) = \left(O(f^1, \dots, f^{k+1}) \right)_\phi \quad \forall f^j \in C^\infty(M). \quad (4.23)$$

Observe that $D_F^{\mathcal{T}_{poly}} F_\phi = 0$ by 4.21 and 4.22 and similarly $D_F^{\mathcal{D}_{poly}} O_\phi = 0$, hence we have

Proposition 4 A section of \mathcal{T}_{poly} is $D_F^{\mathcal{T}_{poly}}$ -horizontal if and only if is a Taylor expansion of a polyvectorfield on M , i.e. iff it is of the form F_ϕ for some $F \in T_{poly}^k(M)$.

Similarly a section of \mathcal{D}_{poly} is $D_F^{\mathcal{D}_{poly}}$ -horizontal if and only if is of the form O_ϕ for some $O \in D_{poly}^k(M)$. The terms in such a flat section are defined by tensors which are universal polynomials (involving concatenations of no-loop type) in the tensors defining the polyvectorfield (or differential operator) on M , the curvature tensor and their iterated covariant derivatives.

Observe also that a D_F closed section $s \in \Omega^q(M, E)$ or $\mathcal{F} \in \Omega^q(M, \mathcal{T}_{poly})$ or $\mathcal{O} \in \Omega^q(M, \mathcal{D}_{poly})$ for $q \geq 1$ is the boundary of a section defined by tensors which are given by universal polynomials (involving concatenations of no-loop type) in the tensors defining the section, the curvature tensor and their iterated covariant derivatives.

The isomorphisms obtained are isomorphisms of differential graded Lie algebras.

5 Construction of a universal star products

The construction of a star product on any Poisson manifold by Cattaneo, Felder and Tomassini proceeds as follows: one quantize the identification of the commutative algebra of smooth functions on M with the algebra of flat sections of E in the following way.

A deformed algebra structure on $\Gamma(M, E)[[\hbar]]$ is obtained through fiberwise quantization of the jet bundle using Kontsevich star product on \mathbb{R}^d . Precisely, one considers the fiberwise Poisson structure on E defined by Λ_ϕ and the fiberwise Kontsevich star product on $\Gamma(M, E)[[\hbar]]$:

$$\sigma *_K^{\Lambda_\phi} \tau = \sigma \tau + \sum_{n=1}^{\infty} \frac{\hbar^n}{n!} U_n(\Lambda_\phi, \dots, \Lambda_\phi)(\sigma, \tau).$$

The operator D_X^G is not a derivation of this deformed product; one constructs a flat connection D which is a derivation of $*_K^{\Lambda_\phi}$. One defines first

$$D_X^1 = X + \sum_{j=0}^{\infty} \frac{v^j}{j!} U_{j+1}(\hat{X}, \Lambda_\phi, \dots, \Lambda_\phi) \quad (5.1)$$

where $\hat{X} := D_X^G - X$ is a vertical vectorfield on E . The formality equations imply that D_X^1 is a derivation of the star product. Using the fact that $U_1(\xi) = \xi$ for any vector field ξ and that, for $n \geq 2$, the maps $U_n(\xi, \alpha_2, \dots, \alpha_n) = 0$ if ξ is a linear vector field, we see that

$$D_X^1 = D_X^G + \sum_{j=1}^{\infty} \frac{v^j}{j!} U_{j+1}(\hat{X}, \Lambda_\phi, \dots, \Lambda_\phi) \quad (5.2)$$

where $\hat{X} = \sum_i X^i (-\partial_{y^i} + \sum_k A_i^k(x; y) \partial_{y^k})$ as defined in equation 4.10, so that it is given by universal polynomials (with no-loop type concatenations) in the tensors defining X , the curvature tensor and their iterated derivatives. The connection D^1 is not flat so one deforms it by

$$D := D^1 + [\gamma, \cdot]_{*_K^{\Lambda_\phi}}$$

so that D is flat. The 1-form γ is constructed inductively using the fact that the cohomology of D^G vanishes.

The next point is to identify series of functions on M with the algebra of flat sections of this quantized bundle of algebras to define the star product on M .

This is done by building a map $\rho : \Gamma(M, E)[[v]] \rightarrow \Gamma(M, E)[[v]]$ so that $\rho \circ D^G = D \circ \rho$. This map is again constructed by induction using the vanishing of the cohomology.

All these points show that the star product constructed in this way is universal.

Similarly, one can show that Dolgushev's formality is universal in the following sense. Dolgushev constructs a L_∞ morphism from the differential graded Lie algebra of polyvectorfields on M to the differential graded Lie algebra of polydifferential operators on M , giving its Taylor coefficients, i.e. a collection of maps U_j^D associating to j multivectorfields F_k on M a multidifferential operator $U_j^D(\alpha_1, \dots, \alpha_j)$. The tensors defining this operator are given by universal polynomials (involving concatenations of no-loop type) in the tensors defining the α_j 's, the curvature tensor and their iterated covariant derivatives.

Indeed this L_∞ -morphism is obtained in two steps from the fiberwise Kontsevich formality from $\Omega(M, \mathcal{T}_{poly})$ to $\Omega(M, \mathcal{D}_{poly})$ building first a twist which depends only on the curvature and its covariant derivatives, then building a contraction using the vanishing of the D_F cohomology.

References

- [1] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz et D. Sternheimer, Deformation theory and quantization, Ann. Phys. 111 (1978) 61-110.
- [2] M. Cahen, S. Gutt, Produits $*$ sur les espaces affins symplectiques localement symétriques, C.R. Acad. Sc. Paris 297 (1983), série I, 417-420.
- [3] A.S. Cattaneo, G.Felder and L.Tommassini, From local to global deformation quantization of Poisson manifolds, Duke Math. J. 115, 2 (2002) 359-352.

- [4] A.S. Cattaneo, G.Felder, On the globalization of Kontsevich's star product and the perturbative Poisson sigma model, hep-th/0111028.
- [5] Simone Gutt, John Rawnsley, Equivalence of star products on a symplectic manifold: an introduction to Deligne's Cech cohomology classes", Journ. of Geom. and Phys.29 (1999) 347–392.
- [6] Simone Gutt, John Rawnsley, Natural star products on symplectic manifolds and quantum moment maps, Lett. in Math. Phys. 66 (2003) 123–139.
- [7] V.Dolgushev, Covariant and equivariant formality theorem, Adv. Math. 191, 1 (2005) 147–177.
- [8] B.V. Fedosov, a simple geometrical construction of deformation quantization. J. Diff. Geom. 40 (1994) 213–238
- [9] M. Flato, A. Lichnerowicz and D. Sternheimer, Crochet de Moyal-Vey et quantification, C.R. Acad. Sc., t. 283, A (1976), 19
- [10] M. Kontsevich, Deformation quantization of Poisson manifold, Lett. Math. Phys. 66, 3 (2003) 157-216.