

Asymptotic behavior of growth functions of D0L-systems

Julien Cassaigne

Christian Mauduit

François Nicolas

January 22, 2020

Abstract

A *D0L-system* is a triple (A, σ, w) where A is a finite alphabet, σ is an endomorphism of the free monoid over A , and w is a word over A . The *D0L-sequence* generated by (A, σ, w) is the sequence of words $(w, \sigma(w), \sigma(\sigma(w)), \sigma(\sigma(\sigma(w))), \dots)$. The corresponding sequence of lengths, that is the function mapping each integer $n \geq 0$ to $|\sigma^n(w)|$, is called the *growth function* of (A, σ, w) . In 1978, Salomaa and Soittola deduced the following result from their thorough study of the theory of rational power series: if the D0L-sequence generated by (A, σ, w) is not eventually the empty word then there exist an integer $\alpha \geq 0$ and a real number $\beta \geq 1$ such that $|\sigma^n(w)|$ behaves like $n^\alpha \beta^n$ as n tends to infinity. The aim of the present paper is to present a short, direct, elementary proof of this theorem.

1 Introduction

1.1 Notation

As usual, \mathbb{N} , \mathbb{R} and \mathbb{C} denote the semiring of natural integers, the field of real numbers, and the field of complex numbers, respectively. For every $a, b \in \mathbb{N}$, $[a, b]$ denotes the set of all integers n such that $a \leq n \leq b$. Let $f, g : \mathbb{N} \rightarrow \mathbb{C}$. We write $f(n) \preceq g(n)$ if there exists a real number $\lambda > 0$ such that $\{n \in \mathbb{N} : |f(n)| > \lambda |g(n)|\}$ is finite. We write $f(n) \asymp g(n)$ if both $f(n) \preceq g(n)$ and $g(n) \preceq f(n)$ hold.

A *word* is a finite string of symbols. Word concatenation is denoted multiplicatively. For every word w , the *length* of w is denoted $|w|$. The word of length zero is called the *empty word*. For every symbol a and every word w , $|w|_a$ denotes the number of occurrences of a in w .

An *alphabet* is a finite set of symbols. Let A be an alphabet. The set of all words over A is denoted A^* . A mapping $\sigma : A^* \rightarrow A^*$ is called a *morphism* if $\sigma(xy) = \sigma(x)\sigma(y)$ for every $x, y \in A^*$. Clearly, σ is completely determined by its restriction to A . For every $n \in \mathbb{N}$, σ^n denotes the n^{th} iterate of σ : for every $w \in A^*$, $\sigma^0(w) = w$, $\sigma^1(w) = \sigma(w)$, $\sigma^2(w) = \sigma(\sigma(w))$, $\sigma^3(w) = \sigma(\sigma(\sigma(w)))$, etc.

A *D0L-system* [2] is defined as a triple (A, σ, w) where A is an alphabet, σ a morphism from A^* to itself, and w is a non-empty word over A . The *growth function* of the D0L-system (A, σ, w) is defined as the integer sequence $(|w|, |\sigma(w)|, |\sigma^2(w)|, |\sigma^3(w)|, \dots)$. For every *D0L-system* (A, σ, w) , either the sequence $(w, \sigma(w), \sigma^2(w), \sigma^3(w), \dots)$ is eventually periodic or $\lim_{n \rightarrow \infty} |\sigma^n(w)| = \infty$.

1.2 Contribution

The aim of the paper is to present a short, elementary proof of the following theorem.

Theorem 1. *Let (A, σ, w) be a D0L-system such that $\sigma^n(w)$ is a non-empty word for every $n \in \mathbb{N}$. There exist a non-negative integer α smaller than the cardinality of A , and a real number $\beta \geq 1$ such that $|\sigma^n(w)| \asymp n^\alpha \beta^n$.*

Theorem 1 plays a crucial role in the proof of an important result: Pansiot's theorem concerning the complexity of pure morphic sequences [3].

In 1978, Salomaa and Soittola laboriously proved a stronger result than Theorem 1.

Theorem 2 (Salomaa and Soittola [6, 1]). *Let (A, σ, w) be a D0L-system such that $\sigma^n(w)$ is a non-empty word for every $n \in \mathbb{N}$. There exist a positive integer q , a non-negative integer α , and a real number $\beta \geq 1$ such that for each $r \in [0, q - 1]$,*

$$\frac{|\sigma^{nq+r}(w)|}{(nq+r)^\alpha \beta^{nq+r}}$$

converges to a positive, finite limit as $n \rightarrow \infty$.

The proof of Theorem 1 presented below cannot likely be refined into a proof of Theorem 2. The original proof of Theorem 2 relies on the theory of rational power series. In particular, two deep results are put to use:

1. Schützenberger's representation theorem [6, 1], and
2. Berstel's theorem concerning the minimum-modulus poles of univariate rational series over the semiring of non-negative real numbers [6, 1].

To conclude this section note that a very interesting particular case of Theorem 2 can be simply deduced from the Perron-Frobenius theory.

Definition 1 (Irreducibility and period). *Let A be an alphabet and let $\sigma : A^* \rightarrow A^*$ be a morphism. We say that σ is irreducible if for each $(a, b) \in A \times A$, there exists $k \in \mathbb{N}$ such that a occurs in $\sigma^k(b)$. For every $a \in A$, the period of a under σ is defined as the greatest common divisor of $\{k \in \mathbb{N} : |\sigma^k(a)|_a \neq 0\}$.*

If the morphism σ is irreducible then all letters in A have the same period under σ . If σ is irreducible and if every letter in A is of period one under σ then σ is called *primitive*: there exists $N \in \mathbb{N}$ such that for each $(a, b) \in A \times A$, a occurs in $\sigma^N(b)$.

Theorem 3 ([4]). *Let A be an alphabet, let $\sigma : A^* \rightarrow A^*$ be an irreducible morphism, and let q denote the period under σ of any letter in A . There exists a real number $\beta \geq 1$ such that for each $(a, b) \in A \times A$ and each $r \in [0, q - 1]$,*

$$\frac{|\sigma^{nq+r}(a)|_b}{\beta^{nq+r}}$$

converges to a positive, finite limit as $n \rightarrow \infty$.

2 Proof of Theorem 1

Our proof of Theorem 1 relies on the equivalence of norms on a finite-dimensional vector space (see Theorem 4 below). For the sake of completeness, the definition of a norm is recalled.

Definition 2 (Norm). *Let V be a real or complex vector space. A norm on V is a mapping $\|\cdot\|$ from V to \mathbb{R} such that the following three properties hold:*

- *for every $x \in V$, $\|x\| = 0$ if and only if x is the zero vector;*
- *for every $x \in X$ and every $\lambda \in \mathbb{R}$, $\|\lambda x\| = |\lambda| \|x\|$;*
- *for every $x, y \in V$, $\|x + y\| \leq \|x\| + \|y\|$.*

For every integer $d \geq 1$, $\mathbb{C}^{d \times d}$ denotes the algebra of d -by- d complex matrices, and $\mathbb{N}^{d \times d}$ denotes the semiring of d -by- d matrices with non-negative, integer entries; the following norms on $\mathbb{C}^{d \times d}$ play a central role in the section.

Definition 3. *Let d be a positive integer and let $X \in \mathbb{C}^{d \times d}$.*

Define $\|X\|$ as the sum of the magnitudes of the entries of X .

For every non-singular matrix $P \in \mathbb{C}^{d \times d}$, define $\|X\|_P$ as the maximum magnitude of the entries of PXP^{-1} .

Next theorem is well-known: it states that, on a finite-dimensional vector space, all norms define the same topology.

Theorem 4. *Let V be a real or complex vector space. If the dimension of V is finite then for any norms $\|\cdot\|_1$ and $\|\cdot\|_2$ on V , there exist positive real numbers λ and μ such that $\lambda \|x\|_1 \leq \|x\|_2 \leq \mu \|x\|_1$ for every $x \in V$.*

Corollary 1. *Let d be a positive integer. For any norms $\|\cdot\|_1$ and $\|\cdot\|_2$ on $\mathbb{C}^{d \times d}$ and for any $M \in \mathbb{C}^{d \times d}$, $\|M^n\|_1 \asymp \|M^n\|_2$ as $n \rightarrow \infty$.*

Next proposition, which is mainly folklore, is the main ingredient of the proof of Theorem 1.

Proposition 1. *Let d be a positive integer and let $M \in \mathbb{C}^{d \times d}$. There exist an integer $\alpha \in [0, d-1]$ and a non-negative real number β such that for every norm $\|\cdot\|$ on $\mathbb{C}^{d \times d}$, $\|M^n\| \asymp n^\alpha \beta^n$ as $n \rightarrow \infty$.*

Proof. Let $P \in \mathbb{C}^{d \times d}$ be a non-singular matrix such that $PM P^{-1}$ is in Jordan normal form: there exist $D, N \in \mathbb{C}^{d \times d}$ such that D is diagonal, N is nilpotent, $PM P^{-1} = D + N$ and $DN = ND$. By Corollary 1, it suffices to study the behavior of $\|M^n\|_P$ as $n \rightarrow \infty$. For every $n \in \mathbb{N}$, the binomial theorem yields:

$$PM^n P^{-1} = (D + N)^n = \sum_{k=0}^n \binom{n}{k} D^{n-k} N^k.$$

Besides, N^k is a zero matrix for every integer $k \geq d$, and thus

$$PM^n P^{-1} = \sum_{k=0}^{d-1} \binom{n}{k} D^{n-k} N^k$$

for every integer $n \geq d-1$. Hence, for each $(i, j) \in [1, d] \times [1, d]$, there exist an eigenvalue λ_i of D and a complex polynomial $f_{i,j}$ with $\deg f_{i,j} \leq d-1$ such that for every integer $n \geq d$, the $(i, j)^{\text{th}}$ entry of $PM^n P^{-1}$ equals $f_{i,j}(n) \lambda_i^n$. The behavior of $\|M^n\|_P$ as $n \rightarrow \infty$ is now clear: there exist indices $i, j \in [1, d]$ such that $\alpha := \deg f_{i,j}$ and $\beta := |\lambda_i|$ satisfy $\|M^n\|_P \asymp n^\alpha \beta^n$ as $n \rightarrow \infty$. (More precisely, if M is not nilpotent then β is positive and the ratio $\frac{\|M^n\|_P}{n^\alpha \beta^n}$ converges to a positive, finite limit as $n \rightarrow \infty$.) \square

Proposition 1 deserves several comments. First, a more precise result is known.

Theorem 5 ([7]). *Let d be a positive integer, let $\|\cdot\|$ denote the spectral norm on $\mathbb{C}^{d \times d}$, and let $M \in \mathbb{C}^{d \times d}$ be such that M is not nilpotent.*

- *Let β denote the spectral radius of M .*
- *Let j denote the maximum size of the Jordan blocks of M with spectral radius β .*

The ratio $\frac{\|M^n\|}{n^{j-1} \beta^n}$ converges to a positive, finite limit as $n \rightarrow \infty$.

Let us also mention that a weak version of Proposition 1 holds in an arbitrary Banach algebra.

Theorem 6 (Gelfand's formula [5]). *Let \mathcal{A} be a complex Banach algebra and let $\|\cdot\|$ denote its norm. For every $M \in \mathcal{A}$, $\sqrt[n]{\|M^n\|}$ converges to the spectral radius of M as $n \rightarrow \infty$.*

Let us now illustrate Proposition 1 with an example. The matrix

$$M := \begin{bmatrix} 4 & -3 \\ 3 & 4 \end{bmatrix}$$

is diagonalizable:

$$PMP^{-1} = \begin{bmatrix} \lambda & 0 \\ 0 & \bar{\lambda} \end{bmatrix}$$

where $\lambda := 4 + 3i$, $\bar{\lambda} := 4 - 3i$, $P := \begin{bmatrix} 1 & i \\ 1 & -i \end{bmatrix}$ and $P^{-1} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ -i & i \end{bmatrix}$. Hence, $\|M^n\|_P = 5^n$ for every $n \in \mathbb{N}$, and thus for every norm $\|\cdot\|$ on $\mathbb{C}^{2 \times 2}$, $\|M^n\| \asymp 5^n$ as $n \rightarrow \infty$.

Noteworthy is that no entry of M^n behaves in the same way as $\|M^n\|$ as $n \rightarrow \infty$. Indeed, for every integer $n \in \mathbb{N}$,

$$M^n = P^{-1} \begin{bmatrix} \lambda^n & 0 \\ 0 & \bar{\lambda}^n \end{bmatrix} P \frac{1}{2} \begin{bmatrix} \lambda^n + \bar{\lambda}^n & i\lambda^n - i\bar{\lambda}^n \\ -i\lambda^n + i\bar{\lambda}^n & \lambda^n + \bar{\lambda}^n \end{bmatrix} = 5^n \begin{bmatrix} \cos(n\theta) & -\sin(n\theta) \\ \sin(n\theta) & \cos(n\theta) \end{bmatrix}$$

where θ is an argument of λ . Besides, both sets $\{\cos(n\theta) : n \in \mathbb{N}\}$ and $\{\sin(n\theta) : n \in \mathbb{N}\}$ are dense subsets of the closed real interval with endpoints -1 and $+1$ (see appendix).

We turn back to the proof of Theorem 1.

Lemma 1. *Let A be an alphabet and let $\sigma : A^* \rightarrow A^*$ be a morphism. Let d denote the cardinality of A . There exists a matrix $M \in \mathbb{N}^{d \times d}$ such that*

$$\sum_{a \in A} |\sigma^n(a)| = \|M^n\| \quad (1)$$

for every $n \in \mathbb{N}$.

Proof. Write A in the form $A = \{a_1, a_2, \dots, a_d\}$. Define the matrix $M \in \mathbb{N}^{d \times d}$ by: for each $(i, j) \in [1, d] \times [1, d]$, the $(i, j)^{\text{th}}$ entry of M^n equals $|\sigma^n(a_j)|_{a_i}$. For every $n \in \mathbb{N}$ and every $i, j \in [1, d]$, the $(i, j)^{\text{th}}$ entry of M equals $|\sigma^n(a_j)|_{a_i}$. Hence, Equation (1) holds. \square

Lemma 2. *Let (A, σ, w) be a DOL-system. If for every $a \in A$ there exists $m \in \mathbb{N}$ such that a occurs in $\sigma^m(w)$ then*

$$|\sigma^n(w)| \asymp \sum_{a \in A} |\sigma^n(a)|$$

as $n \rightarrow \infty$.

Proof. For every $n \in \mathbb{N}$, let

$$S_n := \sum_{a \in A} |\sigma^n(a)|.$$

First, we have

$$|\sigma^n(w)| = \sum_{a \in A} |w|_a |\sigma^n(a)| \leq \left(\max_{a \in A} |w|_a \right) S_n,$$

and thus $|\sigma^n(w)| \preceq S_n$ as $n \rightarrow \infty$. Conversely, let $a \in A$, let $m_a \in \mathbb{N}$ be such that a occurs in $\sigma^{m_a}(w)$, and let $L_a := \max_{b \in A} |\sigma^{m_a}(b)|$. Since $\sigma^n(a)$ occurs in $\sigma^{n+m_a}(w)$, we have

$$|\sigma^n(a)| \leq |\sigma^{n+m_a}(w)| = \sum_{b \in A} |\sigma^n(w)|_b |\sigma^{m_a}(b)| \leq L_a \sum_{b \in A} |\sigma^n(w)|_b = L_a |\sigma^n(w)|.$$

It follows

$$S_n \leq \left(\sum_{a \in A} L_a \right) |\sigma^n(w)|$$

and thus $S_n \preceq |\sigma^n(w)|$ as $n \rightarrow \infty$. □

Proof of Theorem 1. Let \bar{A} denote the set of all symbols $a \in A$ such that a occurs in $\sigma^m(w)$ for some $m \in \mathbb{N}$. Let $\bar{\sigma} : \bar{A}^* \rightarrow \bar{A}^*$ denote the morphism given by: $\bar{\sigma}(x) := \sigma(x)$ for every $x \in \bar{A}^*$. For every $n \in \mathbb{N}$, $\sigma^n(w) = \bar{\sigma}^n(w)$, and thus we may replace the D0L-system (A, σ, w) with $(\bar{A}, \bar{\sigma}, w)$ in the remaining of the proof: now, Lemma 2 applies to (A, σ, w) . Combining Lemma 2, Lemma 1 and Proposition 1, we get that there exists a non-negative integer α smaller than the cardinality of A and a real number $\beta \geq 0$ such that $|\sigma^n(w)| \asymp n^\alpha \beta^n$. Since $|\sigma^n(w)|$ is not less than one for every $n \in \mathbb{N}$, $n^\alpha \beta^n$ does not converge to zero as $n \rightarrow \infty$. From that we deduce $\beta \geq 1$. □

References

- [1] J. Berstel and C. Reutenauer. *Rational series and their languages*, volume 12 of *EATCS Monographs on Theoretical Computer Science*. Springer-Verlag, 1988. The new version is presently available online at Berstel's homepage.
- [2] A. Lindenmayer. Mathematical models for cellular interactions in development. *Journal of Theoretical Biology*, 18(3):280–315, 1968.
- [3] J.-J. Pansiot. Complexité des facteurs des mots infinis engendrés par morphismes itérés. In *Proceedings of the 11th International Colloquium on Automata, Languages and Programming (ICALP'84)*, volume 172 of *Lecture Notes in Computer Science*, pages 380–389. Springer-Verlag, 1984.
- [4] M. Queffélec. *Substitution dynamical systems-spectral analysis*, volume 1294 of *Lecture Notes in Mathematics*. Springer-Verlag, 1987.
- [5] W. Rudin. *Functional analysis*. International Series in Pure and Applied Mathematics. McGraw-Hill, second edition, 1991.
- [6] A. Salomaa and M. Soittola. *Automata-theoretic aspects of formal power series*. Texts and Monographs in Computer Science. Springer-Verlag, 1978.
- [7] R. S. Varga. *Matrix iterative analysis*, volume 27 of *Springer Series in Computational Mathematics*. Springer-Verlag, second edition, 2000.

Appendix

Throughout the section,

- π denotes Archimedes' constant,
- i denotes the imaginary unit,
- $I := \{x \in \mathbb{R} : 0 \leq x \leq +1\}$, and
- $J := \{x \in \mathbb{R} : -1 \leq x \leq +1\}$.

The aim of this appendix is to prove the following proposition.

Proposition 2. *For any argument θ of $4 + 3i$, both sets $\{\cos(n\theta) : n \in \mathbb{N}\}$ and $\{\sin(n\theta) : n \in \mathbb{N}\}$ are dense subsets of J .*

Proposition 2 is a consequence of the following two well-known results.

Proposition 3. *Let ρ be a real number. If both ρ and $\cos(2\pi\rho)$ are rational then $\cos(2\pi\rho) \in \{-1, -\frac{1}{2}, 0, +\frac{1}{2}, +1\}$.*

Proof. Clearly, $2\cos(2\pi\rho)$ is comprised between -2 and $+2$, inclusive.

Assume that ρ is rational. Then, both complex numbers $\exp(2\pi\rho i)$ and $\exp(-2\pi\rho i)$ are algebraic integers. Indeed they are roots of the monic integer polynomial $z^q - 1$, where q is a positive integer such that $q\rho$ is an integer. Since the sum of two algebraic integers is also an algebraic integer, $2\cos(2\pi\rho) = \exp(2\pi\rho i) + \exp(-2\pi\rho i)$ is an algebraic integer.

Assume in addition that $2\cos(2\pi\rho)$ is rational. Since a rational number is an algebraic integer if and only if it is an integer, $2\cos(2\pi\rho)$ belongs to $\{-2, -1, 0, +1, +2\}$. \square

Proposition 4. *For any irrational number $\rho \in \mathbb{R}$, $\{n\rho - \lfloor n\rho \rfloor : n \in \mathbb{N}\}$ is a dense subset of I .*

Proof of Proposition 2. Since the cosine of θ equals $\frac{4}{5}$, $\frac{\theta}{2\pi}$ is irrational by Proposition 3. Hence, $D := \{\frac{n\theta}{2\pi} - \lfloor \frac{n\theta}{2\pi} \rfloor : n \in \mathbb{N}\}$ is a dense subset of I by Proposition 4. Since the function $f : I \rightarrow J$ that maps each $x \in I$ to $\cos(2\pi x)$ is continuous and surjective, $\{\cos(n\theta) : n \in \mathbb{N}\} = f(D)$ is a dense subset of J . In the same way, the function $g : I \rightarrow J$ that maps each $x \in I$ to $\sin(2\pi x)$ is continuous, surjective and such that $\{\sin(n\theta) : n \in \mathbb{N}\} = g(D)$. \square