Localized Solitons of a $(2+1)$ -dimensional Nonlocal Nonlinear Schrödinger Equation

Ken-ichi Maruno¹†

¹ Department of Mathematics, The University of Texas-Pan American, Edinburg, TX 78539-2999

> Yasuhiro Ohta² ² Department of Mathematics, Kobe University,

> > Rokko, Kobe 657-8501, Japan

October 31, 2018

Abstract

An integrable $(2+1)$ -dimensional nonlocal nonlinear Schrödinger equation is discussed. The *N*-soliton solution is given by Gram type determinant. It is found that the localized N-soliton solution has interesting interaction behavior which shows change of amplitude of localized pulses after collisions.

1 Introduction

The nonlinear Schrödinger (NLS) equation,

$$
i\psi_t = \psi_{xx} + \alpha |\psi|^2 \psi, \qquad (1)
$$

is the most important soliton equation which is a widely used model for investigating the evolution of pulses in optical fiber and of surface gravity waves with narrow-banded spectra in fluid [\[1\]](#page-9-0). The study of vector and nonlocal analogues of the NLS equation has received considerable attention recently from both physical and mathematical points of view [\[1,](#page-9-0) [2,](#page-9-1) [3,](#page-9-2) [4,](#page-9-3) [5,](#page-9-4) [6\]](#page-9-5).

In this Letter, we discuss a $(2+1)$ -dimensional nonlocal nonlinear Schrödinger (2DNNLS) equation:

$$
i u_t = u_{xx} + 2u \int_{-\infty}^{\infty} |u|^2 dy, \qquad (2)
$$

where $u = u(x, y, t)$ is a complex function and x, y, t are real. The Gram type determinant solution is presented and localized soliton interactions are studied.

2 Determinant Solution

Using the dependent variable transformation

$$
u(x, y, t) = \frac{g(x, y, t)}{f(x, t)}, \qquad u^*(x, y, t) = \frac{g^*(x, y, t)}{f(x, t)},
$$

where f is real and $*$ is complex conjugate, we have bilinear equations [\[7\]](#page-9-6)

$$
(D_x^2 - iD_t)g \cdot f = 0,\t\t(3)
$$

$$
(D_x^2 + iD_t)g^* \cdot f = 0,\t\t(4)
$$

$$
D_x^2 f \cdot f = 2 \int_{-\infty}^{\infty} g g^* dy. \tag{5}
$$

These bilinear equations have the following Gram determinant solution which is the *N*-soliton solution of the 2DNNLS equation:

$$
f = \begin{vmatrix} \mathcal{A}_N & I_N \\ -I_N & \mathcal{B}_N \end{vmatrix},
$$

$$
g = \begin{vmatrix} \mathcal{A}_N & I_N & \mathbf{e}_N^T \\ -I_N & \mathcal{B}_N & \mathbf{0}^T \\ \mathbf{0} & -\mathbf{a}_N & 0 \end{vmatrix}, \quad g^* = -\begin{vmatrix} \mathcal{A}_N & I_N & \mathbf{0}^T \\ -I_N & \mathcal{B}_N & \mathbf{a}^*_{N}^T \\ -\mathbf{e}_N^* & \mathbf{0} & 0 \end{vmatrix},
$$

where

$$
\mathcal{A}_N=\begin{pmatrix} \frac{e^{\xi_1+\xi_1^*}}{p_1+p_1^*} & \frac{e^{\xi_1+\xi_2^*}}{p_1+p_2^*} & \cdots & \frac{e^{\xi_1+\xi_N^*}}{p_1+p_N^*} \\ \frac{e^{\xi_2+\xi_1^*}}{p_2+p_1^*} & \frac{e^{\xi_2+\xi_2^*}}{p_2+p_2^*} & \cdots & \frac{e^{\xi_2+\xi_N^*}}{p_2+p_N^*} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{e^{\xi_N+\xi_1^*}}{p_N+p_1^*} & \frac{e^{\xi_N+\xi_2^*}}{p_N+p_2^*} & \cdots & \frac{e^{\xi_N+\xi_N^*}}{p_N+p_N^*} \end{pmatrix},
$$

$$
\mathcal{B}_{N} = \begin{pmatrix}\n\frac{\int_{-\infty}^{\infty} a_{1}^{*} a_{1} dy}{p_{1}^{*} + p_{1}} & \frac{\int_{-\infty}^{\infty} a_{1}^{*} a_{2} dy}{p_{1}^{*} + p_{2}} & \cdots & \frac{\int_{-\infty}^{\infty} a_{1}^{*} a_{N} dy}{p_{1}^{*} + p_{N}} \\
\frac{\int_{-\infty}^{\infty} a_{2}^{*} a_{1} dy}{p_{2}^{*} + p_{1}} & \frac{\int_{-\infty}^{\infty} a_{2}^{*} a_{2} dy}{p_{2}^{*} + p_{2}} & \cdots & \frac{\int_{-\infty}^{\infty} a_{2}^{*} a_{N} dy}{p_{2}^{*} + p_{N}} \\
\vdots & \vdots & \ddots & \vdots \\
\frac{\int_{-\infty}^{\infty} a_{N}^{*} a_{1} dy}{p_{N}^{*} + p_{1}} & \frac{\int_{-\infty}^{\infty} a_{N}^{*} a_{2} dy}{p_{N}^{*} + p_{2}} & \cdots & \frac{\int_{-\infty}^{\infty} a_{N}^{*} a_{N} dy}{p_{N}^{*} + p_{N}}\n\end{pmatrix},
$$

and I_N is the $N \times N$ identity matrix, \mathbf{a}^T is the transpose of \mathbf{a} ,

$$
\mathbf{a}_{N} = (a_{1}, a_{2}, \cdots, a_{N}), \quad \mathbf{e}_{N} = (e^{\xi_{1}}, e^{\xi_{2}}, \cdots, e^{\xi_{N}}), \quad \mathbf{0} = (0, 0, \cdots, 0),
$$

$$
\xi_{i} = p_{i}x - \mathbf{i}p_{i}^{2}t, \quad \xi_{i}^{*} = p_{i}^{*}x + \mathbf{i}p_{i}^{*2}t, \qquad 1 \leq i \leq N,
$$

and p_i is a complex wave number of *i*-th soliton and $a_i \equiv a_i(y)$ is a complex phase function of *i*-th soliton.

Here, we show that eq.[\(5\)](#page-1-0) has the above Gram determinant solution.

Let us denote the (i, j) -cofactor of the matrix

$$
M = \begin{pmatrix} \mathcal{A}_N & I_N \\ -I_N & \mathcal{B}_N \end{pmatrix}
$$

as Δ_{ij} . Then the *x*-derivative of $f = \det M$ is given by

$$
f_x = \sum_{i=1}^N \sum_{j=1}^N \Delta_{ij} \frac{\partial}{\partial x} \frac{e^{\xi_i + \xi_j^*}}{p_i + p_j^*} = \sum_{i=1}^N \sum_{j=1}^N \Delta_{ij} e^{\xi_i + \xi_j^*}
$$

=
$$
\begin{vmatrix} \mathcal{A}_N & I_N & \mathbf{e}_N^T \\ -I_N & \mathcal{B}_N & \mathbf{0}^T \\ -\mathbf{e}_N^* & \mathbf{0} & 0 \end{vmatrix}.
$$
 (6)

In the Gram determinant expression of *f*, dividing *i*-th row by e^{ξ_i} and multiplying $(N+i)$ -th column by e^{ξ_i} for $i = 1, \dots, N$, and dividing *j*-th column by $e^{\xi_i^*}$ and multiplying $(N + j)$ -th row by $e^{\xi_j^*}$ for $j = 1, \dots, N$, we obtain another determinant expression of *f*,

$$
f=\det M',
$$

where

$$
M'=\begin{pmatrix} \mathcal{A}'_N & I_N \\ -I_N & \mathcal{B}'_N \end{pmatrix},
$$

$$
\mathcal{A}'_{N} = \begin{pmatrix}\n\frac{1}{p_{1} + p_{1}^{*}} & \cdots & \frac{1}{p_{1} + p_{N}^{*}} \\
\vdots & \ddots & \vdots \\
\frac{1}{p_{N} + p_{1}^{*}} & \cdots & \frac{1}{p_{N} + p_{N}^{*}}\n\end{pmatrix}, \\
\mathcal{B}'_{N} = \begin{pmatrix}\n\frac{e^{\xi_{1}^{*} + \xi_{1}} \int_{-\infty}^{\infty} a_{1}^{*} a_{1} dy}{p_{1}^{*} + p_{1}} & \cdots & \frac{e^{\xi_{1}^{*} + \xi_{N}} \int_{-\infty}^{\infty} a_{1}^{*} a_{N} dy}{p_{1}^{*} + p_{N}} \\
\vdots & \ddots & \vdots \\
\frac{e^{\xi_{N}^{*} + \xi_{1}} \int_{-\infty}^{\infty} a_{N}^{*} a_{1} dy}{p_{N}^{*} + p_{1}} & \cdots & \frac{e^{\xi_{N}^{*} + \xi_{N}} \int_{-\infty}^{\infty} a_{N}^{*} a_{N} dy}{p_{N}^{*} + p_{N}}\n\end{pmatrix}.
$$

Thus the *x*-derivative of *f* is also written as

$$
f_x = \sum_{i=1}^{N} \sum_{j=1}^{N} \Delta'_{N+i,N+j} \frac{\partial}{\partial x} \frac{e^{\xi_i^* + \xi_j} \int_{-\infty}^{\infty} a_i^* a_j dy}{p_i^* + p_j} = \sum_{i=1}^{N} \sum_{j=1}^{N} \Delta'_{N+i,N+j} e^{\xi_i^* + \xi_j} \int_{-\infty}^{\infty} a_i^* a_j dy
$$

=
$$
\int_{-\infty}^{\infty} \sum_{i=1}^{N} \sum_{j=1}^{N} \Delta'_{N+i,N+j} e^{\xi_i^* + \xi_j} a_i^* a_j dy
$$

where Δ'_{ij} is the (i, j) -cofactor of M' . Therefore we have

$$
f_x = \int_{-\infty}^{\infty} \begin{vmatrix} \mathcal{A}'_N & I_N & \mathbf{0}^T \\ -I_N & \mathcal{B}'_N & \mathbf{\tilde{a}}_N^{*T} \\ \mathbf{0} & -\mathbf{\tilde{a}}_N & 0 \end{vmatrix} dy
$$

=
$$
\int_{-\infty}^{\infty} \begin{vmatrix} \mathcal{A}_N & I_N & \mathbf{0}^T \\ -I_N & \mathcal{B}_N & \mathbf{a}^*_{N}^T \\ \mathbf{0} & -\mathbf{a}_N & 0 \end{vmatrix} dy,
$$

where

$$
\widetilde{\mathbf{a}}_N=(e^{\xi_1}a_1,\cdots,e^{\xi_N}a_N).
$$

By differentiating the above f_x by x , we get

$$
f_{xx} = \int_{-\infty}^{\infty} \begin{vmatrix} \mathcal{A}_N & I_N & \mathbf{e}_N^T & \mathbf{0}^T \\ -I_N & \mathcal{B}_N & \mathbf{0}^T & \mathbf{a}^{*}{}_N^T \\ -\mathbf{e}_N^* & \mathbf{0} & 0 & 0 \\ \mathbf{0} & -\mathbf{a}_N & 0 & 0 \end{vmatrix} dy.
$$

On the other hand, using the Jacobi formula for determinant[\[7\]](#page-9-6), we have

$$
gg^* = \begin{vmatrix} \mathcal{A}_N & I_N & \mathbf{e}_N^T & \mathbf{0}^T \\ -I_N & \mathcal{B}_N & \times \begin{vmatrix} -I_N & \mathcal{B}_N & \mathbf{0}^T & \mathbf{a}^*_{N} \\ -\mathbf{e}_N^* & \mathbf{0} & 0 & 0 \\ \mathbf{0} & -\mathbf{a}_N & 0 & 0 \end{vmatrix} \\ -\begin{vmatrix} \mathcal{A}_N & I_N & \mathbf{e}_N^T \\ -I_N & \mathcal{B}_N & \mathbf{0}^T \\ -\mathbf{e}_N^* & \mathbf{0} & 0 \end{vmatrix} \times \begin{vmatrix} \mathcal{A}_N & I_N & \mathbf{0}^T \\ -I_N & \mathcal{B}_N & \mathbf{a}^*_{N}^T \\ \mathbf{0} & -\mathbf{a}_N & 0 \end{vmatrix}
$$

.

Here we note that the *y*-dependence in right-hand side appears only in the last row and last column of the second determinant in each term. Thus we obtain

$$
\int_{-\infty}^{\infty} gg^* dy = f f_{xx} - f_x f_x,
$$

which is a bilinear equation [\(5\)](#page-1-0).

Since eqs.[\(3\)](#page-1-0) and [\(4\)](#page-1-0) are bilinear equations for the NLS equation and do not include *y*, we can prove in the same way in the NLS equation that the above Gram determinant solution satisfies the bilinear identities [\(3\)](#page-1-0) and [\(4\)](#page-1-0), i.e., we can show easily that the bilinear equations [\(3\)](#page-1-0) and [\(4\)](#page-1-0) are made from a pair of Jacobi identities, respectively.

3 Localized Solitons

Using the above formula, we can make 1-soliton solution as follows:

$$
u = \frac{g}{f} = \frac{a_1 e^{\xi_1}}{1 + \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{(p_1^* + p_1)^2} e^{\xi_1 + \xi_1^*}}, \quad u^* = \frac{g^*}{f} = \frac{a_1^* e^{\xi_1^*}}{1 + \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{(p_1^* + p_1)^2} e^{\xi_1 + \xi_1^*}},
$$
(7)

where

$$
f = \begin{vmatrix} \frac{e^{\xi_1 + \xi_1^*}}{p_1 + p_1^*} & 1 \\ -1 & \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{p_1^* + p_1} \end{vmatrix} = 1 + \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{(p_1^* + p_1)^2} e^{\xi_1 + \xi_1^*},
$$

$$
g = \begin{vmatrix} \frac{e^{\xi_1 + \xi_1^*}}{p_1 + p_1^*} & 1 & e^{\xi_1} \\ -1 & \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{p_1^* + p_1} & 0 \\ 0 & -a_1 & 0 \end{vmatrix} = a_1 e^{\xi_1},
$$

$$
g^* = - \begin{vmatrix} \frac{e^{\xi_1 + \xi_1^*}}{p_1 + p_1^*} & 1 & 0 \\ -1 & \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{p_1^* + p_1} & a_1^* \\ -e^{\xi_1^*} & 0 & 0 \end{vmatrix} = a_1^* e^{\xi_1^*}.
$$

If we choose $a_1(y) = \alpha_1 \text{sech}(k(y + \eta_0))$ where α_1 is a complex number and *k* and η_0 are real numbers,

$$
u = \frac{\alpha_1 \mathrm{sech}(k(y + \eta_0))e^{\xi_1}}{1 + \frac{(2/k)|\alpha_1|^2}{(p_1^* + p_1)^2}e^{\xi_1 + \xi_1^*}} = \frac{\alpha_1}{2\sqrt{A}} \mathrm{sech}(ky + \eta_0) \mathrm{sech}\left(\frac{\xi_1 + \xi_1^*}{2} + \frac{1}{2}\log A\right)e^{\frac{\xi_1 - \xi_1^*}{2}},
$$

where $A = \frac{(2/k)|\alpha_1|^2}{(n^*+n)^2}$ $\frac{(2/k)|\alpha_1|^2}{(p_1^*+p_1)^2}$. In this case, we have a localized pulse as shown in figure [1.](#page-6-0) If we choose $a_1(y) = \alpha_1 \text{sech}(k(y + \eta_1)) + \alpha_2 \text{sech}(k(y + \eta_2))$ where α_1 and α_2 are complex numbers and k , η_1 and η_2 are real numbers,

$$
u = \frac{(\alpha_1 \text{sech}(k(y + \eta_1)) + \alpha_2 \text{sech}(k(y + \eta_2)))e^{\xi_1}}{1 + \frac{(2/k)(|\alpha_1|^2 + |\alpha_2|^2) + 4(\eta_1 - \eta_2)(\alpha_1 \alpha_2^* + \alpha_1^* \alpha_2)/(e^{k(\eta_1 - \eta_2)} - e^{-k(\eta_1 - \eta_2)})}{(p_1^* + p_1)^2}e^{\xi_1 + \xi_1^*}
$$

=
$$
\frac{1}{2\sqrt{A}}(\alpha_1 \text{sech}(k(y + \eta_1)) + \alpha_2 \text{sech}(k(y + \eta_2))
$$

× sech $\left(\frac{\xi_1 + \xi_1^*}{2} + \frac{1}{2}\log A\right)e^{\frac{\xi_1 - \xi_1^*}{2}},$

where $A = \frac{(2/k)(|\alpha_1|^2 + |\alpha_2|^2) + 4(\eta_1 - \eta_2)(\alpha_1\alpha_2^* + \alpha_1^*\alpha_2)/(e^{k(\eta_1 - \eta_2)} - e^{-k(\eta_1 - \eta_2)})}{(n^* + n_1)^2}$ $\frac{(p_1^* + p_1)^2}{(p_1^* + p_1)^2}$. We see two localized pulses in figure [2.](#page-6-1) These two localized pulses travel parallel to the *x*axis. With $a_1(y) = \sum_j^M \alpha_j \text{sech}(k(y + \eta_j))$, we can see *M*-localized pulses travelling parallel to the *x*-axis. We call this *M*-localized pulse the $(1, M)$ -localized pulse solution. In the general case of pulse solutions generated from the *N*-soliton formula, it is named by (*N*,*M*)-localized pulse solution.

Figure 1: 1-soliton solution. $\alpha_1 = 1 + 2i$, $p_1 = 2 + 3i$, $k = 3$, $\eta_0 = 0$.

Figure 2: 1-soliton solution. $\alpha_1 = 1 + 2i, \alpha_2 = 1/2 + i, p_1 = 2 + 3i, k = 3, \eta_1 =$ $-6, η₂ = 6.$

Next, we consider the case of $N = 2$, i.e. 2-soliton solution. Using the determinant form of *N*-soliton solution, we have

$$
f = \begin{vmatrix} \frac{e^{\xi_1 + \xi_1^*}}{p_1 + p_1^*} & \frac{e^{\xi_1 + \xi_2^*}}{p_1 + p_2^*} & 1 & 0\\ \frac{e^{\xi_2 + \xi_1^*}}{p_2 + p_1^*} & \frac{e^{\xi_2 + \xi_2^*}}{p_2 + p_2^*} & 0 & 1\\ -1 & 0 & \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{p_1^* + p_1} & \frac{\int_{-\infty}^{\infty} a_1^* a_2 dy}{p_1^* + p_2} \\ 0 & -1 & \frac{\int_{-\infty}^{\infty} a_2^* a_1 dy}{p_2^* + p_1} & \frac{\int_{-\infty}^{\infty} a_2^* a_2 dy}{p_2^* + p_2} \\ \frac{e^{\xi_1 + \xi_1^*}}{p_1^* + p_1} + \frac{c_{12}}{p_1^* + p_2} & e^{\xi_1 + \xi_2} + \frac{c_{21}}{p_2^* + p_1} & e^{\xi_1 + \xi_2^*} + \frac{c_{22}}{p_2^* + p_2} \\ + \left(\frac{c_{12}c_{21} - c_{11}c_{22}}{(p_2^* + p_1)(p_1^* + p_2)} + \frac{c_{11}c_{22} - c_{12}c_{21}}{(p_1^* + p_1)(p_2^* + p_2)} \right) e^{\xi_1 + \xi_2 + \xi_1^* + \xi_2^*},
$$

$$
g = \begin{vmatrix} \frac{e^{\xi_1 + \xi_1^*}}{p_1 + p_1^*} & \frac{e^{\xi_2 + \xi_2^*}}{p_2 + p_1^*} & 1 & 0 & e^{\xi_1} \\ \frac{e^{\xi_2 + \xi_1^*}}{p_2 + p_1^*} & \frac{e^{\xi_2 + \xi_2^*}}{p_2 + p_2^*} & 0 & 1 & e^{\xi_2} \\ -1 & 0 & \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{p_1^* + p_1} & \frac{\int_{-\infty}^{\infty} a_1^* a_2 dy}{p_1^* + p_2} & 0 \\ 0 & -1 & \frac{\int_{-\infty}^{\infty} a_2^* a_1 dy}{p_2^* + p_1} & \frac{\int_{-\infty}^{\infty} a_2^* a_2 dy}{p_2^* + p_2} & 0 \\ 0 & 0 & -a_1 & -a_2 & 0 \\ -a_1 e^{\xi_1} + a_2 e^{\xi_2} + \frac{(c_{12}a_1 - c_{11}a_2)(p_1 - p_2)}{(p_1^* + p_1)(p_1^* + p_2)} e^{\xi_1 + \xi_1^* + \xi_2} \\ + \frac{(c_{22}a_1 - c_{21}a_2)(p_1 - p_2)}{(p_2^* + p_1)(p_2^* + p_2)} e^{\xi_2 + \xi_2^* + \xi_1}, \\ \frac{e^{\xi_1 + \xi_1^*}}{p_1 + p_1^*} & \frac{e^{\xi_2 + \xi_2^*}}{p_2 + p_2^*} & 0 & 1 & 0 \\ e^{\xi_2 + \xi_1^*} & \frac{e^{\xi_2 + \xi_2^*}}{p_2 + p_2^*} & 0 & 1 & 0 \\ 0 & -1 & \frac{\int_{-\infty}^{\infty} a_1^* a_1 dy}{p_1^* + p_1} & \frac{\int_{-\infty}^{\infty} a_1^* a_2 dy}{p_1^* + p_2} & a_1^* \\ -e^{\xi_1^*} & -e^{\xi_2^*} & 0 & 0 & 0 \\ -e^{\xi_1^*} & -e^{\xi_2^*} & 0 & 0 & 0 \\ -a
$$

 $\overline{}$ $\Big\}$ $\Big\}$ $\Big\}$ $\Big\}$ $\Big\}$ $\overline{}$ $\overline{}$ \vert

where $c_{ij} = \int_{-\infty}^{\infty} a_i^* a_j dy / (p_i^* + p_j).$

To make four localized pulses, i.e. (2,2)-localized pulse solution, we consider $a_i(y) = \sum_{j=1}^{2} \alpha_{2(i-1)+j}$ sech $(k(y + \eta_j))$. Then *c_{ij}* is given as follows.

$$
c_{ij} = \frac{(2/k)(\alpha_{2(i-1)+1}^{*}\alpha_{2(j-1)+1} + \alpha_{2(i-1)+2}^{*}\alpha_{2(j-1)+2})}{(p_i^{*} + p_j)} + \frac{4(\eta_1 - \eta_2)(\alpha_{2(j-1)+1}\alpha_{2(i-1)+2}^{*} + \alpha_{2(i-1)+1}^{*}\alpha_{2(j-1)+2})}{(e^{k(\eta_1 - \eta_2)} - e^{-k(\eta_1 - \eta_2)})(p_i^{*} + p_j)}.
$$

Figure [3](#page-8-0) is an example of (2,2)-localized pulse solution. It is observed that 4 localized pulses suddenly change the height of pulses after a collision. Each pair

Figure 3: (2,2)-localized pulse solution. $\alpha_1 = 1 + i, \alpha_2 = 1, \alpha_3 = 1, \alpha_4 = 1, p_1 =$ $3/2-5i/2, p_2 = 3-i, k = 2, \eta_1 = -5, \eta_2 = 5.$

of pulses on lines parallel to the *x*-axis collides, then the total mass of pulses is redistributed. In the case of figure [3,](#page-8-0) the height of a localized pulse become very small after a collision. Although there is a distance between two pulses on a line parallel to the *x*-axis and other two pulses on another line, the collision causes an effect of 4-pulse interaction. As this example, solutions of the 2DNNLS equation have very complicated and interesting properties.

4 Conclusion

We have discussed an integrable 2DNNLS equation and shown that the *N*-soliton solution of the 2DNNLS equation is given by the Gram type determinant and solutions can be localized in *x*-*y* plane.

Note that the integrable 2DNNLS equation discussed in this Letter can be considered as the vector NLS equation with infinitely many components [\[8,](#page-9-7) [9,](#page-9-8) [10,](#page-9-9) [1\]](#page-9-0). This fact suggests that the vector soliton equations can produce nonlocal multi-dimensional soliton equations having localized pulses.

It should be noted that a model for second harmonic generation, i.e., quadratic solitons, was discussed in the paper by Nikolov et al., and they discussed the relationship between a nonlocal soliton equation and a vector soliton system [\[5\]](#page-9-4). Finding physical systems which could be described by the 2DNNLS equation is an interesting problem.

Note added in proof: After the acceptance of this Letter for publication, the

authors noticed the 2DNNLS equation (2) is equivalent to eq.(7.86) in ref.[11]. However, as far as we know, the N-soliton solution has not been obtained so far. The authors thank Dr. Takayuki Tsuchida for letting us know the paper by Zakharov [11].

References

- [1] M. J. Ablowitz, B. Prinari and A. D. Trubatch, *Discrete and Continuous Nonlinear Schrödinger Systems* (Cambridge University Press, 2004).
- [2] D. Pelinovsky, Phys. Lett. A **197** (1995) 401.
- [3] D. Pelinovsky and R. H. J. Grimshaw, J. Math. Phys. **36** (1995) 4203.
- [4] W. Królikowski and O. Bang, Phys. Rev. E **63** (2000) 016610.
- [5] N. I. Nikolov, D. Neshev, W. Królikowski and O. Bang, Phys. Rev. E 68 (2003) 036614.
- [6] B. Deconinck and J. N. Kutz, Phys. Lett. A **319** (2003) 97.
- [7] R. Hirota, *The Direct Method in Soliton Theory* (Cambridge University Press, 2004).
- [8] S. V. Manakov, Sov. Phys. JETP **38** (1974) 248.
- [9] R. Radhakrishnan, M. Lakshmanan, and J. Hietarinta, Phys. Rev. E **56** (1997) 2213.
- [10] P. D. Miller, Phys. Lett. A **101** (1997) 17.
- [11] V. E. Zakharov, in Solitons, ed. Bullough and Caudrey, Topics in Current Physics (Springer, Berlin - New York, 1980) 243.