

On Parallel Sections of a Vector Bundle

Richard Atkins

richard.atkins@twu.ca

Department of Mathematics

Trinity Western University

7600 Glover Road

Langley, BC, V2Y 1Y1 Canada

Abstract

We consider when a smooth vector bundle endowed with a connection possesses non-trivial, local parallel sections. This is accomplished by means of a derived flag of subsets of the bundle. The procedure is algebraic and rests upon the Frobenius Theorem.

1 Introduction

A connection on a vector bundle is a type of differentiation that acts on vector fields. Its importance lies in the fact that given a piecewise continuous curve connecting two points on the underlying manifold, the connection defines a linear isomorphism between the respective fibres over these points. A renowned theorem of differential geometry states that when the Riemann curvature tensor of the connection vanishes, there exist local frames comprised of parallel sections. This paper presents a refinement of this result. That is, given a connection on a vector bundle we determine when there exist local parallel sections and we find the subbundle they generate. This is accomplished by means of an algebraic construction of a derived flag of subsets of the original vector bundle.

This question has been considered in the real analytic case by Trenevski (cf. [3]). Our solution, by contrast, follows from the Frobenius Theorem, which applies to smooth (C^∞) data. Furthermore, while Trenevski's method relies upon power series expansions we take a more geometric approach to the problem.

By applying our methods to the vector bundle of symmetric two-tensors over a manifold we obtain a solution to the problem of determining when a connection is locally a metric connection. For the case of surfaces, this has also been dealt with in [1].

2 The Existence of Parallel Sections

Let $\pi : W \rightarrow M$ be a smooth vector bundle and W' a subset of W with the following two properties:

P1: For each $x \in M$, $W_x \cap W'$ is a linear subspace of $W_x := \pi^{-1}(x)$.

P2: For each $w \in W'$ there exists an open neighbourhood U of $\pi(w)$ in M and a smooth local section $X : U \subseteq M \rightarrow W' \subseteq W$ such that $w = X(\pi(w))$.

Let

$$\nabla : \mathcal{A}^0(W) \rightarrow \mathcal{A}^1(W)$$

be a connection on W , where $\mathcal{A}^n(W)$ denotes the space of local sections $U \subseteq M \rightarrow W \otimes \Lambda^n M$. Define a map

$$\tilde{\alpha} : \mathcal{A}^0(W) \rightarrow \mathcal{A}^1(W/W')$$

by

$$\tilde{\alpha} := \phi \circ \nabla,$$

where W/W' is the quotient of W and W' taken fibrewise and $\phi : W \otimes T^*M \rightarrow (W/W') \otimes T^*M$ denotes the natural projection. For any local section $X : U \subseteq M \rightarrow W' \subseteq W$ and differentiable function $f : U \rightarrow \mathfrak{R}$ we have $\tilde{\alpha}(fX) = f\tilde{\alpha}(X)$. Thus, there corresponds to $\tilde{\alpha}$ a map

$$\alpha_{W'} : W' \rightarrow (W/W') \otimes T^*M$$

acting linearly on each fibre of W' . $\alpha_{W'}$ is the *second fundamental 1-form* of W' .

Let V be any subset of W satisfying P1. Define $\mathcal{S}(V)$ to be the subset of V consisting of all elements v for which there exists a smooth

local section $X : U \subseteq M \rightarrow V \subseteq W$ such that $v = X(\pi(v))$. Then $\mathcal{S}(V)$ satisfies both P1 and P2.

We seek to construct the maximal flat subset \widetilde{W} , of W . \widetilde{W} may be obtained as follows. Set

$$\begin{aligned} V^{(0)} &:= \{w \in W \mid R(\cdot)(w) = 0\} \\ W^{(i)} &:= \mathcal{S}(V^{(i)}) \\ V^{(i+1)} &:= \ker \alpha_{W^{(i)}} \end{aligned}$$

where $R : TM \otimes TM \otimes W \rightarrow W$ denotes the curvature tensor of ∇ . This gives a sequence

$$W \supseteq W^{(0)} \supseteq W^{(1)} \supseteq \dots \supseteq W^{(k)} \supseteq \dots$$

of subsets of W . Note that $W^{(i)}$ is not necessarily a vector bundle over M since the dimension of the fibres may vary from point to point. For some $k \in \mathbb{N}$, $W^{(l)} = W^{(k)}$ for all $l \geq k$. Define $\widetilde{W} = W^{(k)}$, with projection $\tilde{\pi} : \widetilde{W} \rightarrow M$.

In order to extract information from \widetilde{W} we need some concept of regularity. Accordingly, we say that the connection ∇ is *regular at* $x \in M$ if there exists a neighbourhood U of x such that $\tilde{\pi}^{-1}(U) \subseteq \widetilde{W}$ is a vector bundle over U . ∇ is *regular* if \widetilde{W} is a vector bundle over M . The dimension of the fibres of \widetilde{W} , for regular ∇ , shall be denoted $\text{rank } \widetilde{W}$.

Theorem 1 *Let ∇ be a connection on the smooth vector bundle $\pi : W \rightarrow M$.*

(i) If $X : U \subseteq M \rightarrow W$ is a local parallel section then the image of X lies in \widetilde{W} .

(ii) Suppose that ∇ is regular at $x \in M$. Then for every $w \in \widetilde{W}_x$ there exists a local parallel section $X : U \subseteq M \rightarrow \widetilde{W}$ with $X(x) = w$.

Proof:

(i) follows directly from the definition of \widetilde{W} .

(ii) Suppose that ∇ is regular at $x \in M$ and let $w \in \widetilde{W}_x$. By regularity, there exists a neighbourhood U_1 of x and a frame (X_1, \dots, X_n) of $\tilde{\pi}^{-1}(U_1) \subseteq \widetilde{W}$. By choosing a possibly smaller neighbourhood $U_2 \subseteq U_1$ of x we can extend (X_1, \dots, X_n) to a frame $\mathcal{X} := (X_1, \dots, X_n, \dots, X_N)$ of $\pi^{-1}(U_2) \subseteq W$. Let $\omega = \omega_j^i$ denote the connection form of ∇ with respect to \mathcal{X} : $\nabla_X X_j = \sum_{i=1}^N X_i \omega_j^i(X)$. Since \widetilde{W} has zero second fundamental 1-form,

$$\omega = \left(\begin{array}{c|c} \phi & * \\ \hline 0 & * \end{array} \right)$$

where ϕ is an $n \times n$ matrix of 1-forms. The curvature form $\Omega = \Omega_j^i$ of ∇ with respect to \mathcal{X} is

$$\Omega = d\omega + \omega \wedge \omega = \left(\begin{array}{c|c} d\phi + \phi \wedge \phi & * \\ \hline 0 & * \end{array} \right)$$

Since the curvature tensor R is identically zero, when restricted to $TM \otimes TM \otimes \widetilde{W}$, it follows that

$$d\phi + \phi \wedge \phi = 0$$

Therefore, by the Frobenius Theorem, there exists an $n \times n$ matrix of functions $A = A_j^i$ defined in a neighbourhood $U \subseteq U_2$ of x such that $dA = -\phi \wedge A$ and $A(x) = I_{n \times n}$, the $n \times n$ identity matrix (cf. [4],

chp. 7, 2. Proposition 1., pg. 290). Let c^j , $1 \leq j \leq n$, be real scalars satisfying $w = \sum_{j=1}^n X_j(x)c^j$. Define functions f^i on U by

$$f^i = \begin{cases} \sum_{j=1}^n A_j^i c^j & 1 \leq i \leq n \\ 0 & n+1 \leq i \leq N. \end{cases}$$

Let $X : U \rightarrow \widetilde{W}$ be the local section of \widetilde{W} defined by $X := \sum_{i=1}^N X_i f^i$. Since $df + \omega \cdot f = 0$, X is parallel. Moreover, $X(x) = w$.

q.e.d.

Corollary 2 *Let ∇ be a regular connection on the smooth vector bundle $\pi : W \rightarrow M$. Then (\widetilde{W}, ∇) is a flat vector bundle over M .*

Corollary 3 *Let ∇ be a connection on the smooth vector bundle $\pi : W \rightarrow M$, regular at $x \in M$. Then there are $\dim \widetilde{W}_x$ independent local parallel sections in a neighbourhood of $x \in M$.*

Example Consider the symmetric connection ∇ on the 2-sphere, $M = S^2$, defined as follows: $\Gamma_{\phi\phi}^\theta = -\sin\theta\cos\theta$, $\Gamma_{\theta\theta}^\phi = \Gamma_{\phi\theta}^\phi = \cot\theta$ and all other Christoffel symbols are zero. Here θ and ϕ are the polar and azimuthal angles on S^2 , respectively. Let

$$\begin{aligned} X_1 &= d\theta \otimes d\theta \\ X_2 &= d\phi \otimes d\phi \\ X_3 &= d\theta \otimes d\phi + d\phi \otimes d\theta \end{aligned}$$

be a basis of W , the symmetric elements of $T^*M \otimes T^*M$. The curvature terms $R_{\theta\phi} = \nabla_{\frac{\partial}{\partial\theta}} \nabla_{\frac{\partial}{\partial\phi}} - \nabla_{\frac{\partial}{\partial\phi}} \nabla_{\frac{\partial}{\partial\theta}}$ are

$$\begin{aligned} R_{\theta\phi}(X_1) &= -(\sin^2\theta)X_3 \\ R_{\theta\phi}(X_2) &= X_3 \\ R_{\theta\phi}(X_3) &= 2X_1 - 2(\sin^2\theta)X_2 \end{aligned}$$

This gives $W^{(0)} = \text{span}(X_1 + (\sin^2\theta)X_2)$. Non-zero local sections of $W^{(0)}$ are of the form $X = f(X_1 + (\sin^2\theta)X_2)$ where f is a smooth non-vanishing function defined on an open subset of S^2 . The covariant derivative of X is $\nabla X = X \otimes d\log|f|$ and so $W^{(1)} = W^{(0)}$. Thus $\widetilde{W} = W^{(0)}$. Since \widetilde{W} is a rank one vector bundle over S^2 it follows that ∇ is a locally metric connection; in fact, it is the Levi-Civita connection of the induced metric of the standard embedding of the two-sphere in three-dimensional Euclidean space.

References

- [1] R. Atkins and Z. Ge, *An Inverse Problem in the Calculus of Variations and the Characteristic Curves of Connections on $SO(3)$ -Bundles* Can. Math. Bull. (1995)
- [2] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry I* (John Wiley & Sons, 1963)
- [3] K. Trancevski, *On the Parallel Vector Fields in Vector Bundles* Tensor N.S. **60** (1998)
- [4] M. Spivak, *Differential Geometry II* (Publish or Perish, 1970,1979)